

**PRACTICE MANAGEMENT GUIDELINES FOR THE EVALUATION
OF BLUNT ABDOMINAL TRAUMA**

EAST Practice Management Guidelines Work Group

William S. Hoff, MD,¹ Michelle Holevar, MD,² Kimberly K. Nagy, MD,³ Lisa
Patterson, MD,⁴ Jeffrey S. Young, MD,⁵ Abenamar Arrillaga, MD,⁶ Michael P.
Najarian, DO,⁷ Carl P. Valenziano, MD⁸

¹Brandywine Hospital, Coatesville, PA

²Mount Sinai Hospital, Chicago, IL

³Cook County Hospital and Rush University, Chicago, IL

⁴Wright State University, Dayton, OH

⁵University of Virginia Health System, Charlottesville, VA

⁶Greenville Memorial Hospital, Greenville, SC

⁷Lehigh Valley Hospital, Allentown, PA

⁸Morristown Memorial Hospital, Morristown, NJ

Address for Correspondence:

William S. Hoff, MD, FACS
Brandywine Hospital - Department of Traumatology
201 Reeceville Road
Coatesville, PA 19320
Phone: (610) 383-8099 / Fax: (610) 383-8352
E-mail: bill.hoff@uphs.upenn.edu

PRACTICE MANAGEMENT GUIDELINES FOR THE EVALUATION OF BLUNT ABDOMINAL TRAUMA

I. Statement of the problem

Evaluation of patients who have sustained blunt abdominal trauma (BAT) may pose a significant diagnostic challenge to the most seasoned trauma surgeon. Blunt trauma produces a spectrum of injury from minor, single-system injury to devastating, multi-system trauma. Trauma surgeons must have the ability to detect the presence of intra-abdominal injuries across this entire spectrum. While a carefully performed physical examination remains the most important method to determine the need for exploratory laparotomy, there is little Level I evidence to support this tenet. In fact, several studies have highlighted the inaccuracies of the physical examination in BAT.^{1, 2} The effect of altered level of consciousness as a result of neurologic injury, alcohol or drugs, is another major confounding factor in assessing BAT.

Due to the recognized inadequacies of physical examination, trauma surgeons have come to rely on a number of diagnostic adjuncts. Commonly used modalities include diagnostic peritoneal lavage (DPL) and computed tomography (CT). Although not available universally, focused abdominal sonography for trauma (FAST) has recently been included in the diagnostic armamentarium. Diagnostic algorithms outlining appropriate use of each of these modalities individually have been established. Several factors influence the selection of diagnostic testing: (1) type of hospital - i.e., trauma center vs. "non-trauma" hospital; (2) access to a particular technology at the surgeon's institution; (3) the surgeon's individual experience with a given diagnostic modality. As facilities evolve, technologies mature and surgeons gain new experience, it is important that any diagnostic strategy constructed be dynamic.

The primary purpose of this study was to develop an evidence-based, systematic diagnostic approach to BAT utilizing the three major diagnostic modalities: i.e., DPL, CT and FAST. This diagnostic regimen would be designed such that it could be reasonably applied by all general surgeons performing an *initial* evaluation of BAT.

II. Process

A. Identification of references

A MEDLINE search was performed using the key words "abdominal injuries" and the subheading "diagnosis". This search was limited further to (1) clinical research, (2) published in English, (3) publication dates January 1978 through February 1998. The initial search yielded 742 citations. Case reviews, review articles, meta-analyses, editorials, letters to the editor, technologic reports, pediatric series and studies involving a significant number of penetrating abdominal injuries were excluded prior to formal review. Additional references, selected by the individual subcommittee members, were then included to compile the master reference list of 197 citations.

B. Quality of the references

Articles were distributed among subcommittee members for formal review. A review data sheet was completed for each article reviewed which summarized the main conclusions of the study, and identified any deficiencies in the study. Further, reviewers classified each reference by the methodology established by the Agency for Health Care Policy and Research (AHCPR) of the U.S. Department of Health and Human Services as follows:

Class I: Prospective, randomized, double-blinded study

Class II: Prospective, randomized, non-blinded trial

Class III: Retrospective series, meta-analysis

Following review by the subcommittee, references were excluded based on poor design or invalid conclusions. An evidentiary table was constructed using the remaining 101 references: Class I (20); Class II (32); Class III (49). Recommendations were based on studies included in the evidentiary table.

III. Recommendations

A. Level I

1. Exploratory laparotomy is indicated for patients with a positive DPL.
2. CT is recommended for the evaluation of hemodynamically stable patients with equivocal findings on physical examination, associated neurologic injury, or multiple extra-abdominal injuries. Under these circumstances, patients with a negative CT should be admitted for observation.
3. CT is the diagnostic modality of choice for nonoperative management of solid visceral injuries.
4. In hemodynamically stable patients, DPL and CT are complementary diagnostic modalities.

B. Level II

1. FAST may be considered as the initial diagnostic modality to exclude hemoperitoneum. In the presence of a negative or indeterminate FAST result, DPL and CT have complementary roles.
2. When DPL is used, clinical decisions should be based on the presence of gross blood on initial aspiration (i.e., 10 ml) or microscopic analysis of lavage effluent.
3. In hemodynamically stable patients with a positive DPL, follow-up CT scan should be considered, especially in the presence of pelvic fracture or suspected injuries to the genitourinary tract, diaphragm or pancreas.
4. Exploratory laparotomy is indicated in hemodynamically unstable patients with a positive FAST. In hemodynamically stable patients with a positive FAST, follow-up CT permits nonoperative management of select injuries.
5. Surveillance studies (i.e., DPL, CT, repeat FAST) are required in hemodynamically stable patients with indeterminate FAST results.

C. Level III

1. Objective diagnostic testing (i.e., FAST, DPL, CT) is indicated for patient with abnormal mentation, equivocal findings on physical examination, multiple injuries, concomitant chest injury or hematuria.

2. Patients with seatbelt sign (SBS) should be admitted for observation and serial physical examination. Detection of intraperitoneal fluid by FAST or CT in a patient with SBS mandates either DPL to determine the nature of the fluid or exploratory laparotomy.
3. CT is indicated for the evaluation of suspected renal injuries.
4. A negative FAST should prompt follow-up CT for patients at high risk for intraabdominal injuries (e.g., multiple orthopedic injuries, severe chest wall trauma, neurologic impairment).
5. Splanchnic angiography may be considered in patients who require angiography for the evaluation of other injuries (e.g., thoracic aortic injury, pelvic fracture).

IV. Scientific Foundation

A. Diagnostic Peritoneal Lavage (DPL)

DPL was introduced by Root in 1965 as a rapid and accurate method to identify the presence of intra-abdominal hemorrhage following trauma.³ Subsequent studies have confirmed the efficacy of DPL in diagnosing abdominal hemorrhage as well as its superiority over physical examination alone.⁴ The accuracy of DPL has been reported between 92% and 98%.^{5, 6, 7, 8, 9, 10} The high sensitivity of DPL is due to the significant false positive rate of the technique.^{11, 12, 13} Several authors have highlighted the importance of interpreting DPL results in the context of the overall clinical condition of the patient. A positive DPL does not necessarily mandate immediate laparotomy in the hemodynamically stable patient.^{12, 14, 15, 16} DPL has been shown to be more *efficient* than CT scan in identifying patients that require surgical exploration.¹⁷

The complication rate associated with DPL is quite low.¹⁸ The incidence of complications is lower for open DPL compared with to the closed technique. However, closed DPL can be performed more rapidly.^{19, 20, 21, 22} Studies designed to examine the ability of physicians to estimate the red blood cell (RBC) count in DPL fluid have demonstrated the poor sensitivity of visual inspection.^{23, 24, 25} A positive DPL, based on microscopic analysis of lavage fluid, has been defined as $> 10^5$ RBC/mm³. It has been recommended that patients with RBC counts in the equivocal range (i.e., 25,000 – 75,000 RBC/mm³) undergo additional diagnostic testing, such as CT scanning.¹²

The false positive rate for DPL is increased in patients with pelvic fractures.^{26, 27} In order to avoid sampling the retroperitoneal hematoma, a suprapubic approach has been recommended, theoretically reducing the chances of a false positive result.²⁸

The advantages of DPL for detection of hollow visceral injuries have been clearly demonstrated.^{29, 30} Two studies which advocate analysis of DPL fluid for amylase and alkaline phosphatase consistent with enteric injuries have been disputed.^{31, 32, 33} Similarly, the utility of the DPL white blood cell (WBC) count has been questioned.^{34, 35, 36} DPL is sensitive for mesenteric injury and, in fact, has been shown to be superior to CT for the diagnosis of this injury.³⁷

Thus, DPL is a safe, rapid and accurate method for determining the presence of intraperitoneal blood in victims of BAT. It is more accurate than CT for the early diagnosis of hollow visceral and mesenteric injuries, but it does not reliably exclude

significant injuries to retroperitoneal structures. False positive results may occur in the presence of pelvis fractures. Hemodynamically stable patients with equivocal results are best managed by additional diagnostic testing to avoid unnecessary laparotomies.

B. Computed Tomography (CT)

Routine use of CT for the evaluation of BAT was not initially viewed with overwhelming enthusiasm. CT requires a cooperative, hemodynamically stable patient. In addition, the patient must be transported out of the trauma resuscitation area to the radiographic suite. Specialized technicians and the availability of a radiologist for interpretation were also viewed as factors which limited the utility of CT for trauma patients. CT scanners are now available in most trauma centers and, with the advent of helical scanners, scan time has been significantly reduced. As a result, CT has become an accepted part of the traumatologist's armamentarium.

The accuracy of CT in hemodynamically stable blunt trauma patients has been well established. Sensitivity between 92% and 97.6% and specificity as high as 98.7% has been reported in patients subjected to emergency CT.^{38, 39} Most authors recommend admission and observation following a negative CT scan.^{40, 41} In a recent study of 2774 patients, the authors concluded that the negative predictive value (99.63%) of CT was sufficiently high to permit safe discharge of BAT patients following a negative CT scan.⁴²

CT is notoriously inadequate for the diagnosis of mesenteric injuries and may also miss hollow visceral injuries. In patients at risk for mesenteric or hollow visceral injury, DPL is generally felt to be a more appropriate test.^{37, 43} A negative CT scan in such a patient cannot reliably exclude intra-abdominal injuries.

CT has the unique ability to detect clinically unsuspected injuries. In a series of 444 patients in whom CT was performed to evaluate renal injuries, 525 concomitant abdominal and/or retroperitoneal injuries were diagnosed. Another advantage of CT scanning over other diagnostic modalities is its ability to evaluate the retroperitoneal structures.⁴⁰ Kane performed CT in 44 hemodynamically stable blunt trauma patients following DPL. In 16 patients, CT revealed significant intra-abdominal or retroperitoneal injuries not diagnosed by DPL. Moreover, the findings on CT resulted in a modification to the original treatment plan in 58% of the patients.⁴⁴

C. Focused Abdominal Sonography for Trauma (FAST)

In recent years, focused abdominal sonography for trauma (FAST) has emerged as a useful diagnostic test in the evaluation of BAT. The advantages of the FAST examination have been clearly established. FAST is noninvasive, may be easily performed and can be done concurrently with resuscitation. In addition, the technology is portable and may be easily repeated if necessary.^{45, 46, 47, 48} In most cases, FAST may be completed within 3 or 4 minutes.^{49, 50, 51} The test is especially useful for detecting intra-abdominal hemorrhage in the multiply injured or pregnant patient.⁵²

A noted drawback to the FAST examination is the fact that a positive examination relies on the presence of free intraperitoneal fluid. In the hands of most operators, ultrasound will detect a minimum of 200 mL of fluid.⁵³ Injuries not associated with hemoperitoneum may not be detected by this modality.^{49, 54, 55} Thus, ultrasound is not a reliable method for excluding hollow visceral injury.^{47, 49, 56, 57, 58} In addition,

the FAST examination cannot be used to reliably grade solid organ injuries. Therefore, in the hemodynamically stable patient, a follow-up CT scan should be obtained if nonoperative management is contemplated.⁵⁹

FAST compares favorably with more traditionally utilized diagnostic tests. In the hemodynamically stable patient with BAT, FAST offers a viable alternative to DPL.⁶⁰ DPL may also be used as a complementary examination in the hemodynamically stable patient in the presence of an equivocal or negative ultrasound with strong clinical suspicion of visceral injury.^{61, 62} FAST has demonstrated utility in hemodynamically stable patients with BAT.^{58, 60, 63} In addition, ultrasound has been shown to be more cost-effective when compared to DPL or CT.^{45, 47, 60}

Overall, FAST has a sensitivity between 73% and 88%, a specificity between 98% and 100% and is 96% to 98% accurate.^{46, 50, 57, 58, 64, 65} This level of accuracy is independent of the practitioner performing the study. Surgeons, emergency medicine physicians, ultrasound technicians and radiologists have equivalent results.^{46, 53, 64, 65, 66}

D. Other Diagnostic Modalities

As interest in laparoscopic procedures has increased among general surgeons, there has been speculation regarding the role of diagnostic laparoscopy (DL) in the evaluation of BAT. One of the potential benefits postulated is the reduction of nontherapeutic laparotomies. With modification of the technique to include smaller instruments, portable equipment and local anesthesia, DL may be a useful tool in the initial evaluation of BAT. Although there are no randomized, controlled studies comparing DL to more commonly utilized modalities, experience at one institution using minilaparoscopy demonstrated a 25% incidence of positive findings on DL, which were successfully managed nonoperatively and would have resulted in nontherapeutic laparotomies.⁶⁷

Although its ultimate role remains unclear, another modality to be considered in the diagnostic evaluation of BAT is visceral angiography. This modality may have diagnostic value when employed in conjunction with angiography of the pelvis or chest, or when other diagnostic studies are inconclusive.⁶⁸

V. Summary

Injury to intra-abdominal viscera must be excluded in all victims of BAT. Physical examination remains the initial step in diagnosis but has limited utility under select circumstances. Thus, various diagnostic modalities have evolved to assist the trauma surgeon in the identification of abdominal injuries. The specific tests selected are based on the clinical stability of the patient, the ability to obtain a reliable physical examination and the provider's access to a particular modality. It is important to emphasize that many of the diagnostic tests utilized are complementary rather than exclusionary.

Based on the above recommendations, a reasonable diagnostic approach to BAT is summarized in Figures 1 and 2. In hemodynamically stable patients with a reliable physical examination, clinical findings may be used to select patients who may be safely observed. In the absence of a reliable physical examination, the main

diagnostic choice is between CT or FAST (with CT in a complementary role). Hemodynamically unstable patients may be initially evaluated with FAST or DPL.

VI. Future Investigation

Recent literature is replete with studies that emphasize the many advantages of ultrasound in the valuation of BAT. Although this technology is becoming more available to trauma surgeons, for a variety of reasons, it has not become universally available in all centers. Continued research addressing the utility of FAST, with emphasis on its advantages specific to resource utilization, is suggested. In addition, studies should be designed to more closely evaluate the feasibility of FAST as the sole diagnostic test in hemodynamically stable patients. Perhaps safe strategies for nonoperative management of solid visceral injuries could be developed which rely on FAST alone, such that the number of CT scans could be reduced.

VII. References

1. Rodriguez A, DuPriest RW Jr., Shatney CH: Recognition of intra-abdominal injury in blunt trauma victims. A prospective study comparing physical examination with peritoneal lavage. *Am Surg* 48: 457-459, 1982.
2. Schurink GW, Bode PJ, van Luijt PA, et al: The value of physical examination in the diagnosis of patients with blunt abdominal trauma: a retrospective study. *Injury* 28: 261-265, 1997.
3. Root HD, Hauser CW, McKinley CR, et al: Diagnostic peritoneal lavage. *Surg* 57: 633-637, 1965.
4. Bivins BA, Sachatello CR, Daugherty ME, et al: Diagnostic peritoneal lavage is superior to clinical evaluation in blunt abdominal trauma. *Am Surg* 44: 637-641, 1978.
5. Smith SB, Andersen CA: Abdominal trauma: the limited role of peritoneal lavage. *Am Surg* 48: 514-517, 1982.
6. Henneman PL, Marx JA, Moore EE, et al: Diagnostic peritoneal lavage: accuracy in predicting necessary laparotomy following blunt and penetrating trauma. *J Trauma* 30: 1345-1355, 1990.
7. Krausz MM, Manny J, Austin E, et al: Peritoneal lavage in blunt abdominal trauma. *SG & O* 152: 327-330, 1981.
8. Moore JB, Moore EE, Markivchick VJ, et al: Diagnostic peritoneal lavage for abdominal trauma: superiority of the open technique at the infraumbilical ring. *J Trauma* 21: 570-572, 1981.
9. Jacob ET, Cantor E: Discriminate diagnostic peritoneal lavage in blunt abdominal injuries: accuracy and hazards. *Am Surg* 45: 11-14, 1979.
10. Fischer RP, Beverlin BC, Engrav LH, et al: Diagnostic peritoneal lavage: fourteen years and 2586 patients later. *Am J Surg* 136: 701-704, 1978.
11. Bilge A, Sahin M: Diagnostic peritoneal lavage in blunt abdominal trauma. *Eur J Surg* 157: 449-451, 1991.
12. DeMaria EJ: Management of patients with indeterminate diagnostic peritoneal lavage results following blunt trauma. *J Trauma* 31: 1627-1631, 1991.
13. Van Dongen LM, de Boer HH: Peritoneal lavage in closed abdominal injury. *Injury* 16: 227-229, 1985.
14. Day AC, Rankin N, Charlesworth P: Diagnostic peritoneal lavage: integration with clinical information to improve diagnostic performance. *J Trauma* 32: 52-57, 1992.
15. Barba C, Owen D, Fleiszer D, et al: Is positive diagnostic peritoneal lavage an absolute indication for laparotomy in all patients with blunt trauma? The Montreal General Hospital experience. *Can J Surg* 34: 442-445, 1991.
16. Drost TF, Rosemurgy AS, Kearney RE, et al: Diagnostic peritoneal lavage. Limited indications due to evolving concepts in trauma care. *Am Surg* 57: 126-128, 1991.
17. Blow O, Bassam D, Butler K, et al: Speed and efficiency in the resuscitation of blunt trauma patients with multiple injuries: the advantage of diagnostic peritoneal lavage over abdominal computerized tomography. *J Trauma* 44: 287-290, 1998.
18. Davis JW, Hoyt DB, Mackersie RC, et al: Complications in evaluating abdominal trauma: diagnostic peritoneal lavage versus computerized axial tomography. *J Trauma* 30: 1506-1509, 1990.

19. Lopez-Viego MA, Mickel TJ, Weigelt JA: Open versus closed diagnostic peritoneal lavage in the evaluation of abdominal trauma. *Am J Surg* 160: 594-597, 1990.
20. Cue JI, Miller FB, Cryer HM, III, et al: A prospective, randomized comparison between open and closed peritoneal lavage techniques. *J Trauma* 30: 880-883, 1990.
21. Wilson WR, Schwarcz TH, Pilcher DB: A prospective randomized trial of the Lazarus-Nelson vs. the standard peritoneal dialysis catheter for peritoneal lavage in blunt abdominal trauma. *J Trauma* 27: 1177-1180, 1987.
22. Felice PR, Morgan AS, Becker DR: A prospective randomized study evaluating periumbilical versus infraumbilical peritoneal lavage: a preliminary report. A combined hospital study. *Am Surg* 53: 518-520, 1987.
23. Gow KW, Haley LP, Phang PT, et al: Validity of visual inspection of diagnostic peritoneal lavage fluid. *Can J Surg* 39: 114-119, 1996.
24. Wyatt JP, Evans RJ, Cusack SP: Variation among trainee surgeons in interpreting diagnostic peritoneal lavage fluid in blunt abdominal trauma. *J Royal Coll Surg (Edin)* 37: 104-106, 1992.
25. Driscoll P, Hodgkinson D, Mackway-Jones K: Diagnostic peritoneal lavage: It's red but is it positive? *Injury* 23: 267-269, 1992.
26. Mendez C, Gubler KD, Maier RV: Diagnostic accuracy of peritoneal lavage in patients with pelvic fractures. *Arch Surg* 129: 477-482, 1994.
27. Hubbard SG, Bivins BA, Sachatello CR, et al: Diagnostic errors with peritoneal lavage in patients with pelvic fractures. *Arch Surg* 114: 844-846, 1979.
28. Cochran W, Sobat WS: Open versus closed diagnostic peritoneal lavage. A multiphasic prospective randomized comparison. *Ann Surg* 200: 24-28, 1984.
29. Meyer DM, Thal ER, Weigelt JA, et al: Evaluation of computed tomography and diagnostic peritoneal lavage in blunt abdominal trauma. *J Trauma* 29: 1168-1170, 1989.
30. Burney RE, Mueller GL, Coon WW, et al: Diagnosis of isolated small bowel injury following blunt abdominal trauma. *Ann Emerg Med* 12: 71-74, 1983.
31. McAnena OJ, Marx JA, Moore EE: Peritoneal lavage enzyme determinations following blunt and penetrating abdominal trauma. *J Trauma* 31: 1161-1164, 1991.
32. McAnena OJ, Marx JA, Moore EE: Contributions of peritoneal lavage enzyme determinations to the management of isolated hollow visceral abdominal injuries. *Ann Emerg Med* 20: 834-837, 1991.
33. Megison SM, Weigelt JA: The value of alkaline phosphatase in peritoneal lavage. *Ann Emerg Med* 19: 503-505, 1990.
34. D'Amelio LF, Rhodes M: A reassessment of peritoneal lavage leukocyte count in blunt abdominal trauma. *J Trauma* 30: 1291-1293, 1990.
35. Soyka JM, Martin M, Sloan EP, et al: Diagnostic peritoneal lavage: is an isolated WBC count greater than or equal to 500/mm³ predictive of intra-abdominal injury requiring celiotomy in blunt trauma patients? *J Trauma* 30: 874-879, 1990.
36. Jacobs DG, Angus L, Rodriguez A, et al: Peritoneal lavage white count: a reassessment. *J Trauma* 30: 607-612, 1990.
37. Ceraldi CM, Waxman K: Computerized tomography as an indicator of isolated mesenteric injury. A comparison with peritoneal lavage. *Am Surg* 56: 806-810, 1990.

38. Peitzman AB, Makaroun MS, Slasky BS, et al: Prospective study of computed tomography in initial management of blunt abdominal trauma. *J Trauma* 26: 585-592, 1986.
39. Webster VJ: Abdominal trauma: Pre-operative assessment and postoperative problems in intensive care. *Anaest Int Care* 13: 258-262, 1985.
40. Lang EK: Intra-abdominal and retroperitoneal organ injuries diagnosed on dynamic computed tomograms obtained for assessment of renal trauma. *J Trauma* 30: 1161-1168, 1990.
41. Matsubara TK, Fong HM, Burns CM: Computed tomography of abdomen (CTA) in management of blunt abdominal trauma. *J Trauma* 30: 410-414, 1990.
42. Livingston DH, Lavery RF, Passannante MR, et al: Admission or observation is not necessary after a negative abdominal computed tomographic scan in patients with suspected blunt abdominal trauma: Results of a prospective, multi-institutional trial. *J Trauma* 44: 272-282, 1998.
43. Nolan BW, Gabram SG, Schwartz RJ, et al: Mesenteric injury from blunt abdominal trauma. *Am Surg* 61: 501-506, 1995.
44. Kane NM, Dorfman GS, Cronan JJ: Efficacy of CT following peritoneal lavage in abdominal trauma. *J Comp Asst Tomo* 11: 998-1002, 1987.
45. Branney SW, Moore EE, Cantrill SV, et al: Ultrasound based key clinical pathway reduces the use of hospital resources for the evaluation of blunt abdominal trauma. *J Trauma* 42: 1086-1090, 1997.
46. Healey MA, Simons RK, Winchell RJ, et al: A prospective evaluation of abdominal ultrasound in blunt trauma: Is it useful? *J Trauma* 40: 875-883, 1996.
47. Glaser K, Tschmelitsch J, Klingler P, et al: Ultrasonography in the management of blunt abdominal and thoracic trauma. *Arch Surg* 129: 743-747, 1994.
48. Liu M, Lee CH, P'eng FK: Prospective comparison of diagnostic peritoneal lavage, computed tomographic scanning, and ultrasonography for the diagnosis of blunt abdominal trauma. *J Trauma* 35: 267-270, 1993.
49. Boulanger BR, McLellan BA, Brenneman FD, et al: Emergent abdominal sonography as a screening test in a new diagnostic algorithm for blunt trauma. *J Trauma* 40: 867-874, 1996.
50. Boulanger BR, Brenneman FD, McLellan BA, et al: A prospective study of emergent abdominal sonography after blunt trauma. *J Trauma* 39: 325-330, 1995.
51. Ma OJ, Kefer MP, Mateer JR, et al: Evaluation of hemoperitoneum using a single- vs. multiple-view ultrasonographic examination. *Acad Emerg Med* 2: 581-586, 1995.
52. Rozycki GS, Ochsner MG, Jaffin JH, et al: Prospective evaluation of surgeons' use of ultrasound in the evaluation of trauma patients. *J Trauma* 34: 516-527, 1993.
53. Branney SW, Wolfe RE, Moore EE, et al: Quantitative sensitivity of ultrasound in detecting free intraperitoneal fluid. *J Trauma* 39: 375-380, 1995.
54. Sherbourne CD, Shanmuganathan K, Mirvis SE, et al: Visceral injury without hemoperitoneum: a limitation of screening abdominal sonography for trauma. *Emerg Radiol* 4: 349-354, 1997.
55. Tso P, Rodriguez A, Cooper C: Sonography in blunt abdominal trauma: a preliminary progress report. *J Trauma* 33: 39-44, 1992.
56. Buzzas GR, Kern SJ, Smith RS, et al: A comparison of sonographic examinations for trauma performed by surgeons and radiologists. *J Trauma* 44: 604-608, 1998.

57. Smith SR, Kern SJ, Fry WR, et al: Institutional learning curve of surgeon-performed trauma ultrasound. *Arch Surg* 133: 530-536, 1998.
58. McKenney MG, Martin L, Lentz K, et al: 1000 consecutive ultrasounds for blunt abdominal trauma. *J Trauma* 40: 607-612, 1996.
59. Chambers JA, Pilbrow WJ: Ultrasound in abdominal trauma: an alternative to peritoneal lavage. *Arch Emerg Med* 5: 26-33, 1988.
60. McKenney KL, McKenney MG, Nunez DB, et al: Cost reduction using ultrasound in blunt abdominal trauma. *Emerg Radiol* 4: 3-6, 1997.
61. Kimura A, Otsuka T: Emergency center ultrasonography in the evaluation of hemoperitoneum: A prospective study. *J Trauma* 31: 20-23, 1991.
62. Gruessner R, Mentges B, Duber C, et al: Sonography versus peritoneal lavage in blunt abdominal trauma. *J Trauma* 29: 242-244, 1989.
63. Rothlin MA, Naf R, Amgwerd M, et al: Ultrasound in blunt abdominal and thoracic trauma. *J Trauma* 34: 488-495, 1993.
64. Kern SJ, Smith RS, Fry WR, et al: Sonographic examination of abdominal trauma by senior surgical residents. *Am Surg* 63: 669-674, 1997.
65. Rozycki GS, Ochsner MG, Schmidt JA, et al: A prospective study of surgeon-performed ultrasound as the primary adjuvant modality for injured patient assessment. *J Trauma* 39: 492-500, 1995.
66. McKenney M, Lentz K, Nunez D, et al: Can ultrasound replace diagnostic peritoneal lavage in the assessment of blunt trauma? *J Trauma* 37: 439-441, 1994.
67. Berci G, Sackier JM, Paz-Partlow M: Emergency laparoscopy. *Am J Surg* 161: 332-335, 1991.
68. Ward RE: Study and management of blunt trauma in the immediate post-impact period. *Rad Clin NA* 19: 3-7, 1981.

PRACTICE MANAGEMENT GUIDELINES FOR THE EVALUATION OF BLUNT ABDOMINAL TRAUMA

First Author	Year	Reference Title	Class	Conclusions
Bivins BA	1978	Diagnostic exploratory celiotomy: an outdated concept in blunt abdominal trauma.	I	Clinical evaluation alone would have missed 59% of injuries in blunt trauma patients studied. Exploratory laparotomy recommended for (+)DPL. Due to recognized false negative rate, admission and observation recommended for patients with (-)DPL.
Moore JB	1980	Diagnostic peritoneal lavage for abdominal trauma: superiority of the open technique at the intraumbilical ring. <i>J Trauma</i> 21: 570-572	I	Open DPL preferred over closed. Increased time required for open DPL compensated by higher reliability.
Krausz MM	1981	Peritoneal lavage in blunt abdominal trauma. S, G & O 152: 327-330	I	Laboratory study demonstrates safety and reliability of percutaneous (closed) method of DPL. Clinical series documents percutaneous technique to be accurate in the diagnosis of visceral injury and/or hemoperitoneum.
Cochran W	1984	Open versus closed diagnostic peritoneal lavage. A multiphasic prospective randomized comparison. <i>Ann Surg</i> 200: 24-28	I	No significant difference in accuracy between two techniques. Supraumbilical approach more accurate in presence of pelvis fracture. Complication rate higher with open DPL.
Felice PR	1987	A prospective randomized study evaluating periumbilical versus intraumbilical peritoneal lavage: a preliminary report. A combined hospital study. <i>Am Surgeon</i> 53: 518-520	I	Periumbilical peritoneal lavage performed faster and preferred by majority of providers. Safety and sensitivity equivalent between the two techniques.
Wilson WR	1987	A prospective randomized trial of the Lazarus-Nelson vs the standard peritoneal dialysis catheter for peritoneal lavage in blunt abdominal trauma. <i>J Trauma</i> 27: 1177-1180	I	Percutaneous DPL (i.e. Lazarus -Nelson) associated with decreased time to catheter insertion with no significant difference in time to complete lavage, volume of fluid recovered, sensitivity, or specificity compared with open technique. Open DPL recommended for patients with previous abdominal surgery or when percutaneous DPL unsuccessful.
Buechter KU	1990	The use of serum amylase and lipase in evaluating and managing blunt abdominal trauma. <i>Am Surgeon</i> 56: 204-208	I	Serum amylase and lipase are randomly elevated in BAT population. Diagnostic testing is not warranted based on elevated amylase or lipase on initial evaluation.
Howdieshell TR	1989	Open versus closed peritoneal lavage with particular attention to time, accuracy, and cost. <i>Am J Emerg Med</i> 7: 367-371	I	Closed DPL is faster, safer and equally accurate as open DPL.

Kimura A	1991	Emergency center ultrasonography in the evaluation of hemoperitoneum: a prospective study. <i>J Trauma</i> 31: 20-23	I	Recommend US as a screening modality for detection of hemoperitoneum (86. 7% sensitivity; 100% specificity). DPL indicated for neurologically injured patients with (-)US and a high suspicion of visceral injury.
Troop B	1991	Randomized, prospective comparison of open and closed peritoneal lavage for abdominal trauma. <i>Ann Emerg Med</i> 20: 1290-1292	I	Closed DPL superior to open DPL. Open or semi -open technique recommended for patients in whom closed DPL is contraindicated.
Day AC	1992	Diagnostic peritoneal lavage: integration with clinical information to improve diagnostic performance. <i>J Trauma</i> 32: 52-57	I	Combination of clinical evaluation and DPL reduces rate of non - therapeutic laparotomies, but increases the number of missed injuries. The highest accuracy (95%) is obtained by combination of circulatory assessment and DPL.
Tso P	1992	Sonography in blunt abdominal trauma: a preliminary progress report. <i>J Trauma</i> 33: 39-44	I	US sensitive (91%) for detection of free fluid but less sensitive (69%) for identification of free fluid plus organ disruption. US does not rule out organ injury in the absence of hemoperitoneum.
Liu M	1993	Prospective comparison of diagnostic peritoneal lavage, computed tomographic scanning, and ultrasonography for the diagnosis of blunt abdominal trauma. <i>J Trauma</i> 35: 267-270	I	Sensitivity and specificity of US is comparable to CT or DPL. False negatives identified using CT (1) and US (3) in the presence of intestinal perforations. Defined complementary roles of US, CT and DPL in evaluation of BAT.
Rothlin MA	1993	Ultrasound in blunt abdominal and thoracic trauma. <i>J Trauma</i> 34: 488-495	I	US highly sensitive (98.1%) and specific (100%) for identification of intra-abdominal fluid. Specificity remains high (99.6%) but sensitivity decreases (43.6%) for diagnosis of specific organ lesions. Recommend 1) CT to identify specific organ injury, 2) serial US every 1 -2 hrs for first 6 hrs, then every 12 hrs for 2 days.
Rozycki GS	1993	Prospective evaluation of surgeons' use of ultrasound in the evaluation of trauma patients. <i>J Trauma</i> 34: 516-527	I	In mixed blunt (84%) / penetrating (16%) population, US has 79.0% sensitivity and 95.6% specificity. Adjusted sensitivity for blunt trauma is 84.0%. US indicated for 1) blunt thoracoabdominal injury; 2) suspected pericardial tamponade; 3) multi -system injury with unknown etiology of hypotension; 4) pregnant trauma patient.
Goletti O	1994	The role of ultrasonography in blunt abdominal trauma: results in 250 consecutive cases. <i>J Trauma</i> 36: 178-181	I	Overall sensitivity of US 86.7%. Intraperitoneal fluid volumes > 250 ml correlates with high unnecessary laparotomy rate when diagnosed by US; suggest 250 ml as threshold for non -operative management using US. US -guided paracentesis allows safe non -operative management in presence of small volume of fluid.
Huang M	1994	Ultrasonography for the evaluation of hemoperitoneum during resuscitation: a simple scoring system. <i>J Trauma</i> 36: 173-177	I	US 100% specific for diagnosis of hemoperitoneum. Scoring system developed to predict presence of hemoperitoneum and need for surgery; US score > 3 corresponds to > 1000 ml blood with 84% sensitivity, 71% specificity and 71% accuracy.

McKenney M	1994	Can ultrasound replace diagnostic peritoneal lavage in the assessment of blunt trauma? <i>J Trauma</i> 37: 439-441	I	Comparison of US with DPL, CT scan and exploratory laparotomy in 200 patients with BAT. US 83% sensitive, 100% specific and 97% accurate.
Branney SW	1995	Quantitative sensitivity of ultrasound in detecting free intraperitoneal fluid. <i>J Trauma</i> 39: 375-380	I	Prospective study of US performed in patients following DPL with no aspiration of gross blood. US demonstrated 97% sensitivity and detected mean fluid volume of 619 ml. US screen should be initial branch point in BAT algorithm.
Rozycki GS	1995	A prospective study of surgeon - performed ultrasound as the primary adjuvant modality for injured patient assessment. <i>J Trauma</i> 39: 492-500	I	Assessment of surgeon-performed US in 371 patients (295 blunt / 76 penetrating). US is an accurate modality (81.5% sensitivity; 99.7% specificity) which may be performed by surgeons. Recommend repeat US at 12 -14 hrs if initial exam is negative.
Thomas B	1997	Ultrasound evaluation of blunt abdominal trauma: program implementation, initial experience, and learning curve. <i>J Trauma</i> 42: 384-390	I	US examination s performed in 300 patients by surgeons and trauma fellows with review of false(-) and false(+) by radiologist. Demonstrated 81.0% sensitivity and 99.3% specificity. Accuracy plateaus after 100 examinations. Projected cost savings of \$41,000.
McKenney MG	1998	Can surgeons evaluate emergency ultrasound scans for blunt abdominal trauma? <i>J Trauma</i> 44: 649-653	I	Prospective study of 112 FAST examinations performed and initially interpreted by surgeons with final interpretation by radiologist. No false negatives, 2 false positives recorded. Good agreement between interpretation by surgeon and radiologist (99%).
Jacob ET	1979	Discriminate diagnostic peritoneal lavage in blunt abdominal injuries: accuracy and hazards. <i>Am Surgeon</i> 45: 11-14	II	DPL 93.4% accurate in prediction of positive exploratory laparotomy and 96.6% accurate in prediction of negative exploratory laparotomy in patients with mild or equivocal clinical findings.
Buttsworth JF	1980	Detection of occult abdominal trauma in patients with severe head injuries. <i>Lancet</i> 2: 759-762	II	DPL recommended for trauma patients who are unable to obey simple commands secondary to closed head injury to exclude occult intra-abdominal injury.
Alyono D	1982	Reappraisal of diagnostic peritoneal lavage criteria for operation in penetrating and blunt trauma. <i>Surgery</i> 92: 751-757	II	In blunt trauma, the highest level of accuracy is achieved with standard diagnostic criteria: DPL -RBC > 100K/mm ³ ; DPL-WBC > 500/mm ³
Kusminsky RE	1982	The potential value of endotoxin - amylase detection in peritoneal lavage fluid. <i>Am Surgeon</i> 48: 359-362	II	Detection of amylase or endotoxin in DPL fluid is valuable in the detection of pancreatic and gastrointestinal injuries.
Rodriguez A	1982	Recognition of intra -abdominal injury in blunt trauma victims. A prospective study comparing physical examination with peritoneal lavage. <i>Am Surgeon</i> 48: 457-459	II	Findings on PE unreliable in conscious, oriented patients with BAT resulting in potential for missed intra -abdominal injuries. DPL highly accurate and sensitive for detection on intra -abdominal injuries.

Kuminsky RE	1984	The value of sequential peritoneal profile in blunt abdominal trauma. <i>Am Surgeon</i> 50: 248-253	II	Addition of endotoxin in DPL fluid allows safe non -operative management in hemodynamically stable BAT patients with DPL RBC count > 100K/mm ³ .
Davis RA	1985	The use of computerized axial tomography versus peritoneal lavage in the evaluation of blunt abdominal trauma. <i>Surgery</i> 98: 845-850	II	High sensitivity and specificity documented for r DPL compared with CT in BAT: cost of CT 8x cost of DPL. CT as the sole diagnostic modality in hemodynamically stable patients with BAT adds cost, time and risk of missed injury without providing significant additional information.
Pelitzman AB	1986	Prospective study of computed tomography in initial management of blunt abdominal trauma. <i>J Trauma</i> 26: 585-592	II	CT demonstrated to be highly sensitive (97.6%) and specific (98.7%) for the diagnosis of intra -abdominal injuries in hemodynamically stable patients. CT permits safe non -operative management of solid visceral injuries.
Gomez GA	1987	Diagnostic peritoneal lavage in the management of blunt abdominal trauma: a reassessment. <i>J Trauma</i> 27: 1-5	II	DPL is an accurate indicator of significant intra -abdominal injury as documented by exploratory laparotomy in patients with BAT.
Pagliariello G	1987	Abdominopelvic computerized tomography and open peritoneal lavage in patients with blunt abdominal trauma: a prospective study. <i>Can J Surgery</i> 30: 10-13	II	CT less sensitive when compared with DPL. Agreement between DPL and CT demonstrated in 53%. DPL superior to CT for evaluation of BAT.
Chambers JA	1988	Ultrasound in abdominal trauma: an alternative to peritoneal lavage. <i>Arch Emerg Med</i> 5: 26-33	II	US is a reliable diagnostic technique for detection of free intraperitoneal fluid but is unreliable for grading specific injuries.
Frame S	1989	Computed tomography versus diagnostic peritoneal lavage: usefulness in immediate diagnosis of blunt abdominal trauma. <i>Ann Emerg Med</i> 18: 513-516	II	DPL safer and more accurate than CT in the evaluation of BAT.
Gruessner R	1989	Sonography versus peritoneal lavage in blunt abdominal trauma. <i>J Trauma</i> 29: 242-244	II	Ultrasound preferred initial screening method compared to DPL for evaluation of BAT. However, DPL has complementary role in the presence of indeterminate US.
Mckersie RC	1989	Intra-abdominal injury following blunt trauma. Identifying the high -risk patient using objective risk factors. <i>Arch Surgery</i> 124: 809-813	II	Presence of (1) chest injury; (2) base deficit < -3.0; (3) hypotension on arrival; (4) prehospital hypotension; (5) pelvic fracture significantly correlated with intra -abdominal injury. DPL, US, or CT recommended in the presence of one of these risk factors.
Meyer DM	1989	Evaluation of computed tomography and diagnostic peritoneal lavage in blunt abdominal trauma. <i>J Trauma</i> 29: 1168-1170	II	CT scan significantly less sensitive than DPL in BAT patients with equivocal findings on PE (74.3% vs 95.9%). CT unreliable for identification of small intestinal injuries in the acute stage of evaluation.

Cue JI	1990	A prospective, randomized comparison between open and closed peritoneal lavage techniques. <i>J Trauma</i> 30: 880-883	II	Open DPL takes longer to perform with better return of lavage fluid. Time consideration and improved patient tolerance justifies use of closed CPL.
Lopez-Viego MA	1990	Open versus closed diagnostic peritoneal lavage in the evaluation of abdominal trauma. <i>Am J Surgery</i> 160: 594-596	II	Open DPL has fewer complications. Because it may be performed faster, closed DPL is recommended with conversion to open technique if complications occur.
Blige A	1991	Diagnostic peritoneal lavage in blunt abdominal trauma. <i>Eur J Surgery</i> 157: 449-451	II	DPL is highly accurate for the diagnosis of free intraperitoneal blood, but is overly sensitive in that it is unable to distinguish clinically unimportant amounts of intraperitoneal blood. Recommend additional diagnostic studies in hemodynamically stable patients with BAT to reduce incidence of unnecessary laparotomies.
Drost TF	1991	Diagnostic peritoneal lavage. Limited indications due to evolving concepts in trauma care. <i>Am Surgeon</i> 57: 126-128	II	Non-therapeutic exploratory laparotomy performed in 1/3 BAT patients based on DPL results. (+)DPL not a reliable predictor of significant intra-abdominal injury, especially in lieu of non-operative management protocols.
Hoffmann R	1992	Blunt abdominal trauma in cases of multiple trauma evaluated by ultrasonography: a prospective analysis of 291 patients. <i>J Trauma</i> 32: 452-458	II	Documented high sensitivity (89%) and specificity (97%) for US in patients with ISS > 20. False negative results limited by surveillance of indeterminate US with DPL, CT, or exploratory laparotomy.
Boulanger BR	1993	The clinical significance of acute hyperamylasemia after blunt trauma. <i>Can J Surgery</i> 36: 63-69	II	Admission serum amylase levels should not be used to determine clinical or radiographic evaluation of patients with BAT.
Forster R	1993	Ultrasonography in blunt abdominal trauma: influence of the investigators' experience. <i>J Trauma</i> 34: 264-269	II	US performed by surgeons has high sensitivity (96%) and specificity (95%) with a short learning phase.
Jaffin JH	1993	Alkaline phosphatase levels in diagnostic peritoneal lavage fluid as a predictor of hollow visceral injury. <i>J Trauma</i> 34: 829-833	II	Routine measurement of alkaline phosphatase in DPL fluid is not cost-effective.
Glaser K	1994	Ultrasonography in the management of blunt abdominal and thoracic trauma. <i>Arch Surgery</i> 129: 743-747	II	Retrospective review of US performed as the initial diagnostic modality in 1151 patients. US provides results similar to CT and DPL (99% sensitivity, 98% specificity) at less cost and without complications. US inaccurate in diagnosis of small bowel perforations.

Ma OJ	1995	Evaluation of hemoperitoneum using a single - vs multiple -view ultrasonographic examination. <i>Acad Emerg Med</i> 2: 581-586	II	Comparison of single -view (right intercostal oblique) with multiple -view US. Sensitivity greater with multiple -view US (87% vs 51%). Specificity 100% with both techniques.
Nagy KK	1995	Aspiration of free blood from the peritoneal cavity does not mandate immediate laparotomy. <i>Am Surgeon</i> 61: 790-795	II	Comparison of aspiration of gross blood on DPL to actual clinical results in 566 patients who sustained blunt and penetrating abdominal trauma. Aspiration of < 5 ml free blood associated with > 20% non-therapeutic laparotomy rate.
Boulanger BR	1995	A prospective study of emergent abdominal sonography after blunt trauma. <i>J Trauma</i> 39: 325-330	II	Comparison of US with DPL and CT for detection of intraperitoneal fluid. US performed in mean time of 2.6 mins with 81% sensitivity, 98% specificity and 96% accuracy. US is a rapid , accurate examination for initial evaluation of free intraperitoneal following BAT.
Boulanger BR	1996	Emergent abdominal sonography as a screening test in a new diagnostic algorithm for blunt trauma. <i>J Trauma</i> 40: 867-874	II	Description of a diagnostic algorithm using US in BAT. Documented 94% accuracy in 400 patients studied; US exam completed in < 3 mins (82%). US is a rapid and accurate diagnostic modality.
Gow KW	1996	Validity of visual inspection of diagnostic peritoneal lavage fluid. <i>Can J Surgery</i> 39: 114-119	II	Determine predictive value of visual inspection of DPL fluid for identification of intra -abdominal injury. Visual inspection found to have good NPV (98.9%) but poor PPV (52.0%). Hemodynamically stable patients with (+)DPL by visual inspection should have fluid tested before exploratory laparotomy.
Healy MA	1996	A prospective evaluation of abdominal ultrasound in blunt trauma: Is it useful? <i>J Trauma</i> 40: 875-883	II	Assessment of accuracy of technician -performed US in evaluation of 796 patients with BAT. US demonstrated 88.2% sensitive, 97.7% specific, 72.3% PPV and 99.2% NPV. Accuracy of US consistent with other diagnostic modalities.
McKenney MG	1996	1000 consecutive ultrasounds for blunt abdominal trauma. <i>J Trauma</i> 40: 607-610	II	Assessment of utility of US in patients with indications for DPL or CT. US demonstrated 88% sensitivity, 99% specificity, 97% accuracy. (+)US in hemodynamically unstable patients or in the presence of decreasing hematocrit mandates exploratory laparotomy. CT scan following (+)US in hemodynamically stable patients permits selection for non -operative management.
Branney SW	1997	Ultrasound based key clinical pathway reduces the use of hospital resources for the evaluation of blunt abdominal trauma. <i>J Trauma</i> 42: 1086-1090	II	Prospective, non -randomized study of US protocol for evaluation of BAT compared with retrospective controls using CT and DPL. Decreased use of DPL and CT by 74% and 58% respectively without missed injuries. US safe and cost -effective diagnostic modality for evaluation of BAT.
McElveen TS	1997	The role of ultrasonography in blunt abdominal trauma: a prospective study. <i>Am Surgeon</i> 63: 184-188	II	Comparison of surgeon -performed US to CT and DPL in 82 patients with BAT. Ultrasound found to be 88% sensitive, 98% specific, 96% accurate; NPV = 97%, positive predictive value (PPV) = 93%. US accurate and may be performed with minimal training.

Blow O	1998	Speed and efficiency in the resuscitation of blunt trauma patients with multiple injuries: the advantage of diagnostic peritoneal lavage over abdominal computed tomography. <i>J Trauma</i> 44: 287-290	II	Sensitivity and specificity of DPL similar to CT scan in patients with hemodynamic instability, severe TBI or multiple injuries. DPL is more efficient and may be performed with lower cost.
Livingston DH	1998	Admission or observation is not necessary after a negative abdominal computed tomographic scan in patients with suspected blunt abdominal trauma: results of a prospective, multi-institutional trial. <i>J Trauma</i> 44: 273-282.	II	Study demonstrates 99.63% negative predictive value (NPV) for CT scan performed in 2774 patients following blunt abdominal trauma (BAT). CT scan detected 22/25 hollow visceral injuries. Patients with (-) CT scan may be safely discharged.
Fischer RP	1978	Diagnostic peritoneal lavage: fourteen years and 2,586 patients later. <i>Am J Surgery</i> 136: 701-704	III	Organ-specific accuracy of DPL documented for spleen (98.5%), liver (97.1%), small bowel (91.3%), intraperitoneal bladder (66.7%), and diaphragm (59.1%). Compared with historical controls, DPL decreased rate of unnecessary laparotomy from 13% to 6% and decreased mortality from 46.4% to 30%. Decreased mortality presumed due to decreased incidence of missed injury with clinical observation alone.
Hubbard SG	1979	Diagnostic errors with peritoneal lavage in patients with pelvic fractures. <i>Arch Surgery</i> 114: 844-846	III	Accuracy of DPL significantly reduced in the presence of a pelvis fracture. Additional diagnostic tests recommended in hemodynamically stable patients with pelvis fracture and (+)DPL.
Bagwell CE	1980	Blunt abdominal trauma: exploratory laparotomy or peritoneal lavage? <i>Am J Surgery</i> 140: 368-373	III	DPL should be considered mandatory in hemodynamically stable patients with altered mental status or multiple injuries.
Robbs JV	1980	Blunt abdominal trauma with jejunal injury: a review. <i>J Trauma</i> 20: 308-311	III	Clinical findings of pain, tenderness, guarding, absent bowel sounds, and hypovolemia correlate with jejunal injury. Paracentesis (i.e. four-quadrant aspiration) recommended in patients with multiple injuries, concomitant closed head injury, or impaired level of consciousness. If paracentesis is negative, DPL is indicated.
Sherwood R	1980	Minilaparoscopy for blunt abdominal trauma. <i>Arch Surgery</i> 115: 672-673	III	Mini-laparoscopy allows direct visualization of the extent and source of hemorrhage in BAT patients with (1) altered sensorium, (2) multi-system trauma, (3) unexplained hypotension, or (4) equivocal findings on PE. In addition, the clinical importance of intra-abdominal hemorrhage may be determined.
Ward RE	1981	Study and management of blunt trauma in the immediate post-impact period. <i>Rad Clin NA</i> 19: 3-7	III	Splanchnic angiography should be considered as a complement to DPL in patients with (1) pelvis fractures, (2) indications for thoracic aortography, and (3) perplexing abdominal findings.

Alyono D	1982	Significance of repeating diagnostic peritoneal lavage. <i>Surgery</i> 91: 656-659	III	Repeat DPL performed at 1-2 hrs has a high degree of sensitivity, specificity and accuracy in patients with indeterminate initial DPL (ie. DPL-RBC = 50-100 K/mm ³ , DPL-WBC = 100-500/mm ³).
Smith SB	1982	Abdominal trauma: the limited role of peritoneal lavage. <i>Am Surgeon</i> 48: 514-517	III	High degree of accuracy demonstrated with PE in patients capable of a reliable PE. DPL is very sensitive and is associated with a high non-therapeutic laparotomy rate
Berci G	1983	Emergency mini-laparoscopy in abdominal trauma. An update. <i>Am J Surgery</i> 146: 261-265	III	Laparoscopy safer, faster, and more accurate than DPL. Identification of intra-abdominal blood without an identified injury permits non-operative management and decreases the rate of unnecessary exploratory laparotomies.
Burney RE	1983	Diagnosis of isolated small bowel injury following blunt abdominal trauma. <i>Ann Emerg Med</i> 12: 71-74	III	Abdominal pain was a universal symptom in patients who communicate. Other predictive findings on PE included diffuse abdominal tenderness, abdominal rigidity, and absence of bowel sounds. DPL was the most sensitive diagnostic modality for small bowel injury.
Soderstrom CA	1983	The diagnosis of intra-abdominal injury in patients with cervical cord trauma. <i>J Trauma</i> 23: 1061-1065	III	All significant intra-abdominal injuries diagnosed by DPL in patients with cervical cord injuries. Recommend DPL to exclude intra-abdominal injury in BAT patients with concomitant cervical cord injuries.
Berry TK	1984	Diagnostic peritoneal lavage in blunt trauma patients with coagulopathy. <i>Ann Emerg Med</i> 13: 879-880	III	Accuracy of DPL not diminished by presence of coagulopathy. Exploratory laparotomy is indicated in patients with (+)DPL with post-traumatic coagulopathy.
Mustard RA	1984	Blunt splenic trauma: diagnosis and management. <i>Can J Surgery</i> 27: 330-333	III	DPL diagnostic in 86 patients with splenic injury documented by exploratory laparotomy.
McLellan BA	1985	Analysis of peritoneal lavage parameters in blunt abdominal trauma. <i>J Trauma</i> 25: 393-399	III	Based on significant number of therapeutic laparotomies, DPL RBC count > 20K/mm ³ recommended as indication for exploratory laparotomy.
Trooskin SZ	1985	Peritoneal lavage in patients with normal mentation and hematuria after blunt trauma. S, G & O 160: 145-147	III	DPL reveals injuries which require surgery in 45% of BAT patients with normal mentation and hematuria. DPL recommended in patients with BAT who present with hematuria in the presence of normal neurologic examination.
Van Dongen LM	1985	Peritoneal lavage in closed abdominal injury. <i>Injury</i> 16: 227-229	III	DPL may be overly sensitive in evaluation of BAT.
Webster VJ	1985	Abdominal trauma: pre-operative assessment and postoperative problems in intensive care. <i>Anaest & Int Care</i> 13: 258-262	III	CT scan has significantly impacted the use of other diagnostic modalities in the evaluation of hemodynamically stable patients with BAT.

Ryan JJ	1986	Critical analysis of open peritoneal lavage in blunt abdominal trauma. <i>Am J Surgery</i> 151: 221-223	III	False positive rate for peritoneal lavage higher than previously reported (sensitivity = 83%) resulting in 27% non -therapeutic laparotomy rate.
Kane NM	1987	Efficacy of CT following peritoneal lavage in abdominal trauma. <i>J Comp Asst Tomo</i> 11: 998-1002	III	CT revealed substantial intra-abdominal or retroperitoneal injuries in 1/3 patients who underwent CT following DPL. CT recommended when clinical status equivocal regardless of DPL results.
Pattyn P	1989	Peritoneal lavage after abdominal trauma: indications, technique, results. <i>Int Surgery</i> 74: 17-19	III	Based on high sensitivity (false negative rate = 1%) and low incidence of complications (0.5%), DPL recommended for the evaluation of BAT.
Wening JV	1989	Evaluation of ultrasound, lavage, and computed tomography in blunt abdominal trauma. <i>Surg Endoscopy</i> 3: 152-158	III	Demonstrated 84% sensitivity and 97% specificity for US in evaluation of BAT. US is a reliable, fast and repeatable diagnostic modality.
Ceraldi CM	1990	Computerized tomography as an indicator of isolated mesenteric c injury. A comparison with peritoneal lavage. <i>Am Surgeon</i> 561: 806-810	III	Sensitivity of CT inadequate to reliably exclude mesenteric injury. DPL recommended as a more sensitive diagnostic modality.
D'Amelio LF	1990	A reassessment of the peritoneal lavage leukocyte count in blunt abdominal trauma. <i>J Trauma</i> 30: 1291-1293	III	Elevated DPL fluid WBC count (> 500/mm ³) has on diagnostic value in the early (< 4 hrs) post -injury period. Isolated elevation of DPL WBC count may be more useful in delayed setting or in the presence of equivocal PE.
Davis JW	1990	Complications in evaluating abdominal trauma: diagnostic peritoneal lavage versus computerized axial tomography. <i>J Trauma</i> 30: 1506-1509	III	Significantly lower complication rate for DPL compared to CT scan (0.9% vs 3.4%) with no difference in preventable deaths.
Henneman PL	1990	Diagnostic peritoneal lavage: accuracy in predicting necessary laparotomy following blunt and penetrating trauma. <i>J Trauma</i> 30: 1345-1355	III	Semi-open DPL 96% accurate for prediction of need for exploratory laparotomy in BAT and 92% accurate in the presence of pelvic fracture.
Hawkins ML	1990	Is diagnostic peritoneal lavage for blunt trauma obsolete? <i>Am Surgeon</i> 56: 96-99	III	Ease, safety (1% complication rate) and accuracy of DPL (97%) justify continued use in evaluation of BAT.
Jacobs DG	1990	Peritoneal lavage white count: a reassessment. <i>J Trauma</i> 30: 607-612	III	Isolated elevation of DPL WBC count > 500/mm ³ not specific for diagnosis of intra -abdominal injury. Specificity increases with repeat DPL.

Lang EK	1990	Intra-abdominal and retroperitoneal organ injuries diagnosed on dynamic computed tomograms obtained for assessment of renal trauma. <i>J Trauma</i> 30: 1161-1168	III	CT scan helpful in the diagnosis of unsuspected abdominal or retroperitoneal injuries in the evaluation of patients for renal trauma. Patients with (-)CT may be safely observed.
Matsubara TK	1990	Computed tomography of abdomen (CTA) in management of blunt abdominal trauma. <i>J Trauma</i> 30: 410-414	III	CT is a valuable diagnostic modality in hemodynamically stable patients with BAT if performed correctly and interpreted accurately. Patients with (-)CT should be admitted for observation.
Megison SM	1990	The value of alkaline phosphatase in peritoneal lavage. <i>Ann Emerg Med</i> 19: 503-505	III	Measurement of alkaline phosphatase in DPL fluid adds no diagnostic advantage in identification of intestinal injury.
Soyka JM	1990	Diagnostic peritoneal lavage: is an isolated WBC count greater than or equal to 500/mm ³ predictive of intra-abdominal injury requiring celiotomy in blunt trauma patients? <i>J Trauma</i> 30: 874-879	III	Isolated elevation of DPL WBC count > 500/mm ³ should not be an indication for exploratory laparotomy in BAT.
Barba C	1991	Is positive diagnostic peritoneal lavage an absolute indication for laparotomy in all patients with blunt trauma? <i>Can J Surgery</i> 34: 442-445	III	Immediate exploratory laparotomy not necessarily mandated in the presence of (+)DPL. Additional diagnostic studies should be considered in hemodynamically stable patients with (+)DPL.
Berci G	1991	Emergency Laparoscopy. <i>Am J Surgery</i> 161: 332-335	III	Diagnostic laparoscopy (DL) is a viable diagnostic modality in the evaluation of BAT. DL lowers incidence of non -therapeutic exploratory laparotomy.
Davis JW	1991	Base deficit as an indicator of significant abdominal injury. <i>Ann Emerg Med</i> 20: 842-844	III	Base deficit (BD) < - 6.0 is a sensitive indicator of intra -abdominal injury in BAT. DPL or CT recommended for patients with BD < - 6.0.
DeMaria EJ	1991	Management of patients with indeterminate diagnostic peritoneal lavage results following blunt trauma. <i>J Trauma</i> 31: 1627-1631	III	Indeterminate DPL correlates with injuries that may be managed non-operatively. CT recommended following indeterminate DPL rather than repeat DPL.
Fryer JP	1991	Diagnostic peritoneal lavage as an indicator for therapeutic surgery. <i>Can J Surgery</i> 34: 471 -476	III	Sixty-five percent (65%) of patients who underwent exploratory laparotomy for (+) DPL had therapeutic laparotomies.
McAnena OJ	1991	Contributions of peritoneal lavage enzyme determinations to the management of isolated hollow visceral abdominal injuries. <i>Ann Emerg Med</i> 20: 834-837	III	Elevation of DPL fluid amylase is highly specific for isolated small bowel injury. Recommend routine enzyme determinations for DPL effluent as a marker for small bowel injury.


McAnena OJ	1991	Peritoneal lavage enzyme determinations following blunt and penetrating abdominal trauma. <i>J Trauma</i> 31: 1161-1164	III	Elevated amylase and alkaline phosphatase levels in DPL fluid increases index of suspicion for presence of a small bowel injury in patients with (-)DPL by RBC count.
Mure AJ	1991	Serum amylase determination and blunt abdominal trauma. <i>Am Surgeon</i> 57: 210-213	III	Serum amylase has poor sensitivity, specificity and PPV for the diagnosis of intra -abdominal injury. Routine serum amylase has no value in the evaluation of BAT.
Perez FG	1991	Evaluation of the abdomen in intoxicated patients: is computed tomography scan or peritoneal lavage always indicated? <i>Ann Emerg Med</i> 20: 500-502	III	Legally intoxicated patients with normal mentation may be reliably assessed by physical examination. Elevated serum ethanol does not mandate CT or DPL.
Sahdev P	1991	Evaluation of liver function tests in screening for intra -abdominal injuries. <i>Ann Emerg Med</i> 20: 838-841	III	Elevated liver function tests associated with injury to the liver. Patients with elevated liver function tests should undergo CT scan.
Day AC	1992	Diagnostic peritoneal lavage: integration with clinical information to improve diagnostic performance. <i>J Trauma</i> 32: 52-57	I	Combination of clinical evaluation and DPL reduces rate of non - therapeutic laparotomies, but increases the number of missed injuries. The highest accuracy (95%) is obtained by combination of circulatory assessment and DPL.
Driscoll P	1992	Diagnostic peritoneal lavage: it's red but is it positive? <i>Injury</i> 23: 267-269	III	Visual assessment of DPL fluid RBC -count inaccurate.
Knudson MM	1992	Hematuria as a predictor of abdominal injury after blunt trauma. <i>Am J Surgery</i> 164: 482-485	III	Hematuria is a marker for renal or extra -renal intra -abdominal injury. CT recommended in the presence of hematuria with shock.
Pattimore D	1992	Torso injury patterns and mechanisms in car crashes: an additional diagnostic tool. <i>Injury</i> 23: 123-126	III	Injuries to the spleen, liver, pelvis and aorta more likely with side impact compared to front impact collisions.
Wyatt JP	1992	Variation among trainee surgeons in interpreting diagnostic peritoneal lavage fluid in blunt abdominal trauma. <i>J Royal Coll Surg (Edin)</i> 37: 104-106	III	Estimation of DPL RBC -count by visual inspection inaccurate compared to microscopic analysis. Recommend quantitative cell count vs visual assessment of DPL fluid to make decision on management.
Baron BJ	1993	Nonoperative management of blunt abdominal trauma: the role of sequential diagnostic peritoneal lavage, computed tomography, and angiography. <i>Ann Emerg Med</i> 22: 1556-1562	III	Combined modalities of CT scan and an angiography in hemodynamically stable patients with (+)DPL reduces the non - therapeutic laparotomy rate.

Bode PJ	1993	Abdominal ultrasound as a reliable indicator for conclusive laparotomy in blunt abdominal trauma. <i>J Trauma</i> 34: 27-31	III	US demonstrated high sensitivity (92.8%) and specificity (100%). Routine US recommended for 1) abdominal findings not initially felt to warrant immediate laparotomy, 2) equivocal results on initial US, 3) deteriorating clinical situation.
Visvanathan R	1993	Blunt abdominal I trauma - injury assessment in relation to early surgery. <i>J Royal Coll Surg (Edin)</i> 38: 19-22	III	DPL highly sensitive (95%) with 81% specificity and 89% accuracy. Combination of DPL and US facilitates early assessment and management of abdominal injuries .
Mendez C	1994	Diagnostic accuracy of peritoneal lavage in patients with pelvic fractures. <i>Arch Surgery</i> 129: 477-481	III	Registry study of 286 open DPLs performed in patients with BAT in the presence of a pelvis fracture. Open DPL accurate modality (94% sensitivity; 99% specificity) for evaluation of patients with multiple injuries, including pelvis fractures.
Tibbs PA	1980	Diagnosis of acute abdominal injuries in patients with spinal shock: value of diagnostic peritoneal lavage. <i>J Trauma</i> 20: 55-57	III	DPL recommended for exclusion of intra -abdominal injuries in spinal cord injured patients with complete neurologic deficit.
Baron BJ	1993	Nonoperative management of blunt abdominal trauma: the role of sequential diagnostic peritoneal lavage, computed tomography, and angiography. <i>Ann Emerg Med</i> 22: 1556-1562	III	Combined modalities of CT scan and angiography in hemodynamically stable patients with (+)DPL reduces the non - therapeutic laparotomy rate.
Mendez C	1994	Diagnostic accuracy of peritoneal lavage in patients with pelvic fractures. <i>Arch Surgery</i> 129: 477-481	III	Registry study of 286 open DPLs performed in patients with BAT in the presence of a pelvis fracture. Open DPL accurate modality (94% sensitivity; 99% specificity) for evaluation of patients with multiple injuries, including pelvis fractures.
Grieshop NA	1995	Selective use of computed tomography and diagnostic peritoneal lavage in blunt abdominal trauma. <i>J Trauma</i> 38: 727-731	III	Registry review of 956 hemodynamically stable patients with reliable neurologic examinations (GCS > 11). Patients with abnormal PE, chest injury or gross hematuria have high incidence of intra-abdominal injury which require exploratory laparotomy. No CT required in patients with normal PE, no chest injury and no hematuria. Elevated blood alcohol does not alter accuracy of PE.
Noian BW	1995	Mesenteric injury from blunt abdominal trauma. <i>Am Surgeon</i> 61: 501-506	III	Review of 27 patients with mesenteric injury following BAT. CT performed in 10 patients; failed to detect mesenteric injury in 7. High index of suspicion required to identify patients with mesenteric injury. CT scan is insufficient diagnostic modality for this injury and may result in missed injuries to mesentery and small bowel.


Chandler CF	1997	Seatbelt sign following blunt trauma is associated with increased incidence of abdominal injury. <i>Ann Surgeon</i> 63: 885-888	III	Evaluation of seatbelt sign (SBS) as predictor of intra -abdominal injury in 14 patients. Sensitivity for solid visceral injuries was 85% and 100% with CT scan and DPL respectively; sensitivity for hollow visceral injuries was 33% and 100% respectively. Negative CT scan in patients with SBS mandates admission and observation. Free fluid on CT scan warrants further investigation (i.e. DPL or exploratory laparotomy).
Kern SJ	1997	Sonographic examination of abdominal trauma by senior surgical residents. <i>Ann Surgeon</i> 63: 669-674	III	Evaluation of focused abdominal sonography for trauma (FAST) following PE in 518 patients (92.4% blunt / 7.6% penetrating). FAST examination 73.3% sensitive, 97.5% specific with 98.3% NPV and 96.1% PPV. Low sensitivity due to missed hollow visceral injuries.
McKenney KL	1997	Cost reduction using ultrasound in blunt abdominal trauma. <i>Emerg Radiology</i> 4: 3-6	III	Comparison of 626 patients (Group 1) evaluated with CT and DPL with 564 patients (Group 2). Use of DPL and CT decreased by 94% and 63% respectively in Group 2. Decreased cost / patient by \$170. Recommend US as the initial diagnostic test of choice in BAT with unreliable PE. US replaces DPL and allows more resource-efficient use of CT scan.
Porter RS	1997	Use of ultrasound to determine need for laparotomy in trauma patients. <i>Ann Emerg Med</i> 29: 323-330	III	Retrospective review of technician -performed US in 1631 patients without controls. Sensitivity and specificity of US 93% and 90% respectively. US safe and cost -effective diagnostic modality in the evaluation of BAT.
Schurink GW	1997	The value of physical examination in the diagnosis of patients with blunt abdominal trauma: a retrospective study. <i>Injury</i> 28: 261-265	III	Retrospective study of physical examination (PE) in 204 patients with BAT. Patients with isolated TBI, low -impact rib pain, isolated abdominal trauma may be evaluated with PE plus ultrasound (US) with > 85% NPV. Higher incidence of intra -abdominal injury in the presence of low rib fractures and high -energy impact, therefore follow-up CT scan recommended in presence of normal US.
Sherbourne CD	1997	Visceral injury without hemoperitoneum: a limitation of screening abdominal sonography for trauma. <i>Emerg Radiology</i> 4: 349-354	III	Review of 196 patients with intra -abdominal injury: 50/196 (26%) had no hemoperitoneum. Fifteen of 50 patients had (-)FAST examination. US may fail to detect intra -abdominal injuries in the absence of hemoperitoneum.
Takishima T	1997	Serum amylase level on admission in the diagnosis of blunt injury to the pancreas. <i>Ann Surgery</i> 226: 70-76	III	Serum amylase elevated in 84% of patients with pancreatic injury at presentation; elevated in 76% (< 3 hrs post -injury) and 100% (> 3 hrs post-injury). Serum amylase must be measured at least 3 hrs post-injury to avoid missed injuries.

Buzzas GR	1998	A comparison of sonographic examinations for trauma performed by surgeons and radiologists. <i>J Trauma</i> 44: 604-608	III	Comparison of FAST performed by surgical residents (Group A) and US technicians/radiologists (Group B). Sensitivity 73.3% and 79.5% for Group A and B respectively. Specificity 97.5% and 99.3% for Group A and B respectively. Sensitivity improved with exclusion of hollow visceral injuries.
Smith SR	1998	Institutional learning curve of surgeon-performed trauma ultrasound. <i>Arch Surg</i> 133: 530-536	III	Sensitivity and specificity of FAST examination 73% and 98% respectively and may be learned without significant learning curve. Modality unreliable for detection of hollow visceral injuries.

Evaluation of Blunt Abdominal Trauma: Unstable Patient



Evaluation of Blunt Abdominal Trauma: Stable



*CT scan may be elected based on institutional experience or clinical suspicion of intra-abdominal injury.