

This is a peer-reviewed, accepted author manuscript of the following article: Nusurupia, J., Reilly, J. J., & Janssen, X. (2021). Pre-school children's compliance with the WHO 24-hour movement behaviour guidelines: a systematic review of Sub-Saharan African studies. *African Journal of Physical Activity and Health Science*, 26(4), 345-359. <https://doi.org/10.37597/ajphes.2020.26.4.1>

# 1 Pre-school children's compliance with the WHO 24-hour 2 movement behaviour guidelines: A systematic review of Sub- 3 Saharan African studies

4  
5 JACKLINE NUSURUPIA <sup>1,2\*</sup>, JOHN J. REILLY <sup>1</sup> AND XANNE JANSSEN <sup>1</sup>

6  
7 <sup>1</sup>*Faculty of Humanities and Social Sciences, University of Strathclyde, Physical Activity for Health*  
8 *Group, Glasgow G1 1XQ, Scotland. Email: jacklinejema10@gmail.com*

9 <sup>2</sup>*Tanzania Food and Nutrition Centre, 22 Ocean Road, P.O. Box 922 Dar es Salaam, Tanzania*

10  
11 *(Received: ; Revision Accepted: )*

12 *ORCID nos.: JN (N/A), JJR (0000-0001-6165-5471), XJ (0000-0003-1355-0792)*

13 *DOI:*

## 14 15 **Abstract**

16  
17 This review aimed to quantify the extent to which levels of habitual physical activity, sedentary  
18 behaviour, and sleep complied with WHO Guidelines in 3- to 4-year-olds in Sub-Saharan Africa.  
19 The systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-  
20 Analyses (PRISMA). A literature search was conducted in August 2019 from PubMed, Web of  
21 Science, and Medline Ovid databases as well as grey literature. Eligible studies were assessed for  
22 quality using the Joanna Briggs Institute (JBI) tool for prevalence studies. Only three studies were  
23 eligible: two from one South African sample and one from Ghana which did not contain useable  
24 data. The South African studies reported high compliance with physical activity guidelines (n=87),  
25 with a mean total physical activity volume of 446 min/day (SD=67) and a median level of  
26 moderate-to-vigorous physical activity of 109 min/day (SD=37). Results further reflected a lower  
27 compliance with the screen time guidelines (n=56; mean screen time 2 h/day, SD=1.3), with no  
28 studies reported on sleep. There are insufficient data to draw meaningful conclusions on pre-school  
29 children's adherence to the 24-hour movement guidelines in Sub-Saharan Africa; thus warranting  
30 the need for future research in this region.

31  
32 **Keywords:** Pre-school children, physical activity, sedentary behaviour, sleep, Sub-Saharan Africa.

## 33 34 **How to cite this article:**

35 Nusurupia, J., Reilly, J.J. & Janssen, X. (2020). A systematic review of compliance with the WHO  
36 24-hour movement behaviour guidelines in pre-school children in Sub-Saharan Africa. *African*  
37 *Journal for Physical Activity and Health Sciences*, 26(4), . DOI:  
38 <https://doi.org/10.37597/ajphes.2020.26.4>.

## 39 40 **Introduction**

41  
42 Physical inactivity has been identified as one of the most important factors for  
43 global mortality and a contributor to the global rise in overweight and obesity. The  
44 number of overweight or obese infants and young children (age 0 to 5 years)  
45 increased from 32 million globally in 1990 to 40 million in 2016. This increase  
46 has been projected to reach 50 million by 2030 (World Obesity Federation, 2019).

47 In a global effort to counter the rise in overweight and obesity, the World Health  
48 Organisation (WHO) developed 24-hour movement behaviour guidelines for the  
49 early years. These guidelines comprise physical activity, sedentary behaviour, and  
50 sleep (WHO, 2019). The guidelines state that children aged 3-4.9 years should  
51 participate in 180 min of daily physical activity including 60 minutes of moderate-  
52 to-vigorous physical activity (MVPA). In addition, children should spend less than  
53 1 hour behind a screen and less than 1 hour being restrained; and 10-13 hours of  
54 good quality sleep (WHO, 2019).

55  
56 Several studies have found that among school-aged children and adolescents'  
57 levels of physical activity are lower than recommended (Aubert *et al.*, 2018;  
58 Reilly, 2015; Zhu *et al.*, 2019). Only a minority of children and adolescents  
59 globally meet the minimum recommendation of 60 minutes MVPA/day (Aubert  
60 *et al.*, 2018; Guthold *et al.*, 2018). However, there appears to be less global  
61 evidence on levels of physical activity in early childhood (Aubert *et al.*, 2018).  
62 This is because the concept of 24-hour movement behaviours is relatively new and  
63 the first global guidelines for these behaviours in young children were only  
64 published in 2019 (WHO, 2019). Therefore, it is unclear whether habitual levels  
65 of physical activity, sedentary behaviour, and sleep are adequate in children under  
66 5 years of age globally. A few recent studies from high-income countries  
67 suggested that compliance to WHO 2019 guidelines might be low. These studies  
68 indicated that only a minority of children meet combined guidelines for physical  
69 activity, sleep, and sedentary behaviour (screen time) (Cliff *et al.*, 2017; De  
70 Craemer *et al.*, 2018; Lee *et al.*, 2017).

71  
72 Childhood obesity may be a concern for public health in Sub-Saharan Africa  
73 because the childhood population is rapidly increasing and urbanising. Africa's  
74 childhood population is projected to reach 1 billion by 2025, which will make it  
75 the biggest childhood population compared to all the other continents (United  
76 Nations Children's Fund, 2019). A recent study has shown that excessive fatness  
77 was present in nearly a third of children across urban areas in Africa (Diouf *et al.*,  
78 2018). This is consistent with the conclusions of another systematic review which  
79 linked urbanisation with a growing tendency of decreasing physical activity,  
80 increasing sedentary behaviours and decreasing fitness among children and  
81 adolescents in Africa (Muthuri *et al.*, 2014).

82  
83 The study's primary aim, therefore, was to establish the available evidence on  
84 compliance with the 24-hour movement behaviour guidelines among 3- to 5-year-  
85 old children in Sub-Saharan Africa. A secondary aim was to identify gaps in the  
86 available evidence and develop future research questions.

## 87 88 **Methodology**

89  
90 This systematic review was conducted and reported by following the Preferred  
91 Reporting Items for Systematic Reviews (PRISMA) guidelines. PRISMA has

92 been defined as evidence-based guidelines with a minimal set of items for  
93 reporting systematic reviews and meta-analysis (Moher *et al.*, 2009). The protocol  
94 for the review was registered on PROSPERO on the 6<sup>th</sup> of August 2019 with  
95 Prospero registration number: CRD42019140909. The main search terms were  
96 developed following the PECO (population, exposure, comparator, outcome)  
97 format, but only population (Preschool children of 3.0 to 4.9 years in Sub-Saharan  
98 Africa), and exposure (time spent in physical activity, while sleeping and being  
99 sedentary, including screen time) were included.

#### 100 101 *Study inclusion and exclusion criteria*

102 In this review, the following studies were included: studies which included  
103 healthy, typically developing, children aged 3.0 to 4.9 years residing in Sub-  
104 Saharan Africa and reported on objectively or subjectively measured habitual time  
105 spent in physical activity, or while sleeping or during sedentary engagements,  
106 including screen time. Only articles published in English language were  
107 considered. All the articles, which fulfilled the above inclusion criteria, were  
108 examined regardless of the years when published. Articles were excluded if they  
109 were non-English language publications, if participating children were less than  
110 3.0 years and older than 4.9 years, reported studies with children who had a  
111 physical or learning disability, or who were unhealthy.

#### 112 113 *Literature search*

114 Three databases (Medline Ovid, PubMed, and Web of Science) were searched in  
115 August 2019. The searches were independently performed by two researchers, one  
116 of whom was a specialist librarian, using the following search terms: population  
117 (preschool, child) and exposure (time spent in physical activities, sleeping and  
118 time spent in sedentary behaviours including screen time); the search terms were  
119 used in separation or in combination using the boolean operator like "OR" and  
120 "AND". The literature search was restricted to Sub-Sahara Africa. An example of  
121 a search strategy from Web of Science is attached as an appendix (Appendix 1),  
122 which was subsequently adapted for the other two databases. References were  
123 exported, de-duplicated and reviewed using Endnote X9. Two researchers  
124 independently assessed the titles/abstracts (JN and XJ) and full-text articles were  
125 obtained for those papers which met the inclusion criteria. Full-text articles were  
126 screened (JN and XJ) and the two researchers resolved differences in opinion over  
127 the eligibility of papers to be included by discussion, referring to a third author  
128 (JR) where necessary.

#### 129 130 *Grey literature searching*

131 Grey literature was searched through each country's domain website (e.g. Ghana  
132 – com.gh and South Africa – com.za), whereby in the advanced google search box  
133 the following words were entered: Physical activity OR sedentary behaviour OR  
134 Screen OR Sleep AND preschool children, then searched.

135  
136

### *Data extraction, quality assessment and synthesis*

Two researchers independently used the adapted pre-specified excel data sheet to extract all the necessary data from eligible studies (JN and XJ). The necessary data extracted from the studies included the following: study location, design, exposure type (physical activity, sedentary behaviour, and/or sleep), exposure description, exposure measurement type, age and sex of sample, and summary measures of time spent in the behaviours. Any minor disagreements between the two researchers over data extracted were resolved through discussion. The data-extracted excel sheet is provided in Appendix 2.

To test for compliance with recommendations in relation to 24-hour movement behaviours, we used the WHO 2019 guidelines for 3.0 to 4.9-year olds stating that children should, in a 24-hour period: 1) spend at least 180 minutes in physical activity, of which at least 60 minutes should in MVPA; 2) Spend 10.0-13.0 hours asleep; 3) Spend no more than 1 hour per day using a screen. Authors of the present study contacted the corresponding author of one of the eligible studies (Draper *et al.*, 2017) for clarification of data presented in that study and the corresponding author graciously provided the information which was subsequently used in the analysis. Two researchers (JN and JR) independently used the Joanna Briggs Institute (JBI) checklist of the quality of prevalence studies and assessed the quality of individual studies (risk of bias) which were included in this review (The Joanna Briggs Institute, 2009). Minor differences between the two researchers were resolved by discussion.


## **Results**

### *Eligible studies and study selection process*

A total of 2835 and 9 records were found via the database and grey literature searches, respectively. After de-duplication, 2832 papers were screened for inclusion. Subsequent to assessing the title and abstract, 2821 were removed from the record as they did not meet the inclusion criteria. Therefore, 11 articles remained which were considered for full-text screening. Finally, four studies using two unique samples were considered to be eligible and included in this systematic review (2019; Draper *et al.*, Draper *et al.*, 2017; Okyere, 2018; Tomaz, 2018). Of these four studies, one was conducted in Ghana (Okyere, 2018) and the other three in South Africa (Draper *et al.*, 2019; Draper *et al.*, 2017; Tomaz, 2018). A schematic overview of the steps used in the systematic review process is illustrated in the PRISMA flow diagram presented in Figure 1. The South African studies were reported in one PhD thesis (Tomaz, 2018) and two published studies (Draper *et al.*, 2019; Draper *et al.*, 2017). The three studies addressed different research questions and had slightly different sample sizes - for the present synthesis of review findings we selected the published study with the largest sample size of 3.0-4.9y olds for physical activity (Draper *et al.*, 2017), and the only publication which reported screen time (Tomaz, 2018).

182  
183

**Figure 1:** Prisma flow diagram.



184  
185

187 Table 1 summarises the three eligible studies published based on two unique  
 188 samples. Neither study reported the levels of compliance with recommendations  
 189 in relation to all the three behaviours (physical activity, screen time, and sleep),  
 190 though the South African study reported the compliance levels for total volume of  
 191 physical activity and screen time (Draper *et al.*, 2019; Draper *et al.*, 2017; Tomaz,  
 192 2018). The South African study used convenience sampling in urban and rural  
 193 areas and obtained data on the following behaviours: habitual levels of physical  
 194 activity, sedentary behaviour, and sleep. The study from Ghana measured habitual  
 195 levels of physical activity in pre-school children from a single urban municipality,  
 196 but used the parent-reported Global Physical Activity Questionnaire (GPAQ) to  
 197 measure levels of physical activity which is intended for adults, not children  
 198 (Okyere, 2018).  
 199  
 200

**Table 1:** Summary of eligible studies.

Author/ years	Sample size/Age (n)	Data Collection (years)	How behaviours were measured	Levels of movement behaviours	Comments
<b>Okyere (2018)</b>	N: 201 Age: 3-5- year-olds	2018	Children’s physical activity levels using the GPAQ (parent- reported)	54.2% of the sample reported to do at least 30 min/day MVPA 3-5 times/week	Levels of physical activity and compliance with guidelines unclear
<b>Draper et al. (2017)</b>	N: 87 Age: 3-4 year-olds	2012-2014	Physical activity and sedentary behaviour measured by Actigraph GT3X accelerometer	90.8% of the sample complied with the physical activity recommendation; Mean total daily physical activity 446.4 min/day (SD 67.6) Median daily MVPA 109.3 min/day (SD 36.5)	The same sample as Tomaz et al. (2018); Samples were taken from rural and urban areas
<b>Tomaz (2018)</b>	N: 56 (screen time) Age: 3-4 year-olds	2012-2014	Screen-time by parent questionnaire	Mean screen time (n 56) 2.0 hr/day (SD 1.3)	The same sample as Draper et al. (2017); sample were taken from rural and urban areas

203 The study quality appraisal is summarised in Table 2. The eligible studies  
 204 generally did not set out with nationally representative prevalence estimates as  
 205 their primary aims and consequently did not meet some of the quality criteria in  
 206 the JBI checklist. Eligible study samples were not randomly selected or nationally  
 207 representative (Draper *et al.*, 2019; Draper *et al.*, 2017; Okyere, 2018; Tomaz,  
 208 2018).

209  
 210 **Table 2:** Quality appraisal summary using the Joanna Briggs Institute, (JBI) tool (The Joanna  
 211 Briggs Institute, 2009).

<b>Criteria</b>	<b>Okyere (2018)</b>	<b>Draper et al. (2017)</b>	<b>Tomaz et al. (2018)</b>
<b>Sample frame appropriate</b>	NO	YES	YES
<b>Participants sampled appropriately</b>	YES	NO	NO
<b>Is the sample size adequate?</b>	YES for estimating prevalence in the target population only	YES for purposes of an original study, not as prevalence estimate	YES for purposes of an original study, not as prevalence estimate
<b>Subjects &amp; settings described in detail?</b>	YES	NO, but in subsequent publications	NO, but in subsequent publications
<b>Data analysis conducted with sufficient coverage?</b>	NO	YES	YES
<b>Valid methods used for the identification of all the behaviours</b>	NO	NO, not for all behaviours	NO, not for all behaviours
<b>Standard, reliable method of measuring the behaviours in all participants</b>	NO	YES	YES
<b>Appropriate statistical analysis</b>	NO, not for a prevalence estimate	NO, not for a prevalence estimate	NO, not for a prevalence estimate
<b>Response rate adequate/dealt with</b>	NO	YES	YES

212

213

214

## 215 **Discussion**

216

217 The main aim of this systematic review was to quantify the compliance with the  
 218 recently published WHO guidelines (April 2019) on habitual levels of the 24-hour  
 219 movement behaviours (physical activity, screen time, sleep time) in preschool  
 220 children in Sub-Saharan Africa. The results of the systematic review showed that  
 221 data on compliance with 24-hour movement behaviours in Sub-Saharan Africa  
 222 was very limited, even with relatively broad inclusion criteria which comprised  
 223 both subjective and objective measures of the behaviours. No studies examined all  
 224 three 24-hour movement behaviours combined and evidence on adherence to the

individual components of the guidelines was also limited. Only four studies were eligible for inclusion in this systematic review (Draper *et al.*, 2019; Draper *et al.*, 2017; Okyere, 2018; Tomaz, 2018), using two unique samples, and one study could not be used with confidence as the tool used to measure levels of physical activity (GPAQ) were intended for adults, not children. Childhood obesity in general, and the 24-hour movement behaviours in particular, may not have been a high priority for public health research and surveillance in Africa in the past. The burden of undernutrition and infectious disease, as well as the lack of capacity in, and focus on, childhood obesity may explain the paucity of evidence identified in the present review.

Due to the very limited evidence available for inclusion in this review, results cannot be compared between countries in Sub-Saharan Africa. However, several studies have been carried out in high-income countries (HICs) which can be used for comparison. In a study of 151 Canadian children, the majority of toddlers (age: 19.0 months; SD=1.9 months) met the individual physical activity (99.3%; n=141) and sleep (82.1%, n=124) recommendations, while only 23 toddlers met the screen time recommendation (15.2%) (Lee *et al.*, 2017). Similarly, another study in Canada involving 803 children (mean age: 3.5 years; 95% CI = 3.5, 3.6), showed a high proportion of children meeting physical activity (61.8%) and sleep (83.9%) recommendations, but few met the screen time recommendation (24.4 %), while 3.3% did not meet any of the three recommendations (Chaput *et al.*, 2017). Only 12.7% met all three recommendations contained in the guidelines. Results of an Australian study of 248 preschool children (mean age 4.2 years  $\pm$  0.6 years) showed similar results, with a high proportion of the children meeting physical activity (93.1%) and sleep (88.7%) guidelines, but fewer children complying with the screen time guideline (17.3%), and only 14.9% fulfilling all the three recommendations (Cliff *et al.*, 2017). The GET UP! - A study conducted in Australia, found that of the 202 toddlers studied (mean age: 19.7  $\pm$  4.1 months), only 8.9% met the overall 24-hour movement recommendations, high percentages met physical activity (96.5%) and sleep (79.7%) recommendations, but only 11.4% were compliant with the screen time recommendations (Santos *et al.*, 2017). A recent Belgian study of 1082 pre-schoolers (mean age: 4.2years; SD = 0.02years), of whom 595 had data for all the variables, found that only 10% complied with all WHO recommendations on weekdays and 4% on weekends and that compliance with physical activity guidelines among the Belgian pre-schoolers was low (De Craemer *et al.*, 2018).

In summary, recent studies of compliance with the 24-hour movement behaviour guidelines among Canadian and Australian pre-schoolers found that a high proportion of the children complied with the physical activity and sleep recommendations, but there was very low compliance with meeting all the three recommendations; which suggests that screen time was not well complied with. In contrast, a study of pre-schoolers in Singapore noted that the percentage of children who failed to meet the sleep recommendations was particularly high



270 (Chen *et al.*, 2019). It is possible that there is a common trend in HICs, but cultural  
271 differences and disparity between studies in the accelerometry cut-offs used to  
272 measure total volume of physical activity and MVPA might play a part in  
273 explaining the discrepant findings between studies. Future research will be needed  
274 to determine whether discrepancies between studies in level of compliance with  
275 WHO guidelines are due to methodological differences or attributable to real  
276 differences between populations (e.g. cultural or environmental differences).

277  
278 The South African study included in this review showed high levels of physical  
279 activity with 91% of the sample meeting the guideline for total daily physical  
280 activity, but mean levels of daily screen time were well above the guideline, and  
281 no sleep time data were available. There may be differences in the socio-economic  
282 and cultural environment between studies, which complicate data interpretation.  
283 A large cohort study in Brazil, a middle-income country, suggests that pre-school  
284 children (birth-4years) from higher-income earning families go to bed and wake  
285 up later than children of lower-income families (Netsi *et al.*, 2017). On average  
286 though, the times of going to bed and waking up are shifted by around two hours  
287 later in low and middle-income countries (LMIC's) compared to families from  
288 HIC's (Netsi *et al.*, 2017). However, it was also noted that families with lower  
289 income and maternal education tended to have higher rates of co-sleeping (Netsi  
290 *et al.*, 2017). A study (Chaput *et al.*, 2015), suggests that there is a possibility that  
291 in HICs, children's days are more structured and regulated while in LMICs there  
292 is more flexibility and discretionary time, which might mean that children from  
293 LMICs have less sleep than those from HICs.

294  
295 The present review and the research evidence upon which it was based had several  
296 strengths and limitations that are worth mentioning. The main strengths were the  
297 novelty and timeliness of the review as well as the importance of the research  
298 questions. In addition, the review was conducted and reported rigorously,  
299 following PRISMA guidances and a grey literature search. The limitations were;  
300 first, the number of eligible studies was sparse, thus limiting the ability to make  
301 meaningful conclusions about levels of the 24-hour movement behaviour in  
302 preschool children in Sub-Saharan Africa. The main finding of the present review,  
303 which is considered strikingly timely, is that there is a need for more research on  
304 the subject, especially timely since the WHO (2019) guidelines highlight the  
305 importance of the time spent in the behaviours to individual child health and  
306 wellbeing, and to population efforts to tackle obesity and related non-  
307 communicable diseases. Second, the quality of studies included in the systematic  
308 review was somewhat limited, but they did not set out to measure compliance with  
309 all of the 24-hour movement behaviour guidelines as their primary aim. There is a  
310 need for research with this specific aim in future, and an international effort to  
311 improve surveillance of the 24-hour movement behaviours among pre-school  
312 children at an early stage (Sunrise Study, 2020). Finally, the searches from the  
313 database and grey literature were limited to research published in English  
314 language, excluding non-English studies. Therefore, it is plausible that studies

published in other languages which might have partly met the search criteria, were they written in English, have been excluded.

## Conclusions

Few studies have examined the levels of or compliance with 24-hour movement behaviours in pre-school children in Sub-Saharan Africa to date, and the dearth of data from Africa in general is a concern. Future research on levels of the 24-hour movement behaviours among young children in Sub-Saharan Africa should be conducted. Such studies should consider the principles and practice of obtaining prevalence estimates with confidence, utilise valid methods, and consider the possible influence of urban versus rural settings, and socioeconomic status on compliance with WHO guidelines.

## Acknowledgments

The authors would like to thank the Librarian Ms. Sarah Kevill for help with literature searching, Dr. Cathi E. Draper and Dr. Simone Tomaz for providing data and clarifications from their published papers.

## References

- Aubert, S., Barnes, J. D., Abdeta, C., Nader, P. A., Adeniyi, A. F., Aguilar-Farias, N., . . . Cardon, G. (2018). Global matrix 3.0 physical activity report card grades for children and youth: Results and analysis from 49 countries. *Journal of Physical Activity and Health, 15*(s2), S251-S273. <https://doi.org/10.1123/jpah.2018-0472>
- Chaput, J.-P., Colley, R. C., Aubert, S., Carson, V., Janssen, I., Roberts, K. C. & Tremblay, M. S. (2017). Proportion of preschool-aged children meeting the Canadian 24-Hour Movement Guidelines and associations with adiposity: Results from the Canadian Health Measures Survey. *BMC Public Health, 17*(5), 829. <https://doi.org/10.1186/s12889-017-4854-y>
- Chaput, J.-P., Katzmarzyk, P. T., LeBlanc, A. G., Tremblay, M. S., Barreira, T. V., Broyles, S. T., . . . Kurpad, A. (2015). Associations between sleep patterns and lifestyle behaviors in children: an international comparison. *International Journal of Obesity Supplements, 5*(2), S59-S65. <https://doi.org/10.1038/ijosup.2015.21>
- Chen, B., Bernard, J. Y., Padmapriya, N., Yao, J., Goh, C., Tan, K. H., . . . Godfrey, K. M. (2019). Socio-demographic and maternal predictors of adherence to 24-hour movement guidelines in Singaporean children. *International Journal of Behavioral Nutrition and Physical Activity, 16*(1), 70. <https://doi.org/10.1186/s12966-019-0834-1>
- Cliff, D. P., McNeill, J., Vella, S. A., Howard, S. J., Santos, R., Batterham, M., . . . de Rosnay, M. (2017). Adherence to 24-Hour Movement Guidelines for the Early Years and associations with social-cognitive development among Australian preschool children. *BMC Public Health, 17*(5), <https://doi.org/857.10.1186/s12966-019-0834-1>.
- De Craemer, M., McGregor, D., Androustos, O., Manios, Y. & Cardon, G. (2018). Compliance with 24-h movement behaviour guidelines among Belgian pre-school children: the ToyBox-study.

364 *International Journal of Environmental Research and Public Health*, 15(10), 2171.  
365 <https://doi.org/10.1186/s12889-017-4858-7>  
366

367 Diouf, A., Adom, T., Aouidet, A., El Hamdouchi, A., Joonas, N. I., Loechl, C. U., . . . Monyeki,  
368 A. (2018). Body mass index vs deuterium dilution method for establishing childhood obesity  
369 prevalence, Ghana, Kenya, Mauritius, Morocco, Namibia, Senegal, Tunisia and the United  
370 Republic of Tanzania. *Bulletin of the World Health Organization*, 96(11), 772.  
371 <https://doi.org/10.2471/BLT.17.205948>  
372

373 Draper, C. E., Tomaz, S. A., Jones, R. A., Hinkley, T., Twine, R., Kahn, K. & Norris, S. A. (2019).  
374 Cross-sectional associations of physical activity and gross motor proficiency with adiposity in  
375 South African children of pre-school age. *Public Health Nutrition*, 22(4), 614-623.  
376 <https://doi.org/10.1017/S1368980018003579>.  
377

378 Draper, C. E., Tomaz, S. A., Stone, M., Hinkley, T., Jones, R. A., Louw, J., . . . Norris, S. A. (2017).  
379 Developing intervention strategies to optimise body composition in early childhood in South  
380 Africa. *BioMed Research International*, 2017. doi: 10.1155/2017/5283457.  
381

382 United Nations Children's Fund. (2019). Children in Africa: Key statistics on child survival and  
383 population. Retrieved from [https://data.unicef.org/resources/children-in-africa-child-survival-](https://data.unicef.org/resources/children-in-africa-child-survival-brochure/)  
384 [brochure/](https://data.unicef.org/resources/children-in-africa-child-survival-brochure/)  
385

386 Guthold, R., Stevens, G. A., Riley, L. M. & Bull, F. C. (2018). Worldwide trends in insufficient  
387 physical activity from 2001 to 2016: A pooled analysis of 358 population-based surveys with 1·9  
388 million participants. *The Lancet Global Health*, 6(10), e1077-e1086.  
389 [https://doi.org/10.1016/S2214-109X\(18\)30357-7](https://doi.org/10.1016/S2214-109X(18)30357-7)  
390

391 Lee, E.-Y., Hesketh, K. D., Hunter, S., Kuzik, N., Rhodes, R. E., Rinaldi, C. M., . . . Carson, V.  
392 (2017). Meeting new Canadian 24-hour movement guidelines for the early years and associations  
393 with adiposity among toddlers living in Edmonton, Canada. *BMC Public Health*, 17(5), 840.  
394 <https://doi.org/10.1186/s12889-017-4855-x>.  
395

396 Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for  
397 systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine*,  
398 151(4), 264-269. <https://doi.org/10.7326/0003-4819-151-4-200908180-00135>  
399

400 Muthuri, S. K., Wachira, L.-J. M., Leblanc, A. G., Francis, C. E., Sampson, M., Onywera, V. O.  
401 & Tremblay, M. S. (2014). Temporal trends and correlates of physical activity, sedentary behavior,  
402 and physical fitness among school-aged children in Sub-Saharan Africa: A systematic review.  
403 *International Journal of Environmental Research and Public Health*, 11(3), 3327-3359.  
404 <https://doi.org/10.1136/bmj.b2535>  
405

406 Netsi, E., Santos, I. S., Stein, A., Barros, F. C., Barros, A. J. & Matijasevich, A. (2017). A different  
407 rhythm of life: sleep patterns in the first 4 years of life and associated sociodemographic  
408 characteristics in a large Brazilian birth cohort. *Sleep Medicine*, 37, 77-87.  
409 <https://doi.org/10.1016/j.sleep.2017.06.001>  
410

411 Okyere, S. (2018). *Prevalence of Overweight and Obesity in Pre-School Children at La*  
412 *Nkwantanang-Madina Municipality*. University of Ghana, [http://ugspace.ug.edu.gh/](http://ugspace.ug.edu.gh/handle/123456789/25907)  
413 [handle/123456789/25907](http://ugspace.ug.edu.gh/handle/123456789/25907)  
414

415 Reilly, J. (2015). The pandemic of low physical activity in children and adolescents. *Aspetar Sports*  
416 *Medicine Journal*, 4(May), 234-238. [https://www.aspetar.com/](https://www.aspetar.com/journal/viewarticle.aspx?id=236#.XsulhGgzbiU) journal/viewarticle  
417 [.aspx?id=236#.XsulhGgzbiU](https://www.aspetar.com/journal/viewarticle.aspx?id=236#.XsulhGgzbiU)

418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446

Santos, R., Zhang, Z., Pereira, J. R., Sousa-Sá, E., Cliff, D. P. & Okely, A. D. (2017). Compliance with the Australian 24-hour movement guidelines for the early years: Associations with weight status. *BMC Public Health*, 17(5), 867. <https://doi.org/10.1186/s12889-017-4857-8>.

Sunrise Study. (2020). <https://sunrise-study.com/>

The Joanna Briggs Institute (2009). The Joanna Briggs Institute Critical Appraisal tools for use in JBI Systematic Reviews: Checklist for Prevalence Studies. Retrieved from [https://joannabriggs.org/sites/default/files/2019-05/JBI\\_Critical\\_Appraisal-Checklist\\_for\\_Prevalence\\_Studies2017\\_0.pdf](https://joannabriggs.org/sites/default/files/2019-05/JBI_Critical_Appraisal-Checklist_for_Prevalence_Studies2017_0.pdf) <https://open.uct.ac.za/handle/11427/29464>

Tomaz, S.A. (2018). *Physical activity and gross motor skills in rural South African preschool children*. University of Cape Town, <https://doi.org/10.1186/s12889-019-6515-9>

World Health Organisation (2019). Guideline on physical activity, sedentary behavior and sleep for children under 5 years of age. Retrieved from <https://apps.who.int/iris/handle/10665/311664>

World Obesity Federation (2019). Atlas of Childhood Obesity. Retrieved from <https://www.worldobesity.org/nlsegmentation/global-atlas-on-childhood-obesity>

Zhu, Z., Tang, Y., Zhuang, J., Liu, Y., Wu, X., Cai, Y., . . . Chen, P. (2019). Physical activity, screen viewing time, and overweight/obesity among Chinese children and adolescents: An update from the 2017 physical activity and fitness in China—the youth study. *BMC Public Health*, 19(1), 197. <https://doi.org/10.1186/s12889-019-6515-9>

447 **Appendix 1:** An example of a search strategy from Web of Science and it was adapted for the  
448 other two databases.

449

450 1 \*Child, Pre-schooler/

451 2 \*Sleep/

452 3 \*Bed Rest/

453 4 2 or 3

454 5 Physical activity.mp.

455 6 Active play.mp.

456 7 \*Physical Exertion/

457 8 \*running/ or \*swimming/ or \*walking/

458 9 5 or 6 or 7 or 8

459 10 \*Sedentary Behavior/

460 11 sedentary lifestyle.mp.

461 12 "Physical Inactivity".mp.

462 13 \*Restraint, Physical/

463 14 10 or 11 or 12 or 13

464 15 screen time/

465 16 \*Video Games/

466 17 \*television/ or \*videodisc recording/ or \*videotape recording/

467 18 computer.mp.

468 19 "Play Station".mp.

469 20 DVD\*.mp.

470 21 "electronic gam\*".mp.

471 22 \*Smartphone/

472 23 Ipad\*.mp.

473 24 Laptop\*.mp.

474 25 15 or 16 or 17 or 18 or 19 or 20 or 21 or 23 or 24

475 26 Afric\*.mp.

476 27 Sub-sahara.mp.

477 28 26 or 27

478 29 4 or 9 or 14 or 25

479 30 1 and 28 and 29

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

## Appendix 2: Data extraction table

Author/ year	Study Design	Sample (N)	Sample Age (mean);	Behaviour	Exposure Description	Exposure Measurement Type	Measurement Description	Units	Baseline Mean	Baseline SD	NOTES
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	LPA	Actigraph GT3X+ accelerometers	Actigraph 25-420 counts/15s, 20 minutes consecutive zeros, 24h/day, 7 days, 15s epochs	Min/day	337.1	46.8	Emailed author for data 3-4-year-olds
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	MPA	Actigraph GT3X+ accelerometers	Not reported	Min/day	83.3	24.4	Emailed author for data 3-4 year olds
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	VPA	Actigraph GT3X+ accelerometers	Not reported	Min/day	25.9	14	Emailed author for data 3-4 year olds
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	MVPA	Actigraph GT3X+ accelerometers	Actigraph > 420 counts/15s, 20 minutes consecutive zeros, 24h/day, 7 days, 15s epochs	Min/day	109.3	36.5	Emailed author for data 3-4 year olds
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	LMVPA	Actigraph GT3X+ accelerometers	Actigraph > 25 counts/15s, 20 minutes consecutive zeros, 24h/day, 7 days, 15s epochs	Min/day	446.4	67.6	Emailed author for data 3-4 year olds
Draper et al. 2017	Cross-section	N=87	Age 3-4 years; mean age 4.5 y	Physical activity	LMVPA	Actigraph GT3X+ accelerometers	Actigraph > 25 counts/15s, 20 minutes consecutive zeros, 24h/day, 7 days, 15s epochs	% meeting 180min/day guidelines	100% weekly average; 98.2% daily		Emailed author for data 3-4 year olds
Senyo 2018	Cross-section	N=201	Range 2-5 years; mean age 3.9 y	Physical activity	MPA	WHO Global Physical Activity Questionnaire	Categorical at least 30min MVPA per day	%	Not at all: 7.5%; 1-2 times 38.3%; 3-5		MPH thesis

Senyo 2018	Cross-section	N=201	Range 2-5 years; mean age 3.9 y	Physical activity	VPA	WHO Global Physical Activity Questionnaire	Categorical at least 60 min VPA per day	%	times 17.4%; all week 36.8% Not at all: 11.9%; 1-2 times 39.8%; 3-5 times 24.9%; all week 23.4%	MPH thesis	
Tomaz 2018	Cross-section	N=56	Age 3-4 years	Screen time	Screen time	Parent questionnaire	Not validated questionnaire	hr/day	2	1.3	PhD thesis
Draper et al. 2019	Cross-section	Same sample as Draper et al., 2017									

501  
502  
503  
504  
505

