

## **Predictors of gallstone formation after bariatric surgery: a multivariate analysis of risk factors comparing gastric bypass, gastric banding, and sleeve gastrectomy**

**Vicky Ka Ming Li · Nestor Pulido ·  
Patricio Fajnwaks · Samuel Szomstein ·  
Raul Rosenthal · Pedro Martinez-Duarte**

Published online: 3 April 2009  
© Springer Science+Business Media, LLC 2009

**Erratum to: Surg Endosc**  
**DOI 10.1007/s00464-008-0204-6**

Pedro Martinez-Duarte was inadvertently omitted from  
the list of authors.

---

The online version of the original article can be found under  
doi:[10.1007/s00464-008-0204-6](https://doi.org/10.1007/s00464-008-0204-6).

---

V. K. M. Li · N. Pulido · P. Fajnwaks · S. Szomstein ·  
R. Rosenthal (✉) · P. Martinez-Duarte  
Section of Minimally Invasive and Endoscopic Surgery,  
The Bariatric and Metabolic Institute at Cleveland Clinic  
Florida, Cleveland Clinic Florida, 2950 Cleveland Clinic Blvd,  
Weston, FL 33331, USA  
e-mail: [rosentr@ccf.org](mailto:rosentr@ccf.org)