

# Northumbria Research Link

Citation: Faithfull, Sara, Turner, Lauren, Poole, Karen, Joy, Mark, Manders, Ralph, Weprin, Jennifer, Winters-Stone, Kerri and Saxton, John (2019) Prehabilitation for adults diagnosed with cancer: A systematic review of long-term physical function, nutrition and patient-reported outcomes. *European Journal of Cancer Care*, 28 (4). e13023. ISSN 0961-5423

Published by: Wiley-Blackwell

URL: <https://doi.org/10.1111/ecc.13023> <<https://doi.org/10.1111/ecc.13023>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/37845/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Prehabilitation for adults diagnosed with cancer: a  
systematic review of long-term physical function, nutrition  
and psychological outcomes**

Journal:	<i>European Journal of Cancer Care</i>
Manuscript ID	ECC-2017-0523.R1
Manuscript Type:	Feature and Review Paper
Keywords:	Rehabilitation, Exercise, Nutrition, Oncological Outcome, Prehabilitation, Psychological

SCHOLARONE™  
Manuscripts

Review

1  
2  
3 **Title: Prehabilitation for adults diagnosed with cancer: a systematic review of long-term physical**  
4 **function, nutrition and patient reported outcomes**  
5  
6  
7  
8  
9

10 **Abstract:**

11  
12 Prehabilitation is increasingly being used as an intervention to mitigate treatment-related  
13 complications and enhance recovery. An individual's state of health at diagnosis including; obesity,  
14 physical fitness and comorbidities, are influencing factors for the occurrence of adverse effects. This  
15 review explores whether prehabilitation works in improving health outcomes at or beyond **the initial**  
16 30-days post treatment and considers the utility of prehabilitation before cancer treatment. A database  
17 search was conducted for articles published with prehabilitation as a pre cancer treatment intervention  
18 between 2009-2017. Studies with no 30-day post treatment data were excluded. Outcomes post  
19 prehabilitation were extracted for physical function, nutrition and patient reported outcomes. Sixteen  
20 randomised controlled trials **with a combined 2017 participants** and six observational studies **with 289**  
21 **participants** were included. Prehabilitation interventions provided multi-modality components  
22 including exercise, nutrition and psycho educational aspects. Prehabilitation improved gait, cardio  
23 pulmonary function, urinary continence, lung function and mood 30-days post treatment but was not  
24 consistent across studies. When combined with rehabilitation, greater benefits were seen in 30 day  
25 gait and physical functioning compared to prehabilitation alone. Large scale randomized studies are  
26 required to translate what is already known from feasibility studies to improve overall health and  
27 increase long-term cancer patient outcomes.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 **Key words:** Prehabilitation, Rehabilitation, Cancer, Exercise, Nutrition, Survivorship  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Title: Prehabilitation for adults diagnosed with cancer: a systematic review of long-term physical**  
4 **function, nutrition and patient reported outcomes**  
5  
6  
7  
8  
9  
10

## 11 **Introduction**

12 Prehabilitation offers a route to improving patient's physical status and buffering treatment-related  
13 deconditioning between the time of cancer diagnosis and post treatment recovery. Prehabilitation  
14 includes physical and psychological assessments that establish baseline functioning and identify  
15 impairments that can impact on cancer treatment-related morbidity, as well as providing targeted  
16 interventions to maximize patient function prior to treatment onset (Silver and Baima, 2013). The  
17 primary goal of prehabilitation is *"to prevent or reduce the severity of anticipated treatment-related*  
18 *impairments that may cause significant disability (page2)* (Silver and Baima, 2013). There are several  
19 systematic reviews of prehabilitation for those receiving cancer surgery (Boereboom et al., 2016,  
20 Carli et al., 2017, Singh et al., 2013) and all suggest that prehabilitation enhances early discharge from  
21 hospital and reduces surgical adverse effects. **There is a growing requirement to include**  
22 **prehabilitation as part of the cancer pathway with three recent reports advising the value of**  
23 **prehabilitation, two in the USA (National Academies of Sciences, 2018, Stout et al., 2016) and one in**  
24 **the UK (Macmillan Cancer Support, 2018). However evidence that prehabilitation translates into**  
25 **better long-term patient outcomes beyond the initial 30-day post treatment complications is lacking.**  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 **Challenges to providing prehabilitation are that** cancer patients are highly likely to have comorbidities  
45 that complicate treatment delivery and reduce physical fitness (Sarfati et al., 2016, Stairmand et al.,  
46 2015). Comorbid conditions associated **with aging** and particularly excess body weight are common  
47 in patients presenting with cancer (Goodwin and Chlebowski, 2016) and evidence from  
48 epidemiological studies suggests that comorbidities and poorer health are correlates of poorer survival  
49 (Land et al., 2012, Land et al., 2012). There is compelling evidence for the link between obesity and  
50 cancer outcomes (Calle et al., 2003) with particular associations in the following tumour sites breast  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (Jiralerspong and Goodwin, 2016), gastrointestinal (Brown and Meyerhardt, 2016), endometrial  
4 (Onstad et al., 2016), prostate (Vidal et al., 2014) as well as haematological cancers, including  
5 multiple myeloma and leukaemia (Yang et al., 2016). Obesity is an important risk factor for  
6 cardiovascular, kidney disease, diabetes and some musculoskeletal disorders (Collaboration). These  
7 obesity-related comorbidities contribute to the adverse effects of cancer treatment (Bradley et al.,  
8 2014, Sogaard et al., 2013) and combined with an aging demographic, where more than 60% of  
9 cancer patients are over 65, comorbidity and poorer physical and functional health will impact upon  
10 future cancer treatment delivery and outcomes (Greenlee et al., 2016). **These co existing health  
11 problems are strong indicators for providing prehabilitation to maximise cancer treatment outcomes.**

12  
13  
14  
15  
16  
17  
18  
19  
20  
21 Rehabilitation interventions such as exercise, weight reduction and pharmacotherapy are **recognised**  
22 ways of managing comorbidity-related conditions after cancer treatment (Alamuddin et al., 2016) and  
23 there is evidence that **smoking cessation (Sitas et al., 2014) reduces adverse treatment effects and  
24 improves survival.** Preparing patients **prior to** cancer therapy by improving their overall health status  
25 as in prehabilitation could optimise their response to treatment and has important implications for  
26 future service delivery (Silver and Baima, 2013). Prehabilitation has been espoused as a key  
27 component of early recovery in cancer patients and is a term that has been traditionally used to  
28 describe interventions for optimising cardiopulmonary reserve prior to cancer surgery, with the aim of  
29 improving post-operative recovery outcomes (Carli et al., 2017, Silver, 2015, Silver and Baima,  
30 2013). However, prehabilitation programs are also targeting this pre-treatment period to improve  
31 chemotherapy adherence (Le Roy et al., 2016), reduce anxiety (Tsimopoulou et al., 2015), and to  
32 provide a stronger platform for post-treatment rehabilitation aimed at reversing treatment-related side  
33 effects and symptoms, managing comorbidities and enhancing longer-term health-related quality of  
34 life (Alfano et al., 2012, Boereboom et al., 2015, Shun, 2016, Silver, 2014). While fewer studies have  
35 been undertaken outside of the surgical context, a growing number of studies are focusing on different  
36 cancer treatments and modes of prehabilitation using exercise, psychological support and nutritional  
37 interventions. **These studies need to look at longer-term outcomes beyond the traditional enhanced  
38 recovery 30-day post treatment outcomes, to understand treatment adherence, mortality, disease**

1  
2  
3 **prognosis or impact on health economics (Stout et al., 2018).** This is the first systematic review to  
4  
5 critically review the impact of different prehabilitation interventions on long-term health outcomes (at  
6  
7 or beyond 30-days post-treatment completion) in cancer patients and explore the utility of  
8  
9 prehabilitation as a platform for risk management before and after all cancer treatments.

10  
11 **This review addressed two questions:**

- 12  
13  
14 1. **What is the effect of prehabilitation on  $\geq 30$  day post treatment outcomes including; physical**  
15  
16 **functioning, nutrition and patient-reported outcomes ?**
- 17  
18 2. **How can prehabilitation be used to optimise the management of cancer patients with**  
19  
20 **comorbidity or pre-existing risk factors that are associated with poorer cancer treatment**  
21  
22 **outcomes?**

## 23 24 25 26 27 28 29 **Methods**

### 30 31 *Data Sources and search method*

32  
33  
34  
35 The review was registered on PROSPERO (CRD42016050296) international prospective database of  
36  
37 systematic reviews. The search was conducted in two stages. In stage one, studies were identified via  
38  
39 abstracts through a systematic search strategy for Medline (Pub med), CINAHL (with full text)  
40  
41 Embase and Cochrane central register of controlled trials. The databases were chosen to identify  
42  
43 potentially relevant published studies in the field of medicine, exercise, health and psychosocial care.  
44  
45 Search terms were split into two categories “prehabilitation combined with cancer” and terms to  
46  
47 identify the nature of prehabilitation such as “exercise, nutrition, psychology and other **behavioural**  
48  
49 interventions”. The full search strategy and MESH terms are provided in supplementary materials. In  
50  
51 stage two, other relevant publications were retrieved by reviewing the reference lists of these studies  
52  
53 against the eligibility criteria.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Studies selected were published from the period 2000 to February 2017. The following were all  
4 excluded from the review; prehabilitation studies with no reported post-treatment outcomes at 30-days  
5 or longer; studies that combined data from previously published studies; abstracts, case studies,  
6 conference abstracts and those not in English. Participants included were cancer patients who were  
7 treated with any treatment modality and received any form of prehabilitation either in the home or  
8 hospital setting. Prehabilitation was defined as a single or multi-modality intervention that could  
9 include exercise, nutritional support, patient education and/or psychological therapy. Control was  
10 defined as those participant's receiving usual care as defined in the clinical pathway. Identification of  
11 objective clinical, patient reported and delivery outcomes were described at 30-days post treatment  
12 completion. Comorbidity data at **baseline and at completion** were also reviewed. Efficacy in relation  
13 to 30 day post treatment objective physical functioning was explored through meta-analysis but data  
14 was not of sufficient quality to make a comparison. The quality of eligible studies was assessed using  
15 the PRISMA critical appraisal methods (Shamseer L et al., 2015). Risk of bias was assessed by an  
16 interdisciplinary research team using the Cochrane Collaboration's tool for assessing risk of bias  
17 ROBINS-I tool (Sterne et al., 2016). Observational or quasi-experimental studies were included as  
18 they provided additional information as to the use of prehabilitation interventions.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

## 38 **Results**

39  
40 Sixteen randomised controlled trials (RCT) and six observational studies were included in the  
41 narrative synthesis (Figure 1). The quality of the RCT studies varied considerably with 7 of the 16  
42 studies being considered as having a high risk of bias. Studies were not sufficiently consistent in  
43 intervention or outcome data to be included in a meta-analysis. In many studies reporting of the  
44 randomization processes, lack of allocation concealment to those enrolling, blinding of outcome  
45 assessors and poor reporting of missing data may have impacted on study quality (Table 1). Most  
46 studies were single centre studies. However, one of two multi-site studies was a 3-arm trial comparing  
47 psychological prehabilitation strategies, with participants randomized to stress management, a support  
48 group or usual care (Parker et al., 2009). Other studies compared different prehabilitation components  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 head to head as nutritional interventions or psychological approaches. **The number of participants**  
4 **within the RCTs ranged from 48 to 652, with a median of 88 with a total number of subjects in the**  
5 **review of 2017** (Table 1). Most individual RCTs analysed fewer than 60% of the sample originally  
6  
7 recruited in the study, excluding participants due to comorbidity or inability to undergo cardio  
8  
9 pulmonary exercise testing. Participants were adults with colorectal (Carli et al., 2010, Cheville et al.,  
10  
11 2015, Gillis et al., 2014, Gillis et al., 2016, Moriya, 2015) lung (Barlési et al., 2008, Stefanelli et al.,  
12  
13 2013), head and neck (Van Bokhorst-de Van der Schuer et al., 2000), breast (Garssen et al., 2013),  
14  
15 bladder (Jensen et al., 2015, Jensen et al., 2014) and prostate (Bales et al., 2000, Burgio et al., 2006,  
16  
17 Parker et al., 2009) cancer or included individuals with a range of cancers (Schmidt et al., 2015). Trial  
18  
19 designs were primarily feasibility studies and therefore the studies were rarely powered to determine  
20  
21 the efficacy of prehabilitation on post-treatment recovery outcomes. The primary endpoint was  
22  
23 predominantly objective physical function prior to treatment with the secondary endpoints described  
24  
25 at 1 to 6 months post intervention. Only four (25%) of the authors fully reported participant  
26  
27 comorbidities at baseline (Burgio et al., 2006, Jensen et al., 2015, Schmidt et al., 2015, Van Bokhorst-  
28  
29 de Van der Schuer et al., 2000) whilst two actively excluded participants with comorbidities possibly  
30  
31 due to the intensity of the exercise programme (Carli et al., 2010, Stefanelli et al., 2013).  
32  
33

34  
35 The designs of the 6 observational studies were either case controlled cohort, historical controls or  
36  
37 quasi experimental. Studies were primarily feasibility studies and participant numbers were small,  
38  
39 **ranging from 35 to 87 with a total of 289 participants.** Studies included individuals with breast cancer  
40  
41 (Baima et al., 2015), lung cancer (Jones et al., 2007, Peddle et al., 2009, Sekine et al., 2005),  
42  
43 colorectal cancer (Li et al., 2013) and prostate cancer (Sueppel et al., 2001).  
44

45  
46 Comorbidities were only reported in three of the studies **at baseline**, with ill health being cited as a  
47  
48 contributing factor to difficulties with recruitment **rather than this being recorded as an outcome.**  
49  
50 Several studies did not report attrition (Sekine et al., 2005, Sueppel et al., 2001), and among those that  
51  
52 did attrition rates ranged from 0 to 52%. The number and combination of prehabilitation modalities  
53  
54 varied considerably across studies, ranging from 1 to 3 across individual RCTs and observational  
55  
56 studies (Table 2 & 3).  
57  
58  
59  
60



1  
2  
3  
4  
5 Most (16/22) studies included an *exercise modality*, either as a stand-alone prehabilitation intervention  
6 or in combination. Four studies examined the effects of pelvic floor training in men with prostate  
7 cancer over a varying number of weeks before radical prostatectomy (Bales et al., 2000, Burgio et al.,  
8 2006, Centemero et al., 2010, Sueppel et al., 2001). These were predominantly home-based exercise  
9 programmes with some level of instruction and supervision and/or biofeedback training. Two studies  
10 incorporated supervised therapeutic pulmonary exercises (in conjunction with more conventional  
11 conditioning exercise) in lung cancer patients in the 2-3 weeks prior to surgery (Sekine et al., 2005,  
12 Stefanelli et al., 2013). These exercises were performed on 5-7 days per week and included incentive  
13 spirometry, abdominal breathing, huffing and coughing, and respiratory exercises on a bench,  
14 mattress pad and wall bars. Finally, a study in breast cancer patients investigated the feasibility of  
15 therapeutic shoulder mobility exercises in the 2-4 weeks before surgery, comparing in-person teaching  
16 with video-only teaching (Baima et al., 2015). Both methods were shown to be feasible with high  
17 adherence ( $\geq 75\%$ ). Other studies investigated the effects of conventional forms of exercise  
18 conditioning for improving cardiopulmonary fitness and/or muscular strength over durations of 2-8  
19 weeks, though most programmes were of 2-4 weeks duration (Table 2 & 3). All but one of these  
20 studies implemented exercise prehabilitation in the time period before colorectal, lung or bladder  
21 cancer surgery, whereas the remaining study (Cheville et al., 2015) focused on adherence to chemo  
22 radiotherapy in patients with gastrointestinal cancers. Home-based programmes generally consisted of  
23 aerobic and resistance exercise on at least three days per week with varying degrees of face to face  
24 supervision and telephone support (Carli et al., 2010, Gillis et al., 2014, Jensen et al., 2015, Jensen et  
25 al., 2014, Li et al., 2013). Instructions on both the frequency and intensity of aerobic exercise were  
26 generally provided and in some cases participants used heart rate monitors and perceived exertion  
27 scales to self-assess their level of effort (Gillis et al., 2014, Li et al., 2013). Studies of more closely  
28 supervised 2-6 week programs of exercise prehabilitation involved vigorous intensity cycle ergometry  
29 in lung (Jones et al., 2007, Peddle et al., 2009) and rectal cancer patients (West et al., 2015) prior to  
30 surgery and isokinetic muscle strengthening exercises in patients with gastrointestinal cancers during  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 chemo radiotherapy (Cheville et al., 2015). Two further studies included vigorous gym-based aerobic  
4 exercise (Stefanelli et al., 2013) or walking exercise (5000 steps/day)(Sekine et al., 2005) in  
5  
6 combination with therapeutic pulmonary exercises in patients with lung cancer prior to surgery. The  
7  
8 short timelines prior to therapy made a progressive programme difficult to achieve. Although  
9  
10 adherence to the home exercise program was reported in most of these studies, adherence to exercise  
11  
12 at the prescribed intensity and progression of the exercise programme were poorly reported.  
13  
14  
15  
16  
17

18 Only five of the studies provided a *nutritional modality* as part of the prehabilitation package. Some  
19  
20 of the interventions were purely nutrition based (Gillis et al., 2016, Moriya, 2015, Van Bokhorst-de  
21  
22 Van der Schuer et al., 2000), however, two of the studies used nutrition as part of multi-component  
23  
24 prehabilitation intervention (Gillis et al., 2016, Li et al., 2013). The nutritional interventions were  
25  
26 varied with 5-10 days preoperative feeding plus a supplemental arginine formula (Van Bokhorst-de  
27  
28 Van der Schuer et al., 2000) or whey protein (Gillis et al., 2014, Li et al., 2013) or a low or high dose  
29  
30 immune-enhancing diet (Moriya, 2015). Multi-modal prehabilitation interventions provided 90-min of  
31  
32 nutritional counselling with daily whey protein supplementation (Gillis et al., 2016) in comparison to  
33  
34 a control group which received nutritional counselling without supplementation. The timing of  
35  
36 nutritional interventions varied between 5-10 days (Gillis et al., 2016, Moriya, 2015) and 3-6 weeks  
37  
38 pre-operatively (Gillis et al., 2014, Li et al., 2013). The nutritional intervention did not continue  
39  
40 beyond surgery, with one exception (Gillis et al., 2016) which continued the nutritional intervention 4  
41  
42 weeks' post-surgery. Nutritional therapies were primarily targeted on individuals with cancer who  
43  
44 were malnourished, receiving treatment for head and neck (Van Bokhorst-de Van der Schuer et al.,  
45  
46 2000) or colorectal cancer (Gillis et al., 2014, Gillis et al., 2016, Li et al., 2013, Moriya, 2015). Van  
47  
48 Bokhorst (Van Bokhorst-de Van der Schuer et al., 2000) excluded adults from the study if they were  
49  
50 well nourished (10% excluded), whereas Gillis (Gillis et al., 2016) screened for malnutrition using the  
51  
52 Patient Generated – Subjective Global Assessment (PG-SGA) which is a validated tool for nutritional  
53  
54 assessment in oncology. Adherence to nutritional intervention is reported in only one study with  
55  
56 researchers contacting participants on a weekly basis to encourage them to record their whey protein  
57  
58  
59  
60

1  
2  
3 ingestion. This study noted that adherence was higher in the prehabilitation group compared to the  
4  
5 rehabilitation group both pre and post-surgery.  
6  
7  
8  
9

10 Studies involving a *psychoeducation modality* as part of prehabilitation programmes have focused  
11 primarily on anxiety and stress reduction (Cheville et al., 2015, Garssen et al., 2013, Parker et al.,  
12 2009, Schmidt et al., 2015), patient education and lifestyle advice (Baima et al., 2015, Barlési et al.,  
13 2008) (Jensen et al., 2015) and/or counselling (Parker et al., 2009) as part of the intervention;  
14 however, few studies report any detail of the therapeutic components of the intervention. Psycho  
15 educational prehabilitation strategies have been studied as single mode counselling interventions  
16 (Barlési et al., 2008, Cheville et al., 2015) or by comparing a variety of psychological and educational  
17 approaches prior to cancer treatment (Parker et al., 2009). Psycho educational strategies have also  
18 been studied as part of multi component prehabilitation programmes (Gillis et al., 2014, Jensen et al.,  
19 2015, Jensen et al., 2014). Psycho educational interventions prior to surgery for lung and  
20 gastrointestinal cancer provided written and verbal information to participants which described the  
21 disease and associated surgery outcomes (Barlési et al., 2008, Schmidt et al., 2015). An alternative  
22 psycho therapeutic approach involved weekly group sessions with a psychiatrist, focused on  
23 individuals' social, cognitive and emotional care in conjunction with relaxation exercises 30-days  
24 prior to chemotherapy (Cheville et al., 2015). Similarly, Garsen (Garssen et al., 2013) provided 4  
25 sessions over 5 days to women with breast cancer, including stress management, relaxation, guided  
26 imagery techniques and counselling. Parker (Parker et al., 2009) investigated the effects of a similar  
27 programme on post-operative recovery outcomes in men with prostate cancer. These interventions  
28 were all compared to usual or supportive care. Adherence to the intervention was not always reported  
29 in the observational studies with attrition 25-52% respectively (Baima et al., 2015) (Jones et al.,  
30 2007).  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### *Objective clinical outcomes following prehabilitation*

Studies that included an exercise modality investigated the effects of prehabilitation regimens on cancer treatment recovery outcomes and cardiopulmonary fitness (table 4). Three studies reported favourable effects of home-based pelvic floor training on post-operative urinary continence outcomes in prostate cancer patients undergoing radical prostatectomy up to 12 months of follow-up (Burgio et al., 2006, Centemero et al., 2010, Sueppel et al., 2001) and a fourth study (Bales et al., 2000) showed no urinary continence benefits of including biofeedback training. Similarly, a study of female breast cancer patients reported no additional post-operative benefits when home-based shoulder exercise prehabilitation included an in-person teaching session versus video-based instruction (Baima et al., 2015). Supervised exercise prehabilitation programmes in lung cancer patients have generally been more intensive than home-based programmes and have resulted in improvements in pre-operative cardiopulmonary fitness measures, including six minute walk test (6MWT) (Jones et al., 2007) and peak  $VO_2$  (Jones et al., 2007, Stefanelli et al., 2013). However, the improvements in peak  $VO_2$  were modest ( $2-3 \text{ ml kg min}^{-1}$ ) and it is unclear whether improvements of this magnitude translate to improved post-operative recovery outcomes or longer-term outcomes, such as quality of life. It is of interest to note that intensive cycle ergometry prehabilitation had no impact on quality of life pre-surgery or at 2-months post surgery (Peddle et al., 2009). Nevertheless, Sekine (Sekine et al., 2005) reported a reduction in post-operative pulmonary complications and hospital length of stay in lung cancer patients after a prehabilitation programme that involved daily pulmonary therapeutic exercises and walking (5000 steps/day) in the two weeks prior to lobectomy when compared to historical controls. In other studies, prehabilitation programs involving exercise have yielded equivocal results. A supervised programme involving cycling + strengthening exercises in patients with gastrointestinal cancer compared to those in a walking + breathing exercise group showed no differences in 6MWT distance (Carli et al., 2010). Similarly, home-based exercise programs involving aerobic and/or resistance exercise 4 weeks prior to surgery have had minimal impact on post-operative hospital length of stay or severity of complication (Gillis et al., 2014, Jensen et al., 2015, Jensen et al., 2014), although Jensen (Jensen et al., 2015) reported improved post-operative 6MWT distance in bladder

1  
2  
3 cancer patients receiving prehabilitation, 4806m (95%CI 4075m-5536m) compared to 2906m (95%CI  
4 2408-3404m) in those receiving usual care. Gillis (Gillis et al., 2014) reported higher submaximal  
5 cardiopulmonary fitness + 23.4m (6MWT) in a prehabilitation/ rehabilitation group compared to  
6 rehabilitation alone -21.8m (80.7) at 8 weeks after colorectal cancer surgery.  
7  
8  
9

10  
11  
12  
13  
14 Studies that included nutritional outcomes were few and reported no significant differences between  
15 the intervention and control groups at  $\geq 30$  days (Moriya, 2015) on post treatment physical functioning  
16 (table 4). However, preoperative nutritional modality groups showed a significant improvement in  
17 physical functioning and initial symptoms post operatively (Gillis et al., 2014, Li et al., 2013, Van  
18 Bokhorst-de Van der Schuer et al., 2000). Only one study measured upper-body strength (Gillis et al.,  
19 2016) and this improved pre-surgery but was not sustained post-surgery. Participants who received  
20 arginine supplementation with feeding pre-and post-surgery showed reduced appetite at 6 months  
21 (Van Bokhorst-de Van der Schuer et al., 2000) and serum albumin remained stable in a small (n=17)  
22 pre-post intervention study (Li et al., 2013). However, prehabilitation studies nutritional outcomes are  
23 compromised by the lack of consistency in measuring nutritional intake and adherence (mainly  
24 through self-report tools) or objective sarcopenia measures. Such limitations could have important  
25 implications for assessing treatment fidelity and the sensitivity of outcome measures.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

#### 41 *Patient reported outcomes (PRO) of prehabilitation*

42  
43 PRO in the studies reviewed included health related quality of life using the Short Form Health  
44 Survey (SF36) and Prostate Cancer Index (PCI), which incorporate physical and emotional subscales.  
45 Symptom specific measures such as the International Continence Scale for men (ICS male), the  
46 Hospital Anxiety and Depression Scale (HADS) and (PCI) and Activities of Daily Living (ADL) tool  
47 were also reported in some studies (Table 4). Quality of life scores were comparable between  
48 prehabilitation and control groups at 3 months post-intervention in most studies (Barlési et al., 2008,  
49 Burgio et al., 2006, Garssen et al., 2013, Peddle et al., 2009). However, in two studies, self-reported  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 physical function was higher in the prehabilitation group at 1 year (Li et al., 2013, Parker et al., 2009)  
4 and in the study by Li (Li et al., 2013), an increase in self-reported physical activity persisted 8 weeks  
5 after surgery. Post-treatment improvements in mood, anxiety and depression have been reported  
6 immediately post-operatively following prehabilitation involving walking + breathing exercises and  
7 psychological support (Carli et al., 2010, Parker et al., 2009, Schmidt et al., 2015) but effects were  
8 small and between group differences were not sustained long-term (Parker et al., 2009, Schmidt et al.,  
9 2015). **Behavioural change techniques, such as smoking cessation, were rarely reported in studies, this  
10 can impact on radiotherapy side-effects and subsequent post-treatment cancer outcomes (Warren et  
11 al., 2014).**

#### *Patient reported and service outcomes for prehabilitation*

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Complication rates and length of hospital stay post-surgery were the most frequent service delivery  
measures reported for  $\geq 30$ -days post treatment (Table 4). There was no difference in length of stay,  
between prehabilitation and control groups in five studies (Gillis et al., 2014, Gillis et al., 2016,  
Jensen et al., 2014, Li et al., 2013, Schmidt et al., 2015), with the exception of Sekines (Sekine et al.,  
2005), where the intervention group had a reduced length of stay after a 4-6 week prehabilitation  
program. Post-operative complications such as wound healing, seroma formation and bleeding were  
shown to be comparable between intervention and control groups but Moriya (Moriya, 2015) found  
that those receiving a prehabilitation nutritional intervention had fewer post-operative site infections .  
Prehabilitation has been shown to improve initial post-operative mobilisation (Jensen et al., 2015) and  
the number of patients completing chemotherapy (Cheville et al., 2015). Furthermore, in the latter  
study, those receiving the intervention had significantly fewer treatment hospitalisations.

#### **Discussion**

Overall this systematic review suggests prehabilitation impacts on select 30-day outcome measures  
for some people with cancer but few studies have measured or reported overall long-term health

1  
2  
3 benefits. The results of the review are summarised pictorially as a diagram describing the multi-  
4  
5 modality intervention and linked physical function, nutrition and patient reported outcomes used in  
6  
7 the reviewed studies (Figure 2). Many of these studies report service or process data measures such as  
8  
9 length of stay and post-operative complications, but do not consistently capture changes in physical  
10  
11 functioning or patient reported outcomes. The only exception is pre-operative therapeutic pelvic floor  
12  
13 exercises for men undergoing prostatectomy for prostate cancer where prehabilitation improved long-  
14  
15 term urinary continence. This reflects the differentiation between general prehabilitation versus  
16  
17 targeted exercise or nutrition interventions and the greater specificity of their effect. There is  
18  
19 insufficient evidence for demonstration of long-term benefits in other cancer patient populations  
20  
21 beyond the initial 30 day post treatment complications. Even vigorous intensity pre-operative aerobic  
22  
23 exercise conditioning programmes have only resulted in modest improvements in peak oxygen uptake  
24  
25 pre-operatively (of the order of 2-3 ml/kg/min<sup>-1</sup>), possibly a factor of the short duration of programs,  
26  
27 and these gains are lost post-operatively. Not surprisingly then, prehabilitation combined with  
28  
29 rehabilitation was the most effective approach in improving outcomes longer than 30-days.  
30  
31  
32

33  
34 It is now recognised that a physically active lifestyle is inversely related to the risk of certain cancers  
35  
36 and mortality (Brown JC et al., 2012, Schmid and Leitzmann, 2014). Surprisingly few prehabilitation  
37  
38 studies measured or reported participant comorbidities and how they changed over time. Therefore we  
39  
40 were unable to address our second question, how prehabilitation can optimise the management of  
41  
42 cancer patients with comorbidity? Comorbidities in participants in prehabilitation studies were  
43  
44 considered exclusion criteria rather than as predictors of physical functioning that could be mediated  
45  
46 by exercise or nutrition and that could change as a response to intervention (Brown JC et al., 2012).

47  
48 Those participants with high levels of comorbidities and poor fitness were often not eligible to be  
49  
50 included, which suggests those people most in need to improve physical function were less likely to  
51  
52 receive prehabilitation. The multi-modality approach of prehabilitation could optimise the  
53  
54 management of cancer patients with low baseline scores and who have been shown to gain greater  
55  
56 benefits (Minnella et al., 2016). Rather than trying to demonstrate the efficacy of multi-modality  
57  
58  
59  
60

1  
2  
3 prehabilitation on the fittest patients, we should consider using the approach to optimise the  
4 management of the more complex and least fit cancer patients who have most to gain. This requires  
5 more sophisticated tailoring of intervention to personalise and **target prehabilitation**. For example,  
6 current exercise guidelines for cancer survivors recommend muscle strengthening exercises for  
7 overall conditioning (Schmitz KH et al., 2010) but this may not be sufficient to manage specific  
8 deficits. A more task-specific approach incorporating functional movements using strength and  
9 mobility may be optimal for prehabilitation regimens (Winters-Stone et al., 2015).  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 In some studies, the high attrition of participants suggests a balance is required between intensity and  
20 duration of exercise to be able to meet the needs of those with greater limitations. This highlights the  
21 need for continuity and support in establishing exercise habits and expectations around exercise for  
22 people with cancer (Brown JC et al., 2012, Mayo et al., 2011). The content of exercise programmes is  
23 poorly described in some papers, and have not followed the FITT principle of reporting Frequency,  
24 Intensity, Timing and Type of exercise and/or how the exercise programme is personalised or  
25 progressed over time (Thompson et al., 2010). These oversights make it challenging to understand  
26 whether or not the exercise program was insufficiently designed and/or how to revise programs to  
27 optimize adherence and outcomes in the future.  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 The inclusion of nutritional support as part of prehabilitation improved short-term physical function.  
38 The pre-surgical interventions were necessarily short (2-3 weeks) primarily due to treatment target  
39 times. Indications from one study suggest that longer term patient outcomes could benefit with  
40 additional post-surgical rehabilitation. Given the rising proportion of cancer patients who are obese at  
41 diagnosis, the prehabilitative and rehabilitative window is potentially an opportunity to embed new  
42 lifestyle behaviours. Malnutrition is associated with a poorer response to cancer treatment and  
43 hypoalbuminaemia is associated with post-surgical mortality, increased morbidity and length of stay  
44 (Hu W-H et al., 2015). Patients with colorectal cancer are more malnourished than other patients  
45 groups (28% colorectal compared to 4% prostate cancer) (Hu W-H et al., 2015) hence the wide  
46 number of prehabilitation studies in this population. In the nutrition components of prehabilitation  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 programs surrogate measures were used for the combined interventions rather than specific targets  
4 such as, serum values or anthropometric measures. If we are tackling obesity in cancer and its risks,  
5 then a greater focus on adiposity, fat distribution and sarcopenia should be included in prehabilitation  
6 studies. **With emerging therapies and earlier diagnosis techniques, for example of low dose**  
7 **computerised imaging in lung cancer (Smith et al., 2017), the opportunity for prehabilitation becomes**  
8 **more feasible as patients are less likely to be burdened by advanced disease or chronic illness.**  
9  
10  
11  
12  
13  
14  
15  
16

17 Understanding how prehabilitation components work together is a challenge as few studies used a  
18 theoretical or conceptual frameworks to guide design. Exploring how the multi-modality components  
19 work; such as exercise, nutrition, psychoeducational components is essential to maximize outcomes  
20 (Figure 2). The use of factorial research designs in future studies is recommended in evaluating  
21 prehabilitation components (Montgomery et al., 2003). Whilst pre-operative exercise programmes  
22 have incorporated both aerobic and resistance training, most emphasis has been on aerobic exercise.  
23 The effect of resistance exercise on pre-operative muscular function and how this impacts upon post-  
24 operative recovery outcomes has received less attention (Singh et al., 2013). The relationship between  
25 psychological health and exercise behaviour has been well established. Short and long term adherence  
26 may be optimized if anxiety and depression are also addressed during an exercise program, however  
27 the focus on anxiety and depression management at the expense of evidence based behaviour change  
28 strategies may not be the best strategy for long-term adherence (Stacey FG et al., 2015).  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 A limitation of this review is that conclusions have to be considered in the context of a limited  
42 number of studies, the majority of which are underpowered feasibility studies. The importance of  
43 feasibility studies is recognized in the Medical Research Council (MRC) complex interventions  
44 framework and that they should now be used to inform fully powered RCTs. The review highlighted  
45 the need for improved quality of studies for example following consort or strobe reporting guidance  
46 and this has also been described in previous prehabilitation systematic reviews (Singh et al., 2013).  
47 It's imperative that future studies take a more ambitious approach to test efficacy by building on the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 current evidence base using a conceptual framework to guide intervention design and robust  
4  
5 evaluation.

6  
7  
8  
9 Can prehabilitation programmes impact on longer-term cancer health outcomes? The answer is  
10  
11 currently unclear **especially in relation to changing comorbidity**. Prehabilitation is now an integral  
12  
13 part of many cancer surgical preparatory pathways as part of early recovery but there is scope for  
14  
15 greater targeting to include nutrition and psychoeducational components, as well as considering how  
16  
17 prehabilitative interventions may buffer symptoms such as fatigue and pain during adjuvant therapies.  
18  
19 Sophisticated research designs incorporating economic evaluation and longer-term measures are  
20  
21 essential to guide service development and support implementation if the concept of cancer  
22  
23 prehabilitation is to emulate cardiac rehabilitation services.

24  
25  
26  
27 In conclusion, prehabilitation strategies may have an important role to play in addressing the rising  
28  
29 complexity of health needs of those diagnosed with cancer. Forty-percent of all those diagnosed with  
30  
31 cancer have one co-morbid condition and 15% at least two concurrent health problems (Sarfati et al.,  
32  
33 2016). This systematic review highlights that single and multi-modal prehabilitation programmes are  
34  
35 feasible and some approaches confer short-term benefits in the post-surgical recovery period. The  
36  
37 next stage is to design robust efficacy studies to test carefully defined prehabilitative/prehabilitative-  
38  
39 rehabilitative interventions at the time of first cancer treatment (be that surgery, systemic anti-cancer  
40  
41 therapy or radiotherapy) and measure clinical outcome, PRO, patient benefit and service delivery  
42  
43 outcomes throughout the care pathway.

#### 44 45 46 47 **References**

- 48  
49  
50 Alamuddin, N., Bakizada, Z., Wadden, T.A., 2016. Management of Obesity. *Journal of Clinical*  
51 *Oncology* 34 (35), 4295-4305.  
52 Alfano, C.M., Ganz, P.A., Rowland, J.H., Hahn, E.E., 2012. Cancer Survivorship and Cancer  
53 *Rehabilitation: Revitalizing the Link. Journal of Clinical Oncology* 30 (9), 904-906.  
54 Baima, J., Reynolds, S.G., Edmiston, K., Larkin, A., Ward, B.M., O'Connor, A., 2015. Teaching of  
55 *Independent Exercises for Prehabilitation in Breast Cancer. J Cancer Educ.*  
56  
57  
58  
59  
60


- 1  
2  
3 Bales, G.T., Gerber, G.S., Minor, T.X., Mhoon, D.A., McFarland, J.M., Kim, H.L., Brendler, C.B.,  
4 2000. Effect of preoperative biofeedback/pelvic floor training on continence in men  
5 undergoing radical prostatectomy. *Urology* 56, 627-630.
- 6 Barlési, F., Barrau, K., Loundou, A., Doddoli, C., Simeoni, M.C., Auquier, P., Thomas, P., 2008.  
7 Impact of information on quality of life and satisfaction of non-small cell lung cancer  
8 patients: a randomized study of standardized versus individualized information before  
9 thoracic surgery. *J Thorac Oncol* 3 (10), 1146-1152.
- 10 Boereboom, C., Doleman, B., Lund, J., Williams, J., 2016. Systematic review of pre-operative  
11 exercise in colorectal cancer patients. *Tech Coloproctol* 20, 81-89.
- 12 Boereboom, C.L., Williams, J.P., Leighton, P., Lund, J.N., Exercise Prehabilitation in Colorectal  
13 Cancer Delphi Study, G., 2015. Forming a consensus opinion on exercise prehabilitation in  
14 elderly colorectal cancer patients: a Delphi study. *Tech Coloproctol* 19 (6), 347-354.
- 15 Bradley, C.J., Dahman B Fau - Anscher, M., Anscher, M., 2014. Prostate cancer treatment and  
16 survival: evidence for men with prevalent comorbid conditions. (1537-1948 (Electronic)).
- 17 Brown JC, Winters-Stone K, Lee A, KH., S., 2012. Cancer, Physical Activity, and Exercise. .  
18 *Comprehensive Physiology*. 2 (4), 2775-2809.
- 19 Brown, J.C., Meyerhardt, J.A., 2016. Obesity and Energy Balance in GI Cancer. *Journal of Clinical*  
20 *Oncology* 34 (35), 4217-4224.
- 21 Burgio, K.L., Goode, P.S., Urban, D.A., Umlauf, M.G., Locher, J.L., Bueschen, A., Redden, D.T.,  
22 2006. Preoperative biofeedback assisted behavioral training to decrease post-prostatectomy  
23 incontinence: a randomized, controlled trial. *J Urol* 175 (1), 196-201; discussion 201.
- 24 Calle , E.E., Rodriguez , C., Walker-Thurmond , K., Thun , M.J., 2003. Overweight, Obesity, and  
25 Mortality from Cancer in a Prospectively Studied Cohort of U.S. Adults. *New England*  
26 *Journal of Medicine* 348 (17), 1625-1638.
- 27 Carli, F., Charlebois, P., Stein, B., Feldman, L., Zavorsky, G., Kim, D.J., Scott, S., Mayo, N.E., 2010.  
28 Randomized clinical trial of prehabilitation in colorectal surgery. *Br J Surg* 97 (8), 1187-  
29 1197.
- 30 Carli, F., Silver, J.K., Feldman, L.S., McKee, A., Gilman, S., Gillis, C., Scheede-Bergdahl, C.,  
31 Gamsa, A., Stout, N., Hirsch, B., 2017. Surgical Prehabilitation in Patients with Cancer:  
32 State-of-the-Science and Recommendations for Future Research from a Panel of Subject  
33 Matter Experts. *Phys Med Rehabil Clin N Am* 28 (1), 49-64.
- 34 Centemero, A., Rigatti, L., Giraud, D., Lazzeri, M., Lughezzani, G., Zugna, D., Montorsi, F., Rigatti,  
35 P., Guazzoni, G., 2010. Preoperative pelvic floor muscle exercise for early continence after  
36 radical prostatectomy: a randomised controlled study. *European urology* 57 (6), 1039-1043.
- 37 Cheville, A.L., Alberts, S.R., Rummans, T.A., Basford, J.R., Lapid, M.I., Sloan, J.A., Satele, D.V.,  
38 Clark, M.M., 2015. Improving Adherence to Cancer Treatment by Addressing Quality of Life  
39 in Patients With Advanced Gastrointestinal Cancers. *J Pain Symptom Manage* 50 (3), 321-  
40 327.
- 41 Collaboration, N.C.D.R.F., 2016. Trends in adult body-mass index in 200 countries from 1975 to  
42 2014: a pooled analysis of 1698 population-based measurement studies with 19·2 million  
43 participants. *The Lancet* 387 (10026), 1377-1396.
- 44 Garssen, B., Boomsma, M.F., Meezenbroek, E.e.J., Porsild, T., Berkhof, J., Berbee, M., Visser, A.,  
45 Meijer, S., Beelen, R.H., 2013. Stress management training for breast cancer surgery patients.  
46 *Psychooncology* 22 (3), 572-580.
- 47 Gillis, C., Li, C., Lee, L., Awasthi, R., Augustin, B., Gamsa, A., Liberman, A.S., Stein, B.,  
48 Charlebois, P., Feldman, L.S., Carli, F., 2014. Prehabilitation versus rehabilitation: a  
49 randomized control trial in patients undergoing colorectal resection for cancer.  
50 *Anesthesiology* 121 (5), 937-947.
- 51 Gillis, C., Loiselle, S.E., Fiore, J.F., Jr., Awasthi, R., Wykes, L., Liberman, A.S., Stein, B.,  
52 Charlebois, P., Carli, F., 2016. Prehabilitation with Whey Protein Supplementation on  
53 Perioperative Functional Exercise Capacity in Patients Undergoing Colorectal Resection for  
54 Cancer: A Pilot Double-Blinded Randomized Placebo-Controlled Trial. *J Acad Nutr Diet* 116  
55 (5), 802-812.
- 56 Goodwin, P.J., Chlebowski, R.T., 2016. Obesity and Cancer: Insights for Clinicians. *Journal of*  
57 *Clinical Oncology* 34 (35), 4197-4202.


- Greenlee, H., Shi, Z., Molmenti, C.L.S., Rundle, A., Tsai, W.Y., 2016. Trends in Obesity Prevalence in Adults With a History of Cancer: Results From the US National Health Interview Survey, 1997 to 2014. *Journal of Clinical Oncology* 34 (26), 3133-3140.
- Hu W-H, Cajas-Monson LC, Eisenstein S, Parry L, Cosman B, S., R., 2015. Preoperative malnutrition assessments as predictors of postoperative mortality and morbidity in colorectal cancer: an analysis of ACS-NSQIP. *Nutrition Journal* 14 (91).
- Jensen, B.T., Petersen, A.K., Jensen, J.B., Laustsen, S., Borre, M., 2015. Efficacy of a multiprofessional rehabilitation programme in radical cystectomy pathways: a prospective randomized controlled trial. *Scand J Urol* 49 (2), 133-141.
- Jensen, B.T.J., Krintel Petersen, A.K.P., Jensen, J.B., Lausten, S.L., Borre, M.B., 2014. Efficacy of an exercise-based rehabilitation programme in radical cystectomy pathways: A randomised controlled trial. *Eur Urol Suppl* 13, e219.
- Jiralerspong, S., Goodwin, P.J., 2016. Obesity and Breast Cancer Prognosis: Evidence, Challenges, and Opportunities. *Journal of Clinical Oncology* 34 (35), 4203-4216.
- Jones, L.W., Peddle, C.J., Eves, N.D., Haykowsky, M.J., Courneya, K.S., Mackey, J.R., Joy, A.A., Kumar, V., Winton, T.W., Reiman, T., 2007. Effects of presurgical exercise training on cardiorespiratory fitness among patients undergoing thoracic surgery for malignant lung lesions. *Cancer* 110 (3), 590-598.
- Land, L.H., Dalton, S.O., Jensen, M.-B., Ewertz, M., 2012. Impact of comorbidity on mortality: a cohort study of 62,591 Danish women diagnosed with early breast cancer, 1990–2008. *Breast Cancer Research and Treatment* 131 (3), 1013-1020.
- Land, L.H., Dalton, S.O., Jensen, M.B., Ewertz, M., 2012. Influence of comorbidity on the effect of adjuvant treatment and age in patients with early-stage breast cancer. *Br J Cancer* 107 (11), 1901-1907.
- Le Roy, B., Pereira, B., Bouteloup, C., Costes, F., Richard, R., Selvy, M., Petorin, C., Gagniere, J., Futier, E., Slim, K., Meunier, B., Mabrut, J.Y., Mariette, C., Pezet, D., 2016. Effect of prehabilitation in gastro-oesophageal adenocarcinoma: study protocol of a multicentric, randomised, control trial-the PREHAB study. *BMJ Open* 6 (12), e012876.
- Li, C., Carli, F., Lee, L., Charlebois, P., Stein, B., Liberman, A.S., Kaneva, P., Augustin, B., Wongyingsinn, M., Gamsa, A., Kim, D.J., Vassiliou, M.C., Feldman, L.S., 2013. Impact of a trimodal prehabilitation program on functional recovery after colorectal cancer surgery: a pilot study. *Surg Endosc* 27 (4), 1072-1082.
- Macmillan Cancer Support, 2018. Cancer rehabilitation pathways. Macmillan Cancer Support, London.
- Mayo, N., Feldman, L., Scott, S., Zavorsky, G., Kim, D., Charlebois, P., Stein, B., Carli, F., 2011. Impact of preoperative change in physical function on postoperative recovery: argument supporting prehabilitation for colorectal surgery. *Surgery* 150 (3), 505-514.
- Minnella, E.M., Awasthi, R., Gillis, C., Fiore, J.F., Jr., Liberman, A.S., Charlebois, P., Stein, B., Bousquet-Dion, G., Feldman, L.S., Carli, F., 2016. Patients with poor baseline walking capacity are most likely to improve their functional status with multimodal prehabilitation. *Surgery* 160 (4), 1070-1079.
- Montgomery, A.A., Peters, T.J., Little, P., 2003. Design, analysis and presentation of factorial randomised controlled trials. *BMC Medical Research Methodology* 3 (1), 26.
- Moriya, T., 2015. Effects of Preoperative Use of an Immune-Enhancing Diet on Postoperative Complications and Long-Term Outcome: A Randomized Clinical Trial in Colorectal Cancer Surgery in Japanese Patients. *Gastroenterology and Hepatology* 2 (1), 1-8.
- National Academies of Sciences, E.a.M., 2018. Long-term survivorship care after cancer treatment: Proceedings of a workshop. The National Academies Press, Washington, DC: .
- Onstad, M.A., Schmandt, R.E., Lu, K.H., 2016. Addressing the Role of Obesity in Endometrial Cancer Risk, Prevention, and Treatment. *Journal of Clinical Oncology* 34 (35), 4225-4230.
- Parker, P.A., Pettaway, C.A., Babaian, R.J., Pisters, L.L., Miles, B., Fortier, A., Wei, Q., Carr, D.D., Cohen, L., 2009. The Effects of a Presurgical Stress Management Intervention for Men With Prostate Cancer Undergoing Radical Prostatectomy. *J Clin Oncol* 27 (19), 3169-3176.

- 1  
2  
3 Peddle, C.J., Jones, L.W., Eves, N.D., Reiman, T., Sellar, C.M., Winton, T., Courneya, K.S., 2009.  
4 Effects of presurgical exercise training on quality of life in patients undergoing lung resection  
5 for suspected malignancy: a pilot study. *Cancer Nurs* 32 (2), 158-165.
- 6 Sarfati, D., Koczwara, B., Jackson, C., 2016. The impact of comorbidity on cancer and its treatment.  
7 (1542-4863 (Electronic)).
- 8 Schmid, D., Leitzmann, F., 2014. Association between physical activity and mortality among breast  
9 cancer and colorectal survivors: a systematic review and meta-analysis. *Annals of Oncology*  
10 25, 1293-1311.
- 11 Schmidt, M., Eckardt, R., Scholtz, K., Neuner, B., von Dossow-Hanfstingl, V., Sehoul, J., Stief,  
12 C.G., Wernecke, K.D., Spies, C.D., Group, P., 2015. Patient Empowerment Improved  
13 Perioperative Quality of Care in Cancer Patients Aged  $\geq 65$  Years - A Randomized  
14 Controlled Trial. *PLoS One* 10 (9), e0137824.
- 15 Schmitz KH, Courneya KS, Matthews C, Demark-Wahnefried W, Galvão DA, Pinto BM, Irwin ML,  
16 Wolin KY, Segal RJ, Lucia A, Schneider CM, von Gruenigen VE, A, S.A., Medicine.,  
17 m.C.o.S., 2010. American college of sports medicine roundtable on exercise guidelines for  
18 cancer survivors. 2010;42:. *Med Sci Sports Exerc* 42, 1409-1426.
- 19  
20 Sekine, Y., Chiyo, M., Iwata, T., Yasufuku, K., Furukawa, S., Amada, Y., Iyoda, A., Shibuya, K.,  
21 Iizasa, T., Fujisawa, T., 2005. Perioperative rehabilitation and physiotherapy for lung cancer  
22 patients with chronic obstructive pulmonary disease. *Jpn J Thorac Cardiovasc Surg* 53 (5),  
23 237-243.
- 24 Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, LA, S., Group, t.P.-  
25 P., 2015. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols  
26 (PRISMA-P) 2015: elaboration and explanation. *BMJ* 349 (g7647).
- 27 Shun, S.C., 2016. Cancer Prehabilitation for Patients Starting from Active Treatment to Surveillance.  
28 *Asia Pac J Oncol Nurs* 3 (1), 37-40.
- 29 Silver, J.K., 2015. Cancer prehabilitation and its role in improving health outcomes and reducing  
30 health care costs. *Semin Oncol Nurs* 31 (1), 13-30.
- 31 Silver, J.K., 2014. Cancer rehabilitation and prehabilitation may reduce disability and early  
32 retirement. *Cancer* 120 (14), 2072-2076.
- 33 Silver, J.K., Baima, J., 2013. Cancer prehabilitation: an opportunity to decrease treatment-related  
34 morbidity, increase cancer treatment options, and improve physical and psychological health  
35 outcomes. *Am J Phys Med Rehabil* 92 (8), 715-727.
- 36 Singh, F., Netwon, R., Galvao, D., Spry, N., Baker, M., 2013. A systematic review of pre-surgical  
37 exercise intervention studies with cancer patients. *Surgical Oncology* 22, 92-104.
- 38 Sitas, F., Webe, M., Egger, S., Yap, S., Chiew, M., O'Connell, D., 2014. Smoking Cessation After  
39 Cancer. *Journal of Clinical Oncology* 32 (32), 3593-3595.
- 40 Smith, S.R., Khanna, A., Wisotzky, E.M., 2017. An Evolving Role for Cancer Rehabilitation in the  
41 Era of Low-Dose Lung Computed Tomography Screening. *PM&R* 9 (9), S407-S414.
- 42 Sogaard, M., Thomsen, R.W., Bossen, K.S., Sorensen, H.T., Norgaard, M., 2013. The impact of  
43 comorbidity on cancer survival: a review. *Clinical Epidemiology* 5 (Suppl 1), 3-29.
- 44 Stacey FG, James EL, Chapman K, Courneya KS, DR, L., 2015. A systematic review and meta-  
45 analysis of social cognitive theory-based physical activity and/or nutrition behavior change  
46 interventions for cancer survivors. *J Cancer Surviv.* 9 (2), 305-338.
- 47 Stairmand, J., Signal, L., Sarfati, D., Jackson, C., Batten, L., Holdaway, M., Cunningham, C., 2015.  
48 Consideration of comorbidity in treatment decision making in multidisciplinary cancer team  
49 meetings: a systematic review. *Ann. Onc.*, mdv025.
- 50 Stefanelli, F., Meoli, I., Cobuccio, R., Curcio, C., Amore, D., Casazza, D., Tracey, M., Rocco, G.,  
51 2013. High-intensity training and cardiopulmonary exercise testing in patients with chronic  
52 obstructive pulmonary disease and non-small-cell lung cancer undergoing lobectomy. *Eur J*  
53 *Cardiothorac Surg* 44 (4), e260-265.
- 54 Sterne, J.A., Hernán, M.A., Reeves, B.C., Savović, J., Berkman, N.D., Viswanathan, M., Henry, D.,  
55 Altman, D.G., Ansari, M.T., Boutron, I., Carpenter, J.R., Chan, A.-W., Churchill, R., Deeks,  
56 J.J., Hróbjartsson, A., Kirkham, J., Jüni, P., Loke, Y.K., Pigott, T.D., Ramsay, C.R., Regidor,

- 1  
2  
3 D., Rothstein, H.R., Sandhu, L., Santaguida, P.L., Schünemann, H.J., Shea, B., Shrier, I.,  
4 Tugwell, P., Turner, L., Valentine, J.C., Waddington, H., Waters, E., Wells, G.A., Whiting,  
5 P.F., Higgins, J.P., 2016. ROBINS-I: a tool for assessing risk of bias in non-randomised  
6 studies of interventions. *BMJ* 355.
- 7 Stout, N.L., Alfano, C.M., Belter, C.W., Nitkin, R., Cernich, A., Lohmann Siegel, K., Chan, L., 2018.  
8 A Bibliometric Analysis of the Landscape of Cancer Rehabilitation Research (1992-2016). *J*  
9 *Natl Cancer Inst* 110 (8), 815-824.
- 10 Stout, N.L., Silver, J.K., Raj, V.S., Rowland, J., Gerber, L., Cheville, A., Ness, K.K., Radomski, M.,  
11 Nitkin, R., Stubblefield, M.D., Morris, G.S., Acevedo, A., Brandon, Z., Braveman, B.,  
12 Cunningham, S., Gilchrist, L., Jones, L., Padgett, L., Wolf, T., Winters-Stone, K., Campbell,  
13 G., Hendricks, J., Perkin, K., Chan, L., 2016. Toward a National Initiative in Cancer  
14 Rehabilitation: Recommendations From a Subject Matter Expert Group. *Arch Phys Med*  
15 *Rehabil* 97 (11), 2006-2015.
- 16 Sueppel, C., Kreder, K., See, W., 2001. Improved continence outcomes with preoperative pelvic floor  
17 muscle strengthening exercises. *Urol Nurs* 21 (3), 201-210.
- 18 Thompson, W., Gordon, N., Pescatello, L., 2010. ACSM's Guidelines for Exercise Testing and  
19 Prescription. Lippincott, Williams and Wilkins, Philadelphia PA.
- 20 Tsimopoulou, I., Pasquali, S., Howard, R., Desai, A., Gourevitch, D., Tolosa, I., Vohra, R., 2015.  
21 Psychological Prehabilitation Before Cancer Surgery: A Systematic Review. *Ann Surg Oncol*  
22 22 (13), 4117-4123.
- 23 Van Bokhorst-de Van der Schuer, M.A., Langendoen, S.I., Vondeling, H., Kuik, D.J., Quak, J.J., Van  
24 Leeuwen, P.A., 2000. Perioperative enteral nutrition and quality of life of severely  
25 malnourished head and neck cancer patients: a randomized clinical trial. *Clin Nutr* 19 (6),  
26 437-444.
- 27 Vidal, A., Howard, L., Moreira, D., Castro-Santamaria, R., Andriole GL, J., Freedland, S., 2014.  
28 Obesity increases the risk for high-grade prostate cancer: results from the REDUCE study.  
29 *Cancer Epidemiol Biomarkers Prev.* 23 (12), 2936-2942.
- 30 Warren, G.W., Sobus, S., Gritz, E.R., 2014. The biological and clinical effects of smoking by patients  
31 with cancer and strategies to implement evidence-based tobacco cessation support. *The*  
32 *Lancet. Oncology* 15 (12), e568-e580.
- 33 West, M.A., Loughney, L., Lythgoe, D., Barben, C.P., Sripadam, R., Kemp, G.J., Grocott, M.P., Jack,  
34 S., 2015. Effect of prehabilitation on objectively measured physical fitness after neoadjuvant  
35 treatment in preoperative rectal cancer patients: a blinded interventional pilot study. *Br J*  
36 *Anaesth* 114 (2), 244-251.
- 37 Winters-Stone, K., Dieckmann, N., Maddalozzo, G., Bennett JA, Ryan CW, TM., B., 2015.  
38 Resistance Exercise Reduces Body Fat and Insulin During Androgen-Deprivation Therapy for  
39 Prostate Cancer. *Oncology Nursing Forum* 42 (4), 348-356.
- 40 Yang, L., Drake, B.F., Colditz, G.A., 2016. Obesity and Other Cancers. *Journal of Clinical Oncology*  
41 34 (35), 4231-4237.
- 42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1. PRISMA study selection flowchart







1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Table 1 Risk of bias was assessed by an interdisciplinary research team using the Cochrane Collaboration’s tool for assessing risk of bias ROBINS-I tool

		Random Sequence Generation	Allocation concealment	Blinding of participants and personnel Blinding of	Incomplete outcome data	Selective outcome reporting
		random eg. random number table, computer	participants and investigators enrolling participants could not foresee assignment	outcome and the outcome measurement are not likely to be	no missing outcome data, reasons for missing data similar across groups	study protocol available and all pre-specified outcomes are
x	Research Design	Criteria 1	Criteria 2	Criteria 3	Criteria 4	Criteria 5
Bales, 2000	RCT	unclear	high	unclear	unclear	low
Burgio, 2006	RCT	low	high	low	unclear	low
Carli, 2010	RCT Stratified randomisation	unclear	unclear	high	multiple imputations	low
Centemero, 2010	RCT	low	low	low	low	low
Gillis, 2014	RCT	low	low	low	multiple imputations	low
Jensen, 2014	RCT	unclear	unclear	unclear	unclear	high
Jensen, 2015	RCT	low	low	unclear	low	low
Stefanelli, 2013	RCT	unclear	unclear	unclear	unclear	low
Barlesi, 2008	RCT	unclear	unclear	unclear	high	low
Cheville, 2015	RCT	unclear	unclear	low	low	low
Garssen, 2013	RCT Block Randomised	unclear	unclear - block randomiza	low	unclear	high
Parker, 2009	RCT	unclear	unclear	low	unclear/high	low
Schmidt, 2015	RCT Block randomised	low	unclear	unclear	high	low
Gillis, 2015	RCT	low	low	low	low	low
Moriya, 2015	RCT	low	low	unclear	low	low
Van Bokhorst, 2000	RCT	unclear	unclear	low	high	low

Table 2 Randomized Controlled Trials (RCTs) of Prehabilitation and critical analysis

Author and date	Intervention description	Control group description	Sample	Follow-up after intervention	Attrition	Critical analysis
Bales, 2000	<i>Frequency:</i> 2-4 weeks prior to surgery <i>Intensity:</i> nurse-led biofeedback, 10-15 repetitions <i>Advised to practice</i> 4x per day <i>Timing</i> 45 minutes <i>Type:</i> pelvic floor muscle exercise plus biofeedback followed by post-op PFM exercises	Written and brief verbal instructions on how to perform pelvic floor muscle exercises	100 men undergoing radical prostatectomy Mean age intervention 60.9yrs and control 59.3yrs.  Prostate cancer stages T1c-T2c  Comorbidity not reported	Every month for 6-months post-surgery	3% 6%	By 6 months following radical prostatectomy, the incidence of urinary continence in the biofeedback and control groups was 94% and 96%, respectively.  No difference in patients who received biofeedback pre op and those who did not. There was no objective measure, just number of wet diapers (vs weight of diaper); unclear about characteristics of those who dropped out, unknown pelvic floor muscle strength prior to undertaking study
Barlesi, 2008	<i>Frequency:</i> unclear <i>Intensity:</i> unclear  <i>Timing:</i> Prior to surgery <i>Type:</i> Additional oral plus written information including associated symptoms	Oral information only describing the disease and its associated surgery and outcomes	75 patients with NSCLC undergoing thoracic surgery  Comorbidity not reported	Baseline and 3 months	26%	QoL scores (baseline, 3 months) were comparable between both groups. Patients receiving oral plus written information were significantly dissatisfied related to several aspects of care regarding staff as well as the structure.  The information group significantly influenced satisfaction levels at multivariate analysis (standardized beta coefficient, 0.26, $p = 0.04$ ).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Burgio, 2006	<p><i>Frequency:</i> initiated 1 week prior to surgery  <i>Intensity:</i> Daily 45 pelvic floor exercises  <i>Timing:</i> One preoperative session  <i>Type:</i> biofeedback plus assisted behavioural training                  Home based exercise</p>	Usual Care	<p>125 men undergoing surgery</p> <p>Mean age 60.9 +/- 6.9yrs.</p> <p>Prostate cancer</p> <p>Comorbidity reported</p>	6 months' post-surgery	10%	<p>At 6 months:                  Difference between the groups in the proportion of men remaining incontinent was 20.03% (higher in the intervention group) (95% CI 6.02% to 34.63%) (p&lt;0.04).</p> <p>Severe/continual leakage was still present in 19.6% of controls compared to 5.9% of those in intervention group (p &lt;0.04).</p> <p>Intervention group had a</p> <ul style="list-style-type: none"> <li>- higher proportion of dry days (p &lt; 0.04),</li> <li>- lower proportion using pads (p &lt;0.05).</li> </ul> <p>No group differences were found in life-style variables,</p> <ul style="list-style-type: none"> <li>- incontinence impact (p = 0.36,</li> <li>- psychological distress (p = 0.69)</li> <li>- quality of life (p = 0.31 to 0.89).</li> </ul> <p>Unclear how long provided and the intensity of the exercises</p>
Carli, 2010	<p><i>Frequency:</i> average 52 days prior to surgery  <i>Intensity:</i> High intensity exercise  <i>Timing:</i> 3 times per week  <i>Type:</i> Prescribed stationary cycling (daily) with strengthening prescribed</p>	Walk/breathing group: recommendations to walk daily and perform foot and ankle exercises to enhance lower-extremity circulation as well as breathing exercises	<p>112 patients undergoing colorectal surgery.</p> <p>Mean age 60 (SD 16)</p> <p>Participants with comorbidities grade IV or V were excluded from study or if unable to complete testing procedure</p>	10 weeks' post-surgery	16%	<p>Improvement in walking capacity in walk/breathing (47%) vs. bike/strengthening pre-surgery (22%). But not sustained over time</p> <p>Mean peak <math>\dot{V}O_2</math> improved in both groups: Bike/ strengthening 134 ml/min (P = 0.003) versus walk breathing 112 ml/min (P = 0.007) but not over time.</p> <p>Anxiety considerably reduced after surgery but did not change in either group over the prehabilitation period.</p> <p>Depression improved for the bike/strengthening group over the prehabilitation period.</p> <p>Exercise participation bike/strengthening group &gt; walk/breathing group (P = 0.075).</p> <p>Lower exercise had better outcomes because of less dropouts, poor compliance in the higher intensity group may have been too hard for such patients</p>

Centemero, 2010	<p><i>Frequency:</i> started 30 days before surgery 2x per week</p> <p><i>Intensity:</i> Physiotherapist encouragement</p> <p><i>Timing:</i> 30 min</p> <p><i>Type:</i> pelvic floor muscle exercises at hospital and at home</p>	Postoperative pelvic floor muscle exercises 48hrs after catheter removal	<p>118 males undergoing surgery.</p> <p>46-68yrs old.</p> <p>Prostate cancer</p> <p>Comorbidity not reported</p>	3 months	17% didn't start study but were eligible	<p>Continence in preoperative group was 59.3% vs. postoperative group 37.3% (p&lt;0.028)</p> <p>ICS male SF mean score in preoperative group 8.1 vs. postoperative group 12.2.</p> <p>Study found that preoperative PFME improved quality of life (The ICFS is a symptom based tool)</p> <p>No detail re adherence to exercises. The question whether the preoperative or post-operative had most effect is unclear. Differences persisted for up to 6 months at 1 year there was no difference.</p>
Cheville, 2015	<p><i>Frequency:</i> 2-3X per week</p> <p><i>Intensity:</i></p> <p><i>Timing:</i> (6-8 sessions)</p> <p><i>Type:</i> led by psychiatrist included social, cognitive, emotional care via exercise, education and relaxation.</p>	Usual care	<p>61 study sample</p> <p>Women, mean age 61.2</p> <p>Comorbidity not reported</p>	30-day readmission	Not reported	<p>More patients in the intervention completed Chemotherapy (p=0.003) than control</p> <p>Intervention group had significantly fewer treatment hospitalisations (p=0.001)</p> <p>No difference in other measures</p> <p>There is no data capture on medications or functional health status. No PROM outcomes or psychological measures all data capture was through EMR retrospective records.</p>
Garssen, 2013	<p><i>Frequency:</i> 4 sessions 5 and 1 day prior to surgery with session 2 and 30 days post-surgery</p> <p><i>Intensity:</i></p> <p><i>Timing:</i> 40-60 minutes</p> <p><i>Type:</i> Stress management training delivered by clinical psychologist - relaxation, guided imagery techniques, and counselling</p>	Usual care	<p>70 women undergoing surgery for breast cancer</p> <p>Intervention mean age 52yrs control group mean age 54 years</p> <p>Comorbidity partially reported (BMI, alcohol use)</p>	30- 90 days post-surgery	<p>18%</p> <p>Only 57% of eligible patients finished study</p>	<p>At 3 months' post-surgery compared to baseline, the intervention group had:</p> <ul style="list-style-type: none"> <li>- Significant difference at 1 month in depression</li> <li>- no significant difference in quality of life</li> <li>- no significant difference in wellbeing</li> <li>- pain not measured at 3 months</li> <li>- control group more complaints than intervention group (0.001&lt;p&lt;0.01).</li> </ul> <p>At 3-month post-op measures will be affected by start of adjuvant treatment (and no information reported on this). Experience of control measured by author designed 4 item questionnaires (un validated measures). Unable to differentiate effect of intervention content or psychologist intervention. The study power is not clear against its primary outcome. Targeted intervention. The differences between groups at 3 months are not significant.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

<p>Gillis, 2014</p>	<p><i>Frequency:</i> 24 days 3 days per week <i>Intensity:</i> moderate aerobic and resistance exercises, <i>Timing:</i> 50 minutes <i>Type</i> Home-based unsupervised initially personalised to the individual. Nutrition counselling with protein supplementation Coping strategies to reduce anxiety and promote adherence with exercises provided on a CD</p>	<p>Rehabilitation group or 8 weeks post operatively (same as intervention but after surgery)</p>	<p>89 patients undergoing surgery for colorectal cancer</p> <p>prehab mean age = 65.7 (13.6); rehab mean = 66.0 (9.1)</p> <p>Comorbidity fully reported</p>	<p>8 weeks' post-op</p>	<p>13%</p>	<p>Difference between baseline and follow-up in prehabilitation group. Mean different 45.4m (95% CI, 13.9 to 77.0)</p> <p>Baseline: Prehab 421m (SD, 120.0) Rehab 425m (SD, 83.8) Pre-treatment: Prehab +25.2 (50.2) Rehab -16.4 (46.0) P=0.001 50% of patients in both groups remained more than 20 m below baseline</p> <p>At 8 weeks, prehab +23.4 (54.8) rehab -21.8 (80.7). P = 0.020 Complication rates and length of hospital stay were similar in prehabilitation and rehabilitation groups.</p> <p>A change of 20 m is considered clinically meaningful as this is the estimated measurement of community dwelling elderly</p> <p>Limitation of the study is missing data and unclear which modality of prehabilitation responsible for outcomes.</p>
<p>Gillis, 2015</p>	<p><i>Frequency:</i> 4 weeks prior to surgery: <i>Intensity:</i> daily intervention <i>Timing:</i> 90 min per day <i>Type:</i> individualised nutrition counselling with daily whey protein supplementation</p>	<p>Individualised nutrition counselling with a non-nutritive placebo</p>	<p>43 patients undergoing surgical treatment.</p> <p>Mean age 67.6yrs (sd 11.5). Mean age in placebo group (69.1yrs (sd 9.4)</p> <p>Comorbidities partially reported</p>	<p>4-8 weeks post-op</p>	<p>10%</p>	<p>Before surgery improvement in whey group 20.8m (sd 42.6m) and in placebo group (1.2m (sd 65.5m) (p=0.27).</p> <p>Recovery rates were similar between groups in the 4-weeks post-surgery (p=0.81). Comment: intervention focused on building strength. Focus on upper body strength and not lower body. Post intervention lack of nutrition may impact on similar group trajectories. Pre-surgery results are significant but not sustained at 4 weeks</p> <p>6MWT data missing pre-op for 4 patients (2 placebo, 2 whey) and 12 post-op (4 placebo, 8 whey), analysis based on 32 participants. If protein required to improve muscle functional capacity, presumably protein supplementation should be continued post-op for functional capacity to be maintained? This study indicates that effects may only be short-term (i.e. 4 weeks to day of surgery, and associated with duration of intervention)</p>

5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	Jensen, 2014 <i>Frequency:</i> 2 weeks prior to surgery: <i>Intensity:</i> 15-minute step training <i>Timing:</i> 2x per day <i>Type:</i> Written information and motivation home based Pt tailored exercises for 6 muscle strength and endurance  Standardised postoperative mobilisation	Standardised nutritional screening and counselling.  Standardised postoperative mobilisation was encouraged at least 2x at 30 min day	129 patients undergoing radical cystectomy  (analysis based on 100 (intervention 47, standard 53)  Bladder cancer  Comorbidity fully reported	4 months' post-surgery	41%	There was no significant difference in LOS and number of adverse events. Physical capacity was significantly improved (p0.02) and mean walking distance at 7 days' post op.  At follow up both groups had regained physical capacity and no difference was seen.  Patient tailored intervention but unclear as to how this was personalised Abstract has little information re intervention  Just over half (55%) completed the prehab programme at 100%, 59% fulfilled 75%. It would have been useful to compare groups on 6MWT rather than LOS as primary outcome measure.
21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36	Jensen, 2015 <i>Frequency:</i> 2 x daily <i>Intensity:</i> progressive strength and endurance exercises <i>Timing</i> daily <i>Type:</i> Preoperative home-based supervised exercise programme and postoperative and progressive postoperative mobilization	Fast track Patient education counselling on choice of urinary intervention, pre-op preparation, pain control and nutrition	107 patients undergoing surgical treatment for bladder cancer. Mean age intervention group 66yrs, mean age control group 71yrs.  Bladder cancer  Comorbidities fully reported	4 months' post-surgery	7% attrition  55% adherence	Severity of complications: no significant difference was found in the incidence (p = 0.47) or severity (p = 0.64) of complications between the treatment groups at 90 days postoperatively, or in readmission within 30 days (p = 0.49).  Ability to perform ADL: the median time was 3 days in the intervention group, compared with 4 days in the standard group (p < 0.05).  Post-op mobilization: significantly higher in the intervention group, reporting 4806m walked (95% CI 4075 to 5536 m), compared to the standard group with 2906 m walked (95% CI 2408 to 3404 m) (p < 0.001) at 7 days  Significance in 4 month outcomes between groups not seen.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Moriya 2015,	<i>Intervention 1</i> <i>Frequency:</i> 5 days pre-operatively <i>Intensity:</i> High dose 750ml/ day <i>Timing:</i> Daily <i>Type:</i> Immune-enhancing diet (IED) (enriched with arginine, omega-3 fatty acids and RNA) and normal food  <i>Intervention 2.</i> <i>Intensity:</i> Low-dose 250ml/day Immune enhancing diet	Control group – normal food	88 patients undergoing surgery for colorectal cancer  Mean age intervention 64.7 (2.3) control 63.8 (2)  Comorbidity not reported	9-133 months after surgery	Not reported	Incisional SSI rates in the IED groups were significantly lower in the intervention group than in the Control group. (0%*, 0%* and 17%) (*P<0.01 vs. Control).  The incidences of the infections not involving the surgical site (non-SSI) and the lengths of hospital stay were similar among the three groups. No significant differences were observed in RFS or DSS.
--------------	---	-----------------------------	---	----------------------------	--------------	--

Parker, 2009	<p><i>Frequency:</i> 1-2 weeks pre-op  <i>Intensity:</i>  <i>Timing</i> 2x 60-90 minute sessions plus 2 booster sessions on morning of surgery and 48 hours post-surgery)  <i>Type:</i>  <i>Intervention 1.</i> Stress management (SM) including diaphragmatic breathing and guided imagery individual sessions with clinical psychologist,  <i>Intervention 2</i> Supportive attention (SA) group discussed their concerns about the upcoming surgery and had a semi-structured medical interview.</p>	Standard care (no meetings with clinical psychologist)	<p>159 men undergoing surgery for prostate cancer.</p> <p>Mean age 60.9 (5.9)</p> <p>Comorbidities not reported</p>	6-12 months post-surgery	34%	<p>Post treatment improvements of intervention on mood disturbance (p0.02) with the stress management group with no significant differences between groups in any of the assessment times between groups over time.</p> <p>The mixed model analysis targeted therapies. Differences in mood were small and although significant are not clinically significant. At 1 year patients had better physical function (SF36) but this was self-reported. A targeted intervention on those individuals with higher stress may be more beneficial in terms of effect size.</p> <p>Men in SM group had significantly higher physical component summary score on SF35 than men in SC group at one year (p=0.0009), but no difference in mental component summary score or prostate specific Qol in PCI. The study excluded emotionally distressed men who may benefit from such an intervention.</p>
--------------	---	--	---	--------------------------	-----	--



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Schmidt, 2015	<p><i>Frequency:</i> 1 day pre-operatively  <i>Intensity:</i>  <i>Timing:</i> 7 days  <i>Type</i> Information booklet lifestyle advice, mobilisation, nutrition and diary keeping</p>	Standard care - information regarding surgical and anaesthesiology risks and procedures	<p>652 patient's undergoing elective surgery for gastro-intestinal, genitourinary, and thoracic cancer</p> <p>65 years of age</p> <p>Comorbidities reported</p>	3 -12 months after surgery	15%	<p>Complications: Occurrence and severity of complications were comparable in both groups, although severe haemorrhage occurred significantly more often in the intervention group (6.7% vs. 2.5%; p = 0.01).</p> <p>LOS: no significant difference between both groups (p = 0.99).</p> <p>HRQoL: no significant difference between the global HRQoL 12 months after surgery in the intervention and in the control group.</p> <p>Postoperative stress: (mobilization, PONV and postoperative pain). Patients in the intervention group reported less pain on the first postoperative day (75.2% vs. 82.3%, p = 0.03). There were no differences regarding mobilization within the first 24 hours (69.2% vs. 70.4%, p = 0.73), or PONV within the first five days (52.8% vs. 56.4%, p = 0.39).</p> <p>Depression: There was no difference in the geriatric depression scale between intervention and control groups at discharge (p = 0.86).</p> <p>Readmission: The readmission rate within 90 days was slightly higher for patients in the intervention group (p = 0.70). In-hospital length of stay at readmission was shorter than in the standard care group without reaching statistical significance (p = 0.22).</p> <p>Mortality: The overall mortality did not differ significantly between the two groups (Log-Rank-test p = 0.197).</p> <p>Patient empowerment failed to shorten LOS or HRQOL. This type of intervention could enhance quality of care in regards to pain, and since over-treatment of pain is particularly harmful for elderly patients, patient safety can thus be improved. Pre-operative information was received well by patients who were cognitively and physically fit.</p>
---------------	---	---	---	----------------------------	-----	--

Stefanelli 2013	<p><i>Frequency:</i> 3 weeks preoperatively  <i>Intensity:</i> high-intensity training  <i>Timing:</i> 15x3-hour sessions weekdays  <i>Type:</i> outpatient intensive pulmonary rehabilitation (PRP) based on of both upper- and lower-limb muscles</p>	Control surgery according to the normal standard preoperative protocol	<p>40 NSCL and COPD undergoing lobectomy</p> <p>Age &gt;75</p> <p>Excluded patients with comorbidity</p>	60 days after surgery	Not reported	<p>A significant difference was observed both at T1 and T2. In prehabilitation group, peak VO2 improves significantly from T0 to T1, <math>P &lt; 0.001</math> and deteriorates from T1 to T2, <math>P &lt; 0.001</math> in control reverting to a similar value to that at T0.</p> <p>Control group peak VO2 did not change from T0 to T1 and significantly deteriorates from T1 to T2: <math>P &lt; 0.00001</math>. FEV1 NS T0, T1, T2</p> <p>Currently, other studies are needed to demonstrate that the patients who undergo preoperative PRP could have also a better quality of life, less postoperative complications and a longer survival after surgery. Good for CPT VS 6MWT comparison</p>
Van Bokhorst, 2000	<p><i>Frequency:</i> 7-10 days pre-operatively plus 14 days post op  <i>Intensity:</i> 150% of basal energy expenditure  <i>Timing:</i> Daily  <i>Type:</i>  1. Standard pre-and post-op enteral feeding   2. Arginine supplemented pre-op and post-op enteral feeding</p>	No pre-op and standard post-op enteral feeding	<p>49 malnourished head and neck cancer patients</p> <p>Mean age 56.6-61.6</p> <p>Comorbidity fully reported</p>	6 months post-surgery.	37%	<p>Between baseline and the day before surgery, both preoperatively fed groups revealed a positive change for physical and emotional functioning and dyspnoea with significance in arginine group. This was not sustained long term at 6 months.</p> <p>Supplemented group showed a negative change in appetite (<math>P=0.049</math>). Between baseline and 6 months after surgery, There were no differences between control and both pre-fed groups.</p> <p>There were no differences in favour of Arginine supplement compared to other feed group.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Table 3 Observational and Quasi Experimental Prehabilitation Studies with critical analysis

Author, date and research design	Intervention and comparator	Population and Sample	Follow-up after intervention	Attrition	Critical analysis
<p>Baima, 2015</p> <p>Feasibility study with two randomised (by appointment time), non-blinded groups</p>	<p>Timing: 1 month prior to surgery.</p> <p>Type: In-person teaching of exercises plus info sheet, plus link to online video provided</p> <p>Comparator</p> <p>Video-only teaching arm</p>	<p>60 cancer patients undergoing surgery for breast cancer</p> <p>1 male, 59 females</p> <p>Age 35-81.</p> <p>(I=36-C24)</p> <p>Breast cancer</p>	<p>3-mths after surgery</p> <p>2 weeks to 6 months</p>	<p>25%</p>	<p>Exercise compliance:</p> <ul style="list-style-type: none"> <li>- 76% chose to exercise.</li> <li>- No difference in exercise compliance between in-person teaching versus video teaching (OR=1.03).</li> </ul> <p>In person 75% (24/32) compared to Video teaching 77% (10/13)</p> <p>Pain</p> <ul style="list-style-type: none"> <li>- 29% of patients (9/31) had worse shoulder pain than baseline at 1 month post-surgery (24 %, 6/25 exercisers, and 50 %, 3/6 non-exercisers).</li> <li>- 15% percent of patients (4/27) had worse shoulder pain than baseline at 3 months' post-surgery (8 %, 2/23 exercisers, and 100 %, 2/2 non-exercisers).</li> </ul> <p>Shoulder Abduction</p> <ul style="list-style-type: none"> <li>- 66% of patients (20/30) lost greater than 10% shoulder abduction ROM at 1 month post-surgery.</li> </ul> <p>Prehabilitation exercise program inferred no additional risk of seroma formation (Exercisers 21 %, 7/33 vs. non-exercisers 22 %, 2/9, OR=0.94).</p> <p>No strong evidence of difference</p> <p>Single site; Change in intervention based on patient preference, so not random; No control group: every participant received some sort of intervention; Stated feasibility study, but unclear re: efficacy of these home exercises; No explanation given for why study staff did not pursue missing data; Possible social desirability bias with “in-person” arm and thus improved compliance; Compliance with exercises self-reported, possibly participants overestimated; Unclear description regarding seroma formation/evaluation ; Time to follow-up variability (2 weeks – 6 months); Lumpectomy and mastectomy included, possibly allowing for very different outcomes</p>
<p>Jones, 2007</p> <p>Single group design feasibility study</p>	<p>Frequency: 5 endurance sessions per week on consecutive days until surgical resection. .</p> <p>Intensity: Highly individualised and progressive from 60-100% VO<sub>2 peak</sub>, Threshold sessions</p>	<p>25 patients 70% F undergoing surgery for suspected lung cancer.</p> <p>Mean age 65+/-10years</p> <p>Lung cancer</p>	<p>Preoperative assessment</p> <p>30 days post-surgery</p>	<p>Attrition 52%</p> <p>Adherence 70%</p>	<p>For patients who achieved &gt;=80% adherence (n = 12), VO<sub>2</sub> peak increased 3.3 mL/kg/min (P=0.006).</p> <p>Six-minute walk test: Significant improvement in &gt;80% adherence group (p=0.14) compared to &lt;80% adherence (p=1.01).</p> <p>The overall adherence rate was 72% (range, 0%–100%) with patients completing a mean of 30 - 27 sessions (range, 0 - 75).</p> <p>No significant difference. (p&gt;0.1) for all measures of pulmonary function</p> <p>The average duration of hospital stay was 10 _ 8 days with 8 - 5 days in general</p>

	<p>and HIIT sessions. Timing: Carried out for 4-6 weeks. Type: Cycle ergometry</p> <p>Comparator: Individually tailored intervention and hospital based over 4-6 weeks</p>				<p>hospital and 2 - 5 days in the intensive care unit. No difference in complications or length of stay than in routine patient care. The length of intervention may be problematic in a 1-2-week normal wait time for surgery, the authors comment on this re the ability to make significant change in such a short time. There is no recording of how many patients didn't meet the eligibility parameters to reflect the normal practice.</p>
<p>Li, 2013 Pre post intervention study  Historical control group  single centre, cohort</p>	<p>Frequency: Individualized aerobic exercise (30 mins x3 times a week, Intensity: at 50% of max heart rate) and resistance training Timing: 3 times per week, no more detail Type, one or two modifiable dietary behaviours identified and discussed, use of whey protein isolate within 1 hour of exercise at 1.2g/kg body weight per day , 90 minute visit with trained psychologist with a focus on anxiety reduction. Length of prehabilitation determined by wait time for surgery</p>	<p>87 patients undergoing surgery for colorectal cancer (Control =45 intervention= 42)</p> <p>Colorectal cancer</p>	<p>Assessed 1 week pre op, 4 weeks and 8 weeks post-surgery</p> <p>Primary outcome measure 6MWT @ 8 weeks</p>	<p>0%</p>	<p>The patients in the prehabilitation program had better postoperative walking capacity at 8 weeks (mean difference, 84.5 ± 83 m; p&lt;0.01). At 8 weeks, 81 % of the prehabilitated patients were recovered compared with 40 % of the control group (p&lt;0.01). The prehabilitation group also reported higher levels of physical activity before and after surgery. The postoperative complication rates and the hospital length of stay were similar. There were significant emotional &amp; social differences between control and intervention. These are not controlled for in the modelling or analysis of difference. Lack of detail re nutritional component. The question of what's a clinically relevant change NO data on the length of time of wait for surgery and this would determine the amount of time spent on prehabilitation intervention (see column 3). Historical control do not appear to have had baseline assessment, just pre op and 4 and 8 week. Could account for difference between groups on HRQoL measures, because pre-op measures taken at different times (control taken immediately before surgery; intervention after meeting with surgeon). No tool used to capture functional symptoms specific to colorectal cancer e.g. FACT-C, SF35 is too generic. Compliance to prehabilitation intervention reported in discussion section (70% exercising at least 2 days per week, 45% fully compliant). Increase in self-reported physical activity persisted after surgery.</p>
<p>Peddle, 2009 Sub analysis of Jones 2007</p>	<p>Frequency: 5 sessions per week on consecutive days until surgical resection Intensity: Highly individualised and progressive from 60-100% VO<sub>2</sub> peak, Threshold sessions and HIIT sessions. Timing: 4-6 weeks prior to surgery. Type: endurance cycle ergometry</p>	<p>19 patients undergoing lung resection for suspected malignancy were planned to complete baseline to pre-surgery intervention.</p> <p>9 patients will full data set</p> <p>Lung cancer</p>	<p>QoL 2 month post-surgery</p>	<p>0% Subset analysis</p>	<p>Pre-surgical exercise training improved cardiorespiratory fitness, it did not seem to improve QOL from baseline to pre-surgery or mitigate the decline in QOL after surgery. QoL might be influenced by several other factors for exercise to have a meaningful effect. VO<sub>2</sub> Peak did improve.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

	Comparator Individually tailored intervention and hospital based over 4-6 weeks				
Sekine, 2005 Prospective study with usual care control	<p>Frequency: 5x per day Intensity: moderate Timing: 2 weeks prior to surgery Type:                      1. Incentive spirometry                      2. Abdominal breathing and breathing exercises                      3. Bronchodilators 5x per day                      4. Huffing and coughing exercises                      5. 5000 steps per day</p> <p>Comparator historical control group of 60 patients <b>without</b> COPD</p>	<p>N=82 (control=60) Rehab (n=22)  (rehab group had more airflow obstruction FEV1/FVC)  22 patients with COPD.  Lung cancer</p>	<p>30 days' post operation  1 month post-op</p>	<p>0% Not reported</p>	<p>The change in lung function as less diminished in the intervention group (p=0.023)  Post-operative pulmonary complications there were no differences  Postoperative hospital stays were significantly longer in the control group (p=0.003) equivalent to -6.8 days  Study does not mathematically control for the differences between groups re baseline health. Study is underpowered  Less decrease in pulmonary function (FEV1 and predicted decrease post-op) in Rehab group but overall pulmonary function was lower in this group. Note: This is probably because this group had COPD whereas the controls didn't.  Post-op stay was longer in the control group</p>
Sueppel, 2001 Descriptive quasi-experimental	<p>Frequency: Daily Intensity: Low Timing: Several weeks prior and night before surgery  Type: Pelvic floor exercises. Pelvic floor muscle strengthening exercises pre-op with biofeedback confirmation of correct performance, then continue PMEs  Comparator: PMEs via video, then first biofeedback at 6 weeks video for pelvic floor muscle exercises, written instructions, nurse verbal support</p>	<p>16 men pre pre-radical prostatectomy 8 men per group  Prostate cancer</p>	<p>Assessment at 3,6,9,12 months</p>	<p>Not reported</p>	<p>Analysis descriptive no statistical evidence, study underpowered and not controlled. Poorly reported study.  Strengths: consistency of biofeedback instruction by same nurse Limitations: missing data, small sample size; don't know prior voiding patterns of patients  Overall some pre-op information while all other studies have been done post-op</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

--	--	--	--	--	--

For Peer Review

Table 4. Prehabilitation RCT studies with statistically significant outcomes at 30 Days post treatment

Prehabilitation RCT studies with statistically significant outcomes at 30 Days post treatment																							
Ref	Tumour type	Physical functioning						Nutrition	Patient reported outcome				Service benefit										
		Objective					Perceived																
		Gait: 6MWT	V02 Peak	Grip Strength	FEV1*	Pad usage*	CHAMP*	SF36 Physical function	SF36 Role scales and social function	BMI	Self report diet	SF36 mental component	Pain scores*	HAD*	STA	POMS	FACT-L	ICS male SF*	LOS	Less hospitalisation	Pt satisfaction	Fewer complications	
Bales 2000	Prostate cancer					NS																	
Barlesi 2008	NSCLC																					NS	
Burgio 2006	Prostate cancer					+ve																	
Carli 2010	Colorectal cancer	NS	NS																				
Centemero 2010	Prostate cancer					+ve												+ve					
Cheville 2015	GI and gall bladder																			+ve			
Garssen 2013	Breast cancer														NS	+ve					NS		
Gillis 2014	Colorectal cancer	+ve	NS																NS	NS		NS	
Gillis 2015	Colorectal cancer	NS	NS				NS	NS			NS												
Jensen 2014	Bladder cancer																					NS	
Jensen 2015	Bladder cancer																		NS				NS
Moriya 2015	Colorectal cancer																		NS				+ve
Parker 2009	Prostate cancer							+ve			NS												
Schmidt 2015	GI, GU and thoracic cancers												+ve						NS				NS
Stefanelli 2013	NSCLC with COPD		+ve	NS															NS				NS
VanBokhorst 2000	Head and Neck																						

\* Measures targeted to specific diseases or health problems outcomes (+ve) significance >p0.005 (NS)not significant