



# Prehabilitation for esophagectomy

Emre Doganay, Krishna Moorthy

Department of Surgery and Cancer, St Mary's Hospital, Imperial College Healthcare NHS Trust, London, UK

*Contributions:* (I) Conception and design: All authors; (II) Administrative support: All authors; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval: All authors.

*Correspondence to:* Krishna Moorthy. 10<sup>th</sup> Floor, QEOM Wing, St Mary's Hospital, Praed St, London, W2 1NY, UK. Email: k.moorthy@imperial.ac.uk.

**Abstract:** Esophagectomy remains the mainstay treatment of esophageal cancer (EC). Combined with neoadjuvant therapies, the management of EC has deleterious effects on body composition, functional capacity and psychological well-being. Preoperative patient optimisation known as prehabilitation is a novel intervention aimed at reducing morbidity and mortality associated with the trajectory of EC care. There is emerging evidence to suggest that prehabilitation is safe, feasible and efficacious. In addition, there is strong data to infer that prehabilitation has a positive effect on functional capacity through exercise. Nutritional and psychological interventions are less well evaluated. Furthermore, no convincing relationship between prehabilitation and oncological outcomes has been demonstrated. Early studies evaluating prehabilitation are promising however further large scale research is required in order to assess the clinical effectiveness.

**Keywords:** Prehabilitation; exercise; nutrition; preoperative optimization

Submitted Jan 11, 2019. Accepted for publication Feb 12, 2019.

doi: 10.21037/jtd.2019.02.12

View this article at: <http://dx.doi.org/10.21037/jtd.2019.02.12>

## Introduction

Esophagectomy retains an important role in the management of locally advanced esophageal cancer (EC) and remains the mainstay of potentially curative treatment. EC surgery carries significant morbidity and mortality, with 90-day mortality at 3.2% and 1-year survival rates varying between 76% and 78% with morbidity between 20% and 70% (1).

The association between esophagectomy and mortality/morbidity is multifactorial. EC surgery represents complex operations associated with significant weight loss and malnutrition. The implementation of pre-operative chemotherapy and combination chemoradiotherapy, in addition to surgery, has come at a cost on body composition and functional status (2). In addition, patients with EC are typically 'high-risk' because they tend to be frail and elderly. These patients are also susceptible to adverse outcomes due to diminished physiological reserve as a direct consequence of cancer-associated malnutrition (cachexia) (3). As a result patients are at high risk for pulmonary, cardiac and infective

complications as well as prolonged hospital stay associated with increased surgery costs and poor postoperative quality of life (QOL) (4).

The incidence of EC increases with age which means that a higher proportion of elderly patients are being diagnosed with EC and are being considered for curative multimodal therapy. Current statistics show that 30% of patients are over the age of 75 at the time of diagnosis, with a median age of 68–70 (5). This is where the challenge arises as ageing is accompanied with frailty, comorbidities, polypharmacy and invariably a reduction in functional reserve (6). The basic tenet of frailty is that it confers vulnerability to stress, higher risk of treatment toxicities and complications. The perioperative period is when patients are at most risk of developing cardiopulmonary complications with an operative mortality rate of 13.4% in those who are between 70–79 years old and 19.9% in those over the age of 80 (7,8).

Frailty is multi-faceted and accompanied with a deterioration across several physiological domains. Fried *et al.* proposed that frailty is present in the presence of

low grip strength, low energy, slowed walking speed, low physical activity, and/or unintentional weight loss. Some, or all of these are common findings in patients diagnosed with EC and exacerbated by neoadjuvant therapies and/or surgery (9).

There is strong evidence to suggest that baseline frailty infers a greater risk of postoperative complications especially in those with severe frailty (moderate frailty: OR =2.06, 95% CI: 1.18–3.6; severe frailty: OR =2.54, 95% CI: 1.12–5.77). Furthermore, 65–89% of patients with severe frailty had a protracted hospital admissions compared to 44–53% of patients with moderate frailty (10). It is plausible that this is a reflection of decline in cardiorespiratory fitness which accompanies age and previously shown to be linked with postsurgical mortality and morbidity. When clinical outcomes were evaluated, Snowden *et al.* demonstrated that patients over the age of 75 with concomitant poor cardiorespiratory fitness (as reflected by a low anaerobic threshold) spent a median of 11 days longer in hospital (23 *vs.* 12;  $P < 0.0001$ ) and 2 days longer in critical care (2.9 *vs.* 0.9;  $P < 0.00001$ ) (11).

A cause for concern in elderly frail patients with EC is malnutrition. Not only does inadequate nutrition adversely affect QOL it also reduces response to treatment and therefore impacts survival. A reduction in weight of more than 15% off baseline was shown to correlate with higher rates of mortality and morbidity when compared to patients with less weight loss (62% *vs.* 38% respectively). In addition, poor baseline nutrition was a presage of early death in elderly patients who underwent neoadjuvant therapies (12).

The aetiology of malnutrition stems from a combination of either inadequate intake, increased metabolic demands or inflammatory dysregulation that alter nutrient utilization resulting in cachexia and manifesting as a decline in physical fitness and reduced metabolic reserve (13). Cachexia is a pervasive feature of EC resulting in weight loss by reducing both lean and fat mass. EC is among the diseases with the highest known association with cachexia. Luminal obstruction from tumor incursion, toxicity associated with neoadjuvant therapies, and surgical resection, further exacerbates malnutrition (14,15). A recognised phenotypic feature of cachexia is sarcopenia. Sarcopenia is a multifactorial syndrome linked with loss of functional performance and characterized by loss of muscle mass with either low strength or low performance (16). Up to 75% of patients with EC are sarcopenic at diagnosis with associated dose-limiting toxicity (DLT) during neoadjuvant

treatments, disease progression and adverse postoperative outcomes (17). Sarcopenic patients on average have been shown to have a reduced overall survival compared to non-sarcopenic patients (8 *vs.* 26 months). The body composition of cancer patients is highly variable with respect to features of muscle and fat mass as well as the distribution of fat in the abdominal and subcutaneous regions. As a result, sarcopenia should not be limited to patients who outwardly appear thin, and can be found in patients who are overweight or obese; a common finding in patients with adenocarcinoma of the oesophagus (18).

The implementation of neoadjuvant therapies has led to additional oncological and survival benefits, however these have come at a cost to functional capacity when it is needed the most. The relationship between neoadjuvant therapy and their detrimental effects on physiologic and functional capacity is well recognized. Neoadjuvant therapy has been shown to reduce skeletal muscle mass and strength, resulting in a significant increase in the prevalence of sarcopenia from 16% at diagnosis to 31% post-treatment, with an accompanying deterioration in performance status (19). Further studies support this showing that an increase in the prevalence of sarcopenia in patients undergoing neoadjuvant chemotherapy (NAC) has adverse effects on parameters of body composition, such as lean body mass and fat mass, which contribute to morbidity. Body composition, in particular skeletal muscle depletion, has been associated with excess toxicity during neoadjuvant therapy with a recent study demonstrating that DLT was present in 41.6% of patients undergoing NAC, and that sarcopenia was a key contributor to this (20–22).

The risk of toxicity associated with NAC directly influences surgical morbidity. Systemic toxicity results from proteolysis leading to skeletal muscle wasting, oxidative stress and mitochondrial death. Following major surgery, oxygen consumption increases and greater physiological reserve is demanded to allow patients to better withstand the metabolic burden (23). Elderly frail patients tend to have a low baseline level of fitness and in the context of NAC, Jack *et al.* demonstrated reduced one-year survival (3). Physical fitness has been shown to be an important determinant of perioperative outcomes, with less fit patients having higher incidences of morbidity and mortality. Inadequate physical performance is a reflection of reduced exercise capacity, and one method of measuring this is by the  $VO_2$  max (mL/kg/min), which is the measure of a person's individual aerobic capacity. It has been postulated that this measure predicts morbidity, and that increased complications are associated

with a  $\text{VO}_2$  max  $<16$  mL/kg/min (24). The stress of NAC and surgery are important contributors to a reduction in aerobic capacity and therefore poor pre-operative physical performance is linked with all-cause mortality, post-operative complications, length of hospital stay and hospital costs. It is becoming clearer that neoadjuvant treatments are causing harm to physical fitness, which in turn translates into adverse clinical outcomes (25).

Anxiety and depression are not uncommon findings in patients with newly diagnosed EC and inadvertently influences functional capacity during the pre- and postoperative periods due to low adherence. In addition, there is a significant deterioration in postoperative QOL which impacts recovery following surgery (23).

This has led to the introduction of multimodal interventions in the period preceding surgery aimed at optimizing patients and improving post-operative outcomes in EC resectional surgery.

### Prehabilitation

In recent years there has been a shift towards the centralization of cancer surgery, adoption of minimally invasive techniques and widespread uptake of multidisciplinary perioperative care programmes such as enhanced recovery after surgery (ERAS). In combination, these have led to substantial improvements in post-operative outcomes.

Currently ERAS is a well-established component in the trajectory of surgical care but place little emphasis on pre-operative patient optimization. Despite the implementation of ERAS, complications following elective major abdominal surgery are still between 25–55% (23,25).

Prehabilitation is the process of providing patients with a reserve to withstand the stress of major cancer surgery. The principle is to extend beyond just pre-operative physiological and co-morbidity optimization to improvement in functional capacity, nutritional status and psychological well-being in preparation for major surgery (26).

### Presurgical exercise

With frailty and ill-health comes a lack of physical activity which can exacerbate perioperative morbidity (27). Exercise as part of a prehabilitation programme has been proposed to counteract functional decline and enhance patient performance as well as attenuating sarcopenia and limiting deconditioning associated with disease burden (28). Traditionally, the notion of bed rest was advocated in

anticipation for surgery but we now know that sedentary behaviour is associated with loss of lean muscle mass, reduced physical function and aerobic capacity and insulin resistance (29).

This is why emphasis must be placed on aerobic and muscular training to increase endurance, manage weight and build strength. Research highlights that in order to achieve this 3 weeks of exercise may be adequate (30). Several studies on prehabilitation in patients undergoing thoracic and gastrointestinal (GI) cancer resection have demonstrated an increase in preoperative physical fitness, physical activity and decreased postoperative complications with shorter hospital stay (31). In addition, feasibility and safety following neoadjuvant therapies, as well as improvements in long-term physical activity and QOL have also been demonstrated (30,31). Exercise includes consistent physical activity delivered through a structured programme to improve physical fitness. For exercise to be effective it needs to be prescribed at a certain 'dose' and adapted to meet the needs and requirements of the patient in order to achieve the desired outcomes (23,32). The prescription and dose of exercise requires consideration of frequency, intensity, time and type (FITT) of exercise. Although a lower dose of physical activity has clear health benefits, higher doses of exercise will result in greater improvements (33). Several pilot studies in rectal and breast cancer have shown improvements in fitness after supervised interval training and several reviews on exercise as part of a prehabilitation programme have demonstrated similar beneficial effects in patients undergoing major abdominal surgery (34–36). In a review looking at prehabilitation and postoperative clinical outcomes, Moran *et al.* established that aerobic and resistance training reduced the occurrence of complications in patients undergoing abdominal surgery (37). In support of this, one study showed that prehabilitation reduced the incidence of postoperative pulmonary complications [OR 0.27, 95% confidence interval (CI): 0.13–0.57] (13). Several studies have highlighted the positive impact of exercise on measures of functional capacity. West *et al.* reported that  $\text{VO}_2$  increased by an average of  $2.1$  mL·kg<sup>-1</sup>·min<sup>-1</sup> after 6 weeks of exercise ( $P<0.001$ ) and further studies have supported this showing a significant increase in  $\text{VO}_2$  and peak  $\text{VO}_2$  ( $\text{pVO}_2$ ) within exercise groups (38). In addition, several studies looking at postoperative outcomes showed that preoperative exercise reduced the length of hospital stay. Cho *et al.* reported a reduction in length of admission (9 *vs.* 10 days;  $P=0.038$ ) as well as reduced intra-abdominal postoperative complications

(OR 0.12, 95% CI: 0.00–0.89, P=0.033) (39). In a review by Vermillion *et al.*, patients who underwent preoperative exercise training were more likely to return to their baseline fitness after surgical resection. In addition, a combination of aerobic exercise training with resistance training was effective in reversing sarcopenia in elderly individuals with a reduction in postoperative pulmonary complications. As well as enhancing cardiopulmonary fitness, exercise was also proposed to reduce fatigue and have desirable outcomes on QOL (28).

## Nutrition

Malnourished surgical patients are known to have higher postoperative morbidity and mortality, and the risk of malnutrition is most severe immediately before surgery (13). EC patients are chronically malnourished and require nutritional prehabilitation in order to replenish nutrient stores and build a metabolic reserve to buffer the catabolic response to cancer, neoadjuvant therapies and surgery. Early patient engagement is paramount and tailored nutritional goals should be delivered in a pre-emptive rather than reactive method (40). A well-recognized phenomenon in critical illness and surgery is protein catabolism. Protein requirements are elevated in stressed states and therefore it is of paramount importance that provisions of protein are addressed as part of nutritional prehabilitation to maintain lean muscle mass and reduce subsequent risk of sarcopenia and frailty (41). The importance of perioperative nutrition and its influence on prognosis was recently discussed in the European Society for Clinical Nutrition and Metabolism (ESPEN) guidelines. Patients who were nutritionally deplete and had significant weight loss were more likely to have a protracted hospital admission, higher readmission rates and experience morbidity/mortality (42). It is essential that cachectic individuals are identified via early pre-assessment by a specialist dietitian and receive tailored, monitored and ongoing nutritional intervention up to the day of surgery. RCTs and meta-analyses have supported this and have shown that preoperative nutrition in malnourished patients prior to major abdominal surgery reduced postoperative morbidity by 20% (43). Diets aimed at enhancing immune function are called immunonutrition (IN). Several meta-analyses have been published on the clinical effectiveness of preoperative IN showing it to reduce the incidence of postoperative infectious complications and length of hospitalization (44). This is true of patients undergoing surgery for gastrointestinal

malignancy and has been advocated as part of preoperative nutritional optimization. In the context of esophageal surgery however, results are heterogeneous. Current evidence is insufficient to recommend enteral IN in patients undergoing esophagectomy and/or chemoradiotherapy for EC (45).

## Psychological

In the context of major life altering surgery, anxiety and depression is common, and has a negative impact on postoperative pain control, leads to prolonged hospital admission/readmissions and causes functional limitations with poor compliance to exercise (46). Prehabilitation encompasses psychological distress, and addressing anxiety and depression as part of this programme will have a positive influence on adhering to prescribed exercise and in turn promoting psychological well-being.

A number of trials have shown prehabilitation to improve QOL, reduce anxiety and depression as well as pain severity and fatigue (23).

In addition, prehabilitation has been shown to improve functional behaviors such as the ability to perform ambulation and rehabilitation tasks which are key for postoperative recovery. This also has a positive effect on subsequent QOL and the psychological wellbeing of the patient. There is new evidence that patients with distress and low self-confidence experience poorer QOL trajectories in the recovery period (47). Thus interventions to address distress and improve self-efficacy are likely to improve post-operative recovery

## Discussion

There is substantial data which highlights the clear benefits of preoperative exercise and nutritional optimization, at the same time recognizing that psychological wellbeing is a critical aspect of perioperative care. However, the majority of data published on prehabilitation in GI cancer comes from colorectal cancer surgery and at present evidence that prehabilitation works in EC surgery is limited. Soares *et al.* investigated the effects of preoperative physical therapy on pulmonary function and physical performance before and after upper abdominal surgery. In a RCT of 32 participants the authors demonstrated that patients who received physical therapy had higher inspiratory strength and respiratory muscle endurance. Furthermore, patients who were randomly assigned to physical therapy achieved better results in the 6-minute walk distance

(6MWD) than compared to control group. There was also a significant difference between postoperative pulmonary complications with the intervention group being less likely to develop a pulmonary complication ( $P=0.03$ ) (48). In a similar RCT Xu *et al.* observed the effect of a 'walk-and-eat' intervention during neoadjuvant chemoradiotherapy on functional walking capacity and nutritional status. The intervention consisted of weekly supervised walking and nutrition advice. The study demonstrated that the intervention group had significantly less decline in 6MWD during neoadjuvant therapy. Similarly, there was less decline in hand-grip strength, weight and lean muscle mass. There was no difference observed in outcomes (49). More recently Minnella *et al.* looked at the effect of prehabilitation on functional status in patients undergoing EC resection. The authors showed that patients who underwent prehabilitation had improved functional capacity (absolute change in 6MWD) both before surgery and after surgery. However, there was no statistical difference between groups in terms of number and severity of complications, length of hospital stay, and readmission rates (50). A plausible explanation for this is the study being underpowered to detect an association between physical fitness and complications. Although Soares *et al.* did show a reduction in postoperative pulmonary complications, the study was also underpowered.

In line with recently published studies there are a number of ongoing trials centred around prehabilitation in EC. The PREPARE for surgery prehabilitation programme is an ongoing quality improvement project which delivers a personalized, home-based programme consisting of preoperative exercise, nutritional optimization and addressing psychological domains such as anxiety and depression as well as key aspects of QOL. Current data (unpublished) from the PREPARE programme has demonstrated a significant improvement in functional capacity as assessed by  $VO_2$ ,  $mL \cdot kg^{-1} \cdot min^{-1}$  as well as reduced postoperative complications including a reduction in Clavien-Dindo score of  $>2$  (80% to 35%), postoperative pneumonia (70% to 29%) and median length of hospital stay (14 to 11) (26). Further details of trials on prehabilitation in EC can be found at <http://clinicaltrials.gov>. Some of these studies are still in the recruitment process, ongoing or completed and awaiting publication. Majority are RCTs and the focus lies with determining the impact of multimodal prehabilitation on functional exercise capacity, functional recovery both during and after neoadjuvant therapies and following surgery. Some of the studies are also aiming to address key important areas such

as postoperative complications which is currently lacking in the literature. This will be eagerly anticipated as the evidence for the relationship between prehabilitation and postoperative outcomes is limited and where research is available it is not adequately powered.

## Conclusions

Although a novel area of research in EC there is clear evidence that prehabilitation improves functional capacity and supports the patient through nutritional and psychological counselling. However further work is required to determine its effects on overall oncologic outcomes.

## Acknowledgements

Venetia Wynter-Blyth—clinical lead of the PREPARE programme. Laura Halliday, Maria Halley, Hayley Osborn, Alex King and Claudia Rueb—members of the PREPARE programme.

## Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

## References

1. National Oesophago-Gastric Cancer Audit [Internet]. National Oesophago-Gastric Cancer Audit 2018 [cited 26 October 2018]. Available online: <https://www.nogca.org.uk/>
2. Jack S, West MA, Raw D, et al. The effect of neoadjuvant chemotherapy on physical fitness and survival in patients undergoing oesophagogastric cancer surgery. *Eur J Surg Oncol* 2014;40:1313-20.
3. Jack S, West M, Grocott MP. Perioperative exercise training in elderly subjects. *Best Pract Res Clin Anaesthesiol* 2011;25:461-72.
4. Khuri SF, Henderson WG, DePalma RG, et al. Determinants of Long-Term Survival After Major Surgery and the Adverse Effect of Postoperative Complications. *Ann Surg* 2005;242:326-41; discussion 341-3.
5. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin* 2015;65:5-29.
6. Xue QL. The Frailty Syndrome: Definition and Natural History. *Clin Geriatr Med* 2011;27:1-15.
7. Baijal P, Periyakoil V. Understanding Frailty in Cancer Patients. *Cancer J* 2014;20:358-66.

8. Finlayson E, Fan Z, Birkmeyer JD. Outcomes in Octogenarians Undergoing High-Risk Cancer Operation: A National Study. *J Am Coll Surg* 2007;205:729-34.
9. Fried LP, Tangen CM, Watson J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146-56.
10. Makary MA, Segev DL, Pronovost PJ, et al. Frailty as a Predictor of Surgical Outcomes in Older Patients. *J Am Coll Surg* 2010;210:901-8.
11. Snowden CP, Prentis J, Jacques B, et al. Cardiorespiratory Fitness Predicts Mortality and Hospital Length of Stay After Major Elective Surgery in Older People. *Ann Surg* 2013;257:999-1004.
12. Won E. Issues in the management of esophageal cancer and geriatric patients. *Chin Clin Oncol* 2017;6:51.
13. West MA, Wischmeyer PE, Grocott MPW. Prehabilitation and Nutritional Support to Improve Perioperative Outcomes. *Curr Anesthesiol Rep* 2017;7:340-9.
14. Anandavivelan P, Lagergren P. Cachexia in patients with oesophageal cancer. *Nat Rev Clin Oncol* 2016;13:185-98.
15. Evans WJ, Morley JE, Argilés J, et al. Cachexia: A new definition. *Clin Nutr* 2008;27:793-9.
16. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019;48:16-31.
17. Nishigori T, Tsunoda S, Obama K, et al. Optimal Cutoff Values of Skeletal Muscle Index to Define Sarcopenia for Prediction of Survival in Patients with Advanced Gastric Cancer. *Ann Surg Oncol* 2018;25:3596-603.
18. Martin L, Birdsell L, MacDonald N, et al. Cancer Cachexia in the Age of Obesity: Skeletal Muscle Depletion Is a Powerful Prognostic Factor, Independent of Body Mass Index. *J Clin Oncol* 2013;31:1539-47.
19. Elliott JA, Doyle SL, Murphy CF, et al. Sarcopenia: Prevalence, and Impact on Operative and Oncologic Outcomes in the Multimodal Management of Locally Advanced Esophageal Cancer. *Ann Surg* 2017;266:822-30.
20. Awad S, Tan BH, Cui H, et al. Marked changes in body composition following neoadjuvant chemotherapy for oesophagogastric cancer. *Clin Nutr* 2012;31:74-7.
21. Paireder M, Asari R, Kristo I, et al. Impact of sarcopenia on outcome in patients with esophageal resection following neoadjuvant chemotherapy for esophageal cancer. *Eur J Surg Oncol* 2017;43:478-84.
22. Tan BH, Brammer K, Randhawa N, et al. Sarcopenia is associated with toxicity in patients undergoing neoadjuvant chemotherapy for oesophago-gastric cancer. *Eur J Surg Oncol* 2015;41:333-8.
23. Carli F, Gillis C, Scheede-Bergdahl C. Promoting a culture of prehabilitation for the surgical cancer patient. *Acta Oncologica* 2017;56:128-33.
24. McCullough PA, Gallagher MJ, Dejong AT. Cardiorespiratory Fitness and Short-term Complications After Bariatric Surgery. *Chest* 2006;130:517-25.
25. Wilson RJ, Davies S, Yates D, et al. Impaired functional capacity is associated with all-cause mortality after major elective intra-abdominal surgery. *Br J Anaesth* 2010;105:297-303.
26. Wynter-Blyth V, Moorthy K. Prehabilitation: preparing patients for surgery. *BMJ* 2017;358:j3702.
27. Boyce T, Robertson R, Dixon A. Commissioning and Behaviour Change [Internet]. The King's Fund 2019 [cited 9 January 2019]. Available online: <https://www.kingsfund.org.uk/publications/commissioning-and-behaviour-change>
28. Vermillion SA, James A, Dorrell RD, et al. Preoperative exercise therapy for gastrointestinal cancer patients: a systematic review. *Syst Rev* 2018;7:103.
29. Coker RH, Hays NP, Williams RH, et al. Bed Rest Promotes Reductions in Walking Speed, Functional Parameters, and Aerobic Fitness in Older, Healthy Adults. *J Gerontol A Biol Sci Med Sci* 2015;70:91-6.
30. Fearon KC, Jenkins JT, Carli F, et al. Patient optimization for gastrointestinal cancer surgery. *Br J Surg* 2013;100:15-27.
31. Valkenet K, van de Port IG, Dronkers JJ, et al. The effects of preoperative exercise therapy on postoperative outcome: a systematic review. *Clin Rehabil* 2011;25:99-111.
32. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126-31.
33. Haskell WL. Health consequences of physical activity: understanding and challenges regarding dose-response. *Med Sci Sports Exerc* 1994;26:649-60.
34. Westphal T, Rinnerthaler G, Gampenrieder SP, et al. Supervised versus autonomous exercise training in breast cancer patients: A multicenter randomized clinical trial. *Cancer Med* 2018;7:5962-72.
35. Awasthi R, Minnella EM, Ferreira V, et al. Supervised exercise training with multimodal pre-habilitation leads to earlier functional recovery following colorectal cancer resection. *Acta Anaesthesiol Scand* 2019;63:461-7.
36. O'Doherty AF, West M, Jack S, et al. Preoperative aerobic exercise training in elective intra-cavity surgery: a

- systematic review. *Br J Anaesth* 2013;110:679-89.
37. Moran J, Guinan E, McCormick P, et al. The ability of prehabilitation to influence postoperative outcome after intra-abdominal operation: A systematic review and meta-analysis. *Surgery* 2016;160:1189-201.
  38. West MA, Loughney L, Lythgoe D, et al. Effect of prehabilitation on objectively measured physical fitness after neoadjuvant treatment in preoperative rectal cancer patients: a blinded interventional pilot study. *Br J Anaesth* 2015;114:244-51.
  39. Cho H, Yoshikawa T, Oba MS, et al. Matched pair analysis to examine the effects of a planned preoperative exercise program in early gastric cancer patients with metabolic syndrome to reduce operative risk: the Adjuvant Exercise for General Elective Surgery (AEGES) study group. *Ann Surg Oncol* 2014;21:2044-50.
  40. Thomas MN, Kufeldt J, Kissler U, et al. Effects of malnutrition on complication rates, length of hospital stay, and revenue in elective surgical patients in the G-DRG-system. *Nutrition* 2016;32:249-54.
  41. Ferrando AA, Paddon-Jones D, Hays NP, et al. EAA supplementation to increase nitrogen intake improves muscle function during bed rest in the elderly. *Clin Nutr* 2010;29:18-23.
  42. Arends J, Bachmann P, Baracos V, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr* 2017;36:11-48.
  43. Osland E, Yunus RM, Khan S, et al. Early versus traditional postoperative feeding in patients undergoing resectional gastrointestinal surgery: a meta-analysis. *JPEN J Parenter Enteral Nutr* 2011;35:473-87.
  44. Zhang Y, Gu Y, Guo T, et al. Perioperative immunonutrition for gastrointestinal cancer: A systematic review of randomized controlled trials. *Surg Oncol* 2012;21:e87-e95.
  45. Mimatsu K, Fukino N, Ogasawara Y, et al. Effects of Enteral Immunonutrition in Esophageal Cancer. *Gastrointest Tumors* 2018;4:61-71.
  46. Rosenberger PH, Jokl P, Ickovics J. Psychosocial Factors and Surgical Outcomes: An Evidence-Based Literature Review. *J Am Acad Orthop Surg* 2006;14:397-405.
  47. Foster C, Haviland J, Winter J, et al. Pre-Surgery Depression and Confidence to Manage Problems Predict Recovery Trajectories of Health and Wellbeing in the First Two Years following Colorectal Cancer: Results from the CREW Cohort Study. *PloS One* 2016;11:e0155434.
  48. Soares SM, Nucci LB, da Silva MM, et al. Pulmonary function and physical performance outcomes with preoperative physical therapy in upper abdominal surgery: a randomized controlled trial. *Clin Rehabil* 2013;27:616-27.
  49. Xu YJ, Cheng JC, Lee JM, et al. A Walk-and-Eat Intervention Improves Outcomes for Patients With Esophageal Cancer Undergoing Neoadjuvant Chemoradiotherapy. *Oncologist* 2015;20:1216-22.
  50. Minnella EM, Awasthi R, Loiselle SE, et al. Effect of Exercise and Nutrition Prehabilitation on Functional Capacity in Esophagogastric Cancer Surgery. *JAMA Surg* 2018. [Epub ahead of print].

**Cite this article as:** Doganay E, Moorthy K. Prehabilitation for esophagectomy. *J Thorac Dis* 2019;11(Suppl 5):S632-S638. doi: 10.21037/jtd.2019.02.12