

Full Length Research Paper

Preliminary screening of plant essential oils against larvae of *Culex quinquefasciatus* Say (Diptera: Culicidae)

M. Ramar^{1*}, M. Gabriel Paulraj² and S. Ignacimuthu²

¹Department of Zoology, APSA, College, Thiruppattur, Tamil Nadu, India- 626141.

²Entomology Research Institute, Loyola College, Chennai, Tamil Nadu, India -600034.

Accepted 27 August, 2013

Preliminary screenings of 22 plant essential oils were tested for mortality of the mosquito larvae *Culex quinquefasciatus* under laboratory conditions. Percent (%) mortality of the mosquito larvae were obtained for each essential oil. At different exposure periods, viz. 1, 3, 6, 12 and 24 h among the 22 plant oils tested, eight oils viz., aniseed, calamus, cinnamon, clove, lemon, orange, thyme, and tulsi oils gave promising results on larvicidal activity. For larvicidal screening bioassay, the mortality was recorded at different exposure periods viz., 1, 3, 6, 12 and 24 h, and it was found that larval mortality increased when exposure time increased. The clove oil was found to be the most effective treatment. In the preliminary screening, clove oil gave 100% mortality at all exposure periods. Vetiver oil recorded 36.2, 61.2, 76.2, 87.5 and 100% mortality in 1, 3, 6, 12 and 24 h, respectively. Results of this study show that the essential oils may be a potent source of natural larvicides.

Key words: Screening, essential oils, *Culex quinquefasciatus*, Larvicidal.

INTRODUCTION

Culex quinquefasciatus, a domestic mosquito mainly found in urban areas is a vector of human filariasis in India. *C. quinquefasciatus* acts as a vector of *Wuchereria bancrofti*, *Brugia malayi* and *Brugia timori*, which are responsible for lymphatic filariasis, a prevalent disease in India. Filariasis is an endemic disease in many parts of India especially in Kerala, Mysore, Tamil Nadu, Andhra Pradesh and Maharashtra states. Lymphatic filariasis infects 80 million people annually, of which 30 million cases exist in chronic infection. In India, 45 million cases of lymphatic filariasis have been recorded (Bowers et al., 1995).

The plant world comprises a rich storehouse of phytochemicals which are widely used as alternatives to synthetic insecticides. Secondary metabolites obtained from plants have shown proven mosquito control poten-

tial. *Chloroxylons wietenia* DC. (Rutaceae) is a medicinal as well as an aromatic tree of dry deciduous forests. It is commonly known as East Indian Satin Wood. The potential mosquitocidal activity of *C. wietenia*, is widely known in tribal areas of South India, where it is common for people in these forest areas to hang leaf garlands in their houses to eradicate mosquitoes and other insects (Ansari and Razdan, 1995).

Pyrethrin based products have been widely used to protect people from mosquito bites through their repellent and killing effects. Many other material products of botanical origin especially, essential oils hold significant promise in insect vector management (Skinner and Johnson, 1980). Kiso and Hikino (1991) have reported that the compounds vinblastine and vincristine obtained from *Catharanthus roseus* inhibited cell division in

*Corresponding author. Email:ramareri@gmail.com.

Table 1. Plant essential oils selected for the bioassay experiments.

| Botanical name | Family | Common name |
|------------------------------------|-------------|-------------|
| <i>Pimpinella anisum</i> | Apiaceae | Aniseed |
| <i>Citrus bergamia</i> | Rutaceae | Bergamot |
| <i>Acorus calamus</i> | Aeraceae | Calamus |
| <i>Cinnamomum camphora</i> | Lauraceae | Camphor |
| <i>Cedrus atlantica</i> | Pinaceae | Cedarwood |
| <i>Cinnamomum veerum</i> | Lauraceae | Cinnamon |
| <i>Cymbopogon nardus</i> | Poaceae | Citronella |
| <i>Myrtus caryophyllus</i> | Myrtaceae | Clove |
| <i>Eucalyptus globules</i> | Myrtaceae | Eucalyptus |
| <i>Pelargonium graveollens</i> | Geraniaceae | Geranium |
| <i>Citrus limon</i> | Rutaceae | Lemon |
| <i>Cymbopogon flexuosus</i> | Poaceae | Lemongrass |
| <i>Citrus aurantifolia</i> | Rutaceae | Lime |
| <i>Gaultheria frgagrantiissima</i> | Acanthaceae | Luchi |
| <i>Myrtica fragrans</i> | Myristiceae | Nutmeg |
| <i>Citrus sinensis</i> | Rutaceae | Orange |
| <i>Cymbopogon martini</i> | Lamiaceae | Palmarosa |
| <i>Pinus radiata</i> | Pinaceae | Pine |
| <i>Rosmarinus officinalis</i> | Lamiaceae | Rose mary |
| <i>Thymus vulgaris</i> | Labiatae | Thyme |
| <i>Ocimum sanctum</i> | Lamiaceae | Tulsi |
| <i>Vetiveria zizanioides</i> | Poaceae | Vetiver |

fertilized mosquito eggs. Tulsi (*Ocimum sanctum*) is known to have various insectidal properties (Saxena et al., 1979). Extracts of *Vitex* species and *Ocimum* species leaves have been traditionally known as natural insecticides. Indian farmers have been using various plant extracts as insecticides and pesticides (Ghosh, 2000).

Many studies have shown the bioactivity of essential oils against mosquitoes as growth inhibitors and/or larvicides, adulticides, repellents, or oviposition deterrents (Sukumar et al., 1991; Carvalho et al., 2003; Cavalcanti et al., 2004; Ansari et al., 2005). Essential oils from a large number of plants, including *Ocimum* spp, *Cymbopogon* spp), *Eucalyptus maculate citriodon* *Pelargonium citrosum*, *Artemisia vulgaris*, *Lantana camara*, *Mentha piperita*, *Vitex rotundifolia*, *Curcuma* spp. (Pitasawat et al., 2003), *Conyza newii*, *Plectranthus marrubioides*, *Tetradenia riparia*, *Tarhchonanthus camphoratus*, *Lippia javanica* and *Lippia ukambensis* (Omolo et al., 2004), have been demonstrated to exhibit good repellent properties against vector mosquitoes. Paula et al. (2003) reported the chemical composition, toxicity and mosquito repellency of *Ocimum selloi* oil. *Ocimum* essential oils have been traditionally used to kill or repel insects, and also to flavour food products.

In the present investigation, the essential oils were selected in order to find out a new mosquitocidal compound against filarial vector mosquito *C.*

quinquefasciatus. The plant essential oil is largely cultivated throughout India and in all tropical countries.

MATERIALS AND METHODS

Plant essential oils

The plant essential oils were obtained from Tegraj and Co (P) Ltd, Chennai, Tamil Nadu. Based on various biological effects on larvae, the 22 essential oils were selected (Table 1).

Maintenance of larvae

The larvae of laboratory stock culture were reared in plastic troughs (size 18 cm diameter x 19 cm height) and were maintained at 27 ± 1°C, 75 to 85% RH, under 14:10 L/D photoperiod cycles. The larvae were fed with dog biscuits and yeast at 3:1 ratio (per weight). Water was changed alternate days. The breeding medium was regularly checked and dead larvae were removed at sight and the troughs in which larvae were maintained were kept closed with muslin cloth to prevent contamination through foreign mosquitoes.

Screening assay

The selected plant oils were screened for larval toxicity by following standard procedure as follows: In the preliminary screening a single dose (500 ppm) of each oil was tested. Twenty fourth instar larvae of *C. quinquefasciatus* were kept in a 500 ml glass beaker containing 249 ml of dechlorinated water and 1.0 ml of desired plant

Table 2. Preliminary screening of plant essential oils against fourth instar larvae of *Culex quinquefasciatus* at 500 ppm in different exposure periods.

| Essential oil | Larval mortality (%) | | | | |
|--------------------------------------|--------------------------|---------------------------|-------------------------------------|--------------------------|---------------------------|
| | 1 h | 3 h | 6 h | 12 h | 24 h |
| Aniseed | 91.2 ± 0.6 ^g | 100 ± 0.0 ^j | 100 ± 0.0 ^m | 100 ± 0.0 ^k | 100 ± 0.0 ^j |
| Bergamot | 11.2 ± 0.2 ^{bc} | 18.7 ± 0.7 ^{bc} | 30.0 ± 0.5 ^{gh} | 45.0 ± 0.4 ^e | 61.2 ± 0.8 ^{efg} |
| Calamus | 21.2 ± 0.2 ^{de} | 32.5 ± 0.2 ^e | 43.7 ± 0.3 ^j | 66.2 ± 0.4 ^f | 100 ± 0.0 ^j |
| Camphor | 0 ^a | 2.25 ± 0.0 ^a | 7.5 ± 0.2 ^{bc} | 20.0 ± 0.9 ^{bc} | 38.7 ± 0.4 ^c |
| Cedarwood | 18.7 ± 0.4 ^c | 30.0 ± 0.2 ^e | 40.0 ± 0.3 ^j | 58.7 ± 0.6 ^f | 68.7 ± 0.8 ^{hi} |
| Cinnamon | 47.5 ± 0.8 ^g | 83.7 ± 0.5 ^a | 100 ± 0.0 ^m | 100 ± 0.0 ^k | 100 ± 0.0 ^j |
| Citronella | 8.75 ± 0.3 ^c | 20.0 ± 0.5 ^{cd} | 35.0 ± 0.4 ^{hi} | 52.5 ± 0.6 ^f | 93.7 ± 0.4 ⁱ |
| Clove | 100 ± 0.0 ^h | 100 ± 0.0 ⁱ | 100 ± 0.0 ^m | 100 ± 0.0 ^k | 100 ± 0.0 ^j |
| Eucalyptus | 0 ^a | 3.7 ± 0.2 ^a | 12.5 ± 0.5 ^{cd} | 26.2 ± 0.4 ^c | 38.8 ± 0.4 ^c |
| Geranium | 0 ^a | 2.5 ± 0.4 ^a | 15.0 ± 0.5 ^{de} | 25.0 ± 0.4 ^c | 36.2 ± 0.7 ^c |
| Lemon | 23.7 ± 0.2 ^c | 40.0 ± 0.5 ^f | 53.7 ± 0.7 ^k | 86.2 ± 0.8 ⁱ | 100 ± 0.0 ^j |
| Lemongrass | 0 ^a | 1.2 ± 0.3 ^a | 3.75 ± 0.4 ^{ab} | 16.2 ± 0.6 ^b | 28.7 ± 0.5 ^b |
| Lime | 13.7 ± 0.4 ^{bc} | 22.5 ± 0.2 ^b | 38.7 ± 0.6 ^{ij} | 57.5 ± 0.6 ^f | 67. ± 0.6 ^g |
| Luchi | 5.0 ± 0.4 ^c | 21.2 ± 0.4 ^d | 33.7 ± 0.7 ^{hi} | 58.75 ± 0.6 ^f | 73.7 ± 1.1 ⁱ |
| Nutmeg | 10 ± 0.8 ^b | 15.0 ± 0.9 ^{bcd} | 30.0 ± 0.9 ^{gh} | 40.0 ± 0.8 ^d | 56.2 ± 0.2 ^{def} |
| Orange | 28.7 ± 0.2 ^{cd} | 43.7 ± 0.8 ^f | 58.7 ± 0.2 ^k | 75.0 ± 0.4 ^h | 100 ± 0.0 ^j |
| Palmarosa | 5.0 ± 0.8 ^{bc} | 12.5 ± 0.9 ^{bcd} | 28.7 ± 0.9 ^{gh} | 37.5 ± 0.8 ^d | 52.5 ± 0.2 ^d |
| Pine | 2.7 ± 0.2 ^{ab} | 11.2 ± 0.2 ^b | 21.2 ± 0.4 ^{ef} | 35.0 ± 0.7 ^d | 55.0 ± 0.7 ^{de} |
| Rose mary | 1.2 ± 0.4 ^b | 13.7 ± 0.7 ^c | 25.0 ± 0.4 ^{f^g} | 36.2 ± 0.7 ^d | 82.5 ± 0.5 ^{fg} |
| Thyme | 42.5 ± 0.2 ^f | 57.5 ± 0.6 ^g | 68.7 ± 0.2 ^l | 80.0 ± 0.4 ^{hi} | 100 ± 0 ^j |
| Tulsi | 47.5 ± 0.2 ^{ef} | 68.7 ± 0.6 ⁱ | 92.5 ± 0.6 ^m | 100 ± 0.0 ^k | 100 ± 0 ^j |
| Vetiver | 36.2 ± 0.4 ^{ef} | 61.2 ± 0.6 ^g | 76.2 ± 0.8 ^m | 87.5 ± 0.6 ^j | 100 ± 0 ^j |
| Reference pesticide (temephos 1 ppm) | 40.0 ± 0.8 ^{ef} | 65.0 ± 0.4 ^{hi} | 100 ± 0.0 ^m | 100 ± 0 ^k | 100 ± 0 ^j |
| Control (Tween 80 0.01%) | 0 ^a | 0 ^a | 0 ^a | 0 ^a | 0 ^a |

Each value (mean ± S.E.) represents mean of four replicates, Mean values followed by same letters in a column are not significantly different at $P < 0.05$ level (Tukey's test).

essential oil concentrations. The larvae were collected by dropper, placed onto filter paper strips and immediately transferred to the beaker containing test solutions. Five replicates for each concentration were maintained. In the control, 1.0 ml of Tween 80 (0.01%) in 249 ml of dechlorinated water was added. Mortality of larvae was recorded after 1, 3, 6, 12 and 24 h of treatment. While recording the percentage mortalities for each concentration, the moribund and dead larvae in five replicates were combined. It has been described that dead larvae are those that cannot be induced to move when they are probed with a needle in the siphon or the cervical region; moribund larvae are those incapable of rising to the surface (Hasan and Deo, 1994).

Statistical analysis

The larvicidal mortality was corrected by Abbott's formula (Abbott, 1925). Percentages of mortality were determined and transformed to arcsine square root values for analysis of variance (ANOVA). Tukey's test ($P < 0.05$) was used to analyze for significant differences among the test essential oils against mosquitoes.

RESULTS AND DISCUSSION

All the 22 plant oils were screened against the larvae of

C. quinquefasciatus for larvicidal activity. The results are given in Table 2. Among the 22 plant oils tested, eight oils viz., aniseed, calamus, cinnamon, clove, lemon, orange, thyme, and tulsi oils gave promising results on larvicidal activity. For the larvicidal screening bioassay, the mortality was recorded at different exposure periods viz., 1, 3, 6, 12 and 24 h, and it was found that larval mortality increased when exposure time increased. The clove oil was found to be the most effective treatment. In the preliminary screening, clove oil gave 100 percent mortality at all exposure periods.

Vetiver oil recorded 36.2, 61.2, 76.2, 87.5 and 100% mortality in 1, 3, 6, 12 and 24 h, respectively. The lowest mortality of 28.7% was recorded at 500 ppm concentration in lemongrass oil after 24 h exposure. In the temephos treatment, larval mortality of 40, 65 and 100% was recorded at 1 ppm concentration in 1, 3 and 6 h treatments.

Vector control by means of chemicals is creating many unwanted effects including emergence of pesticide resistance in vector mosquitoes (Chandra et al., 1998). Botanical insecticides may serve as suitable alternatives

to synthetic insecticides in future as they are relatively safe, biodegradable and are readily available in many areas of the world (Prabakar and Jebanesan, 2004). Interest in the development of natural products has been revived during the last two decades. Plants are considered as rich source of bioactive chemicals and they may be an alternative source of mosquito control agents. Natural products of plant origin with insecticidal properties have been tried in the recent past for control of variety of insect pests and vectors (Wink, 1993). More than 2000 plant species have been reported to possess chemicals with pest control properties (Ahmed et al., 1984) and among them about 344 species of plants have been known to possess some degree of activity against the mosquitoes (Sukumar et al., 1991).

Though several plants from different families have been reported for insecticidal activity only a few botanicals like neem based insecticides have moved from the laboratory to field use, which might be due to the light and heat stability of neem compounds compared to synthetic insecticides (Green et al., 1991). In the present study, 22 essential oils registered larvicidal, pupicidal, adulticidal, ovicidal, oviposition deterrent, growth regulator regulating and histopathological effects. Plenty of literature is available with regard to bioefficacy of volatile oils against vector mosquitoes.

The most successful method to minimize the incidence of mosquito borne diseases is by means of eradicating the mosquito vector through systematic treatment of their breeding places with the help of larvicides (Cetin et al., 2004). Chemical pesticides that are applied in water bodies are not only killing mosquito larvae but also killing non-target organisms. In addition chemical pesticides select for resistance in mosquitoes. As an alternative to chemical pesticides volatile oils are largely studied against *C. quinquefasciatus* mosquito. In the present study, the larvicidal activity was tested at 500 ppm concentration. Among the 22 essential oils, eight oils registered high larvicidal activity. The larval mortality was found to be directly proportional to the exposure period. Clove oil was found to be the most toxic to the larvae since it registered 100% mortality from 1 h treatment period onwards.

The present investigation revealed that the essential oils possess remarkable larvicidal activity against the tested mosquito *C. quinquefasciatus*. The control of selected mosquito by affecting their survival by essential oils was found evident. Further purification and characterization of the bioactive compound of effective oils are underway in our laboratory.

ACKNOWLEDGEMENTS

We are deeply indebted to the Director, Entomology Research Institute, Loyola College, Chennai, for his interest to promote research activities.

REFERENCES

- Abbott WS (1925). A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 18:265-266.
- Ahmed S, Graivge M, Hylin JW, Mitchell WC, Listering JA (1984). Some promising plant species for use as pest control agents under traditional farming system. In: Schmutterer, H. and K.R.S. Ascher (eds.) *Proc. 2nd International Neem Conference*, Rauscholzhausen, Germany. pp. 565-580.
- Ansari MA, Mittal PK, Razdan RK, Sreehari U (2005). Larvicidal and mosquito repellent activities of Pine (*Pinus longifolia*, Family: Pinaceae) oil. *J. Vect. Borne Dis.* 42:95-99.
- Ansari MA, Razdan RK (1995). Relative efficacy of various oils in repelling mosquitoes. *Ind. J. Malariol.* 32:104-11.
- Barnard, D.R., 1999. Repellency of essential oils to mosquitoes (Diptera: Culicidae). *J. Med. Entomol.* 36:625-629.
- Bowers WS, Sener B, Evans H, Bingal F, Erodogan I (1995). Activity of Turkish medicinal plants against mosquitoes *Aedes aegypti* and *Anopheles gambiae*. *Insect Sci. Appl.* 16:339 - 342.
- Carvalho AFU, Melo VMM, Craveiro AA, Machado MILL, Bantim B, Rabelo EF (2003). In : *Memorias Do Instituto-Oswaldo Cruz.* 11 Fortaleza, CE, Brazil. 98:569-571.
- Cavalcanti ESB, de Moraes CC, Lima MAA, Santana EWP (2004). Larvicidal activity of essential oils from Brazilian plants against *Aedes aegypti* L. *Mem. I. Oswaldo Cruz* 99:541-544.
- Cetin B, Ozer H, Kuscu H (2004). Economics of drip irrigation for apple (*Malus domestica*) orchards in Turkey. *N. Z. J. Crop Hortic. Sci.* 32:349-354.
- Chandra F, Darrit F, Darder M, Cuany A, Doannio JMC, Pasteur N, Guillet P (1998). Prethroid resistance in *Culex quinquefasciatus* form West Africa. *Med. Vet. Entomol.* 12:359-366.
- Ghosh GK (2000). In: *Biopesticide and Integrated pest Management* A.P.H. publishing corporation, New Delhi. pp. 400-446.
- Green MM, Singer JM, Sutherland DJ, Hibben CR (1991). Larvicidal activity of *Tagetes minuta* (Mari gold) towards *Aedes aegypti*. *J. Am. Mosq. Control Assoc.* 7:283-286.
- Hasan SB, Deo PG (1994). *Ocimum sanctum* seeds for mosquito control. *Int. Pest Control* 36:20-21.
- Kiso Y, Hikino H (1991). In: *Plant Biochemistry and it's Bioactivity*. Academic Press, London. pp. 219-233.
- Omolo MO, Okinyo D, Ndiege IO, Lwande W, Hassanali A (2004). Repellency of essential oils of some Kenyan plants against *Anopheles gambiae*. *Phytochemistry* 65:2797-2802.
- Paula JP, Gomes Carneiro MR, Paumgarten FJR, De Paula JP (2003). Studies on effect of *Ocimum selloi* oil against the adult mosquitoes. *J. Ethnopharmacol.* 24(88):253-260.
- Pitasawat B, Choochote W, Tuetun B, Tippawangkosol P, Kanjanapothi D, Jitpakdi A, Riyong D (2003). Repellency of aromatic turmeric *Curcuma aromatica* under laboratory and field conditions. *J. Vect. Ecol.* 28:234-240.
- Prabakar K, Jebanesan A (2004). Larvicidal efficacy of some Cucurbitaceous plant leaf extracts against *Culex quinquefasciatus* (Say). *Bioresour. Technol.* 95(1):113-114.
- Saxena BP, Koul O, Tikku A, Atal CK, Suri, OP, Suri KA (1979). Aristolochic - acid an insect chemosterilant from *Aristolochia bracteata* retz. *Ind. J. Expt. Biol.* 17:354.
- Skinner WA, Johnson HL (1980). The design of insect repellents. In: Ariens, E.T. (Ed.), *Drug Design*, vol. 10. Academic Press, New York. pp. 277-305.
- Sukumar K, Perich MJ, Boobar LR (1991). Botanical derivatives in mosquito control: A review. *J. Am. Mosq. Control Assoc.* 7:210-237.
- Wink M (1993). Production and application of phytochemicals from an agricultural perspective. In van Beek, T. A. and H. Breteleur, editors. *Phytochemistry and agriculture*. Oxford, United Kingdom Clarendon Press. pp. 171-213.