



Published in final edited form as:

J Geriatr Psychiatry Neurol. 2016 November ; 29(6): 320–327. doi:10.1177/0891988716666380.

Preoperative Cognitive Performance Dominates Risk for Delirium among Older Adults

Richard N. Jones, ScD, Edward R. Marcantonio, MD, SM, Jane S. Saczynski, PhD, Douglas Tommet, MS, Alden L. Gross, PhD, Thomas G. Trivison, PhD, David C. Alsop, PhD, Eva M. Schmitt, PhD, Tamara G. Fong, MD PhD, Sevdenur Cizginer, MD, Mouhsin M. Shafi, MD PhD, Alvaro Pascual-Leone, MD PhD, and Sharon K. Inouye, MD, MPH

Abstract

Background—Cognitive impairment is a well-recognized risk factor for delirium. Our goal was to determine if the level of cognitive performance across the non-demented cognitive ability spectrum is correlated with delirium risk, and to gauge the importance of cognition relative to other known risk factors for delirium.

Methods—The SAGES (Successful Aging after Elective Surgery) study enrolled 566 adults age ≥ 70 years scheduled for major surgery. Patients were assessed preoperatively and daily during hospitalization for the occurrence of delirium using the Confusion Assessment Method. Cognitive function was assessed preoperatively with an 11-test neuropsychological battery combined into a composite score for general cognitive performance (GCP). We examined the risk for delirium attributable to GCP, as well as demographic factors, vocabulary ability, and informant-rated cognitive decline, and compared the strength of association to risk factors identified in a previously published delirium prediction rule for delirium.

Results—Delirium occurred in 135 (24%) patients. Lower GCP score was strongly and linearly predictive of delirium risk (RR = 2.0 per each half standard deviation difference in GCP score, 95% confidence interval, 1.5, 2.5). This effect was not attenuated by statistical adjustment for demographics, vocabulary ability, and informant-rated cognitive decline. The effect was stronger than, and largely independent from, both standard delirium risk factors and comorbidity.

Conclusions—Risk of delirium is linearly and strongly related to presurgical cognitive performance level even at levels above the population median, which would be considered unimpaired.

Delirium affects up to half of older adults during surgery or hospitalization, and results in more than \$164 billion in US health care costs per year.¹ Cognitive impairment is a recognized risk factor for delirium. In fact, in a recent systematic review of delirium prediction rules, cognitive impairment was second only to age as the most commonly replicated predictor of delirium, occurring in 10 of 37 published risk prediction models.² A limitation of risk modeling in delirium with regard to cognitive functioning is that the

Corresponding Author: Richard N. Jones, ScD, Department of Psychiatry and Human Behavior, Warren Alpert Medical School, Brown University, Butler Hospital, 345 Blackstone Boulevard, Box G-BH, Providence, Rhode Island 02906-4800, Richard_Jones@Brown.edu, Phone: 401-444-1943, Fax: 401-444-1911.

measurement of cognition is typically accomplished with short mental status instruments originally developed to screen for dementia. Such measures often demonstrate substantial ceiling effects and are insensitive to performance gradients at higher levels of ability.³ Our goal was to overcome this limitation by examining the risk associated with post-operative delirium attributable to cognitive performance level across the full spectrum of cognitive ability. We also aimed to characterize the strength of this association relative to and controlling for demographics, vocabulary ability, informant ratings of cognitive decline, and risk factors identified in a previously published and validated delirium prediction rule.^{4,5}

Methods

The **SAGES** (Successful Aging after Elective Surgery) study is an ongoing prospective cohort study of older adults undergoing elective major non-cardiac surgery. The study design and methods have been described previously.⁶ Briefly, eligible participants were age 70 years and older, English speaking, scheduled to undergo elective surgery at one of two Harvard-affiliated academic medical centers and with an anticipated length of stay of at least 3 days. Eligible surgical procedures were: total hip or knee replacement, lumbar, cervical, or sacral laminectomy, lower extremity arterial bypass surgery, open abdominal aortic aneurysm repair, and colectomy. Exclusion criteria included evidence of dementia, delirium, hospitalization within 3 months, terminal condition, legal blindness, severe deafness, history of schizophrenia or psychosis, and history of alcohol abuse. A total of 566 patients were enrolled between June 18, 2010 and August 8, 2013. Written informed consent was obtained from all participants according to procedures approved by the institutional review boards of Beth Israel Deaconess Medical Center and Brigham and Women's Hospital, the two study hospitals, and Hebrew SeniorLife, the study coordinating center, all located in Boston, Massachusetts.

Participants and assessments

Participants were interviewed in their own homes on average two weeks prior to surgery. During hospitalization, participants received a daily delirium assessment (described below). Interviews were fully structured and conducted by lay research associates who received at least 4 weeks of intensive training and semiannual standardization. Chart reviews were performed by a trained research physician. All data collection staff were kept blind to study hypotheses and to the data collected in previous interviews.⁷

Delirium assessment—Delirium assessments, conducted daily in-hospital beginning with post-operative day 1, involved brief mental status testing with tasks of attention, orientation, and memory.^{6,8} Delirium symptoms were assessed with the Delirium Symptom Interview (DSI)⁹ and interviews with nurses and, if available, family members. Delirium case identification used the Confusion Assessment Method (CAM)¹⁰ and a standardized chart review method for delirium.^{11,12} If either was positive the patient was considered delirious.

Cognitive performance assessment—At preoperative baseline, participants were assessed with a neuropsychological battery that included tests of attention, recall, memory,

language, executive functioning, and visuospatial processing.^{6,13,14} Performance on Trails A & B,¹⁵ Digit Span Forwards & Backwards,¹⁶ Semantic & phonemic fluency,¹⁷ the Boston Naming Test,¹⁸ Digit Symbol Substitution,¹⁹ Hopkins Verbal Learning Test (HVL)²⁰ immediate recall (sum of 3 learning trials) & delayed recall, and the Visual Search and Attention task²¹ were combined into a single unidimensional composite using a graded item response theory model.²² Details of the composite construction have been previously described.²³ The resulting composite -- the general cognitive performance (GCP) score—is calibrated against data from the ADAMS (Aging, Demographics and Memory Study), a sub-study of the Health and Retirement Study. In a sample representative of the community dwelling older adults (age 70+) living in the US, the GCP has a mean of 50 and standard deviation of 10, and scores 45 and higher are consistent with unimpaired cognitive functioning (Mini-Mental State Examination²⁴ scores of 24 and higher).²³

An additional cognitive measure in our preoperative baseline battery, but not included in the GCP composite, was the Wechsler Test of Adult Reading (WTAR).¹⁶ The WTAR includes 50 irregularly spelled words of increasing complexity that the participant is asked to read aloud. The total score is the number of words correctly pronounced (range 0–50). The WTAR, and similar tests such as the Wechsler Adult Intelligence Scale-Vocabulary subtest²⁵ the National Adult Reading Test²⁶ are often used as indicators of premorbid intelligence.²⁷ We also conducted interviews using the IQCODE (Informant Questionnaire for Cognitive Decline in the Elderly²⁸), an informant-rated (family member) measure that records potential cognitive decline observed over the previous 10 years.

Delirium risk factors

The measure of delirium risk we use was published and validated by Inouye and colleagues (1993).^{4,5} This risk index was chosen since it has been validated in both medical^{4,5} and surgical^{4,5} patients. The risk index includes four risk factors. Vision impairment, defined as best-corrected vision worse than 20/70 on both near and distant binocular tests; cognitive impairment, defined as performance of less than 24 of 30 on the Mini-Mental State Examination; severe illness defined as an APACHE II²⁹ score above 16; and dehydration, defined as a blood urea nitrogen/creatinine ratio 18 or more.³⁰ We optimally rescaled the additive sum of risk factors. The motivation for this rescaling was to obtain a linear predictor of delirium risk that could be used as a continuous predictor in regression models, and meaningfully contrast effect estimates against other continuous predictors. To accomplish this, we regressed the risk for postoperative delirium on the count of risk factors as a categorical predictor. Then, we assigned values to the counts based on the log odds ratio (relative to the lowest count value) for delirium estimated for each count level. To minimize risks for a chance association, we repeated this process in 1,001 bootstrap replications and used the mean log odds regression weights as the new scaling metric. The ascending values assigned were: 0 (no risk factors present), 0.20, 1.40, and 2.34 (all four risk factors present).

Additional Study Variables

The baseline interview assessed sex, race, ethnicity, education, and the Modified Mini-Mental State Examination (3MS),³¹ which was equated to the MMSE and used in the

delirium risk index. Age, and Charlson comorbidity score³² were determined from chart review.

Statistical Analysis

Logistic regression models were used to characterize the relationship between GCP scores and the risk of post-operative delirium. Covariates were mean centered, except for age which was centered at 75 years, near the median of the baseline age distribution. Relative risk estimates were obtained as a nonlinear combination of estimated logistic regression parameter estimates, and interval estimates were derived from variance estimates of the nonlinear combinations using the delta method. Continuous predictors were standardized to two standard deviations following the suggestion of Gelman.³³ The two standard deviation standardization metric is useful because this scaling strategy places effect estimates associated with continuous predictors on approximately the same scale as those for binary predictors. Parameter estimates were obtained with Stata software version 14.1 (Stata Corp, College Station TX).

We estimated three models. In the first, we regressed the risk of delirium on GCP score. For descriptive purposes, we also investigated the bivariable association of demographic, vocabulary ability, delirium risk factors, and medical comorbidities. In the second, we add to the model demographic predictors including age, sex, race (white, non-Hispanic participants versus all others), WTAR score, and IQCODE score to the model with GCP predicting delirium. The remaining coefficient for GCP score in this multivariable adjusted model tells us about the remaining explanatory power of GCP after accounting for demographics and WTAR score. In the third, we add risk factors for delirium identified in a published clinical risk prediction rule for delirium, as well as medical comorbidity.

Finally, we evaluated the relative importance of predictors in the risk for delirium model using dominance analysis.^{34–36} Dominance analysis decomposes the average R -square statistic (in our regressions, McFadden's R^2) across all risk factors. The relative importance of a variable in a multivariable regression is its contribution to the overall model R^2 including direct and indirect effects on an outcome. This is a strong approach when predictors are correlated: if we only examined the magnitude of the regression coefficient, we would only be using information regarding the direct effect of a predictor.

Sensitivity analyses

To assess the possibility that our results were sensitive to the distribution of background delirium risk factors in our SAGES sample, we performed two sensitivity analyses. The first involved restricting the sample to SAGES participants with at least 1 of the Inouye et al (1993) delirium risk factors. The second involved re-estimating our predictive models for delirium given GCP and other predictors, and the dominance analysis, within multiple (1001) bootstrap draws from our original sample, where persons were selected with weights producing a distribution on the categorized sum of Inouye et al (1993) delirium risk factors that matches that of the validation sample studied in Inouye et al (1993). This represents an approximate Bayesian³⁷ post-stratification approach to address potential selection bias.³⁸

We under-sample from among patients with 0 and 1–2 of the 4 risk factors, and over-sample from the patients with 3–4 risk factors.

Results

Baseline characteristics of the patients overall and by delirium group are shown in Table 1. The mean age (SD) of the sample was 76.7 (5.2) years and 58% were women. Delirium occurred in 135/566 (24%). At baseline, the delirium group had greater levels of comorbidity and lower GCP scores. The baseline mean GCP values in both groups were above 50, the expected mean for a representative sample of U.S. older adults.²³ It is also important to point out that, as expected, the delirium risk score was higher and most of the four indicators comprising the delirium risk score were more prevalent in the group that developed delirium. The notable exception being BUN/creatinine ratio.

Delirium and cognitive performance: bivariable model

We examined the risk for post-operative delirium associated with GCP score (Figure 1 and Table 2). In Figure 1, the probability of developing post-operative delirium as a function of preoperative GCP score is illustrated. The relationship is plotted with a heavy black line, and is derived from a logistic regression model using a linear function for GCP. The dispersion in this relationship is illustrated with 1,001 separate predicted lines from as many bootstrap resamples drawn from the observed data. In the observed and bootstrap samples, we allowed for a competition between linear and quadratic functions for GCP, and in all cases the linear model was judged to be superior by Bayesian information criteria. The rug illustrates the distribution of GCP scores in the observed sample, and observed proportions of persons with delirium where sample participants have been binned according to ascending GCP score values. This plot demonstrates that the relationship between delirium risk and preoperative cognitive performance is strong and linear across the range of observed cognitive performance. This result implies that across all levels of cognitive function, even high function, persons with better cognitive function are expected to have a lower risk of delirium than persons with lower levels of cognitive function. In Table 2 (Bivariable models column) for each half standard deviation lower GCP score, the relative risk of developing post-operative delirium is 2.0 (95% confidence interval (CI): 1.5, 2.5). This association is stronger than associations identified for age (relative risk per half standard deviation, $RR_{s/2} = 1.3$), education ($RR_{s/2} = 1.1$), WTAR score ($RR_{s/2} = 1.4$), proxy report of cognitive decline (IQCODE score, $RR_{s/2} = 1.6$), and the optimally scaled delirium risk rule of Inouye et al (1993; $RR_{s/2} = 1.7$).

Delirium and cognitive performance: multivariable model including demographics

We examined the risk for post-operative delirium associated with general GCP score and adjusted for demographics and vocabulary ability (Table 2, Multivariable Model 1). The association of GCP and delirium is unchanged ($RR_{s/2} = 2.0$, 95% CI: 1.3, 2.6) implying that the association of GCP and delirium risk is not confounded or mediated by other demographic, vocabulary ability, or proxy report of cognitive decline (IQCODE score). On the other hand, with the exception of proxy-reported cognitive decline, all sociodemographic risk factors significant in the bivariable case (95% confidence intervals do not include 1.0)

have attenuated associations that are not significantly different from what might be expected due to chance when GCP is controlled. Therefore the association of age is fully mediated, and vocabulary ability is partially mediated, by GCP in explaining post-operative delirium risk.

Delirium and cognitive performance: multivariable model including delirium risk score

Table 2, Multivariable Model 2 shows the results of the multivariable model for delirium risk adjusted for demographics, vocabulary ability, and published delirium risk factors. In this model the association of lower GCP score is only slightly attenuated, and probably reflects that cognitive impairment is included in the Inouye 1993 delirium risk score. The association of the delirium risk score is also slightly attenuated relative to the bivariate model, reinforcing this interpretation. Only GCP, IQCODE, and the Inouye 1993 delirium risk score were significant predictors of post-operative delirium as suggested by confidence intervals not including 1.0.

Relative importance (dominance) analysis

The results of the dominance analysis are reported in Table 3. The relative importance column indicates the proportion of the total model R^2 (McFadden's³⁹ pseudo- R^2) that is attributable to the risk factor. The 95% confidence interval is derived from 1,001 bootstrap replications. The dominant in bootstrap replication (%) column indicates the proportion of the 1,001 replicates in which the risk factor dominates all other risk factors. Which 'dominates' is based on the R^2 contribution -- including direct and indirect effects -- for the risk factor is greater than any other risk factor's total effect. In our observed data, the GCP score is the dominant risk factor. It accounts for 29% of the model R^2 (which was relatively small, $R^2 = .09$), and in half of all bootstrap replications was the dominant covariate. The second most-important predictor was the Inouye 1993 risk index, which contributed 25% of the model R^2 and was the dominant risk factor in 35% of the 1,001 bootstrap replications. The IQCODE score was occasionally (12% of 1,001 replications) the dominant risk factor. The remaining risk factors (WTAR score, age, sex, race/ethnicity, education) were relatively unimportant risk factors for post-operative delirium in this analysis.

Sensitivity Analysis

Sensitivity analyses reveal that our inferences are sensitive to the distribution of delirium risk factors in the sample. In the restriction sensitivity analysis -- omitting patients with none of the Inouye et al (1993) delirium risk factors -- our pattern of results in terms of risk ratios was unchanged, with the GCP having the highest relative risk, followed by the Inouye et al (1993) risk score and IQCODE ($RR_{s/2}$ of 1.9, 1.7 and 1.6, respectively). However, the dominance analysis favored the Inouye et al (1993) risk score, followed by the GCP and then IQCODE. The post-stratification sensitivity analysis revealed the Inouye et al (1993) risk factor score, WTAR, and GCP to have as approximately equivalent risk ratios ($RR_{s/2}$ of 1.8, 1.8, and 1.7, respectively) but the dominance analysis clearly favored the Inouye et al (1993) risk score. These sensitivity analyses demonstrate that GCP remains among the most powerful delirium risk factors, but dominance over other risk factors is sample dependent. Caution in interpreting the results of these sensitivity analyses is called for because the restriction sensitivity analysis might attenuate the effect of the Inouye et al (1993) delirium

risk score due to restricted range, and the post-stratification sensitivity analysis makes inordinate use of the 5 patients (1%) falling into the highest risk strata to represent 25% of weighted sample of patients matching the Inouye et al (1993) validation sample.

Discussion

In this prospective cohort study of high functioning older persons undergoing major scheduled non-cardiac surgery, we demonstrated that risk of delirium after surgery is strongly related to presurgical cognitive performance level, even at levels above the population median, which would be considered unimpaired. This risk gradient appears to be linear across the range of cognitive ability included in our sample and measured with a neuropsychological battery capable of measuring very low and very high levels of cognitive performance. Additionally, we show that general cognitive performance is more important than other risk factors from a previously described and validated delirium risk tool, and appears to act directly on delirium risk in a manner robust to control for patient sociodemographic characteristics and additional clinical factors.

There are several plausible interpretations of our finding that the GCP is a strong predictor of delirium risk, even after adjusting for age, sex, race/ethnicity, vocabulary performance, and informant report of long-term cognitive decline. One is that persons with GCP scores that fall below levels predicted by the covariates we included in our models their demographic profiles and markers of cognitive decline are at increased risk for post-operative delirium. Among likely explanations for having general cognitive performance levels lower than expected is the presence of preexisting cognitive decline. Such declines could be due to unmeasured individual difference factors and/or lifetime exposures that contribute to accumulating cognitive impairments (e.g., lead exposure, repeated head trauma⁴⁰), or might reflect the effect of accelerated brain aging or relative decrements in brain plasticity.⁴¹ They might also reflect the presence of a preclinical neurodegenerative or dementing process, potentially predisposing factors for delirium.⁴² These hypotheses would require confirmation in future studies. Studies measuring mechanisms of brain cortical plasticity using noninvasive brain stimulation methods^{43,44} and determinations in the cerebrospinal fluid or neuroimaging studies to identify possible amyloid deposits and other biomarkers of preclinical dementia would be valuable was to extend our findings.

Our findings do not amount to a test of the cognitive or brain reserve hypothesis in delirium.⁴⁵ Reserve is a concept used to explain the observation that some individuals function better than others in the presence of brain pathology.⁴⁶ This analysis does not include measures of brain pathology. Moreover, current (preoperative) cognitive level is not a commonly used indicator of cognitive reserve,⁴⁷ although the strategy has been proposed.⁴⁸

Our study has some important strengths, including a relatively large prospective cohort and state-of-the-art delirium assessments, and high quality data collection with little missing data.⁷ However, several limitations should also be mentioned. First, we only chose to use one predictive model for proof of principle. Several others might have been chosen. The model we chose has been widely used, and validated in both medical and surgical samples. Another limitation is our cognitive battery, while being relatively extensive in the field of

delirium research, would be considered overly brief in the field of developmental psychology and provides limited assessment of individual cognitive domains. Additionally, our study included only persons aged at least 70 years and excluded persons with evidence of dementia by chart review or self-report or severe cognitive impairment. It may be that if we had included younger adults age would emerge as the dominant risk factor, or if we had included persons with dementia the relationship of cognitive performance and delirium risk would be non-linear.

In addition, our sample represents a highly educated population from a single geographic location. While the internal validity is not threatened, generalizability may be limited and the findings will need to be replicated in more diverse populations and settings. We conducted two sensitivity analyses, both addressing the distribution of the Inouye et al (1993) delirium risk factors in our sample. In either restricting to patients with at least one risk factor, or matching the distribution of delirium risk factors to Inouye et al's (1993) validation sample, we continue to see general preoperative cognitive performance as a strong risk factor, but dominance over the Inouye et al (1993) risk score is sample dependent.

Our study finds that preoperative cognitive level is strongly and linearly related to risk of post-operative delirium. This effect is independent of -- and more important than -- other published and validated delirium risk factors, medical comorbidity, and sociodemographic factors. The cause of the association is not well understood. The importance of cognitive level as a risk factor for delirium may have been overlooked in previous research as a consequence of the use of mental status screening instruments with low measurement fidelity at average and higher levels of cognitive ability. Our findings may encourage greater efforts to identify subtle or mild degrees of cognitive impairment preoperatively, since these patients may benefit from institution of delirium prevention strategies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors gratefully acknowledge the contributions of the patients, family members, nurses, physicians, staff members, and members of the Executive Committee who participated in the Successful Aging after Elective Surgery (SAGES) Study. A list of participating personnel of the SAGES Study can be found online as supplementary material.

This work is dedicated to the memory of Dr. Jeffrey Silverstein, a champion of delirium research and a member of the Scientific Advisory Board for this study.

Grant Funding: Supported by Grants No. P01AG031720 (SKI), K07AG041835 (SKI), R01AG044518 (SKI/RNJ) from the National Institute on Aging. Dr. Marcantonio was supported in part by Grants No. R01AG030618 (ERM) and K24AG035075 (ERM) from the National Institute on Aging. Dr. Pascual-Leone was supported by a Sidney R. Baer Jr. Foundation and Harvard Catalyst, The Harvard Clinical and Translational Science Center (NCRR and the NCATS NIH, UL1RR025758). Dr. Inouye holds the Milton and Shirley F. Levy Family Chair. The funding sources had no role in the design, conduct, or reporting of this study.

References

1. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014; 383(9920): 911–922. [PubMed: 23992774]
2. van Meenen LCC, van Meenen DMP, de Rooij SE, ter Riet G. Risk prediction models for postoperative delirium: A systematic review and meta-analysis. *J. Am. Geriatr. Soc.* 2014; 62(12): 2383–2390. [PubMed: 25516034]
3. Ashford JW. Screening for memory disorders, dementia and Alzheimer's disease. *Aging Health*. 2008; 4(4):399–432.
4. Inouye SK, Viscoli CM, Horwitz RI, Hurst LD, Tinetti ME. A predictive model for delirium in hospitalized elderly medical patients based on admission characteristics. *Ann. Intern. Med.* 1993; 119(6):474–481. [PubMed: 8357112]
5. Kalisvaart KJ, Vreeswijk R, De Jonghe JF, Van Der Ploeg T, Van Gool WA, Eikelenboom P. Risk Factors and Prediction of Postoperative Delirium in Elderly Hip-Surgery Patients: Implementation and Validation of a Medical Risk Factor Model. *J. Am. Geriatr. Soc.* 2006; 54(5):817–822. [PubMed: 16696749]
6. Schmitt EM, Marcantonio ER, Alsop DC, et al. Novel risk markers and long-term outcomes of delirium: the successful aging after elective surgery (SAGES) study design and methods. *J Am Med Dir Assoc*. 2012; 13(9):818–810. e811–810.
7. Schmitt E, Inouye S, Jones R, et al. The Successful Aging after Elective Surgery (SAGES) study: Cohort description and data quality procedures. *Journal of the American Geriatrics Society (JAGS)*. 2015 in press.
8. Simon SE, Bergmann MA, Jones RN, Murphy KM, Orav EJ, Marcantonio ER. Reliability of a structured assessment for nonclinicians to detect delirium among new admissions to postacute care. *J Am Med Dir Assoc*. 2006; 7(7):412–415. [PubMed: 16979083]
9. Albert MS, Levkoff SE, Reilly C, et al. The delirium symptom interview: an interview for the detection of delirium symptoms in hospitalized patients. *J Geriatr Psychiatry Neurol*. 1992; 5(1):14–21. [PubMed: 1571069]
10. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med*. 1990; 113(12):941–948. [PubMed: 2240918]
11. Inouye SK, Leo-Summers L, Zhang Y, Bogardus ST Jr, Leslie DL, Agostini JV. A chart-based method for identification of delirium: validation compared with interviewer ratings using the confusion assessment method. *J Am Geriatr Soc*. 2005; 53(2):312–318. [PubMed: 15673358]
12. Saczynski JS, Kosar CM, Xu G, et al. A tale of two methods: chart and interview methods for identifying delirium. *J Am Geriatr Soc*. 2014; 62(3):518–524. [PubMed: 24512042]
13. Gross AL, Jones RN, Fong TG, Tommet D, Inouye SK. Calibration and validation of an innovative approach for estimating general cognitive performance. *Neuroepidemiology*. 2014; 42(3):144–153. [PubMed: 24481241]
14. Fong T, Hshieh T, Wong B, et al. Neuropsychological profiles of an elderly cohort undergoing elective surgery and the relationship of cognitive performance with delirium. *J Am Geriatr Soc*. 2015 In Press.
15. Delis DC. Delis-Kaplan Executive Function System. The Psychological Corporation; San Antonio, TX: 2001.
16. Wechsler D. Wechsler Test of Adult Reading. The Psychological Corporation; San Antonio, TX: 2001.
17. Benton A, Hamsher K. Multilingual Aphasia Examination. University of Iowa; Iowa City: 1976.
18. Goodglass H, Kaplan E. The assessment of aphasia and related disorders. 2nd ed. Lea & Febiger; Malvern, PA: 1983.
19. Randolph C. Repeatable Battery for the Assessment of Neuropsychological Status. The Psychological Corporation; San Antonio: 1998.
20. Brandt J. The Hopkins Verbal Learning Test: Development of a new memory test with six equivalent forms. *Clinical Neuropsychologist*. 1991; 5:125–142.

21. Trenerry MR. Visual Search and Attention Test: Professional Manual. Psychological Assessment Resources; Odessa, FL: 1990.
22. Baker FB, Kim S-H. Item Response Theory: Parameter Estimation Techniques. Second ed. Marcel Dekker, Inc; New York: 2004.
23. Gross AL, Jones RN, Fong TG, Tommet D, Inouye SK. Calibration and validation of an innovative approach for estimating general cognitive performance. *Neuroepidemiology*. 2014; 42(3):144–153. [PubMed: 24481241]
24. Folstein MF, Folstein SE, McHugh PR. 'Mini-mental state.' A practical method for grading the cognitive state of patients for the clinician. *J. Psychiatr. Res.* 1975; 12(3):189–198. [PubMed: 1202204]
25. Wechsler D. Manual for the Wechsler Adult Intelligence Scale. Psychological Corporation; New York: 1955.
26. Nelson H. National adult reading test (NART): Test Manual. NFER-Nelson; Windsor: 1982.
27. Yates AJ. The use of vocabulary in the measurement of intellectual deterioration—a review. *Journal of Mental Science*. 1956; 102(429):409–440. [PubMed: 13367762]
28. Jorm A, Scott R, Cullen J, MacKinnon A. Performance of the informant questionnaire on cognitive decline in the elderly (IQCODE) as a screening test for dementia. *Psychol. Med.* 1991; 21:785–790. [PubMed: 1946866]
29. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit. Care Med.* 1985; 13(10):818–829. [PubMed: 3928249]
30. Seymour D, Henschke P, Cape R, Campbell A. Acute confusional states and dementia in the elderly: the role of dehydration/volume depletion, physical illness and age. *Age Ageing*. 1980; 9(3):137–146. [PubMed: 7457268]
31. Teng EL, Chui HC. The Modified Mini-Mental State (3MS) examination. *The Journal of clinical psychiatry*. 1987; 48(8):314–318. [PubMed: 3611032]
32. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987; 40(5):373–383. [PubMed: 3558716]
33. Gelman A. Scaling regression inputs by dividing by two standard deviations. *Stat. Med.* 2008; 27(15):2865. [PubMed: 17960576]
34. Budescu DV. Dominance analysis: A new approach to the problem of relative importance of predictors in multiple regression. *Psychol. Bull.* 1993; 114(3):542.
35. Azen R, Traxel N. Using dominance analysis to determine predictor importance in logistic regression. *Journal of Educational and Behavioral Statistics*. 2009; 34(3):319–347.
36. Luchman JN. Statistical Software Components. 2014. DOMIN: Stata module to conduct dominance analysis.
37. Hastie T, Tibshirani R, Friedman J. The elements of statistical learning: data mining, inference and prediction. Second ed. Springer; 2013.
38. Geneletti S, Best N, Toledano M, Elliott P, Richardson S. Uncovering selection bias in case-control studies using Bayesian post-stratification. *Stat. Med.* 2013; 32(15):2555–2570. [PubMed: 23303593]
39. McFadden D. Conditional logit analysis of qualitative choice behavior. 1973.
40. Del Ser T, Hachinski V, Merskey H, Munoz DG. An autopsy-verified study of the effect of education on degenerative dementia. *Brain*. 1999; 122(Pt 12):2309–2319. [PubMed: 10581224]
41. Pascual-Leone A, Amedi A, Fregni F, Merabet L. Annual Review Neuroscience. 2005. The plastic human brain cortex.
42. Sperling RA, Aisen PS, Beckett LA, et al. Alzheimer's and Dementia. 2011. Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging and the Alzheimer's Association workgroup.
43. Pascual-Leone A, Freitas C, Oberman L, et al. Characterizing brain cortical plasticity and network dynamics across the age-span in health and disease with TMS-EEG and TMS-fMRI. *Brain Topogr.* 2011; 24(3–4):302–315. [PubMed: 21842407]

44. Freitas C, Perez J, Knobel M, et al. Changes in cortical plasticity across the lifespan. *Frontiers in aging neuroscience*. 2011; 3
45. Christensen H, Anstey KJ, Leach LS, Mackinnon AJ. Intelligence, education, and the brain reserve hypothesis. In: Craik F, Salthouse T, editors *The handbook of aging and cognition*. 3 ed. Psychology Press; New York: 2008. 133–188.
46. Stern Y. Cognitive reserve: implications for assessment and intervention. *Folia Phoniatrica et Logopaedica*. 2013; 65(2):49–54. [PubMed: 23941972]
47. Harrison SL, Sajjad A, Bramer WM, Ikram MA, Tiemeier H, Stephan BC. Exploring strategies to operationalize cognitive reserve: A systematic review of reviews. *J. Clin. Exp. Neuropsychol*. 2015:1–12.
48. Satz P, Cole MA, Hardy DJ, Rassoovsky Y. Brain and cognitive reserve: mediator(s) and construct validity, a critique. *J. Clin. Exp. Neuropsychol*. 2011; 33(1):121–130. [PubMed: 20680883]




Figure 1. Risk for postoperative delirium as a function of GCP score: Results from the SAGES Study (N=566)

This figure demonstrates the relationship between the risk for post-operative delirium and preoperative General Cognitive Performance (GCP) score. The heavy line demonstrates the observed relationship as derived from a logistic regression model. In this figure, dispersion is illustrated with fitted lines from 1,001 bootstrap replications from the observed data. The points depict observed rates of post-operative delirium among groups of patients binned according to GCP score, allowing no fewer than 20 persons in each bin and plotted at the median GCP score within the bin. The rug marks at the bottom of the figure illustrate the distribution of persons according to their GCP score.

Table 1

Baseline characteristics of the SAGES study cohort (N=566)*

| Characteristic | Full Sample N= 566 | Delirium N=135 | No Delirium N=431 |
|---|--------------------|----------------|-------------------|
| Age- mean years (SD) | 76.7 (5.2) | 77.5 (5.0) | 76.4 (5.3) |
| Female- n (%) | 330 (58) | 82 (61) | 248 (58) |
| Nonwhite- n (%) | 43 (8) | 13 (10) | 30 (7) |
| Education- mean years (SD) | 15.0 (2.9) | 14.7 (3.0) | 15.0 (2.9) |
| Married- n (%) | 335 (59) | 79 (59) | 256 (59) |
| Lives Alone- n (%) | 169 (30) | 40 (30) | 129 (30) |
| Charlson Score- n (%) | | | |
| 0 | 260 (46) | 54 (40) | 206 (48) |
| 1 | 140 (25) | 23 (17) | 117 (27) |
| 2+ | 166 (29) | 58 (43) | 108 (25) |
| IQCODE score – mean (SD) | 3.12 (0.25) | 3.20 (0.22) | 3.18 (0.30) |
| GCP score-mean (SD) | 57.5 (7.4) | 54.6 (6.6) | 58.4 (7.4) |
| WTAR score – mean (SD) | 37.7 (10.0) | 35.7 (9.8) | 38.3 (9.9) |
| Inouye (1993) delirium risk score mean (SD) | 0.19 (0.15) | 0.33 (0.55) | 0.15 (0.28) |
| Vision impairment – n (%) yes | 3 (0.5) | 2 (1.5) | 1 (0.2) |
| Cognitive impairment – n (%) yes | 33 (6) | 15 (11) | 18 (4) |
| Severe illness – n (%) yes | 33 (6) | 23 (17) | 10 (2) |
| Dehydration – n (%) yes | 275 (49) | 66 (49) | 209 (49) |

* GCP= General Cognitive Performance; IQCODE, Informant Questionnaire for Cognitive Decline in the Elderly; WTAR, Wechsler Test of Adult Reading; SD= standard deviation. The Charlson comorbidity score ranged from 0–35, with scores of 2 or more indicating higher comorbidity. The Inouye (1993) risk score ranges from 0–2.34 as explained in text, and is the weighted sum of four markers: vision impairment, vision worse than 20/70 on both near and distant binocular tests; cognitive impairment, less than 24 of 30 on the Mini-Mental State Examination; severe illness, APACHE II above 16; dehydration, BUN/creatinine ratio 18 or more.

Table 2

Relative risk of post-operative delirium in bivariable and multivariable models: Results from the SAGES Study (N = 566).

| Predictor | Relative risk | | |
|--|----------------------|--------------------------|--------------------------|
| | (95% CI) | | |
| | Bivariable Models | Multivariable Model 1 | Multivariable Model 2 |
| Lower GCP score (per half SD) | 2.0 (1.5, 2.5) | 2.0 (1.3, 2.6) | 1.9 (1.2, 2.6) |
| Higher age (per half SD) | 1.3 (1.0, 1.7) | 1.0 (0.6, 1.3) | 1.0 (0.6, 1.3) |
| Female (vs male) | 1.1 (0.8, 1.4) | 1.2 (0.8, 1.5) | 1.2 (0.8, 1.7) |
| All other race and ethnicity groups (vs white, non-Hispanic) | 1.0 (0.4, 1.6) | 0.5 (0.1, 1.0) | 0.7 (0.2, 1.1) |
| Lower education (per half SD) | 1.1 (0.9, 1.2) | 0.9 (0.7, 1.1) | 0.9 (0.7, 1.1) |
| Lower WTAR score (per half SD) | 1.4 (1.1, 1.8) | 1.3 (0.8, 1.9) | 1.2 (0.6, 1.7) |
| Higher IQCODE score (per half SD) | 1.6 (1.2, 2.0) | 1.5 (1.1, 1.9) | 1.5 (1.1, 2.0) |
| Higher Inouye 1993 risk score, optimum scaling (per SD) | 1.7 (1.3, 2.1) | -- | 1.6 (1.2, 2.0) |
| Higher Charlson comorbidity (per SD) | 1.2 (1.0, 1.4) | -- | 1.1 (1.0, 1.3) |

GCP= General Cognitive Performance; IQCODE, Informant Questionnaire for Cognitive Decline in the Elderly; WTAR, Wechsler Test of Adult Reading; SD, standard deviation.

Table 3

Dominance analysis of multivariable model for post-operative delirium risk.

| Predictor | Relative Importance | 95 % Confidence Interval | Dominant in Bootstrap Replication (%) |
|--|---------------------|--------------------------|---------------------------------------|
| Lower GCP score (per half SD) | 0.29 | (0.11, 0.51) | 0.50 |
| Higher Inouye 1993 risk score, optimum scaling (per SD) | 0.25 | (0.05, 0.49) | 0.35 |
| Higher IQCODE score (per half SD) | 0.16 | (0.02, 0.37) | 0.12 |
| Higher Charlson comorbidity (per SD) | 0.09 | (0.00, 0.27) | 0.03 |
| Lower WTAR score (per half SD) | 0.07 | (0.02, 0.18) | 0.00 |
| Higher age (per half SD) | 0.04 | (0.01, 0.13) | 0.00 |
| All other race and ethnicity groups (vs white, non-Hispanic) | 0.03 | (0.00, 0.12) | 0.00 |
| Lower education (per half SD) | 0.03 | (0.01, 0.10) | 0.00 |
| Female (vs male) | 0.03 | (0.00, 0.13) | 0.00 |

GCP= General Cognitive Performance; IQCODE, Informant Questionnaire for Cognitive Decline in the Elderly; WTAR, Wechsler Test of Adult Reading; SD, standard deviation. The relative importance column indicated the proportion of the total model R^2 ($R^2 = 0.09$) that is attributable to the risk factor. The 95% confidence interval is derived from 1,001 bootstrap replications of each model. The dominant in bootstrap replication (%) column indicates the proportion of the 1,001 replicates in which the risk factor dominates all other risk factors. Dominates means the R^2 contribution including direct and indirect effects for the risk factor is greater than any other risk factor's total effect.