



Published in final edited form as:

*J Immunother.* 2006 ; 29(6): 616–627.

## Presentation of Tumor Antigens by Dendritic Cells Genetically Modified With Viral and Nonviral Vectors

Michal Lotem<sup>\*,†</sup>, Yangbing Zhao<sup>\*</sup>, John Riley<sup>\*</sup>, Patrick Hwu<sup>‡</sup>, Richard A. Morgan<sup>\*</sup>, Steven A. Rosenberg<sup>\*</sup>, and Maria R. Parkhurst<sup>\*</sup>

<sup>\*</sup>*Surgery Branch, National Cancer Institute, Bethesda, MD 20892, USA*

<sup>†</sup>*Sharett Institute of Oncology, Hadassah University Hospital, Jerusalem, Israel*

<sup>‡</sup>*Department of Melanoma Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, TX.*

### Summary

Genetic modification of dendritic cells (DCs) with recombinant vectors encoding tumor antigens may aid in developing new immunotherapeutic treatments for patients with cancer. Here, we characterized antigen presentation by human DCs genetically modified with plasmid cDNAs, RNAs, adenoviruses, or retroviruses, encoding the melanoma antigen gp100 or the tumor-testis antigen NY-ESO-1. Monocyte-derived DCs were electroporated with cDNAs or RNAs, or transduced with adenoviruses. CD34<sup>+</sup> hematopoietic stem cell-derived DCs were used for retroviral transduction. Genetically modified DCs were coincubated with CD8<sup>+</sup> and CD4<sup>+</sup> T cells that recognized major histocompatibility complex class I- and class II-restricted epitopes from gp100 and NY-ESO-1, and specific recognition was evaluated by interferon $\gamma$  secretion. Cytokine release by both CD8<sup>+</sup> and CD4<sup>+</sup> T cells was consistently higher in response to DCs modified with adenoviruses than cDNAs or RNAs, and maturation of DCs after genetic modification did not consistently alter patterns of recognition. Also, retrovirally transduced DCs encoding gp100 were well recognized by both CD8<sup>+</sup> and CD4<sup>+</sup> T cells. These data suggest that DCs transduced with viral vectors may be more efficient than DCs transfected with cDNAs or RNAs for the induction of tumor reactive CD8<sup>+</sup> and CD4<sup>+</sup> T cells in vitro and in human vaccination trials.

### Keywords

dendritic cells; electroporation; nonviral vectors; adenovirus; retrovirus

---

Dendritic cells (DCs) are potent antigen-presenting cells capable of stimulating both naive CD4<sup>+</sup> T-helper cells and CD8<sup>+</sup> cytotoxic T lymphocytes (CTLs).<sup>1</sup> Therefore, DCs loaded with tumor-associated antigens may facilitate the development of new immunotherapies for the treatment of patients with cancer. DCs can readily be loaded with synthetic peptides, but this strategy is limited to the repertoire of known tumor-associated epitopes and to patients who express particular major histocompatibility complex (MHC) restriction elements. An alternate method for loading DCs with antigen is direct incubation with full-length recombinant protein. However, exogenously loaded protein is predominantly processed by DC via the endocytic compartment, and epitopes are most efficiently presented in the context of MHC class II molecules, eliciting CD4<sup>+</sup> T-cell responses. Through cross-presentation, an exogenously loaded protein may be diverted to the endogenous processing pathway, resulting in presentation

of MHC class I-binding epitopes recognized by CD8<sup>+</sup> T cells.<sup>2,3</sup> However, the extent to which cross-presentation truly results in CTL activation to exogenous self-antigens is unclear.<sup>4,5</sup> In a previous report, we did not detect significant cross-presentation when the melanoma differentiation antigen gp100 was pulsed on DCs as an intact protein or from melanoma cell lysates.<sup>6</sup> Alternatively, in several other reports, efficient cross-presentation was induced using full-length proteins formulated as antigen-antibody immune complexes,<sup>7</sup> with ISCOMATRIX adjuvant,<sup>7,8</sup> or with antibodies against DEC-205.<sup>9</sup>

Insertion of full-length antigens into DC by genetic modification has the potential to overcome the dependence of MHC class I processing on cross-presentation after loading with exogenous protein. A vector-transferred recombinant antigen synthesized in the cytosol may enter the degradation process of intracellular molecules, eventually yielding epitope presentation in the context of MHC class I molecules.<sup>10</sup> Several gene delivery methods have been employed for genetic modification of DC, and these can be divided into viral and nonviral vectors.

The nonviral vectors, including plasmid cDNAs and in vitro transcribed RNAs (IVT RNAs), exclusively confer the antigen of interest and are relatively easy to produce. IVT RNA is an especially attractive vector because it does not integrate into the genome and is not controlled by promoter sequences. RNA transfections are transient, but a short-lived display of peptide on RNA-transfected DCs may be sufficient to prime antigen-specific T cells.<sup>11</sup> DNA and RNA transfections are most efficiently carried out using electroporation.<sup>12</sup> Briefly, high-voltage electrical pulses are applied to cells to induce the formation of transient pores in the cell membrane. DNA requires higher voltages to enter the nuclear membrane and is therefore associated with higher rates of cell damage in comparison to RNA, which only requires passage through the cytoplasmic membrane. DCs transfected with RNA were previously reported to be more efficient than DNA-transfected DCs for antigen-specific T-cell stimulation.<sup>13</sup>

Several viral vectors have been used to transduce DCs, including recombinant adenoviruses,<sup>14</sup> pox viruses,<sup>15</sup> retroviruses,<sup>16</sup> and lentiviruses.<sup>17</sup> In general, viral transductions induce higher levels of transgene expression than their nonviral counterparts. Adenoviral vectors transduce DCs with high efficiency and do not significantly reduce cell viability.<sup>18,19</sup> However, proteins from the adenoviral backbone may dominate the immune response, and attempts to reduce the expression of adenoviral proteins in these vectors have not been successful.<sup>19,20</sup> Retroviruses have the advantage that viral proteins are not expressed after integration of the transgene into the genome, and retrovirally transduced DCs can elicit both CD8<sup>+</sup> and CD4<sup>+</sup> T-cell responses to tumor antigens.<sup>16,21</sup> However, retroviruses can only transduce dividing cells, and therefore, cannot mediate gene transfer into monocyte-derived DC populations. Instead, CD34<sup>+</sup> hematopoietic stem cells (HSCs) must be mobilized and stimulated to proliferate in vitro using a cocktail of cytokines. After retroviral transduction, these cells can then be stimulated to differentiate into DCs expressing the transgene. Lentiviruses are thought to be able to transduce nondividing cells, but levels of transgene expression in DCs is generally not as high as with other viral vectors, and the production of stable high-titer viral supernatants is technically challenging.<sup>17</sup>

In this report, we compared patterns of antigen presentation by DCs genetically modified with viral and nonviral vectors. The use of antigen-specific T-cell lines provided a biologic assay for assessing presentation of specific epitopes because binding of T-cell receptor to relevant peptide-MHC class I or class II complexes on DC surfaces induced measurable interferon (IFN)  $\gamma$  secretion. We selected the human tumor-associated antigens gp100 and NY-ESO-1 as model proteins due to the availability of tumor reactive CD8<sup>+</sup> and CD4<sup>+</sup> T-cell lines with specificities for a variety of antigen-derived epitopes. DCs transduced with viral vectors were consistently recognized more efficiently than those transfected with cDNAs or IVT RNAs by both MHC

class I and class II restricted T cells suggesting that viral vectors may be more efficient for the induction of tumor reactive CD8<sup>+</sup> and CD4<sup>+</sup> T cells in vitro and in human vaccination trials.

## MATERIALS AND METHODS

### Cell Lines and Reagents

Human melanoma cell lines, Epstein-Barr virus-transformed B cells, and T2 cells were routinely cultured in RPMI 1640 supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 2 mM L-glutamine (Invitrogen, Carlsbad, CA). DC cultures were initiated by plating human peripheral blood mononuclear cells (PBMC) in X-Vivo-20 (BioWhittaker, Walkersville, MD). Human lymphocytes and DCs were cultured in complete medium (CM) consisting of RPMI 1640, 2 mM L-glutamine, 50 U/mL penicillin, 50 µg/mL streptomycin (Invitrogen), and 10% heat-inactivated human AB serum (Gemini Bio-Products, Woodland, CA; Valley Biomedical, Winchester, VA). Multiple melanoma-reactive T-cell lines and clones were used to evaluate the presentation of MHC-restricted epitopes by genetically modified DCs<sup>16,22–29</sup> as presented in Table 1.

Expression of gp100 in melanoma cell lines was assessed by fluorescence-activated cell sorting (FACS) using a monoclonal antibody (mAb) (HMB45; Enzo Diagnostics, Farmingdale, NY), and expression of NY-ESO-1 was previously evaluated on the basis of specific IFN $\gamma$  secretion by two T-cell clones, MB4 and M8, respectively, that specifically recognize peptides from these proteins in the context of human leukocyte antigen (HLA)-A\*0201.<sup>26</sup> The F002R melanoma cell line (F002Rmel) did not naturally express gp100, but it had previously been transduced with a vesicular stomatitis virus (VSV)-pseudotyped retroviral vector encoding either gp100 or green fluorescence protein (GFP).<sup>16</sup> Expression of HLA class II molecules on the surfaces of melanoma cells was upregulated by transduction with a retroviral construct encoding the HLA class II transactivator gene (CIITA) as previously described.<sup>30</sup> The expression of cell surface HLA class I and class II molecules on melanoma cells was confirmed by FACS using mAbs against HLA-A2 (One Lambda, Conestoga, CA), HLA-DR (L243; Becton Dickinson Biosciences, Franklin Lakes, NJ), HLA-DR4 (Accurate Chemical and Scientific Corp, Westbury, NY), and HLA-DR7 (Pel-Freez Biologicals, Rogers, AR). In addition, HLA haplotypes of cell lines were determined by DNA sequencing in the HLA Laboratory at the National Institutes of Health. The expression of gp100 and NY-ESO-1, HLA-A\*0201, HLA-DR $\beta$ 1\*0401, and HLA-DR $\beta$ 1\*0701 in melanoma cell lines was as follows: 888mel (gp100<sup>+</sup>, NY-ESO-1<sup>-</sup>, HLA-A\*0201<sup>-</sup>, HLA-DR $\beta$ 1\*0401<sup>-</sup>, HLA-DR $\beta$ 1\*0701<sup>-</sup>), 624mel CIITA (gp100<sup>+</sup>, NY-ESO-1<sup>+</sup>, HLA-A\*0201<sup>+</sup>, HLA-DR $\beta$ 1\*0401<sup>+</sup>, HLA-DR $\beta$ 1\*0701<sup>+</sup>), 526mel CIITA (gp100<sup>+</sup>, NY-ESO-1<sup>-</sup>, HLA-A\*0201<sup>+</sup>, HLA-DR $\beta$ 1\*0401<sup>+</sup>, HLA-DR $\beta$ 1\*0701<sup>-</sup>), F002Rmel CIITA-gp100 (gp100<sup>+</sup>, HLA-A\*0201<sup>+</sup>, HLA-DR $\beta$ 1\*0401<sup>-</sup>, HLA-DR $\beta$ 1\*0701<sup>+</sup>), F002Rmel CIITA-GFP (gp100<sup>-</sup>, HLA-A\*0201<sup>+</sup>, HLA-DR $\beta$ 1\*0401<sup>-</sup>, HLA-DR $\beta$ 1\*0701<sup>+</sup>), and 1088mel CIITA (gp100<sup>+</sup>, HLA-DR $\beta$ 1\*0401<sup>+</sup>, HLA-DR $\beta$ 1\*0701<sup>-</sup>).

### Generation of DC

PBMC and CD34<sup>+</sup> HSCs were obtained from patients with metastatic melanoma who were enrolled in clinical trials in the Surgery Branch of the National Cancer Institute, as part of IRB approved protocols. Immature DCs derived from peripheral blood mononuclear cells were prepared as previously described.<sup>31</sup> Briefly, PBMC were plated in X-Vivo-20 (BioWhittaker), and adherent cells were subsequently cultured in CM containing 1000 IU/mL granulocyte-macrophage colony-stimulating factor (GM-CSF) (Peprotech, Rocky Hill, NJ) and 1000 IU/mL interleukin (IL)-4 (Peprotech). DCs were electroporated with cDNAs or IVT RNAs or were transduced with recombinant adenoviral vectors on day 6. Maturation cytokines were added where indicated as follows: 1 µg/mL prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) (Sigma-Aldrich); 10 ng/

mL tumor necrosis factor (TNF) $\alpha$  (Sigma-Aldrich, St Louis, MO); 1  $\mu$ g/mL soluble trimeric CD40L (Amgen-Immunex, Seattle, WA); 5  $\mu$ g/mL lipo-polysaccharide (LPS) (Sigma-Aldrich); or 50  $\mu$ g/mL IFN $\alpha$ ~(Roche).

For the preparation of HSC-derived DCs, CD34<sup>+</sup> cells were mobilized in peripheral blood of patients by 5 daily subcutaneous injections of 10  $\mu$ g/kg rhG-CSF (Filgrastim; Neupogen; Amgen, Thousand Oaks, CA), followed by lymphocytapheresis to obtain PBMCs on day 6. Mobilized CD34<sup>+</sup> HSCs were then isolated using StemSep CD34-positive selection cocktail and magnetic colloid (StemCell Technologies, Vancouver, Canada). Cells were washed and plated in 6-well plates in CM containing 10 ng/mL TNF $\alpha$  (Sigma-Aldrich, St Louis, MO), 40 ng/mL stem cell factor (SCF) (R&D Systems, Minneapolis, MN), and 1000 IU/mL GM-CSF (Peprotech) as previously described.<sup>32</sup>

## Plasmids

pcDNA3 plasmids containing full-length cDNAs encoding gp100,<sup>33</sup> NY-ESO-1,<sup>34</sup> and eGFP were used for cDNA electroporation. For in vitro transcription of mRNAs, a modified pGEM-4Z plasmid was constructed, to which a 64 bp length poly A tail had been added downstream of T7 promoter and multiple cloning site, as previously described.<sup>35</sup> The full-length NY-ESO-1 was inserted into the pGEM-4Z-64A vector using HindIII and HpaI restriction enzymes. The full length gp100 was inserted using NotI and HindIII restriction digestion and ligation. Plasmids were linearized with SpeI for use as templates for in vitro transcription of RNA with mMACHINE High Yield Capped RNA Transcription Kit (Ambion) according to the manufacturer's instructions.

## DNA Electroporation

DNA electroporation was performed with the AMAXA electroporation system using the Human Dendritic Cell Nucleofactor Kit I (AMAXA Biosystems, Cologne, Germany). Briefly,  $2.5 \times 10^6$  monocyte-derived DCs were resuspended in 100  $\mu$ L Nucleofactor solution and were electroporated in an electroporation cuvette with an electrode gap of 2 mm. DNA was added immediately before electroporation at 5  $\mu$ g/sample. DCs were recovered in prewarmed CM containing GM-CSF and IL-4 at a final concentration of  $1 \times 10^6$  cells/mL. Cell viability ranged from 45% to 70% 24 hours posttransfection, as detected by propidium iodide staining. In some experiments, transfected cells were analyzed by flow cytometry or immunohistochemical staining 24 hours posttransfection.

## RNA Electroporation

RNA electroporation was performed as previously described.<sup>36</sup> Briefly, DCs were washed and gently resuspended in Opti-MEM media (Invitrogen, Grand Island, NY) at  $2.5 \times 10^7$ /mL. Cells were electroporated in 2-mm cuvettes at 300 V for 500  $\mu$ s using an Electro Square Porator ECM 830 (BTX, San Diego, CA). The amount of IVT RNA used was 2  $\mu$ g per  $10^6$  DCs. Cells were immediately transferred to media containing 50% fresh CM and 50% conditioned CM (collected from day 7 cultured DC), supplemented with GM-CSF and IL-4.

## Adenoviral Vectors

Ad2 viruses encoding gp100 and eGFP were commercially supplied (Genzyme, Cambridge, MA) and were used to transduce monocyte-derived DCs as previously described.<sup>14</sup> Briefly, DCs were resuspended in X-Vivo-15 (BioWhittaker) at  $10^7$ /mL and incubated with Ad2 stocks at 37°C at a multiplicity of infection of 300. Transduced DCs were resuspended in CM containing GM-CSF and IL-4 at a final concentration of  $1 \times 10^6$  cells/mL. Coincubation with responding T cells was performed after a 24 hour rest period.

## Retroviral Vectors

The VSV-pseudotyped retroviral system was used as previously described.<sup>16,32</sup> Briefly, the vectors were prepared by first inserting the complete gp100 and NY ESO-1 sequences into the pCLNC retroviral plasmid. The pCLNC-gp100 and pMDG-VSV plasmids were cotransfected into 293-gag-pol packaging cells using Lipofectamine Plus (Invitrogen). The 293-gag-pol packaging cells, provided by I. Verma (The Salk Institute, La Jolla, CA), were cultured in Dulbecco's modified Eagles's medium supplemented with 10% heat-inactivated fetal bovine serum and antibiotics. Media was changed 16 and 48 hours after transfection, and culture supernatants were harvested on days 3 and 4. Producer cells were removed from retroviral supernatants by filtration through a 0.2- $\mu$ m filter (Nalgene, Rochester, NY). Supernatants were immediately frozen at 70°C for future use.

Retroviral supernatants were added to cultured CD34<sup>+</sup> cells on days 2 and 3 at a ratio of 1:1 CM. GM-CSF, SCF, TNF $\alpha$ , and polybrene were added, and cells were spun in the plate at 1000 g for 1 hour. On day 4 (or immediately after the second spinfection), transduced DCs were resuspended in fresh CM supplemented with GM-CSF and IL-4 as described above for monocyte-derived DCs. In some experiments, maturation was induced 16 hours before coculture of DCs and T cells with various cytokine cocktails as indicated.

## FACS Analyses

DCs were characterized for expression of cell surface markers including CD11c, CD14, CD40, CD80, CD86, CD83, HLA-A, B, C, and HLA-DR using PE or fluorescein isothiocyanate-conjugated mAbs (BD PharMingen, San Diego, CA). FACS analyses were performed on a FACSCalibur flow cytometer (BD Biosciences) using CellQuest software (BD Biosciences).

## Cytokine Release Assay

Recognition of target cells by melanoma reactive CD8<sup>+</sup> and CD4<sup>+</sup> T-cell lines and clones was evaluated on the basis of specific IFN $\gamma$  secretion. Responder T cells ( $10^5$ ) were cocultured with 0.5 to  $1 \times 10^5$  genetically modified DCs (250  $\mu$ L total) ~20 hours at 37°C, and the concentration of human IFN $\gamma$  in coculture supernatants was measured by ELISA (Pierce-Endogen, Cambridge, MA). As positive controls for T-cell function, specific IFN $\gamma$  secretion was measured in response to peptide-loaded target cells and melanomas. For HLA-A\*0201 restricted CD8<sup>+</sup> T-cell populations, T2 cells were incubated with 1  $\mu$ M of the appropriate peptide 1 to 3 hours at 37°C. For class II HLA-restricted CD4<sup>+</sup> T cells, Epstein-Barr virus-transformed B cells expressing HLA-DR $\beta$ 1\*0401 or HLA-DR $\beta$ 1\*0701 were incubated with 50  $\mu$ M of the appropriate peptide approximately 3 hours at 37°C. In addition, melanoma cell lines expressing various combinations of HLA-A\*0201, HLA-DR $\beta$ 1\*0401, and HLA-DR $\beta$ 1\*0701 were harvested. Responder T cells ( $10^5$ ) were cocultured with  $10^5$  stimulator cells, and the concentration of human IFN $\gamma$  in coculture supernatants was measured by ELISA (Pierce-Endogen).

## RESULTS

### Efficiencies of Gene Transfer Methods

To estimate transfection or transduction efficiencies, DCs genetically modified with viral and nonviral vectors encoding eGFP were analyzed by FACS (Fig. 1). Using the AMAXA electroporation system for cDNA plasmids, 24 hours after transfection, GFP expression ranged from 18% to 36%, and cell viability was 40% to 75% (data not shown). Transfection efficiencies with IVT RNAs were generally higher than those with cDNAs, and DCs electroporated with IVT RNA expressed detectable levels of GFP in 60% to 96% of transfected cells. Cell viabilities were also generally higher using IVT RNAs in comparison to cDNAs (ie,

80% to 90% vs. 40% to 75%, data not shown). Adenoviral vectors were comparable to IVT RNAs in terms of transduction/transfection efficiencies which generally exceeded 90%. However, expression levels, as evaluated by mean fluorescence intensities (MFIs), were usually higher for adenovirally transduced DCs. The transduction efficiency of the retroviral vector was comparable to plasmid cDNA electroporation (ie, ~23%). However, like the recombinant adenovirus, MFIs for retrovirally transduced DCs were usually higher than those for DCs transfected with cDNA or IVT RNA.

### Expression of Phenotypic Markers on Genetically Modified DCs

In the absence of any maturation cytokines, expression of various phenotypic markers was not significantly different on DCs transfected or transduced with cDNA, IVT RNA, or Ad2 (Fig. 2). All immature DC populations were essentially 100% positive for CD11c and negative for CD14, as is characteristic of monocyte-derived DCs. High levels of the costimulatory molecule CD86 were consistently observed, with much lower levels of CD80. Expression of CD40 on immature DCs was comparable after transfection with cDNA or IVT RNA but was lower after Ad2 transduction. Also, there was a trend for higher levels of the maturation marker CD83 on immature DCs electroporated with IVT RNA. In addition, genetic modification of immature DCs did not induce different patterns of cytokine secretion including IL-12p70, IL-1 $\beta$ , and IL-15 (data not shown). In contrast, significant upregulation of CD80, CD86, and CD40 was observed on RNA-transfected DCs after maturation with CD40L and LPS or TNF $\alpha$  and PGE<sub>2</sub>. However, this was not the case for DCs matured with CD40L and LPS after electroporation with cDNA.

### Presentation of gp100 Epitopes by DCs Genetically Modified With cDNA, IVT RNA, and Ad2

To determine if genetically modified DCs presented relevant MHC class I and class II restricted tumor-associated epitopes, IFN $\gamma$  secretion by multiple CD8<sup>+</sup> and CD4<sup>+</sup> melanoma-reactive T-cell populations (Table 1) was measured in response to transfected or transduced DCs. The melanocyte differentiation antigen gp100 was initially selected as a model system because multiple MHC class I and class II restricted T-cell lines were available that specifically recognized epitopes from this protein.

In multiple experiments, recognition of DCs genetically modified with cDNA, IVT RNA, and Ad2 encoding gp100 by a variety of different MHC class I and class II restricted T-cell lines was directly compared. Data from 2 representative experiments are presented in Tables 2 and 3. In some experiments (Table 3), low amounts of IFN $\gamma$  were secreted by HLA-A\*0201 restricted gp100 reactive CTL in response to DCs transfected with gp100 cDNA. However, in 22 of 26 electroporation experiments (85%), no specific cytokine secretion was detected by CD8<sup>+</sup> T-cell lines in response to immature DCs electro-porated with either gp100 cDNA or IVT RNA (Table 2). MHC class II restricted recognition of gp100 epitopes on electroporated DCs was also inconsistent. In some experiments (Table 3), low amounts of IFN $\gamma$  were secreted by HLA-DR restricted gp100 reactive CD4<sup>+</sup> T cells in response to DCs transfected with gp100 cDNA. In 3 of 4 additional electroporation experiments, DCs transfected with gp100 cDNA or IVT RNA were not well-recognized by HLA-DR $\beta$ 1\*0701 restricted gp100 reactive CD4<sup>+</sup> T cells that recognized either gp100:170–190 or gp100:420–435. However, in most other experiments, the BR-B8 CD4<sup>+</sup> T-cell line clearly recognized gp100:44–59 on electroporated DCs in the context of HLA-DR $\beta$ 1\*0401 (Table 2). In contrast, DCs transduced with Ad2 encoding gp100 were consistently well recognized by both MHC class I and class II restricted T-cell lines as evaluated by the high amounts of specific IFN $\gamma$  secretion observed in multiple experiments (Tables 2 and 3).

### Effect of Maturation on Presentation of gp100 Epitopes by Electroporated DCs

To determine if maturation of DCs after transfection with nonviral vectors encoding gp100 would enhance presentation of MHC class I and class II restricted epitopes, IFN $\gamma$  secretion by gp100 reactive T cells was measured in response to electroporated DCs subsequently stimulated with various cytokine cocktails (Tables 4 and 5). Addition of maturation cytokines did not consistently or significantly effect MHC class I or class II presentation of gp100 epitopes. However, there may have been a slight trend toward enhanced recognition of gp100:209–217 by the L2D8T cell clone upon maturation of DCs with soluble CD40L and LPS. Also, IFN $\gamma$  secretion by the MHC class II restricted T-cell lines BR-B8 and B104 was usually lower after DC maturation with TNF $\alpha$  and PGE $_2$ .

### Presentation of NY-ESO-1 Epitopes by DCs Transfected With IVT RNA

Because of improved DC viability after electroporation with IVT RNA as compared with cDNA in previous experiments, nonviral-mediated transfections of DCs with NY-ESO-1 were performed solely with IVT RNA (Table 6). Also, because of the previously described trends for recognition of gp100 epitopes after maturation of genetically modified DCs, only the combination of CD40L and LPS was evaluated as a maturation reagent for DCs transfected with cDNA or IVT RNA encoding NY-ESO-1. MHC class I restricted presentation of the NY-ESO-1:157-165 epitope was observed by DCs from 4 of 5 patients evaluated using 3 different HLA-A\*0201-restricted T-cell lines. The addition of the maturation cocktail containing CD40L and LPS was necessary for antigen detection by the T-cell clones JH1 and M8 for 1 population of DCs (Donor 15), but no impact was observed for DCs from a second patient (Donor 16). However, DC maturation was not required for the highly avid T-cell line TH1F2L. MHC class II-restricted presentation of the NY-ESO-1:161–180 epitope was also observed using the HLA-DP $\beta$ 1\*0401-restricted T-cell line SG6. However, DC maturation was required for this presentation for both DC populations evaluated.

### Presentation of gp100 and NY-ESO-1 Epitopes by Retrovirally Transduced DCs

Data from the Ad2 vector transductions suggested that virally mediated gene transfer into DCs consistently enabled efficient presentation of both MHC class I and class II-restricted epitopes from tumor-associated antigens. However, adenoviral vectors also enable presentation of epitopes from the adenovirus backbone, and these may dominate the immune response when used to stimulate a heterogeneous population of T cells, such as peripheral blood leukocyte. To avoid this disadvantage, we evaluated epitope presentation by DCs after retroviral transduction because nonreplicating retroviral vectors only induce expression of the transgene after genomic integration into host cells. Retroviral transductions were performed using the Moloney murine leukemia virus (MMLV) retrovirus produced by the 293-gag-pol packaging cell line as previously described.<sup>16,32</sup> Since retroviruses can only transduce dividing cells, we first isolated CD34<sup>+</sup> HSCs and stimulated them to proliferate with TNF $\alpha$ , SCF, and GM-CSF. These cells were then retrovirally transduced and were subsequently differentiated into DCs using GM-CSF and IL-4. DCs transduced with MMLV encoding both gp100 and NY-ESO-1 were well recognized by both MHC class I and class II-restricted T-cell lines (Table 7), and similar results were observed using CD34<sup>+</sup> HSC-derived DCs from a second donor (data not shown). However, it is not possible to compare results directly between retrovirally transduced DCs and other means of genetic modification because of the differences in DC preparation.

## DISCUSSION

Genetic modification of DCs with tumor-associated antigens may facilitate the development of new immunotherapies for the treatment of patients with cancer. The results presented here suggest that genetic engineering of DCs is feasible using both viral and nonviral gene delivery methods. Gene transfer efficiencies of greater than 18% were observed with all methods as

evaluated by FACS after genetic manipulation with viral and nonviral vectors encoding eGFP. High transfection efficiencies did not correlate with efficient antigen processing and presentation of relevant T-cell epitopes. However, higher amounts of protein produced by DCs transduced with viral vectors in comparison with those transfected with cDNA or IVT RNA may have been related to the observation that these cells consistently presented both MHC class I and class II-restricted T-cell epitopes.

Our hypothesis was that antigen present in the cytosol would be degraded by the proteasome, and resulting peptides would be assembled with MHC class I molecules in the endoplasmic reticulum. Peptide-MHC class I complexes would then migrate to the surfaces of the gene-modified DCs and activate CD8<sup>+</sup> T cells as previously described.<sup>37</sup> We also anticipated low levels of CD4<sup>+</sup> T-cell stimulation, because most endogeneously produced proteins do not efficiently enter the endocytic pathway associated with presentation of MHC class II-restricted epitopes.<sup>38</sup> However, we found that DCs transfected with nonviral vectors encoding gp100 did not consistently stimulate tumor reactive CTL, and there may have been a slight trend toward enhanced presentation of MHC class II-restricted T-cell epitopes. In contrast to nonviral vectors, transduction of DCs with recombinant Ad2 and MMLV vectors consistently induced potent presentation of both MHC class I and class II-restricted epitopes. The apparent difference in antigen processing between viral and nonviral vectors could not be attributed to transfection/transduction efficiencies or to differences in phenotypic markers expressed on DCs. The efficiencies of RNA electroporation and Ad2 transduction were similarly high, and both methods maintained high levels of viability of phenotypically similar DCs. However, in multiple experiments, much higher amounts of IFN $\gamma$  were secreted by several different CD8<sup>+</sup> and CD4<sup>+</sup> T-cell lines in response to DCs transduced with Ad2 in comparison with those transfected with IVT RNA encoding gp100. A similar trend of enhanced MHC class I and class II-restricted epitope presentation with a virus compared with a nonviral vector was also observed for NY-ESO-1, although we only evaluated IVT RNA and MMLV for this particular antigen. In contrast, the overall amount of protein produced by DCs transduced with viral vectors was probably higher than those transfected with cDNA or IVT RNA as suggested by higher MFIs after genetic modification with eGFP (Fig. 1). Therefore, enhanced epitope presentation by virally transduced DCs may be related to increased quantities of proteins produced by these cells.

To determine if maturation of DCs after transfection with nonviral vectors would enhance presentation of MHC class I and class II-restricted epitopes, T-cell recognition of electroporated DCs subsequently stimulated with 2 different cytokine cocktails was evaluated. In particular, we evaluated recognition of gp100-transfected DCs after maturation with either CD40L and LPS or TNF $\alpha$  and PGE<sub>2</sub>. Addition of these maturation cytokines did not consistently or significantly effect MHC class I or class II presentation of gp100 epitopes. However, there may have been a slight trend toward enhanced recognition of gp100:209–217 by one particular T-cell clone upon DC maturation with CD40L and LPS and a trend toward decreased recognition of MHC class II-restricted epitopes upon DC maturation with TNF $\alpha$  and PGE<sub>2</sub>. On the basis of these trends for gp100, we only evaluated recognition of NY-ESO-1 epitopes after DC maturation with CD40L and LPS. Likewise, addition of these maturation cytokines did not consistently or significantly effect MHC class I or class II presentation of NY-ESO-1 epitopes. Despite these findings, we did not evaluate the effect of DC maturation on the priming of naïve T cells. Conclusions from such experiments might be very different than those presented here because IL-12 production is believed to be critical in the priming phase of the immune response.

In the experiments presented here, DC viability ranged from 40% to 90% depending on the method of transfection or transduction employed and was lowest using the cDNA electroporation protocol. Therefore, it is possible that antigen released by dead cells may have



contributed, in part, to presentation of epitopes by surviving DCs. For the observed recognition of epitopes by CD4<sup>+</sup> T cells, this phenomena may have contributed significantly because the exogenous pathway is preferred for MHC class II-restricted antigen processing and presentation. However, it is unlikely that this phenomena contributed significantly to cross-presentation. In a previous report, we did not detect any cross-presentation when gp100 was pulsed on DCs as an intact protein or from melanoma cell lysates.<sup>6</sup> Furthermore, in several other reports, efficient cross-presentation of NY-ESO-1 was only observed using full-length protein formulated as antigen-antibody immune complexes<sup>7</sup> or with ISCOMA-TRIX adjuvant,<sup>7,8</sup> not using soluble protein alone.

Presentation of gp100 epitopes in the context of MHC class II molecules after genetic modification of DCs may result from a unique structural characteristic. This glycoprotein, and the other melanosomal membrane glycoproteins, tyrosinase, TRP-1, and TRP-2, contain a dileucine-based sorting motif, the melanosomal transport signal (MTS).<sup>39</sup> This hexapeptide sequence normally directs melanosomal proteins to the melanosome, but it may also facilitate entry of these proteins into the endocytic pathway, on the basis of structural similarities between the endosome and melanosome.<sup>40</sup> The melanosomal transport signal-based targeting of intracellular gp100 to the endocytic pathway could explain its frequent presentation by MHC class II molecules, both in tumor cells and genetically modified DCs.

Lack of a clear distinction between viral and nonviral vectors in terms of gene transfer efficiency and expression of costimulatory molecules directed our focus to the role of the vector itself in determining how the cell degrades and processes the inserted antigen. Viral vectors seem to activate proteasomal processing and MHC class I presentation more efficiently than nonviral vectors. Although our results do not elucidate any cellular mechanisms involved, perhaps cofactors associated with cellular reactions to viral infection are crucial for effective antigen processing and presentation by DC. Also, it seems possible that rapid turnover of defective transgene products, prominent with viruses but not nonviral vectors, contributes to enhanced antigen processing.<sup>41</sup> To address this hypothesis, in a preliminary experiment, we first transduced DCs with Ad2-GFP and subsequently transfected these antigen-presenting cells with IVT RNA encoding gp100 (data not shown). In that experiment, no enhanced recognition by gp100 reactive CD8<sup>+</sup> or CD4<sup>+</sup> T cells was observed, but similar experiments should be repeated in the future to address this issue more thoroughly.

Our results support the use of DCs transduced with recombinant retroviral vectors for the development of new cancer immunotherapies. Although adenoviral vectors are associated with higher transduction efficiencies, the application of these vectors is severely limited by preexisting immunity and by rapidly developing anti-vector immune responses.<sup>42</sup> Despite lower transduction efficiencies with MMLV, this retroviral vector induced efficient presentation of both MHC class I and class II-restricted T-cell epitopes from both gp100 and NY-ESO-1. Since clinical evidence strongly suggests that CD8<sup>+</sup> T cells can mediate tumor regression,<sup>43</sup> MHC class I-restricted presentation of tumor-associated epitopes seems critical for any DC gene modification technique. In addition, MHC class II-restricted epitope presentation may be important for eliciting CD4<sup>+</sup> T cell help which may, in turn, enhance CTL responses.<sup>11,15,36</sup> Therefore, the broad spectrum of antigen presentation in retrovirally transduced DC and the low immunogenicity of the retroviral vector delineate this gene delivery method as a valuable tool both for expanding the spectrum of known tumor-associated antigens, and for potential clinical applications of DC-based anticancer immunotherapy.


## References

1. Banchereau J, Briere F, Caux C, et al. Immunobiology of dendritic cells. *Annu Rev Immunol* 2000;18:767–811. [PubMed: 10837075]

2. Shen Z, Reznikoff G, Dranoff G, et al. Cloned dendritic cells can present exogenous antigens on both MHC class I and class II molecules. *J Immunol* 1997;158:2723–2730. [PubMed: 9058806]
3. Larsson M, Fonteneau JF, Bhardwaj N. Dendritic cells resurrect antigens from dead cells. *Trends Immunol* 2001;22:141–148. [PubMed: 11286729]
4. Ludewig B, McCoy K, Pericin M, et al. Rapid peptide turnover and inefficient presentation of exogenous antigen critically limit the activation of self-reactive CTL by dendritic cells. *J Immunol* 2001;166:3678–3687. [PubMed: 11238607]
5. Melief CJ. Mini-review: regulation of cytotoxic T lymphocyte responses by dendritic cells: peaceful coexistence of cross-priming and direct priming? *Eur J Immunol* 2003;33:2645–2654. [PubMed: 14515248]
6. Parkhurst MR, DePan C, Riley JP, et al. Hybrids of dendritic cells and tumor cells generated by electrofusion simultaneously present immunodominant epitopes from multiple human tumor-associated antigens in the context of MHC class I and class II molecules. *J Immunol* 2003;170:5317–5325. [PubMed: 12734382]
7. Schnurr M, Chen Q, Shin A, et al. Tumor antigen processing and presentation depend critically on dendritic cell type and the mode of antigen delivery. *Blood* 2005;105:2465–2472. [PubMed: 15546948]
8. Maraskovsky E, Sjolander S, Drane DP, et al. NY-ESO-1 protein formulated in ISCOMATRIX adjuvant is a potent anticancer vaccine inducing both humoral and CD8+ t-cell-mediated immunity and protection against NY-ESO-1+ tumors. *Clin Cancer Res* 2004;10:2879–2890. [PubMed: 15102697]
9. Bonifaz L, Bonnyay D, Mahnke K, et al. Efficient targeting of protein antigen to the dendritic cell receptor DEC-205 in the steady state leads to antigen presentation on major histocompatibility complex class I products and peripheral CD8+ T cell tolerance. *J Exp Med* 2002;196:1627–1638. [PubMed: 12486105]
10. York IA, Goldberg AL, Mo XY, et al. Proteolysis and class I major histocompatibility complex antigen presentation. *Immunol Rev* 1999;172:49–66. [PubMed: 10631936]
11. Bonehill A, Heirman C, Tuyaerts S, et al. Messenger RNA-electroporated dendritic cells presenting MAGE-A3 simultaneously in HLA class I and class II molecules. *J Immunol* 2004;172:6649–6657. [PubMed: 15153480]
12. Van Tendeloo VF, Ponsaerts P, Lardon F, et al. Highly efficient gene delivery by mRNA electroporation in human hematopoietic cells: superiority to lipofection and passive pulsing of mRNA and to electroporation of plasmid cDNA for tumor antigen loading of dendritic cells. *Blood* 2001;98:49–56. [PubMed: 11418462]
13. Strobel I, Berchtold S, Gotze A, et al. Human dendritic cells transfected with either RNA or DNA encoding influenza matrix protein M1 differ in their ability to stimulate cytotoxic T lymphocytes. *Gene Ther* 2000;7:2028–2035. [PubMed: 11175315]
14. Linette GP, Shankara S, Longrich S, et al. In vitro priming with adenovirus/gp100 antigen-transduced dendritic cells reveals the epitope specificity of HLA-A\*0201-restricted CD8+ T cells in patients with melanoma. *J Immunol* 2000;164:3402–3412. [PubMed: 10706736]
15. Bonini C, Lee SP, Riddell SR, et al. Targeting antigen in mature dendritic cells for simultaneous stimulation of CD4+ and CD8+ T cells. *J Immunol* 2001;166:5250–5257. [PubMed: 11290810]
16. Lapointe R, Royal RE, Reeves ME, et al. Retrovirally transduced human dendritic cells can generate T cells recognizing multiple MHC class I and class II epitopes from the melanoma antigen glycoprotein 100. *J Immunol* 2001;167:4758–4764. [PubMed: 11591807]
17. Lizée G, Gonzales MI, Topalian SL. Lentivirus vector-mediated expression of tumor-associated epitopes by human antigen presenting cells. *Hum Gene Ther* 2004;15:393–404. [PubMed: 15053864]
18. Dietz AB, Vuk-Pavlovic S. High efficiency adenovirus-mediated gene transfer to human dendritic cells. *Blood* 1998;91:392–398. [PubMed: 9427691]
19. Zhong L, Granelli-Piperno A, Choi Y, et al. Recombinant adenovirus is an efficient and non-perturbing genetic vector for human dendritic cells. *Eur J Immunol* 1999;29:964–972. [PubMed: 10092101]


20. Roth MD, Cheng Q, Harui A, et al. Helper-dependent adenoviral vectors efficiently express transgenes in human dendritic cells but still stimulate antiviral immune responses. *J Immunol* 2002;169:4651–4656. [PubMed: 12370405]
21. Batchu RB, Moreno AM, Szmania S, et al. High-level expression of cancer/testis antigen NY-ESO-1 and human granulocyte-macrophage colony-stimulating factor in dendritic cells with a bicistronic retroviral vector. *Hum Gene Ther* 2003;14:1333–1345. [PubMed: 14503968]
22. Parkhurst MR, Riley JP, Robbins PF, et al. Induction of CD4+ Th1 lymphocytes that recognize known and novel class II MHC restricted epitopes from the melanoma antigen gp100 by stimulation with recombinant protein. *J Immunother* 2004;27:79–91. [PubMed: 14770079]
23. Bakker AB, Schreurs MW, Tafazzul G, et al. Identification of a novel peptide derived from the melanocyte-specific gp100 antigen as the dominant epitope recognized by an HLA-A2.1-restricted anti-melanoma CTL line. *Int J Cancer* 1995;62:97–102. [PubMed: 7541395]
24. Kawakami Y, Eliyahu S, Jennings C, et al. Recognition of multiple epitopes in the human melanoma antigen gp100 by tumor-infiltrating T lymphocytes associated with in vivo tumor regression. *J Immunol* 1995;154:3961–3968. [PubMed: 7706734]
25. Dudley ME, Wunderlich JR, Yang JC, et al. A phase I study of nonmyeloablative chemotherapy and adoptive transfer of autologous tumor antigen-specific T lymphocytes in patients with metastatic melanoma. *J Immunother* 2002;25:243–251. [PubMed: 12000866]
26. Khong HT, Rosenberg SA. Pre-existing immunity to tyrosinase-related protein (TRP)-2, a new TRP-2 isoform, and the NY-ESO-1 melanoma antigen in a patient with a dramatic response to immunotherapy. *J Immunol* 2002;168:951–956. [PubMed: 11777994]
27. Khong HT, Wang QJ, Rosenberg SA. Identification of multiple antigens recognized by tumor-infiltrating lymphocytes from a single patient: tumor escape by antigen loss and loss of MHC expression. *J Immunother* 2004;27:184–190. [PubMed: 15076135]
28. Touloukian CE, Leitner WW, Topalian SL, et al. Identification of a MHC class II-restricted human gp100 epitope using DR4-IE transgenic mice. *J Immunol* 2000;164:3535–3542. [PubMed: 10725708]
29. Bownds S, Tong-On P, Rosenberg SA, et al. Induction of tumor-reactive cytotoxic T-lymphocytes using a peptide from NY-ESO-1 modified at the carboxy-terminus to enhance HLA-A2.1 binding affinity and stability in solution. *J Immunother* 2001;24:1–9. [PubMed: 11211143]
30. Lu Y, Boss JM, Hu SX, et al. Apoptosis-independent retinoblastoma protein rescue of HLA class II messenger RNA IFN-gamma inducibility in non-small cell lung carcinoma cells. Lack of surface class II expression associated with a specific defect in HLA-DRA induction. *J Immunol* 1996;156:2495–2502. [PubMed: 8786310]
31. Sallusto F, Lanzavecchia A. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med* 1994;179:1109–1118. [PubMed: 8145033]
32. Reeves ME, Royal RE, Lam JS, et al. Retroviral transduction of human dendritic cells with a tumor-associated antigen gene. *Cancer Res* 1996;56:5672–5677. [PubMed: 8971174]
33. Kawakami Y, Eliyahu S, Delgado CH, et al. Identification of a human melanoma antigen recognized by tumor-infiltrating lymphocytes associated with in vivo tumor rejection. *Proc Natl Acad Sci USA* 1994;91:6458–6462. [PubMed: 8022805]
34. Wang RF, Johnston SL, Zeng G, et al. A breast and melanoma-shared tumor antigen: T cell responses to antigenic peptides translated from different open reading frames. *J Immunol* 1998;161:3598–3606. [PubMed: 9759882]
35. Boczkowski D, Nair SK, Snyder D, et al. Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J Exp Med* 1996;184:465–472. [PubMed: 8760800]
36. Zhao Y, Boczkowski D, Nair SK, et al. Inhibition of invariant chain expression in dendritic cells presenting endogenous antigens stimulates CD4+ T-cell responses and tumor immunity. *Blood* 2003;102:4137–4142. [PubMed: 12920018]
37. Goldberg AL, Cascio P, Saric T, et al. The importance of the proteasome and subsequent proteolytic steps in the generation of antigenic peptides. *Mol Immunol* 2002;39:147–164. [PubMed: 12200047]
38. Brodsky FM. Antigen processing and presentation: close encounters in the endocytic pathway. *Trends Cell Biol* 1992;2:109–115. [PubMed: 14732015]

39. Wang S, Bartido S, Yang G, et al. A role for a melanosome transport signal in accessing the MHC class II presentation pathway and in eliciting CD4+ T cell responses. *J Immunol* 1999;163:5820–5826. [PubMed: 10570265]
40. Calvo PA, Frank DW, Bieler BM, et al. A cytoplasmic sequence in human tyrosinase defines a second class of di-leucine-based sorting signals for late endosomal and lysosomal delivery. *J Biol Chem* 1999;274:12780–12789. [PubMed: 10212263]
41. Yewdell J. To DRiP or not to DRiP: generating peptide ligands for MHC class I molecules from biosynthesized proteins. *Mol Immunol* 2002;39:139–146. [PubMed: 12200046]
42. Sumida SM, Truitt DM, Kishko MG, et al. Neutralizing antibodies and CD8+ T lymphocytes both contribute to immunity to adenovirus serotype 5 vaccine vectors. *J Virol* 2004;78:2666–2673. [PubMed: 14990686]
43. Dudley ME, Wunderlich JR, Robbins PF, et al. Cancer regression and autoimmunity in patients after clonal repopulation with antitumor lymphocytes. *Science* 2002;298:850–854. [PubMed: 12242449]



**FIGURE 1.**

Gene transfer efficiencies in DCs transfected or transduced with vectors encoding eGFP. For each panel, gene expression was evaluated by FACS ~24 hours after genetic modification. Percentages indicate the percent of cells gated through M1, and reported MFI values are for M1-gated cells. A, Monocyte-derived DCs from 2 different donors were electroporated with IVT RNA or cDNA encoding eGFP. B, Monocyte-derived DCs from 2 different donors were electroporated with IVT RNA or cDNA encoding eGFP or were transduced with an Ad2 vector encoding this protein. C, Monocyte-derived DCs from 1 donor were transduced with an Ad2 vector encoding eGFP, and CD34<sup>+</sup> HSC-derived DCs from the same donor were transduced with a retroviral vector encoding this protein.



**FIGURE 2.**

DC phenotype after genetic modification. Monocyte-derived DCs were electroporated with IVT RNA or cDNA encoding gp100 or were transduced with an adenoviral vector encoding this protein. Approximately, 16 hours after gene modification, maturation cytokines were added as indicated and described in the Materials and Methods section, and approximately 24 hours later, DCs were stained with fluorescein isothiocyanate- (x-axis; FL1-H) and PE- (y-axis; FL2-H) conjugated mAbs as indicated and analyzed by FACS.

**TABLE 1**  
 Characteristics of Antigen Reactive T-cell Lines and Clones

T-cell Clone/Line	CD4/CD8	HLA Restriction Element	Peptide Specificity
TIL 1200 (line)	CD8	HLA-A*0201	gp100:154–162
CL70 (clone)	CD8	HLA-A*0201	gp100:154–162
RB154 (line)	CD8	HLA-A*0201	gp100:154–162
CK3H6 (clone)	CD8	HLA-A*0201	gp100:209–217
L2D8 (clone)	CD8	HLA-A*0201	gp100:209–217
JR1E2 (clone)	CD8	HLA-A*0201	gp100:280–288
JR1A4 (clone)	CD8	HLA-A*0201	gp100:280–288
HT1D11 (clone)	CD8	HLA-A*0201	gp100:280–288
BR-B8 (line)	CD4	HLA-DR $\beta$ 1*0401	gp100:44–59
B104 (line)	CD4	HLA-DR $\beta$ 1*0701	gp100:170–190
JE-D2 (line)	CD4	HLA-DR $\beta$ 1*0701	gp100:420–435
TH1F2L (line)	CD8	HLA-A*0201	NY-ESO-1:157–165
J H1 (line)	CD8	HLA-A*0201	NY-ESO-1:157–165
J H6 (line)	CD8	HLA-A*0201	NY-ESO-1:157–165
M-8 (line)	CD8	HLA-A*0201	NY-ESO-1:157–165
SG6 (line)	CD4	HLA-DP $\beta$ 1*0401	NY-ESO-1:161–180

TABLE 2

Recognition of DCs Genetically Modified With gp100 cDNA, IVT RNA, or Ad2 by HLA-A\*0201 and HLA-DRβ1\*0401 Restricted, gp100-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	Genetic Modification	% GFP Positive <sup>‡</sup>	MTF <sup>§</sup>	CK3H6 (A2)	BR-B8 (DR4)
Donor 3 DC	A2*DR4 <sup>+</sup>	—	gp100 cDNA	—	—	48	1814//
Donor 3 DC	A2*DR4 <sup>+</sup>	—	GFP cDNA	36	586	42	37
Donor 3 DC	A2*DR4 <sup>+</sup>	—	gp100 IVT RNA	—	—	57	>2000
Donor 3 DC	A2*DR4 <sup>+</sup>	—	GFP IVT RNA	96	130	48	62
Donor 3 DC	A2*DR4 <sup>+</sup>	—	gp100 Ad2	—	—	1080	>2000
Donor 3 DC	A2*DR4 <sup>+</sup>	—	GFP Ad2	96	1675	28	27
Donor 4 DC	A2*DR4 <sup>+</sup>	—	gp100 cDNA	—	—	34	569
Donor 4 DC	A2*DR4 <sup>+</sup>	—	gp100 IVT RNA	—	—	34	>2000
Donor 4 DC	A2*DR4 <sup>+</sup>	—	GFP IVT RNA	61	50	51	70
Donor 4 DC	A2*DR4 <sup>+</sup>	—	gp100 Ad2	—	—	>2000	>2000
Donor 4 DC	A2*DR4 <sup>+</sup>	—	GFP Ad2	90	556	35	53
T2 cells	A2 <sup>+</sup>	HBVc:18–26(23Y)	—	—	—	44	—
T2 cells	A2 <sup>+</sup>	gp100:209–217	—	—	—	>2000	—
EBV-B	DR4 <sup>+</sup>	IgK:188–201	—	—	—	—	35
EBV-B	DR4 <sup>+</sup>	gp100:170–190	—	—	—	—	328

\* IFN $\gamma$  secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201 and/or HLA-DRβ1\*0401 with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50 mg/mL gp100:44–59 or Influenza HA:306–324, and T2 cells were preincubated with 1  $\mu$ g/mL gp100:209–217 or HBVc:18–26(23Y).

<sup>‡</sup> Percentage of cells expressing GFP by FACS (Fig. 1B).

<sup>§</sup> MFI of GFP-positive cells by FACS (Fig. 1B).

// Underlined values indicate that IFN $\gamma$  secretion in response to DCs genetically modified with gp100 vectors, or peptide-loaded T2 cells or EBV-B cells, was  $\geq$  50 pg/mL and at least twice background with relevant control target cells.



TABLE 3

Recognition of DCs Genetically Modified With gp100 cDNA or Ad2 by HLA-A\*0201 (A2), HLA-DRβ1\*0401 (DR4), and HLA-DRβ1\*0701 (DR7) restricted, gp100-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	Genetic Modification	TIL1200 (A2)	CK3H6 (A2)	JRIE2 (A2)	B8 (DR4)	B104 (DR7)	D2 (DR7)	JE- (DR7)
Donor 3 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	GFP Ad2	68	25	25	22	23	29	29
Donor 3 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	gp100 Ad2	<u>3677</u> <sup>‡</sup>	<u>6445</u>	<u>6445</u>	<u>2033</u>	≥10000	4476	4476
Donor 3 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	GFP cDNA	105	28	24	22	25	28	28
Donor 3 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	gp100 cDNA	177	92	29	62	486	34	34
Donor 4 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	GFP Ad2	84	23	17	17	21	17	17
Donor 4 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	gp100 Ad2	<u>1162</u>	<u>2317</u>	<u>1218</u>	<u>1814</u>	≥10000	847	847
Donor 4 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	GFP cDNA	67	24	21	18	24	18	18
Donor 4 DC	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	gp100 cDNA	<u>124</u>	<u>201</u>	<u>48</u>	<u>74</u>	<u>238</u>	<u>268</u>	<u>268</u>
media	—	—	—	112	21	19	18	22	19	19
T2 cells	A2 <sup>+</sup>	HBVc:18–26(23Y)	—	162	37	28	—	—	—	—
T2 cells	A2 <sup>+</sup>	gp100:154–162	—	7386	—	—	—	—	—	—
T2 cells	A2 <sup>+</sup>	gp100:209–217	—	—	—	—	—	—	—	—
T2 cells	A2 <sup>+</sup>	gp100:280–288	—	—	≥10000	9214	—	—	—	—
EBV-B	DR4 <sup>+</sup>	HA:306–324	—	—	—	—	—	—	—	—
EBV-B	DR4 <sup>+</sup>	gp100:44–59	—	—	—	—	15	—	—	—
EBV-B	DR7 <sup>+</sup>	IgK:188–201	—	—	—	—	<u>1279</u>	—	—	—
EBV-B	DR7 <sup>+</sup>	gp100:170–190	—	—	—	—	—	65	19	19
EBV-B	DR7 <sup>+</sup>	gp100:420–435	—	—	—	—	—	8583	—	—
F002Rmel CIITA	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	VSV-GFP	81	21	27	17	17	22	22
F002Rmel CIITA	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	VSV-gp100	<u>7321</u>	<u>8668</u>	≥10000	17	≥10000	≥10000	≥10000
1088mel CIITA	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	—	7640	<u>9943</u>	≥10000	<u>824</u>	16	20	20
888mel	A2 <sup>+</sup> DR4 <sup>+</sup> DR7 <sup>+</sup>	—	gp100 cDNA	147	16	18	20	20	22	22

\* IFN $\gamma$  secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201, HLA-DRβ1\*0401, and/or HLA-DRβ1\*0701 with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50  $\mu$ g/mL MHC class II-restricted peptides, and T2 cells were preincubated with 1  $\mu$ g/mL MHC class I-restricted peptides.

<sup>‡</sup> Underlined values indicate that IFN $\gamma$  secretion in response to DCs genetically modified with gp100 vectors, peptide-loaded T2 cells or EBV-B cells, or melanoma cell lines was  $\geq$  50 pg/mL and at least twice background with relevant control target cells.

TABLE 4

Recognition of DCs Transfected with gp100 cDNA in the Presence or Absence of Maturation Cytokines by HLA-A\*0201 (A2), HLA-DRβ1\*0401 (DR4), and HLA-DRβ1\*0701 (DR7) Restricted, gp100-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	Genetic Modification	Maturation Cytokines <sup>‡</sup>	CK3H6 (A2) Exp. 1	JR1E2 (A2) Exp. 1	B104 (DR7) Exp. 1	CL70 (A2) Exp. 2	L2D8 (A2) Exp. 2	BR-B8 (DR4) Exp. 2
Donor 5 DC	A2+ DR7+	—	gp100 cDNA	None	72	76	81	—	—	—
Donor 5 DC	A2+ DR7+	—	GFP cDNA	None	104	71	834 <sup>§</sup>	—	—	—
Donor 5 DC	A2+ DR7+	—	gp100 cDNA	TNFα + PGE <sub>2</sub>	75	67	76	—	—	—
Donor 5 DC	A2+ DR7+	—	GFP cDNA	TNFα + PGE <sub>2</sub>	76	76	71	—	—	—
Donor 6 DC	A2+ DR7+	—	gp100 cDNA	None	72	71	78	—	—	—
Donor 6 DC	A2+ DR7+	—	GFP cDNA	None	<u>212</u>	69	≥ 2000	—	—	—
Donor 6 DC	A2+ DR7+	—	gp100 cDNA	TNFα + PGE <sub>2</sub>	67	68	70	—	—	—
Donor 6 DC	A2+ DR7+	—	GFP cDNA	TNFα + PGE <sub>2</sub>	78	68	77	—	—	—
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	None	—	—	—	20	84	4605
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	None	—	—	—	88	88	250
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	TNFα + PGE <sub>2</sub>	—	—	—	19	58	534
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	TNFα + PGE <sub>2</sub>	—	—	—	130	98	238
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	CD40L+LPS	—	—	—	205	<u>1285</u>	3527
Donor 7 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	CD40L+LPS	—	—	—	251	390	524
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	None	—	—	—	22	263	11737
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	None	—	—	—	33	162	341
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	TNFα + PGE <sub>2</sub>	—	—	—	22	246	2324
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	TNFα + PGE <sub>2</sub>	—	—	—	69	158	313
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	gp100 cDNA	CD40L+LPS	—	—	—	54	603	4973
Donor 8 DC	A2 <sup>†</sup> DR4 <sup>†</sup>	—	GFP cDNA	CD40L+LPS	—	—	—	407	309	381
T2 or EBV-B	T cell matched	control	—	—	65	65	70	18	22	144
T2 or EBV-B	T cell matched	relevant	—	—	≥ 2000	≥ 2000	≥ 2000	<u>3981</u>	<u>11662</u>	<u>6714</u>
F002Rmel CIITA	A2 <sup>†</sup> DR4 <sup>†</sup> DR7 <sup>†</sup>	—	VSV-GFP	—	70	76	93	—	—	—
F002Rmel CIITA	A2 <sup>†</sup> DR4 <sup>†</sup> DR7 <sup>†</sup>	—	VSV-gp100	—	<u>1676</u>	≥ 2000	≥ 2000	—	—	—
526mel CIITA	A2 <sup>†</sup> DR4 <sup>†</sup> DR7 <sup>†</sup>	—	—	—	<u>1403</u>	> 2000	68	—	—	—

\* IFNγ secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201, HLA-DRβ1\*0401, and/or HLA-DRβ1\*0701 with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50 μg/mL MHC class II-restricted peptides, and T2 cells were preincubated with 1 μg/mL MHC class I-restricted peptides. Control peptides were the same as those in Tables 2 and 3, and relevant peptides were those indicated in Table 1 for each specific T-cell line.

<sup>‡</sup> DCs were matured with the indicated cytokines as described in the Materials and Methods section.

<sup>§</sup> Underlined values indicate that IFNγ secretion in response to DCs genetically modified with gp100 vectors, peptide-loaded T2 cells or EBV-B cells, or melanoma cell lines was ≥ 50 pg/mL and at least twice background with relevant control target cells.

**TABLE 5**  
 Recognition of DCs Transfected With gp100 IVT RNA in the Presence or Absence of Maturation cytokines by HLA-A\*0201 (A2) and HLA-DRβ1\*0401 (DR4) Restricted, gp100-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	Genetic Modification	Maturation Cytokines <sup>‡</sup>	CL70 (A2)	L2D8 (A2)	HTIE11 (A2)	B8 (DR4)	BR- B8 (DR4)
Donor 9 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	None	12	38	12	12	<u>49904</u> <sup>§</sup>
Donor 9 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	None	9	27	17	17	147
Donor 9 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	TNFα + PGE <sub>2</sub>	25	51	17	17	<u>24572</u>
Donor 9 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	TNFα + PGE <sub>2</sub>	16	21	18	18	51
Donor 9 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	CD40L+LPS	15	<u>144</u>	14	14	<u>19725</u>
Donor 9 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	CD40L+LPS	24	41	28	28	72
Donor 10 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	None	36	46	28	28	<u>57778</u>
Donor 10 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	None	26	29	27	27	209
Donor 10 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	TNFα + PGE <sub>2</sub>	23	57	33	33	<u>24617</u>
Donor 10 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	TNFα + PGE <sub>2</sub>	23	22	28	28	57
Donor 10 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	CD40L+LPS	38	<u>105</u>	34	34	<u>22456</u>
Donor 10 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	CD40L+LPS	25	26	29	29	132
Donor 11 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	None	18	<u>191</u>	9	9	<u>39000</u>
Donor 11 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	None	12	26	15	15	48
Donor 11 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	TNFα + PGE <sub>2</sub>	28	89	13	13	<u>12721</u>
Donor 11 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	TNFα + PGE <sub>2</sub>	12	21	21	21	53
Donor 11 DC	A2*DR4 <sup>+</sup>	—	gp100 RNA	CD40L+LPS	8	42	15	15	<u>36278</u>
Donor 11 DC	A2*DR4 <sup>+</sup>	—	GFP RNA	CD40L+LPS	17	22	28	28	39
T2 or EBV-B	T cell matched	control	—	—	36	84	791	82	82
T2 or EBV-B	T cell matched	relevant	—	—	<u>7450</u>	<u>45424</u>	<u>9156</u>	<u>5604</u>	<u>5604</u>

\* IFN $\gamma$  secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201 and/or HLA-DRβ1\*0401, with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50 μg/mL MHC class II-restricted peptides, and T2 cells were preincubated with 1 μg/mL MHC class I-restricted peptides. Control peptides were the same as those in Tables 2 and 3, and relevant peptides were those indicated in Table 1 for each specific T-cell line.

<sup>‡</sup> DCs were matured with the indicated cytokines as described in the Materials and Methods section.

<sup>§</sup> Underlined values indicate that IFN $\gamma$  secretion in response to DCs genetically modified with gp100 vectors, peptide-loaded T2 cells or EBV-B cells, or melanoma cell lines was  $\geq$  50 pg/mL and at least twice background with relevant control target cells.

**TABLE 6**  
 Recognition of DCs Transfected with NY-ESO-1 IVT RNA in the Presence or Absence of Maturation Cytokines by HLA-A\*0201 (A2) and HLA-DPβ1\*0401 (DP4) Restricted, NY-ESO-1-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	Genetic Modification	Maturation Cytokines <sup>‡</sup>	TH1F2L (A2) Exp. 1	SG6 (DP4) Exp. 1	J-H1 (A2) Exp. 2	M8 (A2) Exp. 2	SG6 (DP4) Exp. 2
Donor 12 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	377 <sup>§</sup>	14	—	—	—
Donor 12 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	None	52	0	—	—	—
Donor 13 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	279	13	—	—	—
Donor 13 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	None	86	5	—	—	—
Donor 14 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	897	13	—	—	—
Donor 14 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	None	215	5	—	—	—
Donor 15 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	—	—	14	28	17
Donor 15 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	None	—	—	14	15	18
Donor 15 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	CD40L+LPS	—	—	166	267	605
Donor 15 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	CD40L+LPS	—	—	15	16	17
Donor 16 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	—	—	16	63	19
Donor 16 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	None	—	—	18	33	17
Donor 16 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	None	—	—	32	81	211
Donor 16 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	GFP RNA	CD40L+LPS	—	—	24	56	74
Donor 16 DC	A2 <sup>+</sup> DP4 <sup>+</sup>	—	ESO RNA	CD40L+LPS	—	—	17	14	—
T2 cells	A2 <sup>+</sup>	HBVc:18–26(23Y)	GFP RNA	—	388	—	—	—	—
T2 cells	A2 <sup>+</sup>	ESO:157–165	—	—	>2000	—	—	—	—
EBV-B	DP4 <sup>+</sup>	HA:306–324	—	—	—	—	—	—	62
EBV-B	DP4 <sup>+</sup>	ESO:161–180	—	—	—	—	—	—	1366
624mel	A2 <sup>+</sup> DP4 <sup>+</sup>	—	—	—	1766	21	—	—	—
888mel	A2 <sup>+</sup> DP4 <sup>+</sup>	—	—	—	39	9	—	—	—

\* IFN $\gamma$  secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201 and/or HLA-DPβ1\*0401 with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50 μg/mL MHC class II-restricted peptides, and T2 cells were preincubated with 1 μg/mL MHC class I-restricted peptides.

<sup>‡</sup> DCs were matured with the indicated cytokines as described in the Materials and Methods section.

<sup>§</sup> Underlined values indicate that IFN $\gamma$  secretion in response to DCs genetically modified with NY-ESO-1 vectors, peptide-loaded T2 cells or EBV-B cells, or melanoma cell lines was  $\geq$  50 pg/mL and at least twice background with relevant control target cells.

**TABLE 7**  
 Recognition of DCs Genetically Modified with Retroviral Vectors Encoding gp100 or NY-ESO-1 by HLA-A\*0201 (A2) and HLA-DRβ1\*0701 (DR7) Restricted, Antigen-reactive T-cell Populations\*

Target Cells	HLA Expression	Preloaded Peptide <sup>†</sup>	MMLV Modification	RB-154 (A2)	L2D8 (A2)	HT1E11 (A2)	B104 (DR7)	D2 (DR7)	JE-DR7	JH1 (A2)	M8 (A2)
Donor 17 HSC-DC	A2 <sup>+</sup> DR7 <sup>+</sup>	—	gp100	>1000 <sup>‡</sup>	>1000	783	745	>1000	—	—	—
Donor 17 HSC-DC	A2 <sup>+</sup> DR7 <sup>+</sup>	—	NY-ESO-1	—	—	—	—	—	—	>1000	287
Donor 17 HSC-DC	A2 <sup>+</sup> DR7 <sup>+</sup>	—	GFP	29	19	272	30	24	—	19	30
T2 cells	A2 <sup>+</sup>	HBVc:18–26(23Y)	—	22	16	25	—	—	—	17	17
T2 cells	A2 <sup>+</sup>	gp100:154–162	—	>1000	—	—	—	—	—	—	—
T2 cells	A2 <sup>+</sup>	gp100:209–217	—	—	>1000	—	—	—	—	—	—
T2 cells	A2 <sup>+</sup>	gp100:280–288	—	—	—	>1000	—	—	—	—	—
T2 cells	A2 <sup>+</sup>	ESO:157–165	—	—	—	—	—	—	—	>1000	>1000
EBV-B	DR7 <sup>+</sup>	lgK:188–201	—	—	—	—	15	16	—	—	—
EBV-B	DR7 <sup>+</sup>	gp100:170–190	—	—	—	—	>1000	>1000	—	—	—
EBV-B	DR7 <sup>+</sup>	gp100:420–435	—	—	—	—	—	—	—	—	—
F002Rmel CIITA	A2 <sup>+</sup> DR7 <sup>+</sup>	—	GFP	16	15	18	14	60	—	—	—
624mel CIITA	A2 <sup>+</sup> DR7 <sup>+</sup>	—	—	588	>1000	26	445	534	—	76	198
526mel (ESO <sup>-</sup> )	A2 <sup>+</sup> DR7 <sup>-</sup>	—	—	>1000	>1000	27	18	16	—	15	14

\* IFN $\gamma$  secretion (pg/mL) in 20 h coculture supernatants of target cells expressing HLA-A\*0201 and/or HLA-DRβ1\*0701 with T lymphocytes.

<sup>†</sup> EBV-B cells were preincubated with 50  $\mu$ g/mL MHC class II-restricted peptides, and T2 cells were preincubated with 1  $\mu$ g/mL MHC class I-restricted peptides.

<sup>‡</sup> Underlined values indicate that IFN $\gamma$  secretion in response to DCs genetically modified with gp100 or NY-ESO-1 vectors, peptide-loaded T2 cells or EBV-B cells, or melanoma cell lines was  $\geq$  50 pg/mL and at least twice background with relevant control target cells.