



Published in final edited form as:

*AIDS Behav.* 2017 April ; 21(4): 1138–1148. doi:10.1007/s10461-016-1584-6.

## Prevalence and predictors of substance use disorders among HIV care enrollees in the United States

**Bryan Hartzler<sup>1</sup>, Julia C. Dombrowski<sup>2</sup>, Heidi M. Crane<sup>2</sup>, Joseph J. Eron<sup>3,4</sup>, Elvin H. Geng<sup>5</sup>, W. Christopher Mathews<sup>6</sup>, Kenneth H. Mayer<sup>7,8</sup>, Richard D. Moore<sup>9,10,11</sup>, Michael J. Mugavero<sup>12</sup>, Sonia Napravnik<sup>3</sup>, Benigno Rodriguez<sup>13</sup>, and Dennis M. Donovan<sup>1,14</sup>**

<sup>1</sup>Alcohol & Drug Abuse Institute, University of Washington, Seattle WA

<sup>2</sup>Division of Allergy and Infectious Disease, University of Washington, Seattle WA

<sup>3</sup>Department of Medicine, University of North Carolina, Chapel Hill NC

<sup>4</sup>Department of Epidemiology, University of North Carolina, Chapel Hill NC

<sup>5</sup>School of Medicine, University of California, San Francisco CA

<sup>6</sup>Department of Medicine, University of California, San Diego CA

<sup>7</sup>School of Medicine, Harvard University, Boston MA

<sup>8</sup>School of Public Health, Harvard University, Boston MA

<sup>9</sup>Department of Medicine, Johns Hopkins University, Baltimore MD

<sup>10</sup>Department of Epidemiology, Johns Hopkins University, Baltimore MD

<sup>11</sup>Center for Global Health, Johns Hopkins University, Baltimore MD

<sup>12</sup>Department of Medicine, University of Alabama, Birmingham AL

<sup>13</sup>Department of Medicine, Case Western Reserve University, Cleveland OH

<sup>14</sup>Psychiatry and Behavioral Sciences, University of Washington, Seattle WA

### Abstract

Prior efforts to estimate U.S. prevalence of substance use disorders (SUDs) in HIV care have been undermined by caveats common to single-site trials. The current work reports on a cohort of 10,652 HIV-positive adults linked to care at seven sites, with available patient data including geography, demography, and risk factor indices, and with substance-specific SUDs identified via

---

Corresponding author: Bryan Hartzler, Ph.D., Alcohol & Drug Abuse Institute, University of Washington, Box 354805, 1107 NE 45<sup>th</sup> Street, Suite 120, Seattle, WA 98105-4631, U.S.A., 206.543.8369, 206.543.5473 (fax), hartzb@u.washington.edu.

**Conflicts of Interest:** Julia C. Dombrowski has conducted STD clinical research unrelated to this work supported by grants to the University of Washington from Genentech, ELITech, Melinta Therapeutics, Curatek Pharmaceuticals, Quidel, and Hologic. Among the authorship group, no other conflicts of interest were declared.

**Disclaimer:** The content of this report is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Health.

**Ethical Approval:** All procedures involving human participants were in accordance with institutional review boards at the CNICS-affiliate universities, and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

**Informed Consent:** Informed consent was obtained from all individuals during their initial enrollment in CNICS.

self-report instruments with validated diagnostic thresholds. Generalized estimating equations also tested patient indices as SUD predictors. Findings were: 1) a 48% SUD prevalence rate (between-site range of 21–71%), with 20% of the sample evidencing polysubstance use disorder; 2) substance-specific SUD rates of 31% for marijuana, 19% alcohol, 13% methamphetamine, 11% cocaine, and 4% opiate; and 3) emergence of younger age and male gender as robust SUD predictors. Findings suggest high rates at which SUDs occur among patients at these urban HIV care sites, detail substance-specific SUD rates, and identify at-risk patient subgroups.

## Resumen

Los esfuerzos previos para estimar la prevalencia de los trastornos por uso de sustancias (TUS) de Estados Unidos en la atención del VIH han sido socavados por los problemas comunes de la investigación realizada en un solo sitio. Este documento informa sobre un estudio de una cohorte de 10,652 adultos con VIH que reciben atención en siete sitios, con los datos del paciente disponibles sobre la geografía, la demografía y los índices de factores de riesgo, y con trastornos por uso de sustancias para sustancias específicas identificadas con los instrumentos de autoinforme con umbrales de diagnóstico que han sido validados. Ecuaciones de estimación generalizadas también evaluaron los índices de pacientes como predictores de TUS. Los resultados fueron: 1) una tasa de prevalencia de TUS de 47% (entre-ubicación gama de 21 a 71%), con 20% de la muestra que demuestra un trastorno que implica múltiples sustancias; 2) las tasas SUD por sustancia específica de 31% para la marihuana, 19% para el alcohol, 13% de la metanfetamina, 11% de la cocaína, y 4% de los opiáceos; y 3) el surgimiento de menor edad y el sexo masculino como predictores robustos de los trastornos por uso de sustancias. Los resultados sugieren que los pacientes en las clínicas urbanas VIH tienen altas tasas de TUS, describen las tasas de sustancias específicas, e identifican subgrupos de pacientes en situación de riesgo.

## Keywords

HIV care settings; substance use disorders; patient demography; United States

---

## Introduction

Prior reports suggest 80% of HIV+ Americans effectively engaged in care reach viral suppression (1, 2), though consequent optimism is tempered as this applies to a subset of those living with HIV. Estimates suggest 14–21% of HIV+ Americans are unaware of their status, and up to half of those linked to care ineffectively engage in services (3). While health policies, delivery systems, and providers may all influence patient engagement in HIV care (4), clinical attributes of the HIV+ population also play a key role. One such attribute is substance use disorders (SUDs), defined by a set of adverse physiological and behavioral consequences (i.e., tolerance, withdrawal, role failure, craving, unsuccessful quitting). Increased care access among persons with SUD due to the Affordable Care Act (5, 6) and strong inter-rater reliability for the singular DSM-V conceptualization of SUD (7) are recent developments suggesting this as an opportune time for reporting SUD prevalence estimates among HIV care enrollees.

From a public health perspective, SUDs and HIV comprise a health syndemic for which deleterious impacts are observed throughout the HIV Care Continuum (8). With regard to HIV transmission, effectiveness of universal *test-and-treat* approaches is diminished among persons with SUD (9, 10). Post-diagnosis linkage to care occurs less often among persons with SUD (11), likely due to a complex mix of system, provider, and patient factors (12). Even after care linkage, persons with SUD visit clinic inconsistently, initiate antiretroviral medication at later stages of illness, and display poor adherence (13–18). Though definitions of HIV care retention may vary (4, 19), research suggests the presence of an SUD has a detrimental influence (20–22). Comparatively less effective HIV diagnosis, care linkage, antiretroviral medication adherence, and retention in services would be expected to diminish likelihood of eventual viral suppression; however, those with SUD respond no differently to antiretroviral medication when regimens are followed (23). Further, adherence and consequent viral suppression are achievable if appropriate health services are in place (24, 25). Thus, clarity of the scope of SUD prevalence may inform service needs of substance-using populations along the HIV Care Continuum.

To date, nearly all efforts to estimate SUD prevalence in U.S.-based HIV care have been limited to single-site trial data. Inherent geographic isolation and selection bias common to such trials contribute to diverse estimates, ranging from 21–65% (26–45). Caveats are compounded by a lack of diagnostic specificity, as trial sample sizes have typically precluded substance-specific examination even as individual substances of abuse pose differential risk in HIV transmission, course, and outcome (15, 46–49). Alternative data sources, if generated via continuous and coordinated multisite collection, may address apparent gaps in extant literature to offer more comprehensive, detailed estimation of SUD prevalence. Patient geography and demography (i.e., age, gender, race/ethnicity, sexual identity) predict both substance use among HIV care enrollees (50–52) and SUD rates in community sampling (53, 54), and thereby merit inclusion in such analytic work. The Center for AIDS Research Network of Integrated Clinical Systems (CNICS) (55) is a U.S.-based data source offering a multi-regional, continuous cohort of HIV care enrollees, with demographic information and capacity to delineate substance-specific SUDs.

Aims of the current work were to report prevalence estimates for SUDs among HIV care enrollees, and identify demographic predictors that increase likelihood that an SUD is present. Corresponding examination of a 10,000+ cohort, drawn from seven urban university-affiliated care centers, enabled derivation of multiregional, substance-specific SUD prevalence estimates. Patient geography, demography, and HIV transmission risk factors were explored as potential SUD predictors, in effort to identify patient subgroups at greater consequent risk to prematurely disengage from HIV care. Increased understanding of the scope of the SUD-HIV syndemic may spur implementation of addiction-focused services that respond to needs of HIV care enrollees.

## Methods

### Data Sources

Prevalence of SUDs was examined via CNICS (55), a network initiated in 1995 for longitudinal observation of patients enrolled at its affiliated sites. Continual integration of

clinical data from these sites affords opportunity to explore questions not readily addressed by sampling biases and surrogate endpoints inherent in clinical trials (56). Available data include information documented by staff at clinic visits, standard HIV-related risk factor data obtained at enrollment, medication/laboratory data from electronic medical records, and patient-reported outcomes collected since 2007 by personal computer or touch-screen tablet (55). Approval of a university-based institutional review board (IRB) at each CNICS site governs data collection, and the University of Washington IRB approved analytic procedures with de-identified data provided to the principal investigator by the CNICS Data Management Core.

The current work is restricted to patient demography/background indices as well as the patient-reported Alcohol Use Disorders Identification Test-Version C [AUDIT-C(57)] and Alcohol, Smoking, and Substance Involvement Test [ASSIST(58)]. Demography/background indices were patients' CNICS enrollment site, age, gender, ethnicity, race, transgender status, and sexual orientation (the latter collected among recent enrollees at three CNICS sites). Two HIV risk factor indices were examined, based on clinical notation kept at CNICS sites: a history of injection drug use (IDU), and a history of men who have sex with men (MSM).

The AUDIT-C is governed by a 'past 12 months' reporting interval, for which a summary score is generated for which a diagnostic threshold identified alcohol use disorder. In a national sample, Dawson et.al (59) demonstrated its utility as a diagnostic screening instrument with 84% sensitivity and 83% specificity for DSM-V diagnosis derived in clinical interviews. The CNICS assessment battery limits ASSIST measurement to four drug categories: cocaine, marijuana, methamphetamine, and non-prescription opioids. The ASSIST is governed by a 'past 3 months' reporting interval, and results in an 'involvement score' for each drug category. Humeniuk et.al (58) documented cross-cultural utility of the ASSIST as a diagnostic screening instrument, with the four CNICS-relevant involvement score thresholds demonstrating 91–97% sensitivity and 87–96% specificity for DSM-IV diagnoses of the four corresponding SUDs in clinical interviews.

### SUD Identifications

Applying AUDIT-C (59) and ASSIST (58) diagnostic thresholds, cases were identified for five substance-specific SUDs (hereafter referenced as *alcohol UD*), *cocaine UD*), *marijuana UD*), *methamphetamine UD*, and *opioid UD*). Persons for whom the AUDIT-C summary score and four ASSIST involvement scores failed to reach diagnostic threshold comprise a *no SUD* subgroup. Polysubstance UD was tallied for persons who exceeded diagnostic thresholds for multiple substances. Models testing patient indices as predictors utilized a binary outcome (any SUD, no SUD).

### Participants

The aggregate sample (N=10,652) were HIV+ adults linked to care at one of seven urban sites who completed a patient-reported outcome assessment between 01/01/2007 – 12/31/2014. Site locations were at Harvard University, Johns Hopkins University, University of Alabama-Birmingham, University of California-San Diego, University of California-San

Francisco, University of North Carolina-Chapel Hill, and the University of Washington. All patients were aged 18+ years, provided demography/background information upon clinic enrollment, and completed a patient-reported outcome assessment at a routine clinic visit. Per CNICS policy, persons deemed medically unstable, appearing intoxicated, evidencing significant cognitive impairment, or unable to speak English or Spanish did not complete the assessment.

### Analytic Strategy

Descriptive rates of alcohol UD, cocaine UD, marijuana UD, methamphetamine UD, and opioid UD were computed for the aggregate sample and by CNICS site. Preliminary review of distributional properties for demography data prompted decisions to: 1) create five age groups (18–29 years, 30–39 years, 40–49 years, 50–59 years, 60+ years), 2) transform race and ethnicity to a single categorical ‘race/ethnicity’ variable (non-Hispanic Caucasian, non-Hispanic Black, Hispanic, Other) as in prior CNICS reporting (57), 3) retain transgender status for descriptive analyses only, given low base rate of affirmative response, and 4) retain sexual orientation for descriptive analyses only, as preliminary analyses revealed poor subsample representativeness.

In this multisite design, generalized estimating equations (GEEs) examined population-average models for a binary logistic outcome (any SUD, no SUD). This approach describes change in this target outcome due to variance in patient demography and HIV risk factor indices, while accounting for nonindependence in observations within sites (60). Initial bivariate models defined categorical patient-based indices as independent variables, with specification of robust covariance structure due to the large aggregate sample size. Preliminary models examined geographic and historical clustering of SUDs, with respective dummy-coding of CNICS site and assessment timing. Though SUD rates did not vary as a function of assessment timing, site was a robust predictor (as later detailed) and was consequently a covariate in subsequent bivariate models respectively testing age-group, gender, race/ethnicity, IDU history, and MSM history (the latter model restricted to the 8882 male patients in the aggregate sample) as SUD predictors.

Patient-based indices identified as SUD predictors in the aggregate sample were included as independent variables in an eventual multivariable model, with similar inclusion of CNICS site as a covariate and model specification of robust covariance structure.

### Results

In the aggregate sample, age ranged from 18–84 years ( $M=43.7$ ,  $S.D.=10.6$ ). Table 1 lists SUD prevalence of patient subgroups defined by age, gender, race/ethnicity, transgender status, and IDU history. Prevalence of SUD is also noted for MSM history among males ( $n=8882$ ), and for sexual orientation among the persons for whom this self-report data was available ( $n=1716$ ).

### Prevalence of Substance-Specific SUDs

Prevalence of SUD in the aggregate sample was 48%, ranging from 21–71% at the seven CNICS sites. Substance-specific SUD prevalence was: 31% for marijuana UD (site-specific

range, 4–52%); 19% for alcohol UD (site-specific range, 13–27%), 13% for methamphetamine UD (site-specific range, 1–21%), 11% for cocaine UD (site-specific range, 7–18%), and 4% of opiate UD (site-specific range, 1–8%). Multiple diagnostic thresholds were met by 20% of the aggregate sample (site-specific range, 3–37%), commensurate with polysubstance UD. Table 2 lists site-specific rates of any SUD and each substance-specific SUD, with de-identification of individual CNICS-affiliate care sites (as stipulated by institutional review board agreements).

### Modeling Patient-based Predictors of SUD Prevalence

An initial bivariate GEE model identified substantial between-site variability in SUD prevalence, Wald  $X^2(6) = 723.71$ ,  $p < .000$ . This prompted inclusion of site as a covariate in all subsequent models, for which Tables 3 and 4 present beta values with 95% confidence interval limits as well as the corresponding standard errors, Wald  $X^2$  values, their statistical significance, and odds-ratios with 95% confidence interval limits for comparison to the referent group.

For age-group, model statistics noted significant prediction of SUD prevalence, Wald  $X^2(4) = 35.31$ ,  $p < .000$  (see Table 3). The age-group X site interaction was nonsignificant, Wald  $X^2(4) = .55$ ,  $p = .968$ . Relative to referent 60+ year-olds, SUD prevalence was progressively greater in younger groups: 59% among 18–29 year-olds, 54% among 30–39 year-olds, 48% among 40–49 year-olds, 41% among 50–59 year-olds, and 30% among 60+ year-olds.

For gender, model statistics indicated significant prediction of SUD prevalence, Wald  $X^2(1) = 41.11$ ,  $p < .000$  (see Table 3). The gender X site interaction was significant, Wald  $X^2(1) = 6.83$ ,  $p < .01$ . Relative to referent females, SUD prevalence was more likely among males (50% vs. 36%) with the gender X site interaction prompting site-level examination. At two sites, SUD prevalence among males and females was not appreciably different, but was greater among males than females at the remaining five sites.

For race/ethnicity, model statistics suggested significant prediction of SUD prevalence, Wald  $X^2(3) = 49.86$ ,  $p < .000$  (see Table 3). However, the race/ethnicity X site interaction failed to reach statistical significance, Wald  $X^2(3) = 4.91$ ,  $p = .178$ . Further, subgroup comparisons to those of the referent ‘Other’ race/ethnicity failed to reach statistical significance. In descriptive terms, SUD prevalence was 54% among non-Hispanic Caucasians, 48% among both Hispanic and ‘Other’ race/ethnicity subgroups, and 39% among non-Hispanic Blacks.

For IDU history, model statistics noted significant prediction of SUD prevalence, Wald  $X^2(1) = 16.41$ ,  $p < .000$  (see Table 3). The IDU history X site interaction was significant, Wald  $X^2(1) = 53.64$ ,  $p < .000$ . Relative to no IDU history referents, prevalence was greater among those with IDU history (58% vs. 46%) with the IDU history X site interaction prompting site-level examination. Prevalence of SUD among those with an IDU history did not appreciably differ from that of those without an IDU history at a single site, but was higher among those with IDU history than those without an IDU history at the remaining six sites.

Finally, MSM history was examined in the subsample of 8882 male patients. Model statistics revealed significant prediction of SUD prevalence, Wald  $X^2(1) = 42.06$ ,  $p < .000$

(see Table 3). The MSM history X site interaction was significant, Wald  $X^2(1) = 12.77$ ,  $p < .000$ . Relative to no MSM history referents, prevalence was greater among those with MSM history (53% vs. 41%). The MSM history X site interaction prompted site-level examination, revealing inconsistent direction of effects. Prevalence of SUD did not appreciably differ between males with vs. without MSM history at one site, was greater among males with MSM history at four sites, and was greater among males without MSM history at the remaining two sites.

A multivariable model tested relative influences of four patient indices (i.e., age-group, gender, race/ethnicity, IDU history) demonstrated in bivariate models to predict SUD in the aggregate sample, with site again included as a covariate. Model statistics revealed an expected attenuation of influences, albeit with prediction of SUD persisting for: 1) age-group, Wald  $X^2(4) = 33.79$ ,  $p < .000$ ; 2) gender, Wald  $X^2(1) = 10.81$ ,  $p = .001$ ; and 3) race/ethnicity, Wald  $X^2(3) = 37.43$ ,  $p < .000$ . IDU history failed to predict SUD, only trending toward statistical significance in this model, Wald  $X^2(1) = 3.69$ ,  $p = .055$  (see Table 4). Interactions of each SUD predictor with site were nonsignificant [age-group X site, Wald  $X^2(4) = .48$ ,  $p = .975$ ; gender X site, Wald  $X^2(1) = .10$ ,  $p = .749$ ; race/ethnicity X site, Wald  $X^2(3) = 4.82$ ,  $p = .185$ ]. In age-group comparisons, SUDs were more prevalent among 18–29 year-olds, 30–39 year-olds, and 40–49 year-olds relative to 60+ year-olds. Likewise, SUD prevalence was greater among males than females. Specific subgroup differences were not indicated in SUD prevalence for race/ethnicity or IDU history.

## Discussion

Utilizing CNICS to estimate SUD prevalence at seven HIV care sites in the U.S., the current work advances understanding of the corresponding health syndemic. Study findings include: 1) 48% SUD prevalence, encompassing substantial geographic variability (21–71%); 2) substance-specific prevalence topped at 31% for marijuana UD, 19% for alcohol UD, 13% for methamphetamine UD, 11% for cocaine UD, and 4% for opioid UD; and 3) emergence of younger age and male gender as robust SUD predictors in a multivariable model. Collective study findings offer multiregional prevalence estimates for substance-specific SUDs at these urban HIV care sites, identify patient subgroups at greater relative risk for evidencing an SUD, and lay groundwork for future comparative investigation of SUD as an influence on virologic outcomes, clinical processes, and indices of health and well-being.

The 48% SUD prevalence rate in CNICS falls amid an aforementioned range of prior single-site estimates. The lone multisite study to previously estimate SUD prevalence was a cross-sectional HIV Cost and Services Utilization Study (HCSUS) of 2864 adults completing an HIV-focused health visit in a two-month period in 1996. Based on clinical interviews (albeit with incomplete diagnostic questioning), HCSUS estimated prevalence of ‘heavy drinking’ at 15% (52) and ‘drug dependence’ at 12% (51). The current work identified a higher collective SUD rate—and did so in this much larger, continuous, and multiregional CNICS cohort via SUD screening instruments with concurrent validity established for DSM diagnoses (58, 59). Given the university affiliation of CNICS sites and their urban location in large U.S. cities, definitive report of SUD prevalence in HIV care in the U.S. may require

future recruitment of a nationally-representative sample of patients receiving services from the HIV care community.

The specificity of prevalence estimates reported herein for alcohol, cocaine, marijuana, methamphetamine, and opioid UD are an advance over the conglomerated reporting in extant research. A surprisingly high rate of marijuana UD is at the upper end of wide-ranging (12–36%) population estimates (61, 62), and suggests this is an area meriting future attention. Community sampling efforts, employing a ‘past 12 months’ reporting interval, offer additional contemporary points of comparison. For alcohol UD, community prevalence was 14% (53), whereas 19% is noted of this CNICS cohort. When drug-based UDs were aggregated across the four CNICS drug categories plus sedatives/tranquilizers, solvents/inhalants, hallucinogens, and club drugs, community prevalence was 4% (54). In the CNICS cohort—with measurement limited to cocaine, marijuana, methamphetamine, and opioids as well as a ‘past 90-day’ reporting interval (58)—29% prevalence is noted. These comparatively elevated rates of alcohol and (particularly) drug-based UDs hold preventative implications. A high SUD prevalence broadens transmission pathways among HIV+ persons involving the sharing of drug injection equipment, intoxicated involvement in unprotected sex, and sexual violence and victimization. These pathways are amplified among those with SUD—whether due to unawareness of HIV+ status, unsuppressed viral load, or both (8). As earlier noted, these are among the continual and recursive challenges that presence of an SUD poses for patients along the HIV Care Continuum.

Multivariable model findings—specifically, regional differences and greater relative risk of SUD among males and young adults—broadly replicate HCSUS patterns of two decades ago (51, 52). Replication of HCSUS findings, as well as those of national epidemiological studies (53, 54), in this CNICS cohort lends credence to emphases given such predictors in SUD detection efforts. With respect to age, available CNICS data precluded examination of SUD chronicity in this cohort. Nevertheless, persistent substance abuse among aging HIV+ persons is a progressive risk for mortality (63, 64). While improved clinical management of HIV infection has contributed to domestic decline in deaths due to AIDS-related causes, substance-related causes of death (i.e., drug overdose, mental disorders resulting from substance abuse) continue to increase among HIV+ persons of all ages (65). Less robust race/ethnicity findings in this CNICS cohort coincide with equivocal HCSUS reporting in this area (51, 52). Weak SUD prediction by patient IDU history may reflect effective use of harm reduction strategies, historically promoted among HIV+ populations (66). Likewise, SUD prediction by MSM history in the CNICS male subsample, albeit with site differences in strength and direction, suggests context may be critical to interpret how this risk factor for HIV transmission influences the prevalence of SUDs (67).

Strengths and caveats of this work bear further mention. The former include: a large, multiregional cohort of HIV care enrollees; inclusion of established SUD screening instruments with validated diagnostic thresholds; and use of secure, private means to collect patient-reports of recent substance use behavior. One noteworthy caveat is setting representativeness, given the prominent size, resources, university affiliation, and urban location of the seven CNICS sites.



Definitive reporting of SUD prevalence among domestic HIV care enrollees awaits future study with a nationally-representative sample. Regarding this cohort of 10,652 patients, potential selection bias is acknowledged given enrollment of 32,000+ persons since CNICS inception (68). The reported 48% SUD prevalence may reflect an underestimation, due to: 1) omission of licit (i.e., tobacco) and illicit (i.e., sedatives/tranquilizers, solvents/inhalants, hallucinogens, club drugs) substance categories in the abbreviated CNICS version of the ASSIST, and 2) CNICS data collection policy precluding completion of the assessment battery by patients appearing intoxicated. Despite these paired caveats of measurement and sampling, SUD prevalence in this cohort is safely within the range of prior published estimates for U.S.-based HIV care samples. Another caveat relates to potential influences of unassessed 3<sup>rd</sup>-variables, like socioeconomic (i.e., employment, income, education) and historical (i.e., family history of SUD, exposure to sexual trauma/victimization) patient background indices. Socioeconomic indices may interact with SUD and other health conditions to influence course and outcome of HIV infection (69), whereas historical indices are linked in HIV+ populations to substance use, treatment failure, morbidity, and mortality (70, 71).

## Conclusions

Caveats notwithstanding, the current study advances understanding of the scope of the American SUD-HIV syndemic. Nearly half of a large, multiregional CNICS cohort met diagnostic threshold for an SUD, and 20% met thresholds for two or more of five substance categories evaluated. Given detrimental impacts that SUDs have on patient care throughout the HIV Care Continuum (3), more effective disease control efforts may derive through arming of HIV care settings with greater capacity to offer their patients addiction-focused services. In addition to established pharmacotherapies for alcohol UD (i.e., acamprosate, naltrexone) and opioid UD (i.e., buprenorphine, methadone, extended release naltrexone), behavior therapies offer a useful response to many clinical challenges posed by persons with SUD. For example, efficacy to improve antiretroviral medication adherence among substance-misusing patients has been demonstrated in multiple randomized controlled trials for cognitive-behavior therapy (72, 73), contingency management (74, 75), and motivational interviewing (76, 77). Further, the considerable empirical support documented for each of these behavior therapies in the addictions field has prompted recommendation of their broad application across adult patient populations and substances of abuse (78). As in other health settings, a key to effective implementation of these behavior therapies may be in maximizing their compatibility with existing service provisions in HIV care. Findings reported herein suggest there is need for addiction-focused services in U.S.-based HIV care settings, and specify young adults and men as patient subgroups for whom they will most often be applicable.

## Acknowledgements

The funding source for this analytic work was National Institute on Drug Abuse R03DA039719 (*Informing Dissemination of Behavior Therapies to Enhance HIV Care Among Substance Abusers*, Hartzler PI). The authors thank the CNICS sites for contributing data to this endeavor. CNICS is funded by R24 AI067039 with sites at University of Alabama at Birmingham (P30 AI027767), University of Washington (P30 AI027757), University of California San Diego (P30 AI036214), University of California San Francisco (P30 AI027763), Case Western

Reserve University (P30 AI036219), John Hopkins University (P30 AI094189, U01 DA036935), Fenway Health/Harvard (P30 AI060354), and University of North Carolina Chapel Hill (P30 AI50410).

## References

1. Moore RD , Bartlett JG . Dramatic decline in the HIV-1 RNA level over calendar time in a large urban HIV practice. *Clinical Infectious Diseases*. 2011;53:600–4.21844006
2. Althoff KN , Buchacz K , Hall HI , et al. U.S. trends in antiretroviral therapy use, HIV RNA plasma viral loads, and CD4 T-lymphocyte cell counts among HIV-infected persons. *Annals of Internal Medicine*. 2012;157:325–35.22944874
3. Gardner EM , McLees MP , Steiner JF , del Rio C , Burman WJ . The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clinical Infectious Diseases*. 2011;52(6):793–800.21367734
4. Mugavero MJ , Amico KR , Horn T , Thompson MA . The state of engagement in HIV care in the United States: From cascade to continuum to control. *Clinical Infectious Diseases*. 2013;57(8):1164–71.23797289
5. McLellan AT , Woodworth AM . The affordable care act and treatment for ‘substance use disorders’: Implications of ending segregated behavioral healthcare. *Journal of Substance Abuse Treatment*. 2014;46:541–5.24679908
6. McCance-Katz EF , Rabiner CA , Rivers JLA . The Affordable Care Act: Implementation and implications for addiction specialty care. *The American Journal on Addictions*. 2014;23:429–30.25065262
7. Denis CM , Gelernter J , Hart AB , Kranzler HR . Inter-observer reliability of DSM-5 substance use disorders. *Drug and Alcohol Dependence*. 2015;153:229–35.26048641
8. Hall HI , Holtgrave DR , Tang T , Rhodes P . HIV transmission in the United States: Considerations of viral load, risk behavior, and health disparities. *AIDS and Behavior*. 2013;17:1632–36.23456577
9. Dieffenbach WW , Fauci AS . Universal voluntary testing and treatment for prevention of HIV transmission. *JAMA*. 2009;301:2380–2.19509386
10. Granich RM , Gilks CF , Dye C , De Cock KM , Williams BG . Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: A mathematical model. *Lancet*. 2009;373(48–57).19038438
11. Giordano TP , Visnegarwala F , White ACJ , Troisi CL , Frankowski RF , Hartman CM , et al. Patients referred to an urban HIV clinic frequently fail to establish care: Factors predicting failure. *AIDS Care*. 2005;17(6):773–83.16036264
12. Meyer JP , Althoff AL , Altice FL . Optimizing care for HIV-infected people who use drugs: Evidence-based approaches to overcoming healthcare disparities. *Clinical Infectious Diseases*. 2013;57(9):1309–17.23797288
13. DeLorenze GN , Weisner C , Tsai AL , Satre DD , Quesenberry CP , Excess mortality among HIV-infected patients diagnosed with substance use dependence or abuse receiving care in a fully integrated medical care program. *Alcoholism: Clinical and Experimental Research*. 2011;35(2):203–10.
14. Arnsten JH , Demas PA , Grant RW , et al. Impact of active drug use on antiretroviral therapy adherence and viral suppression in HIV-infected drug users. *Journal of General Internal Medicine*. 2002;17(5):377–81.12047736
15. Chander G , Lau B , Moore RD . Hazardous alcohol use: A risk factor for non-adherence and lack of suppression in HIV infection. *Journal of Acquired Immune Deficiency Syndromes*. 2006;43(4):411–7.
16. Reback CJ , Larkins S , Shoptaw S . Methamphetamine abuse as a barrier to HIV medication adherence among gay and bisexual men. *AIDS Care*. 2003;15(6):775–85.14617499
17. Pence BW , Miller WC , Gaynes BN , Eron JJ , Psychiatric illness and virologic response in patients initiating highly active antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes*. 2007;44(2):159–66.
18. King WD , Larkins S , Hucks-Ortiz C , et al. Factors associated with HIV viral load in a respondent driven sample in Los Angeles. *AIDS and Behavior*. 2009;13(1):145–53.18064555

19. Giordano TP , Gifford AL , White ACJ , et al. Retention in care: A challenge to survival with HIV infection. *Clinical Infectious Diseases*. 2007;44:1493–9.17479948
20. Torian LV , Wiewel EW . Continuity of HIV-related medical care, New York City, 2005–2009: Do patients who initiate care stay in care? *AIDS, Patient Care, and STDs*. 2011;25:79–88.21284498
21. Hall HI , Gray KM , Tang T , Li J , Shouse L , Mermin J . Retention in care of adults and adolescents living with HIV in 13 U.S. areas. *Journal of Acquired Immune Deficiency Syndromes*. 2012;60:77–82.
22. Rebeiro P , Althoff KN , Buchacz K , Gill J , Horberg M , Krentz H , et al. Retention among North American HIV-infected persons in clinical care, 2000–2008. *Journal of Acquired Immune Deficiency Syndromes*. 2013;62(3):356–62.
23. Werb D , Mills EJ , Montaner JS , Wood E . Risk of resistance to highly active antiretroviral therapy among HIV-positive injecting drug users: A meta-analysis. *Lancet Infectious Diseases*. 2010;10:464–9.20610328
24. Malta M , Strathdee SA , Magnanini MM , Bastos FI . Adherence to antiretroviral therapy for human immunodeficiency virus/acquired immune deficiency syndrome among drug users: A systematic review. *Addiction*. 2008;103:1242–57.18855813
25. Roux P , Carrieri MP , Cohen J , Ravaux I , Poizot-Martin I , Dellamonica P , et al. Retention in opioid substitution treatment: A major predictor of long-term virological success for HIV-infected injection drug users receiving antiretroviral treatment. *Clinical Infectious Diseases*. 2009;49(9):1433–40.19807275
26. Malee KM , Mellins CA , Huo Y , Tassiopoulos K , Smith R , Sirois PA , et al. Prevalence, incidence, and persistence of psychiatric and substance use disorders among mothers living with HIV. *Journal of Acquired Immune Deficiency Syndromes*. 2014;65(5):526–34.
27. Rabkin JG , McElhinney MC , Ferrando SJ . Mood and substance use disorders in older adults with HIV/AIDS: Methodological issues and preliminary evidence. *AIDS*. 2004;18(Supplement #1):S43–S8.15075497
28. Rosenberg SD , Drake RE , Brunette MF , Wolford GL , Marsh BJ . Hepatitis C virus and HIV coinfection in people with severe mental illness and substance use disorders. *AIDS*. 2005;19(Supplement #3):S26–S33.16251824
29. Gaynes BN , Pence BW , Erron JJ , Miller WC . Prevalence and comorbidity of psychiatric diagnoses based on reference standard in an HIV+ patient population. *Psychosomatic Medicine*. 2008;70(4):505–11.18378865
30. Pence BW , Miller WC , Whetten K , Erron JJ , Gaynes BN . Prevalence of DSM-IV-defined mood, anxiety, and substance use disorders in an HIV clinic in the southeastern United States. *Journal of Acquired Immune Deficiency Syndromes*. 2006;42(3):298–306.
31. Atkinson JH , Grant I , Kennedy CJ , Richman DD , Spector SA , McCutchan JA . Prevalence of psychiatric disorders among men infected with human immunodeficiency virus: A controlled study. *Archives of General Psychiatry*. 1988;45:859–64.3415427
32. Dew MA , Becker JT , Sanchez J , Caldararo R , Lopez OL , Wess J , et al. Prevalence and predictors of depressive, anxiety, and substance use disorders in HIV-infected and uninfected men: A longitudinal evaluation. *Psychological Medicine*. 1997;27:395–409.9089832
33. Perry S , Jacobsberg LB , Fishman B , Frances A , Bobo J , Jacobsberg BK . Psychiatric diagnosis before serological testing for the human immunodeficiency virus. *American Journal of Psychiatry*. 1990;147:89–93.2293794
34. Catalan J , Klimes I , Bond A , Day A , Garrod A , Rizza C . The psychosocial impact of HIV infection in men with haemophilia: Controlled investigation and factors associated with psychiatric morbidity. *Journal of Psychosomatic Research*. 1992;36:409–16.1619581
35. Catalan J , Klimes I , Day A , Garrod A , Bond A , Gallwey J . The psychosocial impact of HIV infection in gay men: A controlled investigation and factors associated with psychiatric morbidity. *British Journal of Psychiatry*. 1992;161:774–8.1483162
36. Chuang HT , Jason GW , Pajurkova EM , Gill MJ . Psychiatric morbidity in patients with HIV infection. *Canadian Journal of Psychiatry*. 1992;37:109–15.1562954
37. Williams JB , Rabkin JG , Remien RH , Gorman JM , Ehrhardt A . Multidisciplinary baseline assessment of homosexual men with and without human immunodeficiency virus infection:

- Standardized clinical assessment of current and lifetime psychopathology. *Archives of General Psychiatry*. 1991;48:124–30.1671198
38. Brown GR , Rundell JR , McManis SE , Kendall SN , Zachary R , Temoshok L . Prevalence of psychiatric disorders in early stages of HIV infection. *Psychosomatic Medicine*. 1992;54:588–601.1438661
39. Lipsitz JD , Williams JB , Rabkin JG , Remien RH , Bradbury M , el Sadr W , et al. Psychopathology in male and female intravenous drug users with and without HIV infection. *American Journal of Psychiatry*. 1994;151:1662–8.7943458
40. Rosenberger PH , Bornstein RA , Nasrallah HA , Para MF , Whitacre CC , Fass RJ , et al. Psychopathology in human immunodeficiency virus infection: Lifetime and current assessment. *Comprehensive Psychiatry*. 1993;34:150–8.8339532
41. Maj M , Strarace F , Sartorius N . Mental disorders in HIV-I infection and AIDS In: Sartorius N , Prilipko LL , editors. WHO Expert Series on Biological Psychiatry. Seattle, WA: Hogrefe & Huber Publishers; 1993.
42. Lyketsos CG , McHugh PR , Hanson A , Treisman GJ , Fishman M . Screening for psychiatric morbidity in a medical outpatient clinic for HIV infection: The need for a psychiatric presence. *International Journal of Psychiatry in Medicine*. 1994;24:103–13.7960418
43. Bix BC , Glosser G , Holmes W , Ballas C , Meritz M , Hutelmeyer C , et al. Relationship between psychiatric disease and neurological impairment in HIV seropositive individuals. *Journal of the International Neuropsychological Society*. 1995;1:581–8.9375246
44. Johnson JG , Williams JB , Rabkin JG , Goetz R , Remien RH . Axis I psychiatric symptoms associated with HIV infection and personality disorder. *American Journal of Psychiatry*. 1995;152:551–4.7694903
45. Summers J , Zisook S , Atkinson JH , Sciolla A , Whitehall W , Brown S , et al. Psychiatric morbidity associated with acquired immune deficiency syndrome-related grief resolution. *Journal of Nervous and Mental Disease*. 1995;183:384–9.7798087
46. Sohler NL , Wong MD , Cunningham WE , Cabral H , Drainoni ML , Cunningham CO . Type and pattern of illicit drug use and access to health care services for HIV-infected people. *AIDS Patient Care and STDs*. 2007;21(Supplement 1):S68–S76.17563292
47. Rondinelli AJ , Ouellet LJ , Strathdee SA , Latka MH , Hudson SM , Hagan H , et al. Young adult injection drug users in the United States continue to practice HIV risk behaviors. *Drug and Alcohol Dependence*. 2009;104(1–2):167–74.19559543
48. Engstrom M , Shibusawa T , El-Bassel N , Gilbert L . Age and HIV sexual risk among women in methadone treatment. *AIDS and Behavior*. 2011;15(1):103–13.19936912
49. Strathdee SA , Sherman SG . The role of sexual transmission of HIV infection among injection and non-injection drug users. *Journal of Urban Health*. 2003;80(Supplement 3):iii7–iii14.14713667
50. Mimiaga MJ , Reisner SL , Grasso C , Crane HM , Safren SA , Kitahata MM , et al. Substance use among HIV-infected patients engaged in primary care in the United States: Findings from the Centers for AIDS Research Network of Integrated Clinical Systems cohort. *American Journal of Public Health*. 2013;103(8):1457–67.23763417
51. Bing EG , Burnam MA , Longshore D , et al. Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Archives of General Psychiatry*. 2001;58(8):721–8.11483137
52. Galvan FH , Bing EG , Fleishman JA , et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: Results from the HIV Cost and Services Utilization Study. *Journal of Studies on Alcohol*. 2002;63(2):179–86.12033694
53. Grant BF , Goldstein RB , Saha TD , Chou C , Jung J , Zhang H , et al. Epidemiology of DSM-5 alcohol use disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry*. 2015;72(8):757–66.26039070
54. Grant BF , Saha TD , Ruan WJ , Goldstein RB , Chou SP , Jung J , et al. Epidemiology of DSM-5 drug use disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry*. 2016;73(1):39–47.26580136

55. Kitahata MM , Rodriquez B , Haubrick R , Boswell S , Mathews WC , Lederman MM , et al. Cohort profile: The Centers for AIDS Research Network of Integrated Clinical Systems. *International Journal of Epidemiology*. 2008;37(948–955).18263650
56. Hughes MD . Initial treatment of HIV infection: Randomized trials with clinical endpoints are still needed. *Journal of Infectious Diseases*. 2006;194:542–4.16897649
57. Dombrowski JC , Kitahata MM , Von Rumpalay SE , Crane HM , Mugavero MJ , Eron JJ , et al. High levels of antiretroviral use and viral suppression among persons in HIV care in the United States, 2010. *JAIDS*. 2013;63(3):299–306.23572013
58. Humeniuk R , Ali R , Babor TF , Farrell M , Formigoni ML , Jittiwuikarn J , et al. Validation of the alcohol, smoking, and substance involvement screening test (ASSIST). *Addiction*. 2008;103(6):1039–47.18373724
59. Dawson DA , Smith SM , Saha TD , Rubinsky AD , Grant BF . Comparative performance of the AUDIT-C in screening for DSM-IV and DSM-5 alcohol use disorders. *Drug and Alcohol Dependence*. 2012;126:384–8.22728044
60. Hubbard AE , Ahern J , Fleischer NL , Van der Laan M , Lippman SA , Jewell N , et al. To GEE or not to GEE: Comparing population average and mixed models for estimating the associations between neighborhood risk factors and health. *Epidemiology*. 2010;21:467–74.20220526
61. Hasin DS , Kerridge BT , Saha TD , Huang B , Pickering RP , Smith SM , et al. Prevalence and correlates of DSM-5 cannabis use disorder, 2012–2013: Findings from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *American Journal of Psychiatry*. 2016;173(6):588–99.26940807
62. Grucza RA , Agrawal A , Krauss MJ , Cavazos-Rehg PA , Bierut LJ . Recent trends in the prevalence of marijuana use and associated disorders in the United States. *JAMA Psychiatry*. 2016;73(3):300–1.26864618
63. Reif S , Pence BW , Hall I , Hu X , Whetten K , Wilson E . HIV diagnoses, prevalence, and outcomes in nine southern states. *Journal of Community Health*. 2015;40(4):642–51.25524210
64. Cima M , Parker RD , Ahmed Y , Cook S , Dykema S , Dukes K , et al. Cause of death in HIV-infected patients in South Carolina (2005–2013). *International Journal of STD and AIDS*. 2016;27(1):25–32.25691444
65. Schwarcz SK , Vu A , Hsu LC , Hessol NA . Changes in causes of death among persons with AIDS: San Francisco, California, 1996–2011. *AIDS Patient Care and STDs*. 2014;28(10):517–23.25275657
66. Remien RH , Goetz R , Rabkin JG , Williams JB , Bradbury M , Ehrhardt A , et al. Remission of substance use disorders: Gay men in the first decade of AIDS. *Journal of Studies on Alcohol*. 1995;56:226–32.7760570
67. Santos GM , Do T , Beck J , Makofane K , Arreola S , Pyun T , et al. Syndemic conditions associated with HIV risk in a global sample of men who have sex with men. *Sexually Transmitted Diseases*. 2014;90(3):250–3.
68. CNICS. CFAR Network of Integrated Clinical Systems. University of Alabama-Birmingham; 2016 [cited 2016 May 16]; Available from: [www.uab.edu/cnics](http://www.uab.edu/cnics).
69. Oldenburg CE , Perez-Brumer AG , Reisner SL . Poverty matters: Contextualizing the syndemic condition of psychological factors and newly diagnosed HIV infection in the United States. *AIDS*. 2014;28(18):2763–69.25418633
70. Pence BW . The impact of mental health and traumatic life experiences on antiretroviral treatment outcomes for people living with HIV/AIDS. *Journal of Antimicrobial Chemotherapy*. 2009;63:636–40.19153077
71. Kalichman SC , Sikkema KJ , DiFonzo K , Luke W , Austin J . Emotional adjustment in survivors of sexual assault living with HIV/AIDS. *Journal of Traumatic Stress*. 2002;15(4):286–96.
72. Parsons JT , Rosof E , Punzalan JC , Di Maria L . Integration of Motivational Interviewing and Cognitive Behavioral Therapy to Improve HIV medication adherence and reduce substance use among HIV-positive men and women: Results of a pilot project. *AIDS Patient Care and STDs*. 2005;19(1):31–9.15665633
73. Safren SA , O’Cleirigh CM , Bullis JR , Otto MW , Stein MD , Pollack MH . Cognitive behavioral therapy for adherence and depression (CBT-AD) in HIV-infected injection drug users: A

- randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2012;80(3):404–15.22545737
74. Petry NM , Weinstock J , Alessi SM , Lewis MW , Dieckhaus K . Group-based randomized trial of contingencies for health and abstinence in HIV patients. *Journal of Consulting and Clinical Psychology*. 2010;78(1):89–97.20099954
75. Rosen MI , Dieckhaus K , McMahon TJ , Valdes B , Petry N , Cramer JA , et al. Improved adherence with contingency management. *AIDS Patient Care and STDs*. 2007;21(1):30–40.17263651
76. Diiorio C , McCarty F , Resnicow K , McDonnell-Holstad M , Soet J , Yeager K , et al. Using motivational interviewing to promote adherence to antiretroviral medications: A randomized controlled study. *AIDS Care*. 2008;20(3):273–83.18351473
77. Williams AB , Fennie KP , Bova CA , Burgess JD , Danvers KA , Dieckhaus K . Home visits to improve adherence to highly active antiretroviral therapy: A randomized controlled trial. *JAIDS*. 2006;42(3):314–21.16770291
78. NIDA. Principles of drug addiction treatment: A research-based guide (3rd ed.). Services USDoHaH, editor 2012.

**Table 1.**

## Substance Use Disorder UD Prevalence by Patient Demography and HIV Risk Factors

|  | <u>Subsample Size (%)</u> | <u>'Any SUD' Prevalence<sup>a</sup></u> |
|--|---------------------------|---|
| Aggregate Sample <sup>b</sup>                      | 10,652 (100%)             | 48%                                     |
| <u>Patient Demography</u>                          |                           |   |
| Age-Group  |                           |   |
| 18–29 years  | 1254 (12%)                | 59%                                     |
| 30–39 years  | 2310 (22%)                | 54%                                     |
| 40–49 years  | 3901 (36%)                | 48%                                     |
| 50–59 years  | 2535 (24%)                | 41%                                     |
| 60+ years  | 652 (6%)                  | 30%                                     |
| Gender   |                           |   |
| Male   | 8882 (83%)                | 50%                                     |
| Female   | 1770 (17%)                | 36%                                     |
| Race/Ethnicity                                     |                           |   |
| Non-Hispanic White                                 | 5278 (49%)                | 54%                                     |
| NonHispanic Black                                  | 3632 (34%)                | 39%                                     |
| Hispanic   | 1270 (13%)                | 48%                                     |
| Other  | 472 (4%)                  | 48%                                     |
| Sexual Orientation (subsample n=1716) <sup>c</sup> |                           |   |
| Lesbian, Gay, or Homosexual                        | 1280 (75%)                | 59%                                     |
| Straight or Heterosexual                           | 278 (16%)                 | 52%                                     |
| Bisexual   | 95 (6%)                   | 65%                                     |
| 'Something Else'                                   | 34 (2%)                   | 76%                                     |
| 'Don't Know'                                       | 29 (2%)                   | 41%                                     |
| Transgender  |                           |   |
| Yes  | 87 (1%)                   | 52%                                     |
| No   | 10,565 (99%)              | 48%                                     |
| <u>HIV Risk Factors<sup>d</sup></u>                |                           |   |
| History of MSM (subsample of n=8882)               |                           |   |
| Yes  | 7039 (79%)                | 53%                                     |
| No   | 1843 (21%)                | 41%                                     |
| History of IDU                                     |                           |   |
| Yes  | 1718 (16%)                | 58%                                     |
| No   | 8934 (84%)                | 46%                                     |

<sup>a</sup>Any SUD ' identification based on substance-specific diagnostic thresholds from the AUDIT-C (alcohol UD) and the ASSIST (cocaine UD, marijuana UD, methamphetamine UD, opioid UD).

<sup>b</sup>Sample consists of HIV+ persons enrolling in services 01/01/2007 – 12/31/2014

<sup>c</sup>Patient-reported sexual orientation collected only at three CNICS sites since 2012

<sup>d</sup>MSM history and IDU history per chart notation at CNICS care sites

**Table 2.**

## Substance Use Disorder Prevalence by Geographic Site

|                    | Site<br>#1<br>(818) | Site<br>#2<br>(852) | Site<br>#3<br>(2580) | Site<br>#4<br>(3179) | Site<br>#5<br>(1161) | Site<br>#6<br>(706) | Site<br>#7<br>(1356) | Aggregate<br>(10,652) |
|--------------------|---------------------|---------------------|----------------------|----------------------|----------------------|---------------------|----------------------|-----------------------|
| Any SUD            | 60%                 | 21%                 | 39%                  | 48%                  | 71%                  | 34%                 | 61%                  | 48%                   |
| Alcohol UD         | 27%                 | 13%                 | 16%                  | 18%                  | 21%                  | 14%                 | 22%                  | 19%                   |
| Cocaine UD         | 13%                 | 7%                  | 11%                  | 8%                   | 17%                  | 7%                  | 18%                  | 11%                   |
| Marijuana UD       | 36%                 | 4%                  | 26%                  | 29%                  | 52%                  | 24%                 | 42%                  | 31%                   |
| Methamphetamine UD | 14%                 | 1%                  | 4%                   | 17%                  | 31%                  | 2%                  | 21%                  | 13%                   |
| Opioid UD          | 3%                  | 1%                  | 3%                   | 3%                   | 8%                   | 1%                  | 7%                   | 4%                    |

Site de-identification stipulated by institutional review board of one or more CNICS university-affiliate care sites Corresponding sample/subsample sizes listed in parentheses

SUD identification based on substance-specific diagnostic thresholds from the AUDIT-C (alcohol UD) and the ASSIST (cocaine UD, marijuana UD, methamphetamine UD, opioid UD).



**Table 3.****Bivariate Prediction of Substance Use Disorders by Patient Demography and HIV Risk Factors**

|  | <b>Beta value</b> | <b>95% CI (lower, upper)</b> | <b>Standard Error</b> | <b>Wald <math>\chi^2</math> (1)</b> | <b>Odds Ratio</b> | <b>95% CI (lower, upper)</b> |
|--|-------------------|------------------------------|-----------------------|-------------------------------------|-------------------|------------------------------|
| <b>Age-Group</b>                                 |                   |                              |                       |                                     |                   |                              |
| 18–29 years                                      | 1.36              | (.80, 1.91)                  | .28                   | 23.23***                            | 3.88              | (2.24, 6.73)                 |
| 30–39 years                                      | 1.10              | (.59, 1.61)                  | .26                   | 17.87***                            | 3.01              | (1.80, 5.00)                 |
| 40–49 years                                      | .84               | (.35, 1.33)                  | .25                   | 11.30***                            | 2.31              | (1.42, 3.77)                 |
| 50–59 years                                      | .57               | (.06, 1.08)                  | .26                   | 4.87***                             | 1.77              | (1.07, 2.93)                 |
| 60+ years (reference)                            | 0                 |                              |                       |                                     | 1.00              |                              |
| <b>Gender</b>                                    |                   |                              |                       |                                     |                   |                              |
| Male   | .92               | (.64, 1.20)                  | .14                   | 41.11***                            | 2.51              | (1.90, 3.33)                 |
| Female (reference)                               | 0                 |                              |                       |                                     | 1.00              |                              |
| <b>Race/Ethnicity</b>                            |                   |                              |                       |                                     |                   |                              |
| Nonhispanic Caucasian                            | .47               | (–.12, 1.05)                 | .30                   | 2.47, ns                            | 1.59              | (.89, 2.85)                  |
| Nonhispanic Black                                | –.32              | (–.91, .27)                  | .30                   | 1.10, ns                            | .73               | (.41, 1.32)                  |
| Hispanic   | .01               | (–.74, .77)                  | .38                   | .00, ns                             | 1.02              | (.48, 2.15)                  |
| Other (reference)                                | 0                 |                              |                       |                                     | 1.00              |                              |
| <b>IDU History</b>                               |                   |                              |                       |                                     |                   |                              |
| Yes  | .60               | (.31, .88)                   | .15                   | 16.41***                            | 1.81              | (1.36, 2.42)                 |
| No (reference)                                   | 0                 |                              |                       |                                     | 1.00              |                              |
| <b>MSM history (male subsample only, n=8882)</b> |                   |                              |                       |                                     |                   |                              |
| Yes  | .94               | (.66, 1.22)                  | .14                   | 42.06***                            | 2.56              | (1.93, 3.40)                 |
| No (reference)                                   | 0                 |                              |                       |                                     | 1.00              |                              |

Analyses based on aggregate sample (N=10,652), except where otherwise indicated, and include enrollment site as a covariate.

SUD identification based on substance-specific diagnostic thresholds from the AUDIT-C (alcohol UD) and the ASSIST (cocaine UD, marijuana UD, methamphetamine UD, opioid UD).

Odds-ratios reflect the likelihood of SUD relative to the reference category

\*\*\*  
 $p < .001$

\*\*  
 $p < .01$

\*  
 $p < .05$ .

**Table 4.**

Multivariable Prediction of Substance Use Disorders by Patient Demography and HIV Risk Factors

|                       | <b>Beta value</b> | <b>95% CI (lower, upper)</b> | <b>Standard Error</b> | <b>Wald <math>\chi^2</math> (1)</b> | <b>Odds Ratio</b> | <b>95% CI (lower, upper)</b> |
|-----------------------|-------------------|------------------------------|-----------------------|-------------------------------------|-------------------|------------------------------|
| Age-Group             |                   |                              |                       |                                     |                   |                              |
| 18–29 years           | 1.27              | (.72, 1.82)                  | .28                   | 20.48***                            | 3.55              | (2.05, 6.15)                 |
| 30–39 years           | 1.07              | (.56, 1.58)                  | .26                   | 16.82***                            | 2.90              | (1.75, 4.83)                 |
| 40–49 years           | .73               | (.24, 1.21)                  | .25                   | 8.60**                              | 2.06              | (1.27, 3.35)                 |
| 50–59 years           | .49               | (-.01, .99)                  | .26                   | 3.72, ns                            | 1.64              | (.99, 2.70)                  |
| 60+ years (reference) | 0                 |                              |                       |                                     | 1.00              |                              |
| Gender                |                   |                              |                       |                                     |                   |                              |
| Male                  | .51               | (.21, .81)                   | .15                   | 10.81**                             | 1.66              | (1.23, 2.25)                 |
| Female (reference)    | 0                 |                              |                       |                                     | 1.00              |                              |
| Race/Ethnicity        |                   |                              |                       |                                     |                   |                              |
| Nonhispanic Caucasian | .53               | (-.06, 1.11)                 | .30                   | 3.10, ns                            | 1.69              | (.94, 3.05)                  |
| Nonhispanic Black     | -.19              | (-.78, .41)                  | .31                   | .37, ns                             | .83               | (.46, 1.51)                  |
| Hispanic              | .01               | (-.74, .77)                  | .39                   | .00, ns                             | 1.01              | (.47, 2.15)                  |
| Other (reference)     | 0                 |                              |                       |                                     | 1.00              |                              |
| IDU History           |                   |                              |                       |                                     |                   |                              |
| Yes                   | .29               | (-.01, .59)                  | .15                   | 3.69, ns                            | 1.34              | (.99, 1.81)                  |
| No (reference)        | 0                 |                              |                       |                                     | 1.00              |                              |

Analyses based on aggregate sample (N=10,652), except where otherwise indicated, and include enrollment site as a covariate.

SUD identification based on substance-specific diagnostic thresholds from the AUDIT-C (alcohol UD) and the ASSIST (cocaine UD, marijuana UD, methamphetamine UD, opioid UD).

Odds-ratios reflect the likelihood of SUD relative to the reference category;

\*\*\*  
 $p < .001$ ,

\*\*  
 $p < .01$ ,

\*  
 $p < .05$ .