

Prevalence of and Factors Associated with Migraine in Medical Students at BandarAbbas, Southern Iran, in 2012

Maryam Yazdanparast^{1,2}, Ali Asghar Abrishamizadeh³, Hamidreza Mahboobi^{4,5}, Aria Omrani⁶, Mahsa Ghasemi¹, Monir Ghorashi¹, Milad Zahed¹, Abbass Mohammadi¹, Zahra Karimi¹

¹ Student of Medicine of Hormozgan University of Medical Science (HUMS), Bandarabbass, Iran.

² Research Center for Behavioral and Neurosciences of Hormozgan University of Medical Science (HUMS), Bandarabbass, Iran.

³ Assistant Professor of Neurology Department, Neurologist. Hormozgan University of Medical Science (HUMS), Bandarabbass, Iran.

⁴ General Physician, Infectious and Tropical Diseases Research Center, Hormozgan University of Medical Sciences (HUMS), Iran.

⁵ Payam Noor University (PNU), Tehran, Iran.

⁶ Resident of Clinical Neurology, Institute of Neurology, National Hospital of Neurology and Neurosurgery (UK) & UCL, Belgium.

Corresponding Author:

Hamidreza Mahboobi, General Physician, Infectious and Tropical Diseases Research Centre, Hormozgan University of Medical Sciences (HUMS), Hormozgan, Iran. Tel: +989172121995, Email: hamidrezamahboobi@yahoo.com

Abstract:

Background: Migraine is one of the most common etiologies for headache. This very common neurological disorder has a significant impact on patients' quality of life. The aim of the current study is to evaluate the prevalence of migraine among medical students in the Hormozgan University of Medical Sciences (HUMS).

Methods: A total of 350 medical students were enrolled in our descriptive study. Data were collected using the standard questionnaire of the International Headache Association. The data were analyzed by SPSS 20.0 software using descriptive statistics, Chi-Square, and Independent Samples T-Test. A P-value of ≥ 0.05 was considered statistically significant, since most public health professionals use this value as a standard.

Results: Among the medical students in our study, 24.6% had experienced frequent, severe headaches. The underlying causes of the headaches were diagnosed in 19.8% of the students. The prevalence of migraine in our study was 16.3% (mean age=21.28±2.71years). The prevalence varied by gender, and it was greater among male students.

Conclusion: Our findings indicated that there was a high prevalence of migraine among the medical students in our study, and these findings were consistent with those of previous studies in Iran and other countries.

Keywords: Headache; Migraine; Medical students

Additional Information for citing this article:

Title of Journal: Electronic physician; Abbreviated title of journal: Electron. Physician
doi: 10.14661/2013.679-684

Editorial information:

Type of article: Original

Published: August.01.2013

© 2013 The Authors. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

1. Introduction

Migraine is the most common type of headache (1), and it is a prevalent, neurological disorder in young people. It has a significant, detrimental impact on their quality of life (2-4). Migraine is characterized by unilateral, pulsatile episodes of headache, which is more prominent in the temporal area and lasts from 4 to 72 hours (5-7). Migraine is more prevalent among females which may be due to hormonal changes (8). The literature suggests a different prevalence for this neurological disorder dependent on study populations, geographical factors, and age groups (9-14). One study in Iran reported a prevalence of 12.3% in adolescents who were 12-14 years old (15).

The main risk factors include sleep disorders, high temperature, anxiety, and stress. Migraine is usually associated with nausea, vomiting, and photophobia, and it is worsened by physical activity (12, 16-20). Migraine is more prevalent among people who have higher educational levels (21, 22). Migraine affects patients' daily activities and quality of life (3, 14, 23). The literature indicates that 31-53% of migraine patients are prevented from doing their daily activities (24, 25). Students are more susceptible to severe episodes of migraine, especially those who are experiencing high levels of stress, such as medical students. Thus, the aim of the current study was to assess the prevalence of migraine among medical students in Bandar Abbas, southern Iran.

2. Material and Methods

This descriptive study was performed at the Hormozgan University of Medical Sciences in 2012. We selected 350 students in the medical sciences, including students in various medical fields, including nursing, obstetrics, radiology, health Sciences, the operating room, anesthesiology, laboratory, medical records, emergency, and health information technology (HIT).

Our questionnaire had two parts. In the first part, we focused on demographic information, including age, gender, entrance year, and the field of education. The second part was designed on the basis of the headache questionnaire developed by the International Headache Society (IHS) (26). The aims of the study were explained in detail to the students prior to their being enrolled as participants. Students who declined to participate in the study were excluded. Data were analyzed using SPSS 20.0 software (27) with descriptive statistics (frequency, percentage, mean, and standard deviation), Chi-Square Test, and Independent Samples T-Test. A P-value of ≥ 0.05 was considered to be statistically significant.

3. Results

The mean age of the 350 medical science participants was 21.28 ± 2.78 (95% CI: 20.98-21.57) years. Among the participants, 114 (32.6%) were male and 236 (67.4%) were female. The mean age of the males was 21.90 ± 3.41 (95% CI: 21.27-22.53), and the mean age of the females was 20.97 ± 2.37 (95% CI: 20.67-21.28), which suggested a statistically significant difference ($T=2.604$; $P=0.01$). Among the study participants, 305 (87.1%) were single and 45 (12.9%) were married. The distribution of patients based on the field of study was as follows: 131 (37.4%) in medicine, 37 (10.6%) in nursing, 37 (10.6%) in obstetrics, 21 (6%) in anesthesiology, 18 (5.1%) in operating room, 21 (6%) in radiology, 31 (8.9%) in laboratory, 27 (7.7%) in health and 27 (7.7%) in other fields. Among the 350 participants, only 70 (20%) lived in a dormitory at the University.

Eighty-six of the 350 participants (24.6%) had experienced frequent episodes of severe headaches, and the definite causes of the headaches were diagnosed in 17 cases (19.8%) before the study. In total, 57 (16.3%) had migraine, among which 28 (49.1%) were female and 29 (50.9%) were male. The mean age of the patients who had migraine was 21.28 ± 2.71 (95% CI: 20.56-22) years. Among the 57 patients with migraine, 32 (56.1%) had experienced episodes of migraine that lasted from 4 to 72 hours. Pulsatile headaches were reported in 33 (57.9%) of the patients with migraine. Unilateral headache was reported in 24 (42.1%) of the patients with migraine. Forty-nine (87.5%) of the patients with migraine reported that their headaches worsened with exercise. Among the patients with migraine, 20 (35.1%) reported associated nausea and vomiting, 35 (61.4%) had reported photophobia, and 47 (82.5%) reported that noise exacerbated their headaches.

Migraine was more prevalent among entomology students (28.6%) and was less frequent among midwifery students (8.1%) (Chi-Square=8.403; $P=0.677$). Migraine was more prevalent among males (25.4%) in comparison to females (11.9%) (Chi-Square=10.389; $P=0.001$). Migraine was more prevalent among non-married students (17%) than married students (11.1%), but the difference was not statistically significant (Chi-Square=1.014; $P=0.314$). Migraine was more prevalent among dormitory students (16.1%) than it was in students who did not live in a dormitory (17.1%), but this difference was not statistically significant (Chi-Square=0.047; $P=0.828$).

Our findings suggested that there was no relationship between the prevalence of migraine and the field of education ($P=0.677$), entrance year ($P=0.1$), marital status ($P=0.314$), and place of residence ($P=0.828$). More details are shown in Table 1. The mean age of the patients with migraine in this study was 21.28 ± 2.71 years (95% CI: 20.56-22.00), whereas it was 21.27 ± 2.80 (95% CI: 20.95-21.60) for students who did not have migraine ($T=0.015$; $P=0.998$).

Table 1: Prevalence of migraine according to students' characteristic

Classification		Migraine	Non-migraine	Chi-square	P Value
Field of education	Medicine	23 (17.6%)	108 (82.4%)	8.403	0.677
	Nursing	8 (21.6%)	29 (78.4%)		
	Midwifery	3 (8.1%)	34 (91.9%)		
	Anesthesiology	5 (23.8%)	16 (76.2%)		
	Operation room	5 (27.8%)	13 (72.2%)		
	Radiology	2 (9.5%)	19 (90.5%)		
	Laboratory sciences	3 (9.7%)	28 (90.3%)		
	Hygiene	3 (11.1%)	24 (88.9%)		
	Entomology	2 (28.6%)	5 (71.4%)		
	Health Information Technology (HIT)	1 (14.3%)	6 (85.7%)		
	Medical records	1 (14.3%)	6 (85.7%)		
	Medical emergencies	1 (16.7%)	5 (83.3%)		
Entrance year	2004	0 (0%)	1 (100%)	12.028	0.100
	2005	0 (0%)	11 (100%)		
	2006	3 (42.9%)	4 (57.1%)		
	2007	4 (19%)	17 (81%)		
	2008	12 (25%)	36 (75%)		
	2009	3 (6.5%)	43 (93.5%)		
	2010	17 (15.6%)	92 (84.4%)		
	2011	18 (16.8%)	89 (83.2%)		
Marital Status	Single	52 (17%)	253 (83%)	0.314	0.314
	Married	5 (11.1%)	40 (88.9%)		
Place of residence	Dormitory	45 (16.1%)	235 (83.9%)	0.828	0.828
	Non-dormitory	12 (17.1%)	58 (82.9%)		

4. Discussion

Migraine is the most prevalent type of headache. Its high prevalence among students and its negative effects on their lives and daily activities show the importance of migraine as a case study. This study assessed the prevalence of migraine among medical students. Two-thirds of the students in our study were females, and about 87% of the participants were single. The prevalence of migraine was 16.3% in our study. The high prevalence of migraine among medical students in our study can be related to stress, inadequate sleep, and their heavy workload. Some earlier studies in Iran have confirmed the higher rates of migraine among medical students (28).

Prevalence rates of 7.3 to 23.8% have been reported among medical students in these studies (28). Also, one study in Turkey reported a prevalence of 12.6% among students (22). Conversely, one study in Iran reported the lowest prevalence of migraine among medical students in comparison to other students (29). Only 17 students (19.8%) in our study had visited a physician for their headache, which may be due to their workload and inattention to their own health issues. This percentage was reported as 17.7% in a study in Tabriz and 18% by Wang et al. (2000) (30, 31).

The prevalence of migraine was higher in males than in females in our study. Some studies have reported higher rates of migraine among women (14, 25, 31, and 32), while others reported that there is association between migraine and gender (33, 34). Different from some other studies conducted in Iran (28, 31), we found no difference in the rates of migraine between single and married students, and our findings were in agreement with the results of a study conducted by Vukovic et al. (35). In our study, the prevalence of migraine was not significantly greater among dormitory students than students who lived elsewhere.

5. Conclusion

In our study, the prevalence of migraine was high among students of the medical sciences, which was in agreement with the findings of other studies. This may be due to their stress, duties, and workload. Solutions should be considered since headaches reduce the quality of their work and have detrimental effects on their health.

Acknowledgements

We want to thank to Hormozgan University of Medical Sciences (HUMS) research committee for their supports and helps for conduction this project and preparing the manuscript. The article is prepared based on the scientific writing guidelines (38, 39).

Conflict of Interest:

There is no conflict of interest to be declared.

Authors' contributions:

All of authors contributed to this project and article equally. All authors read and approved the final manuscript.

References

1. Domingues RB, Cezar PB, Schmidt Filho J, de Moraes Filho MN, Pinheiro MN, Marchiori JG, et al. Prevalence and impact of headache and migraine among Brazilian Tupiniquim natives. *Arquivos de neuro-psiquiatria*. 2009;67(2B):413-5. Epub 2009/07/23.
2. Steiner TJ, Scher AI, Stewart WF, Kolodner K, Liberman J, Lipton RB. The prevalence and disability burden of adult migraine in England and their relationships to age, gender and ethnicity. *Cephalalgia : an international journal of headache*. 2003;23(7):519-27. Epub 2003/09/03.
3. Houinato D, Adoukonou T, Ntsiba F, Adjien C, Avode DG, Preux PM. Prevalence of migraine in a rural community in south Benin. *Cephalalgia : an international journal of headache*. 2010;30(1):62-7. Epub 2009/06/06.
4. Lu SR, Fuh JL, Juang KD, Wang SJ. Migraine prevalence in adolescents aged 13-15: a student population-based study in Taiwan. *Cephalalgia : an international journal of headache*. 2000;20(5):479-85. Epub 2000/10/19.
5. Bigal ME, Liberman JN, Lipton RB. Age-dependent prevalence and clinical features of migraine. *Neurology*. 2006;67(2):246-51. Epub 2006/07/26.
6. Piovesan EJ, Kowacs PA, Lange MC, Pacheco C, Piovesan LR, Werneck LC. [Prevalence and semiologic aspects of the idiopathic stabbing headache in a migraine population]. *Arquivos de neuro-psiquiatria*. 2001;59(2-A):201-5. Epub 2001/06/13. Prevalencia e características da cefaleia idiopática em punhaladas em uma população de migranosos.
7. Lusic I. Population variation in migraine prevalence--the unsolved problem. *Collegium antropologicum*. 2001;25(2):695-701. Epub 2002/01/29.
8. Lucchetti G, Peres MF. The prevalence of migraine and probable migraine in a Brazilian favela: results of a community survey. *Headache*. 2011;51(6):971-9. Epub 2011/06/03.
9. Winkler AS, Dent W, Stelzhammer B, Kerschbaumsteiner K, Meindl M, Kaaya J, et al. Prevalence of migraine headache in a rural area of northern Tanzania: a community-based door-to-door survey. *Cephalalgia : an international journal of headache*. 2010;30(5):582-92. Epub 2009/09/09.
10. Ertas M, Baykan B, Orhan EK, Zarifoglu M, Karli N, Saip S, et al. One-year prevalence and the impact of migraine and tension-type headache in Turkey: a nationwide home-based study in adults. *The journal of headache and pain*. 2012;13(2):147-57. Epub 2012/01/17.
11. Radtke A, Neuhauser H. Prevalence and burden of headache and migraine in Germany. *Headache*. 2009;49(1):79-89. Epub 2009/01/08.
12. Lampl C, Buzath A, Baumhackl U, Klingler D. One-year prevalence of migraine in Austria: a nation-wide survey. *Cephalalgia : an international journal of headache*. 2003;23(4):280-6. Epub 2003/04/29.
13. Moens G, Johannik K, Verbeek C, Bulterys S. The prevalence and characteristics of migraine among the Belgian working population. *Acta neurologica Belgica*. 2007;107(3):84-90. Epub 2007/12/13.
14. Ofovwé GE, Ofili AN. Prevalence and impact of headache and migraine among secondary school students in Nigeria. *Headache*. 2010;50(10):1570-5. Epub 2010/10/16.
15. Fallahzadeh H, Alihaydari M. Prevalence of migraine and tension-type headache among school children in Yazd, Iran. *Journal of pediatric neurosciences*. 2011;6(2):106-9. Epub 2012/03/13.
16. Kowacs PA, Piovesan EJ, Lange MC, Werneck LC, Tatsui CE, Ribas LC, et al. Prevalence and clinical features of migraine in a population of visually impaired subjects in Curitiba, Brazil. *Cephalalgia : an international journal of headache*. 2001;21(9):900-5. Epub 2002/03/21.
17. Houinato D, Adoukonou T, Ntsiba F, Adjien C, Avode DG, Preux PM. Prevalence of migraine in a rural community in south Benin. *Cephalalgia : an international journal of headache*. 2010;30(1):62-7. Epub 2009/06/06.
18. Vukovic V, Plavec D, Pavelin S, Janculjak D, Ivankovic M, Demarin V. Prevalence of migraine, probable migraine and tension-type headache in the Croatian population. *Neuroepidemiology*. 2010;35(1):59-65. Epub 2010/06/05.
19. Neut D, Fily A, Cuvellier JC, Vallee L. The prevalence of triggers in paediatric migraine: a questionnaire study in 102 children and adolescents. *The journal of headache and pain*. 2012;13(1):61-5. Epub 2011/11/02.
20. Pareja-Angel J, Campo-Arias A. [The prevalence of symptoms of anxiety and depression in female migraine sufferers]. *Revista de neurologia*. 2004;39(8):711-4. Epub 2004/10/30. Prevalencia de sintomas de ansiedad y de depresion en mujeres con migraña.
21. Muniz R, Macia C, Montiel I, Gonzalez O, Martin R, Asensio M, et al. [Prevalence of migraine in the medical student population as determined by means of the 'Alcoi 1992' questionnaire]. *Revista de*

- neurologia. 1995;23(122):870-3. Epub 1995/07/01. Prevalencia de migrana en una poblacion de estudiantes de medicina mediante el cuestionario 'Alcoi-1992'.
22. Balaban H, Semiz M, Senturk IA, Kavakci O, Cinar Z, Dikici A, et al. Migraine prevalence, alexithymia, and post-traumatic stress disorder among medical students in Turkey. *The journal of headache and pain*. 2012. Epub 2012/04/27.
 23. Wahab KW, Ugheoke AJ. Migraine: prevalence and associated disability among Nigerian undergraduates. *The Canadian journal of neurological sciences Le journal canadien des sciences neurologiques*. 2009;36(2):216-21. Epub 2009/04/22.
 24. Patel NV, Bigal ME, Kolodner KB, Leotta C, Lafata JE, Lipton RB. Prevalence and impact of migraine and probable migraine in a health plan. *Neurology*. 2004;63(8):1432-8. Epub 2004/10/27.
 25. Lipton RB, Stewart WF, Diamond S, Diamond ML, Reed M. Prevalence and burden of migraine in the United States: data from the American Migraine Study II. *Headache*. 2001;41(7):646-57. Epub 2001/09/14.
 26. Michel P, Dartigues J, Henry P, Tison S, Auriacombe S, Brochet B, Et al. Validity of the International Headache Society Criteria for Migraine. *Neuroepidemiology* 1993;12:51-57.
 27. IBM Corp. Released 2011. IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp.
 28. Hashemilar M , Aminisani N , Savadioskouei D , Yousefian M. The Prevalence of Migraine Among Students of Ardebil University Of Medical Sciences (JAUMS). *March-June 2004*; 3(11):64-70.
 29. Modara F , Rostamkhani M . Prevalence of tension and migraine headaches among the students of Ilam Medical University . *Journal of Ilam University of Medical Sciences* , 2008;15(4): 17-19.
 30. Wang SJ, Fuh JL, Young YH, Lu SR, Shia BC. Prevalence of migraine in Taipei, Taiwan: a population-based survey. *Cephalalgia : an international journal of headache*. 2000;20(6):566-72. Epub 2000/11/15.
 31. Sadreddini SA , Pashapour A , Talebi M. Assessment of Prevalence and Type of Headache in Medical Students. *Medical journal of Tabriz university of medical sciences*. 2006 ;28 (3):73-79.
 32. Jimenez-Sanchez S, Fernandez-de-Las-Penas C, Jimenez-Garcia R, Hernandez-Barrera V, Alonso-Blanco C, Palacios-Cena D, et al. Prevalence of Migraine Headaches in the Romany Population in Spain: Sociodemographic Factors, Lifestyle and Co-Morbidity. *Journal of transcultural nursing : official journal of the Transcultural Nursing Society / Transcultural Nursing Society*. 2012. Epub 2012/07/18.
 33. Unalp A, Dirik E, Kurul S. Prevalence and clinical findings of migraine and tension-type headache in adolescents. *Pediatrics international : official journal of the Japan Pediatric Society*. 2007;49(6):943-9. Epub 2007/11/30.
 34. Shahrakai MR, Mirshekar H, Tajik Ghanbari A, Shahraki AR, Shahraki E. Prevalence of Migraine Among Medical Students in Zahedan Faculty of Medicine (Southeast of Iran). *Basic and clinical Neuroscience*. 2011; 2 (2):20-25.
 35. Vukovic V, Plavec D, Pavelin S, Janculjak D, Ivankovic M, Demarin V. Prevalence of migraine, probable migraine and tension-type headache in the Croatian population. *Neuroepidemiology*. 2010;35(1):59-65. Epub 2010/06/05.
 36. Jalalian M, Danial A. H. Writing for academic journals: A general approach. *Electronic Physician*. 2012; 4(2):474-476. doi: 10.14661/2012.474-476
 37. Jalalian M. Writing an eye-catching and evocative abstract for a research article: A comprehensive and practical approach. *Electronic Physician*. 2012; 4(3):520-524. doi: 10.14661/2012.520-524