

Prevalence of parasitic infections in HIV-positive patients in southern Ethiopia: A cross-sectional study

Sintayehu Fekadu¹, Kefyalew Taye², Wondu Teshome³, Solomon Asnake¹

¹Department of Medical Microbiology and Parasitology, College of Medicine and Health Sciences, Hawassa University, Ethiopia

²Department of Pediatrics and Child Health, College of Medicine and Health Sciences, Hawassa University, Ethiopia.

³Department of Public and Environmental Health, College of Medicine and Health Sciences, Hawassa University, Ethiopia

Abstract

Introduction: Intestinal parasitic infections are a major public health burden in tropical countries. Although all HIV/AIDS patients are susceptible to parasitic infections, those having lower immune status are at greater risk. The aim of this study was to determine the prevalence of intestinal parasitic infections in patients living with HIV/AIDS.

Methodology: This was a facility-based cross-sectional study. A total of 343 consecutively sampled HIV/AIDS patients from the HIV care clinic of Hawassa University Referral Hospital were included. Subjects were interviewed for demographic variables and diarrheal symptoms using structured questionnaires. Stool examinations and CD4 cells counts were also performed.

Results: The prevalence of intestinal parasitic infection was 47.8% among HIV/AIDS patients; single helminthic infection prevalence (22.7%) was higher than that the prevalence of protozoal infections (14.6%). About 54% of study participants had chronic diarrhea while 3.4% had acute diarrhea. The prevalence of intestinal parasites in patients with chronic diarrhea was significantly higher than in acute diarrhea ($p < 0.05$). Non-opportunistic intestinal parasite infections such as *Ascaris lumbricoides*, *Taenia* spp., and hookworm were commonly found, regardless of immunestatus or diarrheal symptoms. Opportunistic and non-opportunistic intestinal parasitic infection were more frequent in patients with a CD4 count of $< 200/\text{mm}^3$ (OR=9.5; 95% CI: 4.64-19.47) when compared with patients with CD4 counts of ≥ 500 cells/ mm^3 .

Conclusions: Intestinal parasitic infections should be suspected in HIV/AIDS-infected patients with advanced disease presenting with chronic diarrhea. Patients with low CD4 counts should be examined critically for intestinal parasites, regardless of diarrheal status.

Key words: CD4; HIV/AIDS; intestinal parasite; opportunistic parasite

J Infect Dev Ctries 2013; 7(11):868-872. doi:10.3855/jidc.2906

(Received 01 August 2012– Accepted 20 June 2013)

Copyright © 2013 Fekadu *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Intestinal parasitic infections are among the most widespread of all chronic human infections worldwide and remain an important cause of morbidity and mortality in developing countries [1]. The rate of infection is remarkably high in sub-Saharan Africa, where the majority of human immunodeficiency virus (HIV) infections and AIDS cases are concentrated [2]. In Ethiopia, the ecology of intestinal parasitism is very wide [3]. Further, it has been estimated that over two million people have been infected [4]. Lack of sanitary facilities and health education are major reasons for the transmission of intestinal parasites [1].

HIV infection is a worldwide problem, having infected approximately 33.4 million people globally;

sub-Saharan Africa accounts for more than half (22.4million) of this number [5]. In Ethiopia, the prevalence of HIV/AIDS is 2.4%, with variances across regions[6]. Intestinal parasitic infections, one of the major health problems in sub-Saharan Africa, are common in these patients.

The involvement of HIV/AIDS in gastrointestinal infection is common and is almost universal; significant disease occurs in 50-90% of patients, while diarrhea can be a presenting manifestation or a life-threatening complication in the course of the disease [7]. Several factors such as autonomic dysfunction, medication, immunedysregulation, and nutritional supplementation play a major role in diarrhea related to HIV/AIDS [8]. Microorganisms and parasites are

also the main etiological agents of diarrhea in HIV/AIDS patients. Parasitic protozoa such as *Cryptosporidium parvum*, *Isospora belli*, *Cyclospora* species, *Giardia intestinalis*, *Entamoebahistolytica*, *Blastocystishominis*, and *Microsporidia* have been associated with acute and chronic diarrhea in HIV patients. The nematode species *Strongyloides stercoralis* can also cause diarrhea and overwhelming infestation in patients with suppressed immune status mainly due to HIV/AIDS. In addition, helminths such as *Ascaris lumbricoides*, *Taenia* spp., *Ancylostoma duodenale*, and *Necator americanus* are commonly diagnosed in HIV/AIDS patients [7].

Investigations into the relationship between HIV/AIDS and enteric parasites in Asia and South America have indicated that several species of protozoa and parasites are involved in acute and chronic diarrhea in HIV/AIDS patients [9,10]. In Malaysia, Thailand, Cuba, and Brazil, reports have shown that opportunistic coccidian intestinal parasites and *S. stercoralis* were significantly more frequent in the low-immunity group of HIV-infected patients with diarrhea, while non-opportunistic intestinal parasite infections such as hookworms, *Opisthorchis viverrini*, and *Ascaris lumbricoides* were commonly found in HIV-infected people, regardless of immune status or diarrheal symptoms [10,11,16]. Findings from African countries such as Ivory Coast, Tanzania, and Ethiopia indicate a high prevalence of intestinal parasites in HIV/AIDS patients. *Cryptosporidium parvum* had the highest prevalence, followed by *Giardia intestinalis*, *Entamoeba histolytica*, *Isospora belli*, and *Blastocystis hominis* [9,13-15]. Helminth infection rate was also higher among HIV-seropositive subjects, with mainly *Strongyloides stercoralis* causing overwhelming infestation in such patients [17,18].

Since the types of intestinal parasites infecting humans vary from one locality to another and such a study has not been conducted in the Ethiopian region, there is a need to clarify the magnitude of the problem and to develop a plan for intervention. Hence, this cross-sectional study was designed to investigate the magnitude of intestinal parasitic infection in HIV/AIDS patients attending the HIV care clinic of Hawassa Teaching and Referral Hospital in south Ethiopia.

Methodology

Study area

Hawassa University Referral Hospital (HURH) is located in Hawassa City, the capital of the Southern Nations and Nationalities People's Regional State

(SNNPRS), located 275 kilometres south of Addis Ababa, the capital of Ethiopia. Established in 2005, it is the only referral and teaching hospital in the region. The hospital provides various types of health services.

Study design

After being approved by the institutional ethical committee, this cross-sectional study was conducted between August and December 2008, examining consecutively selected 343 HIV/AIDS patients visiting the hospital. Informed consent and socio-demographic, clinical, and laboratory data were obtained from the study subjects using a pre-tested and structured questionnaire and laboratory analysis of stool and blood samples. The main variables included in the questionnaires were age, sex, and presence of diarrheal diseases.

Sample collection and processing

Patients with confirmed HIV/AIDS who had not received any anti-protozoal and anti-helminthic in the past three weeks were enrolled in the study. Both blood and stool samples were collected from these participants. Five milliliters of EDTA blood from each subject was collected for the CD4 cells count. The CD4 cells count determination was performed by an automated flow cytometry analyzer, FACS Calibur (Becton Dickinson, Franklin Lakes, USA). All subjects were categorized by their immune status according to the 1993 revised classification system for HIV infection by CD4 T-cell categories. Stool samples for parasitological investigation were collected from each study subject in leak-proof plastic containers and were processed by both simple direct mount and concentration technique in the Microbiology and Parasitology laboratory unit of HURH. In the direct mount technique, match stick head stool was mixed with a drop of normal saline and examined microscopically. Concentration of parasites was made using formol-ether concentration technique followed by microscopic examination.


Statistical analysis

The data were entered and statistical analyses were done using SPSS version 16.0. Prevalence and odds ratios were calculated. Observed differences in data were considered significant and noted in the text if $p < 0.05$.

Results

During the six-month study period, a total of 343 subjects were enrolled and examined for intestinal

Figure 1: Prevalence of intestinal parasites among HIV/AIDS patients in Hawassa Teaching and Referral Hospital, 2008



parasites. The age range of the study subjects was 15 to 60 years. There were 113 (32.94%) males (Table 1). The majority of study participants (n = 290; 84.5 %) had low incomes (earned less than 1.25 USD per day). Among the study population, 164 (47.8%) had at least a single parasitic infection. Thirty-six (10.6%) had dual infections (Figure 1).

More than half of the study participants (n = 185; 53.9%) had chronic diarrhea, and 11 (4.3%) had acute diarrhea. The prevalence of parasitic infection was higher in patients with chronic diarrhea than in patients with acute diarrhea (55.7% vs. 17.3%).

Patient classification by the CD4 cells count categories in relation to intestinal parasite infection showed a relatively higher prevalence of intestinal parasitic infection in those patients having lower CD4 cells counts (CD4 < 200/μL) (Table 2). The odds of having any parasitic infections were 2.08 (95% CI 1.09, 4.00) times higher among patient with low CD4 (< 200 cells/mm³) counts when compared with those who had CD4 counts between 200 and 349 cells/mm³. Chi square for the linear trend revealed that infection with any kind of parasites significantly increased as the CD4 cells declined (χ² for linear trend test = 60.71, p < 0.001).

Discussion

HIV infection is a major global public health problem; the sub-Saharan region of Africa has the highest rates of the disease [2]. A large number of adults with HIV/AIDS have chronic diarrhea [11]. In

patients infected with HIV/AIDS, intestinal parasitic infections are among the leading causes of morbidity and mortality.

Deterioration of the immune system due to HIV infection predisposes patients to numerous opportunistic infections, among which gastrointestinal parasitic infection, which usually presents with diarrheal symptoms, is universally recognized.

The results of this study indicated that pathogenic and opportunistic intestinal parasites are common among HIV/AIDS patients with chronic diarrhea. This finding was in agreement with other studies done in Ethiopia [14,15,17] and other African countries [21,22,18].

The overall prevalence of intestinal parasites in HIV/AIDS patients in this study was 47.8%, which was in agreement with results obtained in southwestern Ethiopia (41% and 44.8%) [3,14], and in Brazil (40%) [25], but lower than the prevalence found in northwestern Ethiopia (67.6%) and central Ethiopia (70%) [26,23] and in Thailand (50%) [11]. Among patients with chronic diarrhea, the opportunistic protozoan parasite *C. parvum* was isolated; this finding was in agreement with studies done in Ethiopia (11%), Cuba (11.9%), and India (11%) [14,10,27]. Other opportunistic parasites such as *C. catyensis* were not identified in this study, due possibly to the use of a single stool specimen.

The prevalence of *Entamoeba histolytica/dispar* (7%) was similar to the prevalence found in Nigeria

Table 1: Age and sex distribution of study participants.

Variable	No	%	
Age in years	15-24	64	18.70%
	25-34	149	43.40%
	35-44	91	26.50%
	45-60	39	11.40%
Sex	Male	113	32.94%
	Female	230	67.06%
Total	343	100.00%	

Table 2: Univariate analysis of CD4 cells count level as determining factor of parasitic infections.

	Parasite		OR (95% CI)
	Present	Absent	
	18 (25.35%)	53 (74.65%)	I(reference)
	29 (28.15%)	74 (71.85%)	1.15 (0.58, 2.29)
	46 (60.52%)	30 (39.48%)	4.51 (2.23, 9.14)
	71 (76.34%)	22 (23.66%)	9.5 (4.64, 19.47)

(5.7%) and Malaysia (6.7%) [18,24], greater than the prevalence found in Brazil (3.3%) [25], but lower than found in Ethiopia in other studies (10.3% and 24.6%) [3,23]. Similarly, the prevalence rate of *Giardia lamblia* (4.1%) was in agreement with rates found in Ethiopia (3.8%) and Brazil (3.5%) in other studies [3,25]. The prevalence was greater than that found in Nigeria (2.9%) [18], and lower than that found in Malaysia (8.3%) and Brazil (16%) [16,25]. In general, such prevalence of these non-opportunistic parasites in the present study could have been the effect of water and food contamination, or it could have been due to poor personal hygiene.

The 10.5% prevalence of the nematode *S. stercoralis* was similar to the prevalence found in other studies in Ethiopia (11.5% and 13.0%) [18], but lower than the prevalence found in Brazil (30.1%) [25], and greater than found in other studies in Ethiopia (5.1%) and Brazil (2%) [23,25]. The high prevalence might be because *S. stercoralis* has been shown to be endemic in developing countries, where it is associated with situations that lead to immunodeficiency, including HIV infection [28]. The life cycle of this helminth favors self-infection, with the possibility of hyper infection and spreading, both of which are aggravated in conditions of immunosuppression [29]. The prevalence of *Ascaris lumbricoides* (6.7%), was lower than that found in other studies in Ethiopia (22.2%), Nigeria (20%), and Brazil (15.6%), [23,18,24] but greater than the

prevalence found in another study in Brazil (2.5%) [25].

From the findings, chronic infection with helminthes explains some of the elements of chronic immune activation that may also be found in HIV infection [12]. This type of immune activation has been suggested as a major factor for the increased susceptibility and progression of HIV infection in Africa and other developing countries [13].

The CD4 cells counts of the study's subjects negatively correlated to the parasite count. Individuals with a CD4 cells count of less than or equal to 200/ μ L had the highest prevalence of parasitic infection.

Conclusions

In conclusion, both opportunistic and non-opportunistic intestinal parasite infections are highly prevalent in HIV-infected patients in tropical Africa, particularly in patients with lower immune statuses. Routine examination of stool samples for intestinal parasites is recommended, irrespective of diarrheal status, in the management of HIV-infected patients, as it can reduce morbidity and mortality associated with intestinal parasitic infection.

The study had some potential limitations. First, the stool examinations were done only once and may not have accurately depicted the parasite prevalence. Second, the study was facility-based and thus could be difficult to generalize to the general public.

Acknowledgements

We would like to thank Netherlands Program for the Institutional Strengthening of Post-secondary Education and Training Capacity (NPT/ETH/107) project for financial support. Our thanks also go to Hawassa University Referral Hospital ART clinic case team who provided necessary information and recruitment of study subjects.

References

- Chan MS (1997) The global burden of intestinal nematode infections-fifty years on. *Parasitology Today* 13: 438-443.
- Grant AD, De Cock KM (1998) The growing challenge of HIV/AIDS in developing countries. *Br Med Bull* 54: 369-381.
- Hailemariam G, Kassu A, Abebe G, Abate E, Damte D, Mekonnen E, Ota F (2004) Intestinal parasitic infections in HIV/AIDS and HIV seronegative Individuals in a Teaching Hospital, Ethiopia. *Jpn J Infect Dis* 57: 41-43.
- Joint United Nations Programme on HIV/AIDS/World Health Organization (2002): HIV epidemic update Geneva.
- Joint United Nations Programme on HIV/AIDS (UNAIDS) (2005) World Health Organisation (WHO) HIV/AIDS epidemic update: UNAIDS/WHO 2-41.
- Ministry of health and national HIV/AIDS prevention and control office (2006) AIDS in Ethiopia sixth report.
- Framm SR, Soave R (1997) Agents of Diarrhea. *Med Clin North Am* 81:427-447.
- Dallabetta GA, Miotti PG (1992) Chronic diarrhoea in AIDS patients in the tropics: a review. *Trop Doct* 22: 3-9.
- Fleming AF (1990) Opportunistic infections in AIDS in developed and developing countries. *Trans R Soc Trop Med Hyg* 84 Suppl 1: 1-6.
- Escobedo AA, Nunez FA (1999) Prevalence of intestinal parasites in Cuban acquired immunodeficiency syndrome (AIDS) patients. *Acta Trop* 72: 125-130.
- Wiwantit V (2001) Intestinal parasitic infections in Thai HIV-infected patients with different immunity status. *BMC Gastroenterology* 1: 3.
- Moura H, Fernandes O, Viola JP, Silva SP, Passos RH, Lima DB (1989) Enteric parasites and HIV infection: occurrence in AIDS patients in Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz* 84: 527-533.
- Tarimo DS, Killewo JZJ, Mnjas JN and Msamanga GI (1996) Prevalence of intestinal parasites in patient with enteropathic AIDS in North-Eastern Tanzania. *E Afr Med J* 73: 397-399.
- Awole M, Gebre-Selassie S, Kassa T, Kibru G (2003) Prevalence of Intestinal Parasites in HIV-Infected adult patients in Southwestern Ethiopia. *Ethiop J Health Dev* 17: 71-78.
- Mengesha B (1994) Cryptosporidiosis among medical Patients with AIDS in TikurAnbessa Teaching hospital, Ethiopia. *E Afr Med J* 71: 376-378.
- Oguntibeju OO (2006) Prevalence of Intestinal Parasites In HIV-Positive/AIDS Patients. *Malays J Med Sci* 13: 68-73.
- Fisseha B, Petros B, Woldemichael T. (1998) Cryptosporidium and other parasites in Ethiopia AIDS Patients with chronic diarrhea. *East Afr Med J* 75: 100-101.
- Okodua M, Adeyeba OA, Tafeng YM, Okpala HO (2003) Age and Sex Distribution of Intestinal Parasitic Infection Among HIV Infected Subjects in Abeokuta, Nigeria. *Online J Health Allied Scs* 4: 3.
- Fincham J (2003) HIV/AIDS and Tuberculosis. *Science in Africa. AfricaFirst On-Line Science Magazine* 19: 44-58.
- Conlon CP, Pinching AJ, Perera CU, Moody A, Luo NP, Lucas SB (1990) HIV-related enteropathy in Zambia: a clinical, microbiological, and histological study. *Am J Trop Med Hyg* 42: 83-88.
- Sarfati C, Bourgeois A, Menotti J, Liegeois F, Moyou-Somo R, Delaporte E, Derouin F, Ngole EM, Molina JM (2006) Prevalence of intestinal parasites including microsporidia in human immunodeficiency virus-infected adults in Cameroon: a cross-sectional study. *Am J Trop Med Hyg* 74: 162-164.
- Sewankambo N, Mugerwa RD, Goodgame R, Carswell JW, Moody A, Lloyd G, Lucas SB (1987) Enteropathic AIDS in Uganda, An endoscopic, histologic and microbiologic study. *AIDS* 1: 9-13.
- Fontanet AL, Sahlut, Rinke DWT, Messele T, Masho W, Woldemichael T, Yeneneh H, Coutinho RA (2000) Epidemiology of infections with intestinal parasites and human immunodeficiency virus (HIV) among sugar-estate residents in Ethiopia. *Ann Trop Med Parasitol* 94: 269-278.
- Bachur TP, Vale JM, Coelho IC, Queiroz TR, Chaves Cde S (2008) Enteric parasitic infections in HIV/AIDS patients before and after the highly active antiretroviral therapy. *Braz J Infect Dis* 12: 115-122.
- Cimerman S, Cimerman B, Lewi DS (1999) Prevalence of intestinal parasitic infections in patients with acquired immunodeficiency syndrome in Brazil. *Int J Infect Dis Summer* 3: 203-206.
- Moges F, Kebede Y, Kassa A, Degu G, Tiruneh M, Gedefaw M (2006) Infection with HIV and intestinal parasites among street dwellers in Gondar city, northwest Ethiopia. *Jpn J Infect Dis* 59: 400-403.
- Prasad KN, Nag VL, Dhole TN, Ayyagari A (2000) Identification of enteric pathogens in HIV-positive patients with diarrhoea in northern India. *J Health Popul Nutr* 18: 23-26.
- Keiser PB, Nutman TB (2004) *Strongyloides stercoralis* in the Immunocompromised Population. *Clin Microbiol Rev* 17: 208-217.
- Gomez Morales MA, Atzori C, Ludovisi A, Rossi P, Scaglia M, Pozio E (1995) Opportunistic and non-opportunistic parasites in HIV-positive and negative patients with diarrhoea in Tanzania. *Trop Med Parasitol* 46: 109-114.
- Mariam ZT, Abebe G, Mulu A (2008) Opportunistic and other intestinal parasitic infections in AIDS patients, HIV seropositive healthy carriers and HIV seronegative individuals in southwest Ethiopia. *East Afr J Public Health* 5: 169-173.

Corresponding author

Sintayehu Fekadu Kebede
 Department of Medical Microbiology and Parasitology
 College of Medicine and Health Sciences
 Hawassa University, P.O. box 1560, Ethiopia
 Phone: +251-911-790-197/+251-920-568-817
 Fax: +251-462-208-755
 Email: sintayehufekadukebede@gmail.com

Conflict of interests: No conflict of interests is declared.