

Research Article

Prevalence of *Salmonella* Infection in Dogs in Maiduguri, Northeastern Nigeria

Saleh Mohammed Jajere,¹ Samson Amali Onyilokwu,²
Nuhu Bala Adamu,¹ Naphtali Nayamanda Atsanda,¹ Adamu Saleh Saidu,³
Shuaibu Gidado Adamu,¹ and Fatima Bukar Mustapha¹

¹ Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, University of Maiduguri, PMB 1069, Maiduguri, Borno State, Nigeria

² Veterinary Microbiology Research Laboratory, Department of Veterinary Microbiology and Parasitology, Faculty of Veterinary Medicine, University of Maiduguri, PMB 1069, Maiduguri, Borno State, Nigeria

³ Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, Ahmadu Bello University, PMB 1096, Zaria, Kaduna State, Nigeria

Correspondence should be addressed to Saleh Mohammed Jajere; drmsalehjajere@gmail.com

Received 23 August 2014; Accepted 7 October 2014; Published 23 October 2014

Academic Editor: Toni L. Poole

Copyright © 2014 Saleh Mohammed Jajere et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The prevalence and antimicrobial sensitivity of *Salmonella* from dogs in Maiduguri Metropolis were determined using standard bacteriological methods to assess the risk of possible transmission of *Salmonella* infection from dogs to humans. Of 119 samples, *Salmonella* was isolated from 52 (43.7%). Males had higher prevalence of 50.0% compared with 34.7% in females ($P < 0.05$). Dogs older than 24 months had higher prevalence of 61.0% and the lowest was seen in dogs aged 13–24 months ($P < 0.05$). The prevalence of 31.8%, 41.2%, and 58.8% was observed in dogs aged 3–6, 10–12, and 7–9 months, respectively. High prevalence of 49.5% was observed in Mongrels, while Terrier and Alsatian breeds had 30.0% and 8.3%, respectively. *Salmonella* isolates from Alsatian and Terrier breeds showed about 100% susceptibility to all the tested antimicrobials. Higher percentage of the *Salmonella* isolates from Mongrels also showed susceptibility to ciprofloxacin (89.7%), amoxicillin (87.6%), vancomycin (86.6%), and chloramphenicol (84.5%). However about 50% of these isolates showed resistance to ofloxacin. The carrier status of *Salmonella* is high among dogs especially Mongrels. Therefore good environmental hygiene, discouraging straying coupled with feeding of dogs with properly cooked and uncontaminated feeds was recommended to mitigate risk of human salmonellosis.

1. Introduction

Salmonellosis is a broad term applied to enteric infections caused by a group of gram-negative bacteria of the genus *Salmonella*, belonging to the family Enterobacteriaceae [1]. *Salmonella enterica*, the causative agent of salmonellosis, is known to be a primary inhabitant of the gastrointestinal tract of numerous animal species including humans and mostly affects a wide host range with the exception of a few *Salmonella* serovars that are host-specific: *Salmonella* Typhi in humans, *S. Gallinarum* and *S. Pullorum* in poultry, and *S. Dublin* in cattle [1]. They are common contaminants of wide range of food, eggs, vegetables, and water and are therefore considered as the most common causes of

foodborne zoonotic infection worldwide [2]. Consequently, they are recognized as a major public health problem in both developed and developing countries [3]. Feaces of nearly all-animal species including dogs may serve as a potential source of *Salmonella* infection to humans and even to other animals [1]. Several studies have documented the isolation of *Salmonella* spp. from healthy dogs and subsequent transmission of the infection to humans [4–7]. Other studies have recognized shedding of *Salmonella* by dogs as a possible source of infection for dog owners and their communities [8, 9]. Dogs generally remain resistant to *Salmonella* infections and infected ones may remain carriers and fecal shedding therefore may serve as sources of infection to man and other animals [8]. Due to the increase in dog keeping among

the elites especially those living within the metropolitan cities in Nigeria, there is an increased risk of transmission of *Salmonella* infection to humans. There is paucity of information regarding the role of dogs as sources of *Salmonella* infection to humans in Maiduguri Metropolis and therefore this study aims to determine the carrier status of *Salmonella* in dogs and their role as potential sources of infection to humans.

2. Materials and Methods

2.1. Study Area. Maiduguri is located in the Sahel Savannah region of Northeastern Nigeria at 11°50'–11.83° North Latitude and 13°09'–13.15° East Longitude. It shares an extensive border with Niger to the North, Chad to the Northeast, and Cameroon to the East. Its altitude is about 350 metres above sea level and has an area of 75,540.9 km². Its population is about 4,171,104 with a population density of 60 people per km² (2006 census figures). The climate is favourable, with a mean annual rainfall and temperature of about 650 mm and 32°C, respectively. The months of March and April are the hottest periods of the year with temperatures ranging between 30°C and 40°C. It is usually cold and dry during the harmattan, November to January being the coldest months (<http://en.wikipedia.org/wiki/Maiduguri>).

2.2. Sources, Baseline Characteristics, and Inclusion Criteria of Dogs. The dogs investigated in this study were from Maiduguri Metropolis and its environs. Samples were obtained only from dogs kept at home for security/guard or other purposes and from dogs at the Veterinary Teaching Hospital ($n = 20$), University of Maiduguri staff quarters ($n = 11$), Mairi village ($n = 23$), Gwange ward ($n = 15$), Bulunkutu Polo ($n = 18$), Custom area ($n = 10$), Gommari ($n = 13$), and Ummalari ($n = 9$) wards. A total of 119 faecal samples, comprising 70 males and 49 females, were collected from three breeds of dogs (Nigerian local breed Mongrel, Alsatian, and Terrier breed). The age of the dogs was determined by dental eruption at the time of sample collection. Only one dog was sampled per household visited. Most of the household visited had 3 to 4 dogs. A total of 97 Nigerian local breed Mongrel and 12 Alsatian and 10 Terrier breeds were sampled. All the ages of the sampled dogs ranged from <3 months to older than 2 years. All the dogs were apparently healthy dogs with only 5 (4.2%) having diarrhoea as at the time of sample collection.

2.3. Sample Collection and Bacteriological Analysis. Rectal swabs were collected aseptically using sterile swab sticks (Oxoid) and placed in tubes containing Carry-Blair transport medium (Oxoid, Basingstoke, UK). The samples were immediately transported ice cooled within 8 hours of collection to the Microbiology Laboratory, Faculty of Veterinary Medicine, University of Maiduguri, for further processing and analysis. All the samples collected were then inoculated into 2 mL of Selenite-Faeces-broth (Oxoid, Biotec, Suffolk, UK) for enrichment and incubated aerobically at 37°C for 24 hours. Subcultures were then made from each

broth culture by streaking onto Brilliant Green Agar (BGA) (Oxoid), McConkey agar (Oxoid), *Salmonella-Shigella* Agar (SSA) (Oxoid), and Xylose Lysine deoxycholate (XLD) agar (Oxoid). The cultured plates were incubated at 37°C for 24–48 hours [10]. The cultured plates were then examined for the presence of typical colonies of *Salmonella* based on cultural and morphological characteristics: presumptive *Salmonella* colonies appearing colourless and nonlactose fermenting on McConkey agar, dome-shaped colonies with central black spot on XLD agar, transparent colonies with black centre on SSA, and pink colonies surrounded by a red medium were selected as presumptive *Salmonella* colonies and subjected to further biochemical tests and gram staining as described by Barrow and Feltham [11].

2.4. Antibiotic Sensitivity Testing. Antibiotic susceptibility pattern of isolates to various routine antibiotics was carried out using the Kirby-Bauer disk diffusion method [12] on Mueller-Hinton Agar (Oxoid) as recommended by Anonymous [13]. The susceptibility of the isolates to the following antimicrobials was determined: tetracycline (30 µg), gentamycin (30 µg), chloramphenicol (30 µg), ciprofloxacin (10 µg), ofloxacin (5 µg), streptomycin (10 µg), ampicillin (30 µg), amoxicillin-clavulanic acid (AMC) (5 µg), vancomycin (30 µg), neomycin (25 µg), trimethoprim (25 µg), and erythromycin (15 µg).

2.5. Statistical Analysis. Prevalence of *Salmonella* infection was expressed in percentages and proportions in Microsoft Excel version 2010 before finally importing the data into Statistical Package for Social Sciences (SPSS) software version 16.0 to determine the statistical association between dependent and independent variables. The association between infection and other risk factors was determined using Chi-square test at 5% significance level [14].

3. Results

Of the 119 sampled dogs, 70 (58.8%) and 49 (41.2%) were males and females, respectively; 97 (81.5%), 12 (10.1%), and 10 (8.4%) were Nigerian local breed Mongrel, Alsatian, and Terrier, respectively; 2 (1.2%), 22 (18.5%), 17 (14.3%), 17 (14.3%), 20 (16.8%), and 40 (34.5%) were from dogs aged < 3, 3–6, 7–9, 10–12, 13–24, and older than 24 months, respectively (Table 1).

Of the 119 dogs examined, 52 (43.7%) were positive for *Salmonella*. Of these (52), 50.0% and 34.7% were the male and female infection rates, respectively (Table 2). There was significant difference ($P < 0.05$) in the prevalence of *Salmonella* infection in either sex. Highest prevalence was observed in dogs aged older than 24 months with 61.0% (25/41) and except for dogs < 3 months, the lowest was seen in dogs 13–24 months with 15.0% (3/20). While the prevalence of 31.8%, 41.2%, and 58.8% was observed in dogs aged 3–6, 10–12, and 7–9 months in this order of infection rates (Table 2). There was statistically significant difference ($P < 0.05$) in the prevalence of *Salmonella* infection across the age groups. Highest prevalence of 49.5% (48/97) was observed

TABLE 1: Baseline characteristics of the sampled dogs in Maiduguri, Nigeria ($n = 119$).

Risk factors	Number sampled	% of total population
Sex		
Male	70	58.8
Female	49	41.2
Age (months)		
<3	2	1.7
3–6	22	18.5
7–9	17	14.3
10–12	17	14.3
13–24	20	16.8
>24	41	34.5
Breeds		
Mongrel	97	81.5
Alsatian	12	10.1
Terrier	10	8.4
Overall total	119	100

in the Nigerian local breed Mongrel, while Terrier and Alsatian breeds had 30.0% (3/10) and 8.3% (1/12), respectively (Table 2). This was statistically significant ($P < 0.05$).

The antimicrobial sensitivity testing of the *Salmonella* isolates reveals that *Salmonella* isolates from the Terrier and Alsatian breeds had high susceptibility to all the antimicrobials used, with small pockets of resistance (Table 3). While a higher percentage of the *Salmonella* isolates from the Mongrels showed susceptibility to ciprofloxacin (89.7%), amoxicillin (87.6%), vancomycin (86.6%), chloramphenicol (84.5%), trimethoprim (81.4%), ampicillin (80.4%), neomycin (77.3%), and gentamycin (71.1%) (Table 3), however, about 50% of the isolates from Mongrels were resistant to ofloxacin.

4. Discussion

Dogs as carriers of *Salmonella* spp. worldwide have been incriminated as the potential sources of *Salmonella* infection to humans especially children due to close contact and interaction that exists between them [15]. Cases of *Salmonella* transmission from dogs to humans resulting into severe infection in humans have been documented [16]. It was also reported that the intestinal carriage of *Salmonella* by dogs is more common than the prevalence of clinical disease [9]. The frequency of fecal isolation of *Salmonella* spp. from clinically healthy dogs was reported to be between 0.0% and 43.0% [9, 15], while other studies reported prevalence of *Salmonella* in fecal samples from clinically healthy or hospitalized dogs to range from 1% to 36% [17]. However, recent studies suggested that the prevalence is probably decreasing because more pets are fed commercially processed foods [17]. In the present study, *Salmonella* was isolated from 52 (43.7%) of the dogs examined. This prevalence is within the reported 0.0% and 43.0% range of prevalence of *Salmonella* isolated from clinically healthy dogs [9, 15]. However, this finding

TABLE 2: Prevalence of *Salmonella* among exotic and local breeds of dogs according to sex, age, and breeds in Maiduguri, Nigeria ($n = 119$).

Risk factors	Number sampled	Number (%) positive
Sex		
Male	70	35 (50.0) ^a
Female	49	17 (34.7)
Age (months)		
<3	2	—
3–6	22	7 (31.8) ^a
7–9	17	10 (58.8)
10–12	17	7 (41.2)
13–24	20	3 (15.0)
>24	41	25 (61.0)
Breeds		
Mongrel	97	48 (49.5) ^a
Alsatian	12	1 (8.3)
Terrier	10	3 (30.0)
Total	119	52 (43.7)

^aStatistically significant difference in infection rates ($P < 0.05$).

is consequently higher than the results obtained from other parts of Nigeria and other countries in the world. In similar studies by Britt et al. [18] in Vom, Nigeria reported that 18.3% of 120 domestic dogs were excreting *Salmonella* serotypes. Another study by Khan [19] in Sudan reveals that about 23.5% of 442 dogs were positive for *Salmonella*. The differences in the sample sizes, period of study, type of fecal samples, geographical areas, and isolation methods employed in the various studies above may account for the differences in the prevalence and may all affect the prevalence [20]. The high prevalence of 43.7% in dogs in Maiduguri Metropolis might be associated with poor management conditions, compromised good and hygienic environmental standards, contaminated kennels, and food sources. Mongrels in the study area are mostly kept under less restraint and are left to wander and scavenge for food. This makes them highly exposed to *Salmonella* contaminated food materials in the environment (Table 2).

The absence of *Salmonella* in dogs within the age of <3 months could be attributed to the small number of dogs from which samples were obtained (Table 2). It could also result probably from the protection by the maternal antibodies in dogs within this age bracket. There is seemingly high prevalence of *Salmonella* in dogs within the age groups of 3–6, 7–9, and 10–12 months. This is in concordance with the works of Britt et al. [18] in Vom, Plateau State, that *Salmonella* infections are acquired during the early stages of life in dogs. This could result from the low resistance of the immune system to infection by *Salmonella* during this stage of life in dogs. The low prevalence of 15.0% observed in dogs within 13–24 months could be the result of risen immunity and possibly good environmental and management conditions provided by the pet owners. Morse et al. [4] reported that fecal shedding could last for a period of 6 weeks and this could possibly be the reason why the prevalence was

TABLE 3: Antibiotic susceptibility pattern of *Salmonella* isolates from exotic and local breeds of dogs in Maiduguri, Nigeria ($n = 119$).

Antibiotic (amount in μg)	Mongrel ^a	Alsatian ^b	Terrier ^c	Total
	Number (%) <i>Resistant</i>	Number (%) <i>Resistant</i>	Number (%) <i>Resistant</i>	Number (%) <i>Resistant</i>
Gentamycin (30)	28 (28.9)	0 (0.0)	0 (0.0)	28 (23.5)
Amoxicillin-clavulanic acid (5)	11 (11.3)	0 (0.0)	0 (0.0)	11 (9.2)
Erythromycin (15)	32 (33.0)	1 (8.3)	0 (0.0)	33 (27.7)
Ciprofloxacin (10)	10 (10.3)	0 (0.0)	0 (0.0)	10 (8.4)
Ampicillin (30)	19 (19.6)	2 (16.0)	0 (0.0)	21 (17.6)
Streptomycin (10)	30 (30.9)	0 (0.0)	0 (0.0)	30 (25.2)
Tetracycline (30)	36 (37.1)	0 (0.0)	0 (0.0)	36 (30.3)
Chloramphenicol (30)	15 (15.6)	0 (0.0)	0 (0.0)	15 (12.6)
Ofloxacin (5)	48 (49.5)	1 (8.3)	0 (0.0)	49 (41.2)
Vancomycin (30)	13 (13.4)	0 (0.0)	0 (0.0)	13 (10.9)
Neomycin (25)	22 (22.7)	0 (0.0)	0 (0.0)	22 (18.5)
Trimethoprim (25)	18 (18.6)	0 (0.0)	0 (0.0)	18 (15.1)

^aMongrels ($n = 97$); ^bAlsatian ($n = 12$); ^cTerrier ($n = 10$).

lower in dogs within this age bracket. A high prevalence of 61.0% observed in dogs within the age of >24 months might possibly be attributed to the fact that older dogs are usually immunosuppressed and therefore at high risk of infection [17]. This renders dogs asymptomatic carriers of the infection and consequently contaminates human environment, food, and water resulting in outbreaks of human salmonellosis.

Males had higher prevalence of *Salmonella* infection compared with females (50.0% versus 34.7%). This could be due to the differing number of samples in each sex (Table 2). According to breeds, Nigerian local breed Mongrel showed the highest prevalence of 49.5%. This is in agreement with the work of Britt et al. [18] who highlighted the fact that due to the poor management and compromised sanitary conditions, a large number of Mongrels tend to acquire the infection and shed the organism in the environment. A large population of the Mongrels in the study area is normally kept under less restraint and left to wander around wider range of distances scavenging for food and therefore get exposed to many contaminated materials compared with the Terrier and Alsatian breeds kept under good hygienic conditions. Scavengers are likely to harbor more *Salmonella* serovars than nonscavengers kept under strict restraint and good hygienic conditions [1]. Compromised sanitary conditions could lead to fecal contamination of the food and water of the Mongrels in the environment and therefore could serve as a source of infection in this breed. Finley et al. [21] also observed that when dogs are fed with *Salmonella*-contaminated feed, raw food, and commercially prepared dry foods, they can become infected and consequently shed the organism in their feces to contaminate the environment, domestic animals, other dogs, and even man. The local breed Mongrels are provided with poor environmental conditions and are less cared for compared with the exotic breeds of Terrier and Alsatian. The Alsatian and Terrier breeds are kept at home mostly by the elites (restrained) and fed with commercial food and this

makes them less exposed to *Salmonella* infection compared with the Mongrels (Table 2).

The antimicrobial sensitivity testing results reveal that high percentages of the *Salmonella* isolates from the Alsatian and Terrier breeds were susceptible to the antimicrobials used, while a high percentage of the isolates from Mongrels showed susceptibility to ciprofloxacin (89.7%), amoxicillin-clavulanic acid (AMC) (87.6%), vancomycin (86.6%), and chloramphenicol (84.5%). However, about 50% of these isolates were resistant to ofloxacin (Table 3). In the past decades, the emergence of antibiotic resistant strains of *Salmonella* has become a major public health concern. Mongrels are more likely to harbor *Salmonella* serovars due to their scavenging nature and therefore are more likely to be treated with antibiotics than the exotic breeds in the study area. Alsatian and Terrier breeds are mostly restricted and fed at home and are not allowed to scavenge or stray in the environment. This makes them less likely to acquire *Salmonella* infection and therefore not likely to be treated with antibiotics. Indiscriminate use of antibiotics in animals could be responsible for the emergence of antimicrobial resistant strains of bacteria [1]. Therefore, the use of antibiotics should be well regulated and used only when it is indicated. Ciprofloxacin and amoxicillin-clavulanic acid (AMC) could be useful antibiotics in the treatment of majority of Salmonellosis cases as revealed by their effectiveness against a high percentage of *Salmonella* isolates from the study area.

In conclusion, this study revealed a high *Salmonella* carrier status in dogs kept in households in Maiduguri Metropolis, Nigeria. Mongrels had high *Salmonella* carrier status compared with the other breeds in the study area. Most of the households visited keep Mongrels, where they are left to wander and scavenge for food in the environment and only return back home in the evening. This has public health significance as dogs may pose a risk to humans where close contact between dogs and their owners occurs in households

and children and immunosuppressed individuals may be particularly at risk [6]. As the infection is acquired during the early stage of their life, pet keepers or dog keepers should maintain a high personal and environmental standard hygiene of the dogs to mitigate the potential zoonotic transmission to humans. Properly cooked and uncontaminated feed should be fed to dogs and scavenging and straying of dogs should be discouraged. It is recommended that complete bacteriological identification be carried out on the isolates to determine the serotypes found in the study area.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The authors would like to thank all the technical staff of the Veterinary Microbiology Laboratory, Department of Veterinary Microbiology and Parasitology, Faculty of Veterinary Medicine, University of Maiduguri, Nigeria, for their assistance during this research work.

References

- [1] O. E. Ojo and A. I. Adetosoye, "Salmonella typhimurium infection in diarrhoeic and non-diarrhoeic dogs in Ibadan, Nigeria," *Veterinarski Arhiv*, vol. 79, no. 4, pp. 371–377, 2009.
- [2] E. Scallan, R. M. Hoekstra, F. J. Angulo et al., "Foodborne illness acquired in the United States-Major pathogens," *Emerging Infectious Diseases*, vol. 17, no. 1, 2011.
- [3] K. O. Akinyemi, W. Phillip, W. Beyer, and R. Bohm, "In-vitro antimicrobial susceptibility patterns of *Salmonella enterica* serovars and emergence of *Salmonella* phage type DT071 in a suspected community-associated outbreak in Lagos, Nigeria," *Journal of Infection in Developing Countries*, vol. 1, no. 1, pp. 48–54, 2007.
- [4] E. V. Morse, M. A. Duncan, D. A. Estep, W. A. Riggs, and B. O. Blackburn, "Canine salmonellosis: a review and report of dog to child transmission of *Salmonella enteritidis*," *American Journal of Public Health*, vol. 66, no. 1, pp. 82–84, 1976.
- [5] M. Birnbaum, K. Miller, and G. T. Stratigos, "Salmonella typhi facial infection. Report of case," *Oral Surgery Oral Medicine and Oral Pathology*, vol. 49, no. 3, pp. 219–220, 1980.
- [6] Y. Sato, T. Mori, T. Koyama, and H. Nagase, "Salmonella virchow infection in an infant transmitted by household dogs," *Journal of Veterinary Medical Science*, vol. 62, no. 7, pp. 767–769, 2000.
- [7] R. F. Kahrs, D. N. Holmes, and G. C. Poppensiek, "Diseases transmitted from pets to man: an evolving concern for veterinarians," *The Cornell Veterinarian*, vol. 68, no. 4, pp. 442–459, 1978.
- [8] M. Kozak, K. Horosova, V. Lasanda, J. Bilek, and J. Kyselova, "Do dogs and cats present a risk of transmission of salmonellosis to humans?" *Bratislavske Lekarske Listy*, vol. 104, no. 10, pp. 323–328, 2003.
- [9] S. Sanchez, C. L. Hofacre, M. D. Lee, J. J. Maurer, and M. P. Doyle, "Animal sources of salmonellosis in humans," *Journal of the American Veterinary Medical Association*, vol. 221, no. 4, pp. 492–497, 2002.
- [10] International Organization of Standardization, "Microbiology general guidelines on methods for the detection of *Salmonella*," ISO 6579, International Organization of Standardization, Geneva, Switzerland, 2002.
- [11] G. I. Barrow and R. K. A. Feltham, *Cowan and Steel's Manual for Identification of Medical Bacteria*, Cambridge University Press, Cambridge, UK, 3rd edition, 1993.
- [12] A. W. Bauer, W. M. Kirby, J. C. Sherris, and M. Turck, "Antibiotic susceptibility testing by a standardized single disk method," *American Journal of Clinical Pathology*, vol. 45, no. 4, pp. 493–496, 1966.
- [13] National Committee for Clinical Laboratory Standards, "Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals," Approved Standard M31-A2, National Committee for Clinical Laboratory Standards, Wayne, Pa, USA, 1999.
- [14] G. W. Snedecor and W. G. Cochran, *Statistical Methods*, Iowa State University Press, Ames, Iowa, USA, 8th edition, 1989.
- [15] M. E. Carter and J. P. Quinn, "Salmonella infections in dogs and cats," in *Salmonella in Domestic Animals*, C. Wray and A. Wray, Eds., pp. 231–244, CAB International, Wallingford, UK, 2000.
- [16] E. V. Morse and M. A. Duncan, "Canine salmonellosis: prevalence, epizootiology, signs, and public health significance," *Journal of the American Veterinary Medical Association*, vol. 167, no. 9, pp. 817–820, 1975.
- [17] C. E. Greene, "Salmonellosis," in *Infectious Diseases of the Dog and Cat*, C. E. Greene, Ed., pp. 355–360, Saunders, 3rd edition, 2006.
- [18] D. P. Britt, T. A. Cole, and C. R. Shipp, "Salmonellae from dogs in Vom, Northern Nigeria," *Tropical Animal Health and Production*, vol. 10, no. 4, pp. 215–218, 1978.
- [19] A. Q. Khan, "Salmonella infections in dogs and cats in the Sudan," *British Veterinary Journal*, vol. 126, no. 11, pp. 607–612, 1970.
- [20] N. Seepersadsingh, A. A. Adesiyun, and R. Seebarsingh, "Prevalence and antimicrobial resistance of *Salmonella* spp. in non-diarrhoeic dogs in Trinidad," *Journal of Veterinary Medicine Series B: Infectious Diseases and Veterinary Public Health*, vol. 51, no. 7, pp. 337–342, 2004.
- [21] R. Finley, R. Reid-Smith, C. Ribble, M. Popa, M. Vandermeer, and J. Aramini, "The occurrence and antimicrobial susceptibility of *Salmonellae* isolated from commercially available canine raw food diets in three Canadian cities," *Zoonoses and Public Health*, vol. 55, no. 8–10, pp. 462–469, 2008.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

