

Review Article

Prevalence, Risk Factors, and Antimicrobial Resistance Profiles of Thermophilic *Campylobacter* Species in Humans and Animals in Sub-Saharan Africa: A Systematic Review

Noel Gahamanyi ¹, Leonard E. G. Mboera,¹ Mecky I. Matee,² Diudonné Mutangana,³ and Erick V. G. Komba¹

¹SACIDS Foundation for One Health, College of Veterinary Medicine and Biomedical Sciences, Sokoine University of Agriculture, P.O. Box 3015, Chuo Kikuu, Morogoro, Tanzania

²School of Medicine, Muhimbili University of Health and Allied Sciences, P.O. Box 65001, Dar es Salaam, Tanzania

³College of Science and Technology, University of Rwanda, P.O. Box 3900, Kigali, Rwanda

Correspondence should be addressed to Noel Gahamanyi; gahamanyi.noel@sacids.org

Received 29 May 2019; Revised 25 October 2019; Accepted 26 December 2019; Published 14 January 2020

Academic Editor: Barbara H. Iglewski

Copyright © 2020 Noel Gahamanyi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Thermophilic *Campylobacter* species are clinically important aetiologies of gastroenteritis in humans throughout the world. The colonization of different animal reservoirs by *Campylobacter* poses an important risk for humans through shedding of the pathogen in livestock waste and contamination of water sources, environment, and food. A review of published articles was conducted to obtain information on the prevalence and antimicrobial resistance (AMR) profiles of thermophilic *Campylobacter* species in humans and animals in sub-Saharan Africa (SSA). Electronic databases, namely, PubMed, Google Scholar, Research4life-HINARI Health, and Researchgate.net, were searched using the following search terms “thermophilic *Campylobacter*,” “*Campylobacter jejuni*,” “*Campylobacter coli*,” “diarrhea/diarrhoea,” “antimicrobial resistance,” “antibiotic resistance,” “humans,” “animals,” “Sub-Saharan Africa,” and “a specific country name.” Initially, a total of 614 articles were identified, and the lists of references were screened in which 22 more articles were identified. After screening, 33 articles on humans and 34 on animals and animal products were included in this review. In humans, Nigeria reported the highest prevalence (62.7%), followed by Malawi (21%) and South Africa (20.3%). For *Campylobacter* infections in under-five children, Kenya reported 16.4%, followed by Rwanda (15.5%) and Ethiopia (14.5%). The country-level mean prevalence in all ages and under-five children was 18.6% and 9.4%, respectively. The prevalence ranged from 1.7%–62.7% in humans and 1.2%–80% in animals. The most reported species were *C. jejuni* and *C. coli*. The AMR to commonly used antimicrobials ranged from 0–100% in both humans and animals. Poultry consumption and drinking surface water were the main risk factors for campylobacteriosis. The present review provides evidence of thermophilic *Campylobacter* occurrence in humans and animals and high levels of AMR in SSA, emphasizing the need for strengthening both national and regional multisectoral antimicrobial resistance standard surveillance protocols to curb both the campylobacteriosis burden and increase of antimicrobial resistance in the region.

1. Introduction

Diarrhoea remains the main cause of morbidity and mortality in low- and middle-income countries (LMICs) [1–3]. Worldwide, under-five children experience approximately 1.4 billion episodes of diarrhoea each year, with several medical checks, hospitalizations, and around two million

deaths. Over 78% of diarrhoea cases are found in the LMICs [4]. The burden of diarrhoeal diseases is complicated by the lack of appropriate case management [5], limited ability to detect the aetiologies [6], and antimicrobial resistance [7].

The most common aetiologies of diarrhoea include bacteria such as *Escherichia coli*, *Vibrio cholerae*, *Campylobacter jejuni*, *Salmonella* spp., *Aeromonas* spp., and

Yersinia enterocolitica; viruses mainly rotavirus, norovirus, sapovirus, and adenovirus; and protozoa largely *Entamoeba histolytica*, *Giardia* spp., and *Cryptosporidium* spp. [8, 9]. Of the bacterial aetiologies, *Campylobacter* is a leading cause of gastroenteritis in both high-, middle-, and low- income countries, responsible for 400–500 million cases of diarrhoea each year [10]. The clinically important *Campylobacter* species are *C. jejuni* and *C. coli*, which are responsible for about 98% of all human *Campylobacter* gastroenteritis cases [11, 12].

In most cases, campylobacteriosis does not require any antimicrobial therapy except in severe cases, especially in immune-deficient or immune-suppressed individuals [13, 14]. The recommended drugs are macrolides (mostly erythromycin), fluoroquinolones (mainly ciprofloxacin), and tetracycline [10, 15, 16]. Nevertheless, there is an escalating number of *Campylobacter* isolates resistant to these drugs [17, 18] due to the immeasurable and misuse of antimicrobials [19], not only in animals but also in humans [20]. Several factors have been associated with occurrence of *Campylobacter* infections. They include consumption of different food items like undercooked poultry meat and pork, red meat at barbecue, grapes, and drinking unpasteurized milk, having a chronic illness [21–23], drinking contaminated water, type of water source, animal contact, young age, eating prepared salad, latrine usage, bottle feeding, and nutritional status [24–26]. There is a wide range of natural reservoirs for *Campylobacter* including chicken and other poultry, wild birds, pigs, dogs, cats, sheep, and cows [27, 28]. Consequently, colonization of different reservoirs by *Campylobacter* poses an important risk for humans through shedding of the pathogen in livestock waste and water sources contamination, environment, and food [29, 30].

In LMICs, studies on thermophilic *Campylobacter* species are few due to limited capacity in laboratory diagnosis [31] and lack of surveillance of enteric diseases [32]. The objective of this review was to gather information on the prevalence, risk factors, and antimicrobial resistance profiles of thermophilic *Campylobacter* species in humans and animals in SSA. The findings of this review are expected to provide evidence for policy formulation, prevention, and control of *Campylobacter* infections and increase awareness of the AMR issue.

2. Methods

The data were collected by searching articles published in English from electronic databases, namely, PubMed, Google Scholar, Research4life-HINARI Health, and Researchgate.net. The search terms were “thermophilic *Campylobacter*,” “*Campylobacter jejuni*,” “*Campylobacter coli*,” “diarrheal diarrhoea,” “antimicrobial resistance,” “antibiotic resistance,” “humans,” “animals,” “Sub-Saharan Africa,” and “a specific country name.” Initially, a total of 614 articles were identified, and the lists of references were screened in which 22 more articles were identified. After screening, 33 articles on humans and 34 on animals and animal products were included in this review (Figure 1). The reviewed articles

were those published from 1997 to 2018. During the review process, the data extracted included title, country, sex and age distribution, sample size, isolation and identification methods, isolation rates, and antimicrobial resistance profiles. Articles for which the sample size was not shown or which used archived *Campylobacter* cultures were excluded from this review.

3. Results

3.1. *Campylobacter* Infections in Humans. Of the 47 SSA countries [33], data on human campylobacteriosis were available from 15 (31.9%) countries. The prevalence of thermophilic *Campylobacter* in humans was reported in 33 articles (Table 1). Nigeria reported the highest overall prevalence of thermophilic *Campylobacter* (62.7%); followed by Malawi (21%) and South Africa (20.3%). Kenya reported the highest prevalence (16.4%) of *Campylobacter* infections in under-five children; followed by Rwanda (15.5%) and Ethiopia (14.5%). The mean prevalence in all ages and under-five children was 18.6% and 9.4%, respectively. Burkina Faso and Mozambique had the lowest prevalence of campylobacteriosis for all ages (2.3%) and under-five (1.7%), respectively. Of the 33 articles reviewed, 16 (48.5%) presented data on distribution of *Campylobacter* infections by sex but the difference was not statistically significant. Of these 16 articles, campylobacteriosis was more prevalent among males (22.7%; $n = 3966$) than females (17.7%; $n = 3705$). Culture methods on selective media, biochemical tests, molecular, and biotyping techniques were used for identification of *Campylobacter* (Table 1). Of the 33 articles, 27 studies were carried out at clinical settings (hospitals and health centres) while 6 were community-based studies. Probability sampling methods were adopted in 5 articles while the remaining used convenience sampling. Although *C. jejuni* and *C. coli* were isolated in the mentioned articles, 15 articles reported other enteric pathogens as probable aetiologies of diarrhoea. Furthermore, more than 85% of the articles considered diarrhoeic cases while the remaining included even asymptomatic participants.

Of the 33 articles, only four reported on risk factors of campylobacteriosis in humans. In Tanzania, *Campylobacter* infections were associated with sex, young age, poultry meat consumption, and eating of salads [26, 38]. In Ethiopia, human campylobacteriosis was significantly associated with nonuse of latrines, water source, drinking unboiled water, bottle feeding, nutritional status, and exposure to domestic animals including cats, dogs, poultry, and pigeons [25]. In Burkina Faso, *Campylobacter* infections were most common among under-fives and those aged 21–40 years with more pet contacts [57].

3.2. *Campylobacter* spp. in Animals and Contamination of Animal Products. Of the 34 articles from which data on animals were extracted, 17 collected faeces from live animals, while 16 collected samples from meat or caeca at abattoirs. In 2 articles, samples were collected from both markets and




FIGURE 1: Flowchart showing article selection process.

TABLE 1: Prevalence of thermophilic *Campylobacter* spp. in humans in sub-Saharan Africa, 1997–2018.

Country	Age group (sample size)	Number of articles	Prevalence (%)	Detection method	References
Uganda	Children <5 (226)	1	9.3 (<i>C. jejuni</i> : 80.9%; <i>C. coli</i> : 4.8%)	Culture, biochemical	[34]
Tanzania	Children <5 (1,512)	5	8.8 (2.6–19) (<i>C. jejuni</i> : 89.2%; <i>C. coli</i> : 9.8%)	Culture, biochemical, Gram staining, molecular	[8, 35–38]
Kenya	Children <5 (2,550)	1	16.4	Culture, biochemical, serotyping	[39]
Rwanda	Children <5 (706)	1	15.5 (<i>C. jejuni</i> : 100%)	Molecular	[40]
Madagascar	Children <5 (5,620)	2	9.4 (9.3–9.5) (<i>C. jejuni</i> : 73.6; <i>C. coli</i> : 24.3%)	Culture, serotyping, molecular	[41, 42]
Burkina Faso	Children <5 (283)	1	2 (<i>C. jejuni</i> : 60%; <i>C. coli</i> : 40%)	Culture, molecular	[43]
Ethiopia	Children <5 (670)	2	14.5 (12.7–16.7) (<i>C. jejuni</i> : 71.1%; <i>C. coli</i> : 21.1%)	Culture, biochemical, Gram staining	[25, 44, 45]
Nigeria	Children <5 (1,311)	3	4.4 (0.5–8.2) (<i>C. jejuni</i> : 28%; <i>C. coli</i> : 72%)	Culture, biochemical, biotyping, Gram staining	[46–48]
Niger	Children <5 (260)	1	11.4 (<i>C. jejuni</i> : 100%)	Culture, biochemical, Gram staining	[49]
Mozambique	Children <5 (529)	1	1.7	Culture, biochemical, Gram staining	[50]
Cameroon	Children <5 (260)	1	9.6 (<i>C. jejuni</i> : 100%)	Culture, biochemical, Gram staining	[51]
Botswana	Under 15 years	1	14	Molecular	
Tanzania	All ages (2,487)	4	11.1 (1.9–21.6) (<i>C. jejuni</i> : 93.3%; <i>C. coli</i> : 6.1%)	Culture, biochemical, Gram staining, molecular	[26, 52–54]
Kenya	All ages (4,274)	2	9.2 (8.5–9.8) (<i>C. jejuni</i> : 76.2; <i>C. coli</i> : 12.7%)	Culture	[55, 56]
Burkina Faso	All ages (1,246)	1	2.3 (<i>C. jejuni</i> : 51.8%; <i>C. coli</i> : 13.8%)	Culture, biochemical, Gram staining	[57]
Ethiopia	All ages (640)	2	9.8 (8–11.6) (<i>C. jejuni</i> : 94.1%; <i>C. coli</i> : 5.9%)	Culture, biochemical, Gram staining	[58, 59]
Nigeria	All ages (150)	1	62.7 (<i>C. jejuni</i> : 24.5%; <i>C. coli</i> : 62.3%)	Culture, biochemical, Gram staining	[60]
Ghana	All ages (202)	1	17.3 (<i>C. jejuni</i> : 42.8%; <i>C. coli</i> : 37%)	Culture, biochemical, Gram staining	[61]
Malawi	All ages (1,941)	1	21 (<i>C. jejuni</i> : 85%; <i>C. coli</i> : 14%)	Molecular	[62]
South Africa	All ages (565)	1	20.3 (<i>C. jejuni</i> : 85%; <i>C. coli</i> : 15%)	Culture, biochemical, molecular	[63]

abattoirs. Probability sampling methods were used in 6 articles while the remaining used convenience sampling.

Data on *Campylobacter* in cattle were obtained from ten articles published from studies conducted in six countries. The overall mean prevalence was 17.6% and *C. jejuni* had higher prevalence (70%) than *C. coli* (23.5%). The highest [64] and the lowest overall prevalence [52] were reported from Tanzania. Furthermore, Tanzania and Ghana showed higher prevalence for *C. jejuni* and *C. coli*, respectively (Table 2).

Data on *Campylobacter* in goats were reported in three articles from three different countries. The overall mean prevalence was 31.2%, and *C. jejuni* presented with a higher prevalence (56.2%) than *C. coli* (38.5%). The highest and lowest prevalence were reported from the Democratic Republic of Congo (DRC) [83] and Ghana [83], respectively. Ethiopia [70] and DRC [83] had the highest frequencies for *C. jejuni* and *C. coli*, respectively (Table 2). For sheep, data were reported in four articles from three countries. The overall mean prevalence was 31.8%, with *C. jejuni* being reported at a higher frequency (56.7%) than *C. coli* (35.4%). The highest and lowest prevalence were reported from Ethiopia [70] and Ghana [70], respectively. Ethiopia [70] and Tanzania [75] had the highest prevalence for *C. jejuni* and *C. coli*, respectively (Table 2).

Data on presence of thermophilic *Campylobacter* in pigs were available from six articles from five countries. The overall mean prevalence was 45.5% and contrary to other animals, *C. coli* occurred at a higher prevalence (70.1%) than *C. jejuni* (27.2%). The highest and lowest prevalence were reported from Nigeria [60] and South Africa [78], respectively. Ethiopia [69] had both higher and lower values for *C. jejuni* and *C. coli* (Table 2).

Data on thermophilic *Campylobacter* in chickens were obtained from 11 articles from five different countries. In this review, the number of articles on chickens was the highest compared to other reservoirs. The overall mean prevalence was 62.6% which was the highest in all animal reservoirs documented in this review. *Campylobacter jejuni* was reported in higher prevalence (81.0%) than *C. coli* (18.1%). The highest and lowest prevalence rates were reported in Ethiopia [70] and South Africa [18], respectively (Table 2).

As regards to animal products, data on cattle meat were reported in three articles from three countries. The overall mean prevalence was 5.5%, and *C. jejuni* had higher prevalence (95.2%) than *C. coli* (4.8%). The highest and lowest prevalence rates were reported in Ethiopia [72] and Kenya [73], respectively. For cattle carcasses, data were reported by two articles from two countries with a mean prevalence of 15.9%. Ghana [68] reported a higher prevalence of *C. jejuni* while Tanzania [74] observed a higher prevalence of *C. coli* (Table 2).

Data on sheep meat were reported by a single article from Ethiopia [72] with the prevalence of 10.5%. In sheep carcasses, the mean prevalence was 23.3% computed using two articles from two countries. Ghana [68] showed a higher prevalence of *C. jejuni* while Ethiopia [76] reported a higher prevalence of *C. coli*. In pork, the prevalence was 8.5% in one

article from Ethiopia [72] with *C. coli* being more prevalent than *C. jejuni*. In pig carcasses, the prevalence was 36.3% from one article reporting a study carried out in Ghana [68]. In chicken meat, the mean prevalence was 49.4% reported by two articles from two countries. Dadi and Asrat in a study conducted in Ethiopia [72] indicated a higher prevalence for *C. jejuni* while a study in Kenya [73] found a higher prevalence for *C. coli*. For chicken carcasses, the prevalence was 50% from one article in Burkina Faso [79] and all isolates were *C. jejuni*. In goat meat, the mean prevalence was 22.5% reported by only one article from Ethiopia [72]. In goat carcasses, the mean prevalence was 16.7% reported by two articles from two countries. A study conducted in Ghana [76] reported a higher prevalence for *C. jejuni* while that in Ethiopia [76] found a higher prevalence for *C. coli* (Table 2).

The overall prevalence of thermophilic *Campylobacter* in cats [84] and dogs [84, 85] were 18.3% and 20%, respectively. Of the reviewed articles, some presented data on companion, wild, and other animals (Table 3).

3.3. Antimicrobial Resistance Profiles of *C. jejuni* and *C. coli* in Humans and Animals. In humans, the AMR profiles, determined using disk diffusion, were available in 4 articles from four different countries (Figure 2), while the remaining did not specify the species. The antimicrobials considered in this review for the ease of comparison were ampicillin (AMP), erythromycin (ERY), tetracycline (TET), cefalotin (CF), nalidixic acid (NAL), azithromycin (AZM), gentamicin (GEN), ciprofloxacin (CIP), chloramphenicol (CHL), and trimethoprim-sulfamethoxazole (TM-SFX).

The percentage of antimicrobial resistant isolates ranged from 2–100% for *C. jejuni* and 0–100% for *C. coli*. The AMR data for CIP and ERY, which are drugs of choice for treating *Campylobacter* infections, showed that Ghana [61] and Tanzania [26] reported higher values for both *C. jejuni* and *C. coli*. Resistance of *Campylobacter jejuni* to GEN was similar for both Tanzania and Ghana while for *C. coli*, it was higher in Tanzania compared with that of Ghana [26, 61]. Higher frequencies of resistance were also reported for TET and AMP which have been in use for many years. In general, higher levels of AMR were reported in *C. jejuni* than *C. coli*.

In animal and animal products, the following antimicrobials were used in the reviewed articles: chloramphenicol (CHL), ampicillin (AMP), erythromycin (ERY), ciprofloxacin (CIP), nalidixic acid (NAL), streptomycin (STR), tetracycline (TET), gentamicin (GEN), and trimethoprim-sulfamethoxazole (TM-SFX) (Figure 3).

In animals, the percentage of resistant isolates varied from 0–100%. Resistance to CIP was in the range of 0–80.5% and 0–68.8% for *C. jejuni* and *C. coli*, respectively. Resistance to ERY varied from 0–99.5% and 0–100% for *C. jejuni* and *C. coli*, respectively. Resistance to GEN was <55.6% for both *C. jejuni* and *C. coli*. The highest resistance to most of the drugs was seen in Ghana [68] while the lowest resistance was observed in Tanzania [74, 89]. Resistance to nalidixic acid was high for both *C. jejuni* and *C. coli* in a study conducted in Tanzania [75]. Data on multidrug resistance were available from three studies in which values ranged from 23.3% to

TABLE 2: Prevalence of *Campylobacter* spp. in domestic animals and animal products.

Animal type	Sample type	Country	Overall prevalence	<i>C. jejuni</i> (%)	<i>C. coli</i> (%)	References
Cattle	Faeces	South Africa	19.3	72.4	27.6	[18]
		Nigeria	18.5	80	20	[65]
			12.9	65.1	23	[66]
			2.3	100	0	[52]
		Tanzania	5.6	83.3	16.7	[67]
			32.5	65.5	27.3	[64]
		Ghana	13.2	25	43.8	[68]
		Ethiopia	12.7	53.8	38.5	[69]
		Ethiopia	48	75.3	17.6	[70]
		Mozambique	11	80	20	[71]
<i>Average</i>			17.6	70	23.5	
Cattle	Meat	Tanzania	2.8	100	0	[67]
		Ethiopia	6.2	85.7	14.3	[72]
		Kenya	2	100	0	[73]
		<i>Average</i>	5.5	95.2	4.8	
Cattle	Carcasses	Tanzania	3.7	75	25	[67]
		Ethiopia	9.5	62.5	29.2	[74]
		Ghana	34.5	84.2	13.1	[68]
		<i>Average</i>	15.9	73.9	22.4	
Sheep	Faeces	Tanzania	31.6	55.6	44.4	[75]
		Ethiopia	38	59.3	40.7	[69]
		Ethiopia	39	84.6	15.4	[70]
		Ghana	18.6	27.2	40.9	[68]
<i>Average</i>			31.8	56.7	35.4	
Sheep	Carcasses	Ethiopia	10.6	73.9	26.1	[76]
		Ghana	35.9	92.8	0	[68]
<i>Average</i>			23.3	83.4	13.1	
Sheep	Meat	Ethiopia	10.5	83.3	0	[72]
		Nigeria	92.7	14	78.7	[60]
		Ethiopia	50	0	100	[69]
Pig	Faeces	Ethiopia	50	0	100	[69]
		Tanzania	66.7	81.8	18.2	[77]
		Tanzania	32.5	2.7	91.9	[64]
		Ghana	28.7	48.2	48.2	[68]
		South Africa	2.3	16.7	83.3	[78]
		<i>Average</i>			45.5	27.2
Pig	Carcasses	Ghana	36.3	28.4	10.8	[68]
Pig	Pork	Ethiopia	8.5	25	50	[72]
		Burkina Faso	68	70	30	[79]
Chicken	Faeces	Tanzania	69.8	91.2	8.8	[53]
			42.5	87.1	12.9	[38]
			77.8	91.1	7.3	[54]
			35.3	84.9	15.1	[18]
			49.7	100	0	[80]
		South Africa	54.8	54.8	40.2	[81]
			72.7	92.5	7.5	[59]
			Ethiopia	68.1	80.8	16.2
		Ethiopia	86.6	86.9	11.9	[70]
			Ivory Coast	63.8	51.3	48.7
<i>Average</i>			62.6	81	18.1	
Chicken	Colon	South Africa	14.2	68.8	31.2	[78]
Chicken	Carcasses	Burkina Faso	50	100	0	[79]
Chicken	Meat	Ethiopia	21.7	84	8	[72]
		Kenya	77	59	39	[73]
<i>Average</i>			49.4	71.5	23.5	
Goat	Faeces	DRC	41.7	32.7	59.4	[83]
		Ghana	18.5	36	56	[68]
		Ethiopia	33.3	100	0	[70]
<i>Average</i>			31.2	56.2	38.5	

TABLE 2: Continued.

Animal type	Sample type	Country	Overall prevalence	<i>C. jejuni</i> (%)	<i>C. coli</i> (%)	References
Goat	Carcasses	Ethiopia	9.4	70.6	29.4	[76]
		Ghana	23.9	81.3	0	[68]
<i>Average</i>			16.7	76	14.7	
Goat	Meat	DRC	37.3	21.3	74.7	[83]
		Ethiopia	7.6	71.4	28.6	[72]
<i>Average</i>			22.5	46.4	51.7	
Cattle	Milk	Tanzania	13.4	55.3	31.6	[74]

TABLE 3: Prevalence of *Campylobacter* spp. in companion, wild, and other animals.

Animal type	Specimen	Country	Overall prevalence	<i>C. jejuni</i> (%)	<i>C. coli</i> (%)	References
<i>Companion animals</i>						
Cat	Faeces	Nigeria	18.3	21.1		[66]
Dog	Faeces	Nigeria	27.7	23.1	0	[66]
		<i>Average</i>	12.3	53.8	30.8	[85]
<i>Other animals</i>						
Crow	Faeces	Tanzania	72.8	93.8	6.2	[53]
Duck	Faeces	Tanzania	80	81.5		[86]
Greater crested tern	Faeces	South Africa	16	15	1	[87]
Kelp gull	Faeces	South Africa	12.4	11.6	0.8	[87]
Quail	Caeca	Nigeria	31.1	81	19	[88]
Horse	Faeces	Tanzania	60	66.7	33.3	[75]
Guinea pig	Faeces	Tanzania	26.7	50	50	[75]
Rat	Faeces	Tanzania	1.2	66.7	33.3	[75]




FIGURE 2: Antimicrobial resistance data in humans by the disk diffusion method.

63.3% for *C. jejuni* [18, 59, 74] and from 0–25% for *C. coli* [59, 70]. There were variations in resistance levels to commonly used antimicrobials in animal species depending on the species tested.

4. Discussion

The overall mean prevalence of thermophilic *Campylobacter* in humans ranged from 9.6–18.5% and is within the ranges reported elsewhere in LMICs [31] and in Poland [90]. However, the prevalence was higher than that reported from

Korea [91], and was lower than that reported from the USA [92]. This variation may be attributed to the fact that campylobacteriosis is hyperendemic in LMICs probably due to poor sanitation and close proximity of humans and domestic animals [31]. The risk factors for human infections highlighted in this review partly explain this. They include consumption of poultry meat, drinking surface water, and animal contact, which is in agreement with other studies with consumption of poultry being the major risk factor [24, 93].

The prevalence of thermophilic *Campylobacter* in animals varied between 1.2% and 80%. The mean prevalence




FIGURE 3: Antimicrobial resistance data in animals by the disk diffusion method. 1: duck, 2: sea birds, 3: chicken, 4: raw meat, 5: laboratory and farm animals, 6: pig, 7: food animals, 8: cattle; TZ: Tanzania, SA: South Africa, ET: Ethiopia, IC: Ivory Coast, GH: Ghana.

recorded in chickens (60.3%) concurs with findings from other LMICs such as Thailand [94], Sri Lanka [95], and Vietnam [96]. The mean prevalence of thermophilic *Campylobacter* in pigs was comparable to what was reported in Spain and Vietnam [30, 96] but lower than those reported in Norway and the Netherlands [97, 98]. The prevalence of *Campylobacter* in goats and sheep was slightly higher than the prevalence reported in Germany and Trinidad [99, 100] but lower than the prevalence reported in Spain [30]. The prevalence in cattle (17.6%) was lower than those reported in the USA and Iran [101, 102] but higher than the prevalence reported in another paper in the USA [103].

Although thermophilic *Campylobacter* species are frequently isolated from animal faeces, this review showed that they are also present in considerable amounts in a number of animal products. The reported prevalence of *Campylobacter* in cattle and goat carcasses in sub-Saharan Africa was higher compared to the prevalence in Poland for cattle [104] and in Canada for goat [105]. The contamination of carcasses may result from contact with gut contents during manual skin removal, cleaning, and processing in the slaughter house [106]. The prevalence rates in beef, pork, and mutton were slightly higher compared to those observed in other countries [107–109]. The variation could be influenced by the differences in husbandry practices which determine exposure of the animals to the bacteria. Partly, this could also be attributable to slaughter and animal product handling practices which enhance the contamination of the products.

Campylobacter jejuni and *C. coli* were the most frequently encountered species from both human and animals. Similar observations have been reported by other authors [30, 110]. The predominance of *C. jejuni* in various animals, other than pigs, in sub-Saharan Africa has been previously reported [31, 111]. The possible explanation is that most of the studies rely on culture and biochemical tests which may

not correctly identify some species. Another reason is the use of selective media containing antibiotics to which some other *Campylobacter* species are sensitive to. Furthermore, higher incubation temperatures may limit the growth of some thermophilic *Campylobacter* species like *C. lari* and *C. upsaliensis* [112, 113].

In pigs, *C. coli* showed higher prevalence (67.4%) than *C. jejuni* (27.2%) which is in agreement with reports in Canada and the USA that *C. coli* is a normal flora of pigs' intestines [114, 115]. Furthermore, some studies show that *C. jejuni* and *C. coli* may cohabit in pigs but usually *C. jejuni* is always present in lower frequencies than *C. coli* [116, 117].

The results on AMR in both humans and animals highlight that resistance to mostly used antimicrobials is frequent. The resistance ranged from 0 to 100%, and higher resistance rates were reported in *C. jejuni* than in *C. coli*. The antimicrobials to which resistance was high included AMP, TET, ERY, and TET. The findings concur with the reports from other studies in both LMICs and high-income countries showing an increment in the number of *Campylobacter* strains resistant to most of the antimicrobials used in treating human campylobacteriosis [118–120]. The increase in resistance to most antimicrobial agents and emergence of MDR isolates could be associated with extensive use of antimicrobials not only as therapeutic agents for human infections [20] but also for prophylaxis and growth promotion in animal husbandry [68]. However, there are challenges in surveillance, differences in design and predominance of the disk diffusion method and not using globally accepted methods. These may cause differences within and between countries and certainly limit comparability with data reported in other parts of the world. The resistance to TET was comparable with the findings reported from Poland [121] and the USA [122] and the pooled estimate prevalence worldwide (94.3%) [120]. This resistance

may be due to wide use of tetracycline in both human and veterinary medicine [20]. The proportion of isolates resistant to macrolides (ERY) ranged from 0 to 100% in both humans and animals for *C. jejuni* while the range was from 0 to 92.3% for *C. coli*. The frequency of isolates resistant to fluoroquinolone was relatively lower in humans which is comparable to rates described in Western Europe [118, 121]. The resistance to both erythromycin and ciprofloxacin is of public health concern as there are currently limited options in the choice of treatment of *Campylobacter* infections. The proportion of multidrug resistance (MDR) isolates varied between 23.3 and 63.3% (Figure 3) which falls within the range of 37–90% from studies in China, Korea, and France [123–125].

There are no internationally agreed criteria of susceptibility testing and breakpoint assessment for *Campylobacter* spp. [126]. Therefore, it is difficult to interpret the available data and draw conclusion. Several laboratory standards have been applied for the susceptibility testing of *Campylobacter* species. Although disk diffusion was used in some studies, it should be used only as a screening method for resistance to erythromycin and ciprofloxacin [127].

5. Conclusion

This review indicates that *C. jejuni* and *C. coli* are frequently isolated from humans, food animals, and animal products in sub-Saharan Africa. Isolates from the different sources display varying degrees of resistance to commonly used antimicrobial agents. The findings of this review suggest that the disease burden due to thermophilic *Campylobacter* species in SSA is of public and economic importance. Therefore, routine diagnosis of *C. jejuni* and *C. coli*, appropriate use of antimicrobials, educating communities on hygienic practices, establishment of both national and regional multisectoral antimicrobial resistance standard surveillance protocols are necessary to curb both the campylobacteriosis burden, and increase of antimicrobial resistance in the region.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References





- [1] C. Bern, J. Martines, I. de Zoysa, and R. I. Glass, "The magnitude of the global problem of diarrhoeal disease: a ten-year update," *Bulletin of the World Health Organization*, vol. 70, no. 6, pp. 705–714, 1992.
- [2] C. L. F. Walker, I. Rudan, L. Liu et al., "Global burden of childhood pneumonia and diarrhoea," *The Lancet*, vol. 381, no. 9875, pp. 1405–1416, 2013.
- [3] L. Liu, S. Oza, D. Hogan et al., "Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis," *The Lancet*, vol. 385, no. 9966, pp. 430–440, 2015.
- [4] J. W. Ahs, W. Tao, J. Löfgren, and B. C. Forsberg, "Diarrheal diseases in low- and middle-income countries: incidence, prevention and management," *The Open Infectious Diseases Journal*, vol. 4, no. 1, pp. 113–124, 2010.
- [5] L. Carvajal-Vélez, A. Amouzou, J. Perin et al., "Diarrhea management in children under five in sub-Saharan Africa: does the source of care matter? A countdown analysis," *BMC Public Health*, vol. 16, no. 1, p. 830, 2016.
- [6] C. A. Petti, C. R. Polage, T. C. Quinn, A. R. Ronald, and M. A. Sande, "Laboratory medicine in Africa: a barrier to effective health care," *Clinical Infectious Diseases*, vol. 42, no. 3, pp. 377–382, 2006.
- [7] T. V. Nguyen, P. V. Le, C. H. Le, and A. Weintraub, "Antibiotic resistance in diarrheagenic *Escherichia coli* and *Shigella* strains isolated from children in Hanoi, Vietnam," *Antimicrobial Agents and Chemotherapy*, vol. 49, no. 2, pp. 816–819, 2005.
- [8] M. Vargas, J. Vila, C. Casals et al., "Etiology of diarrhea in children less than five years of age in Ifakara, Tanzania," *The American Journal of Tropical Medicine and Hygiene*, vol. 70, no. 5, pp. 536–539, 2004.
- [9] O. G. Gómez-Duarte, J. Bai, and E. Newell, "Detection of *Escherichia coli*, *Salmonella* spp., *Shigella* spp., *Yersinia enterocolitica*, *Vibrio cholerae*, and *Campylobacter* spp. enteropathogens by 3-reaction multiplex polymerase chain reaction," *Diagnostic Microbiology and Infectious Disease*, vol. 63, no. 1, pp. 1–9, 2009.
- [10] G. M. Ruiz-Palacios, "The health burden of *Campylobacter* infection and the impact of antimicrobial resistance: playing chicken," *Clinical Infectious Diseases*, vol. 44, no. 5, pp. 701–703, 2007.
- [11] B. M. Allos and M. J. Blaser, "*Campylobacter jejuni* and the expanding spectrum of related infections," *Clinical Infectious Diseases*, vol. 20, no. 5, pp. 1092–1101, 1995.
- [12] D. Gilliss, A. B. Cronquist, M. Cartter et al., "Incidence and trends of infection with pathogens transmitted commonly through food—foodborne diseases active surveillance network, 10 U.S. Sites, 1996–2012," *Morbidity and Mortality Weekly Report*, vol. 62, no. 15, pp. 283–287, 2013.
- [13] E. Guévremont, É. Nadeau, M. Sirois, and S. Quessy, "Antimicrobial susceptibilities of thermophilic *Campylobacter* from humans, swine, and chicken broilers," *Canadian Journal of Veterinary Research*, vol. 70, no. 2, pp. 81–86, 2006.
- [14] S. Thakur, S. Zhao, P. F. McDermott et al., "Antimicrobial resistance, virulence, and genotypic profile comparison of *Campylobacter jejuni* and *Campylobacter coli* isolated from humans and retail meats," *Foodborne Pathogens and Disease*, vol. 7, no. 7, pp. 835–844, 2010.
- [15] R. L. Guerrant, T. Van Gilder, T. S. Steiner et al., "Practice guidelines for the management of infectious diarrhea," *Clinical Infectious Diseases*, vol. 32, no. 3, pp. 321–351, 2001.
- [16] F. M. Aarestrup, P. F. McDermott, and H. C. Wegener, *Transmission of Antibiotic Resistance from Food Animals to Humans*, American Society for Microbiology, Washington, DC, USA, 3rd edition, 2008.
- [17] F. M. Aarestrup and J. R. Engberg, "Antimicrobial resistance of thermophilic *Campylobacter*," *Veterinary Research*, vol. 32, no. 3/4, pp. 311–321, 2001.
- [18] Uaboi-Egbenni, "Potentially pathogenic *Campylobacter* species among farm animals in rural areas of Limpopo province, South Africa: a case study of chickens and cattle," *African Journal of Microbiology Research*, vol. 6, no. 12, 2012.
- [19] P. Padungton and J. B. Kaneene, "*Campylobacter* spp. in human, chickens, pigs and their antimicrobial resistance," *Journal of Veterinary Medical Science*, vol. 65, no. 2, pp. 161–170, 2003.

- [20] N. M. Iovine, "Resistance mechanisms in *Campylobacter jejuni*," *Virulence*, vol. 4, no. 3, pp. 230–240, 2013.
- [21] J. Neimann, J. Engberg, K. Mølbak, and H. C. Wegener, "A case-control study of risk factors for sporadic campylobacter infections in Denmark," *Epidemiology and Infection*, vol. 130, no. 3, pp. 353–366, 2003.
- [22] Y. Doorduyn, W. E. Van Den Brandhof, Y. T. H. P. Van Duynhoven, B. J. Breukink, J. A. Wagenaar, and W. Van Pelt, "Risk factors for indigenous *Campylobacter jejuni* and *Campylobacter coli* infections in the Netherlands: a case-control study," *Epidemiology and Infection*, vol. 138, no. 10, pp. 1391–1404, 2010.
- [23] J. Mossong, L. Mughini-Gras, C. Penny et al., "Human campylobacteriosis in Luxembourg, 2010–2013: a case-control study combined with multilocus sequence typing for source attribution and risk factor analysis," *Scientific Reports*, vol. 6, no. 1, 2016.
- [24] G. Kapperud, "Factors associated with increased and decreased risk of *Campylobacter* infection: a prospective case-control study in Norway," *American Journal of Epidemiology*, vol. 158, no. 3, pp. 234–242, 2003.
- [25] A. Lengerh, F. Moges, C. Unakal, and B. Anagaw, "Prevalence, associated risk factors and antimicrobial susceptibility pattern of *Campylobacter* species among under five diarrheic children at Gondar University Hospital, Northwest Ethiopia," *BMC Pediatrics*, vol. 13, no. 1, p. 82, 2013.
- [26] E. V. G. Komba, R. H. Mdegela, P. L. M. Msoffe, L. N. Nielsen, and H. Ingmer, "Prevalence, antimicrobial resistance and risk factors for thermophilic *Campylobacter* infections in symptomatic and asymptomatic humans in Tanzania," *Zoonoses and Public Health*, vol. 62, no. 7, pp. 557–568, 2015.
- [27] M. B. Skirrow, "Epidemiology of *Campylobacter enteritis*," *International Journal of Food Microbiology*, vol. 12, no. 1, pp. 9–16, 1991.
- [28] S. N. Workman, G. E. Mathison, and M. C. Lavoie, "Pet dogs and chicken meat as reservoirs of *Campylobacter* spp. in Barbados," *Journal of Clinical Microbiology*, vol. 43, no. 6, pp. 2642–2650, 2005.
- [29] N. W. Luechtefeld, L. B. Reller, M. J. Blaser, and W. L. Wang, "Comparison of atmospheres of incubation for primary isolation of *Campylobacter fetus* subsp. *jejuni* from animal specimens: 5% oxygen versus candle jar," *Journal of Clinical Microbiology*, vol. 15, no. 1, pp. 53–57, 1982.
- [30] B. Oporto, J. I. Esteban, G. Aduriz, R. A. Juste, and A. Hurtado, "Prevalence and strain diversity of thermophilic campylobacters in cattle, sheep and swine farms," *Journal of Applied Microbiology*, vol. 103, no. 4, pp. 977–984, 2007.
- [31] A. O. Coker, R. D. Isokpehi, B. N. Thomas, K. O. Amisu, and C. L. Obi, "Human campylobacteriosis in developing countries," *Emerging Infectious Diseases*, vol. 8, no. 3, pp. 237–243, 2002.
- [32] WHO/FAO/OIE, *The Global View of Campylobacteriosis: Report of an Expert Consultation*, World Health Organization, Geneva, Switzerland, 2013.
- [33] F. Mullan and S. Frehywot, "Non-physician clinicians in 47 sub-Saharan African countries," *The Lancet*, vol. 370, no. 9605, pp. 2158–2163, 2007.
- [34] S. Mshana, M. Joloba, A. Kakooza, and D. Kaddu-Mulindwa, "*Campylobacter* spp among children with acute diarrhea attending Mulago hospital in Kampala-Uganda," *African Health Sciences*, vol. 9, no. 3, pp. 201–205, 2009.
- [35] R. Kingamkono, E. Sjögren, and U. Svanberg, "Enteropathogenic bacteria in faecal swabs of young children fed on lactic acid-fermented cereal gruels," *Epidemiology and Infection*, vol. 122, no. 1, pp. 23–32, 1999.
- [36] R. Oketcho, C. Nyaruhucha, S. Taybali, and E. Karimuribo, "Influence of enteric bacteria and parasite infection and nutritional status on diarrhoea occurrence in six to 60 month old children admitted at Morogoro regional hospital of Tanzania," *Tanzania Journal of Health Research*, vol. 14, no. 2, 2012.
- [37] A. P. Deogratias et al., "Prevalence and determinants of *Campylobacter* infection among under five children with acute watery diarrhea in Mwanza, North Tanzania," *Archives of Public Health*, vol. 72, no. 1, 2014.
- [38] I. S. Chuma, H. E. Nonga, R. H. Mdegela, and R. R. Kazwala, "Epidemiology and RAPD-PCR typing of thermophilic campylobacters from children under five years and chickens in Morogoro Municipality, Tanzania," *BMC Infectious Diseases*, vol. 16, no. 1, pp. 1–11, 2016.
- [39] M. Beatty et al., "Sporadic paediatric diarrhoeal illness in urban and rural sites in Nyanza province, Kenya," *East African Medical Journal*, vol. 86, no. 8, 2010.
- [40] J.-C. Kabayiza, M. E. Andersson, S. Nilsson, T. Bergström, G. Muhirwa, and M. Lindh, "Real-time PCR identification of agents causing diarrhea in Rwandan children less than 5 years of age," *The Pediatric Infectious Disease Journal*, vol. 33, no. 10, pp. 1037–1042, 2014.
- [41] R. Randremanana, F. Randrianirina, M. Gousseff et al., "Case-control study of the etiology of infant diarrheal disease in 14 districts in Madagascar," *PLoS One*, vol. 7, no. 9, Article ID e44533, 2012.
- [42] R. V. Randremanana, F. Randrianirina, P. Sabatier et al., "*Campylobacter* infection in a cohort of rural children in Moramanga, Madagascar," *BMC Infectious Diseases*, vol. 14, no. 1, 2014.
- [43] I. J. O. Bonkougou, K. Haukka, M. Österblad et al., "Bacterial and viral etiology of childhood diarrhea in Ouagadougou, Burkina Faso," *BMC Pediatrics*, vol. 13, no. 1, p. 36, 2013.
- [44] B. Tafa, T. Sewunet, H. Tassew, and D. Asrat, "Isolation and antimicrobial susceptibility patterns of *Campylobacter* species among diarrheic children at Jimma, Ethiopia," *International Journal of Bacteriology*, vol. 2014, Article ID 560617, 7 pages, 2014.
- [45] M. Getamesay, B. Getenet, and Z. Ahmed, "Prevalence of *Shigella*, *Salmonella* and *Campylobacter* species and their susceptibility patterns among under five children with diarrhea in Hawassa town, South Ethiopia," *Ethiopian Journal of Health Sciences*, vol. 24, no. 2, p. 101, 2014.
- [46] O. Adekunle, A. Coker, and D. Kolawole, "Incidence, isolation and characterization of *Campylobacter* species in Osogbo," *Biology and Medicine*, vol. 1, no. 1, pp. 24–27, 2009.
- [47] S. O. Samuel, A. O. Aboderin, A. A. Akanbi, B. Adegboro, S. I. Smith, and A. O. Coker, "*Campylobacter enteritis* in Ilorin, Nigeria," *East African Medical Journal*, vol. 83, no. 9, pp. 478–484, 2006.
- [48] A. O. Aboderin, S. I. Smith, A. O. Oyelese, A. O. Onipede, S. B. Zailini, and A. O. Coker, "Role of *Campylobacter jejuni/coli* in Ile-Ife, Nigeria," *East African Medical Journal*, vol. 79, no. 8, pp. 423–426, 2002.
- [49] C. Langendorf, S. Le Hello, A. Moumouni et al., "Enteric bacterial pathogens in children with diarrhea in Niger: diversity and antimicrobial resistance," *PLoS One*, vol. 10, no. 3, Article ID e0120275, 2015.
- [50] I. M. Mandomando, P. L. Alonso, E. V. Macete et al., "Etiology of diarrhea in children younger than 5 years of age

- admitted in a rural hospital of southern Mozambique,” *The American Journal of Tropical Medicine and Hygiene*, vol. 76, no. 3, pp. 522–527, 2007.
- [51] H. B. N. Yongs, “Pathogenic microorganisms associated with childhood diarrhea in low-and-middle income countries: case study of Yaoundé-Cameroon,” *International Journal of Environmental Research and Public Health*, vol. 5, no. 4, pp. 213–229, 2008.
- [52] L. J. M. Kusiluka, E. D. Karimuribo, R. H. Mdegela et al., “Prevalence and impact of water-borne zoonotic pathogens in water, cattle and humans in selected villages in Dodoma rural and Bagamoyo districts, Tanzania,” *Physics and Chemistry of the Earth, Parts A/B/C*, vol. 30, no. 11–16, pp. 818–825, 2005.
- [53] R. H. Mdegela, H. E. Nonga, H. A. Ngowi, and R. R. Kazwala, “Prevalence of thermophilic *Campylobacter* infections in humans, chickens and crows in Morogoro, Tanzania,” *Journal of Veterinary Medicine Series B*, vol. 53, no. 3, pp. 116–121, 2006.
- [54] P. Jacob, R. H. Mdegela, and H. E. Nonga, “Comparison of Cape Town and Skirrow’s *Campylobacter* isolation protocols in humans and broilers in Morogoro, Tanzania,” *Tropical Animal Health and Production*, vol. 43, no. 5, pp. 1007–1013, 2011.
- [55] R. L. Shapiro, L. Kumar, P. Phillips-Howard et al., “Antimicrobial-resistant bacterial diarrhea in rural western Kenya,” *The Journal of Infectious Diseases*, vol. 183, no. 11, pp. 1701–1704, 2001.
- [56] J. T. Brooks, J. B. Ochieng, L. Kumar et al., “Surveillance for bacterial diarrhea and antimicrobial resistance in rural western Kenya, 1997–2003,” *Clinical Infectious Diseases*, vol. 43, no. 4, pp. 393–401, 2006.
- [57] L. Sangaré, A. K. Nikiéma, S. Zimmermann et al., “*Campylobacter* spp. epidemiology and antimicrobial susceptibility in a developing country, Burkina Faso (West Africa),” *African Journal of Clinical and Experimental Microbiology*, vol. 13, no. 2, pp. 110–117, 2012.
- [58] G. Beyene and A. Haile-Amlak, “Antimicrobial sensitivity pattern of *Campylobacter* species among children in Jimma university specialized hospital, Southwest Ethiopia,” *Ethiopian Journal of Health Development*, vol. 18, no. 3, pp. 185–189, 2004.
- [59] D. Ewnetu and A. Mihret, “Prevalence and antimicrobial resistance of *Campylobacter* isolates from humans and chickens in bahir dar, Ethiopia,” *Foodborne Pathogens and Disease*, vol. 7, no. 6, pp. 667–670, 2010.
- [60] P. B. Gwimi, O. O. Faleke, M. D. Salihu et al., “Prevalence of *Campylobacter* species in fecal samples of pigs and humans from Zuru Kebbi State, Nigeria,” *International Journal of One Health*, vol. 1, pp. 1–5, 2015.
- [61] A. B. Karikari, K. Obiri-Danso, E. H. Frimpong, and K. A. Krogfelt, “Antibiotic resistance in *Campylobacter* isolated from patients with gastroenteritis in a teaching hospital in Ghana,” *Open Journal of Medical Microbiology*, vol. 7, no. 1, pp. 1–11, 2017.
- [62] J. Mason, M. Iturriza-Gomara, S. J. O’Brien et al., “*Campylobacter* infection in children in Malawi is common and is frequently associated with enteric virus co-infections,” *PLoS One*, vol. 8, no. 3, Article ID e59663, 2013.
- [63] A. Samie, J. Ramalivhana, E. O. Igumbor, and C. L. Obi, “Prevalence, haemolytic and haemagglutination activities and antibiotic susceptibility profiles of *Campylobacter* spp. isolated from human diarrhoeal stools in Vhembe district, South Africa,” *Journal of Health, Population and Nutrition*, vol. 25, no. 4, pp. 406–413, 2007.
- [64] I. P. Kashoma, I. I. Kassem, A. Kumar et al., “Antimicrobial resistance and genotypic diversity of *Campylobacter* isolated from pigs, dairy, and beef cattle in Tanzania,” *Frontiers in Microbiology*, vol. 6, pp. 1–11, 2015.
- [65] S. S. Ngulukun, S. I. Oboegbulem, I. O. Fagbamila, W. Bertu, and M. O. Odugbo, “Prevalence and molecular characterization of thermophilic *Campylobacter* species isolated from cattle in Plateau State, Nigeria,” *Nigerian Veterinary Journal*, vol. 32, no. 4, pp. 349–356, 2011.
- [66] M. D. Salihu, J. U. Abdulkadir, S. I. Oboegbulem et al., “Isolation and prevalence of *Campylobacter* species in cattle from Sokoto State, Nigeria,” *Veterinaria Italiana*, vol. 45, no. 4, pp. 501–505, 2009.
- [67] H. E. Nonga, P. Sells, and E. D. Karimuribo, “Occurrences of thermophilic *Campylobacter* in cattle slaughtered at Morogoro municipal abattoir, Tanzania,” *Tropical Animal Health and Production*, vol. 42, no. 1, pp. 73–78, 2010.
- [68] A. B. Karikari, K. Obiri-Danso, E. H. Frimpong, and K. A. Krogfelt, “Antibiotic resistance of *Campylobacter* recovered from faeces and carcasses of healthy livestock,” *BioMed Research International*, vol. 2017, Article ID 4091856, 9 pages, 2017.
- [69] T. Kassa, S. Gebre-selassie, and D. Asrat, “The prevalence of thermotolerant *Campylobacter* species in food animals in Jimma Zone, Southwest Ethiopia,” *Ethiopian Journal of Health Development*, vol. 19, no. 3, pp. 225–229, 2005.
- [70] A. Abamecha, G. Assebe, B. Tafa, and B. Wondafrash, “Prevalence of thermophilic *Campylobacter* and their antimicrobial resistance profile in food animals in Lare District, Nuer Zone, Gambella, Ethiopia,” *Journal of Drug Research and Development*, vol. 1, no. 2, 2016.
- [71] S. Achá, I. Kühn, P. Jonsson, G. Mbazima, M. Katouli, and R. Möllby, “Studies on calf diarrhoea in Mozambique: prevalence of bacterial pathogens,” *Acta Veterinaria Scandinavica*, vol. 45, no. 1, pp. 27–36, 2004.
- [72] L. Dadi and D. Asrat, “Prevalence and antimicrobial susceptibility profiles of thermotolerant *Campylobacter* strains in retail raw meat products in Ethiopia,” *Ethiopian Journal of Health Development*, vol. 22, no. 2, pp. 195–200, 2008.
- [73] O. Osano and S. M. Arimi, “Retail poultry and beef as sources of *Campylobacter jejuni*,” *East African Medical Journal*, vol. 76, no. 76, pp. 141–143, 1999.
- [74] I. P. Kashoma, I. I. Kassem, J. John et al., “Prevalence and antimicrobial resistance of *Campylobacter* isolated from dressed beef carcasses and raw milk in Tanzania,” *Microbial Drug Resistance*, vol. 22, no. 1, pp. 40–52, 2016.
- [75] E. V. G. Komba, R. H. Mdegela, P. L. M. Msoffe, D. E. Matowo, and M. J. Maro, “Occurrence, species distribution and antimicrobial resistance of thermophilic *Campylobacter* isolates from farm and laboratory animals in Morogoro, Tanzania,” *Veterinary World*, vol. 7, no. 8, pp. 559–565, 2014.
- [76] T. Woldemariam, D. Asrat, and G. Zewde, “Prevalence of thermophilic *Campylobacter* species in carcasses from sheep and goats in an abattoir in Debre Zeit area, Ethiopia,” *Ethiopian Journal of Health Development*, vol. 23, no. 3, pp. 229–233, 2009.
- [77] R. H. Mdegela, K. Laurence, P. Jacob, and H. E. Nonga, “Occurrences of thermophilic *Campylobacter* in pigs slaughtered at Morogoro slaughter slabs, Tanzania,” *Tropical Animal Health and Production*, vol. 43, no. 1, pp. 83–87, 2011.

- [78] A. Jonker and J. A. Picard, "Antimicrobial susceptibility in thermophilic *Campylobacter* species isolated from pigs and chickens in South Africa," *Journal of the South African Veterinary Association*, vol. 81, no. 4, pp. 228–236, 2010.
- [79] A. Kagambèga, A. Thibodeau, V. Trinetta et al., "Salmonella spp. and *Campylobacter* spp. in poultry feces and carcasses in Ouagadougou, Burkina Faso," *Food Science & Nutrition*, vol. 6, no. 6, pp. 1601–1606, Jul. 2018.
- [80] L. A. Bester and S. Y. Essack, "Prevalence of antibiotic resistance in *Campylobacter* isolates from commercial poultry suppliers in KwaZulu-Natal, South Africa," *Journal of Antimicrobial Chemotherapy*, vol. 62, no. 6, pp. 1298–1300, 2008.
- [81] L. A. Bester and S. Y. Essack, "Observational study of the prevalence and antibiotic resistance of *Campylobacter* spp. from different poultry production systems in KwaZulu-Natal, South Africa," *Journal of Food Protection*, vol. 75, no. 1, pp. 154–159, 2012.
- [82] G. Gblossi Bernadette, A. Eric Essoh, K.-N. G. Elise Solange et al., "Prevalence and antimicrobial resistance of thermophilic *Campylobacter* isolated from chicken in côte d'Ivoire," *International Journal of Microbiology*, vol. 2012, Article ID 150612, 5 pages, 2012.
- [83] R. K. A Mpalang, R. Boreux, P. Melin, K. M. Akir Ni Bitiang, G. Daube, and P. De Mol, "Prevalence of *Campylobacter* among goats and retail goat meat in Congo," *The Journal of Infection in Developing Countries*, vol. 8, no. 2, pp. 168–175, 2014.
- [84] M. D. Salihu, A. A. Magaji, J. U. Abdulkadir, and A. Kolawale, "Survey of thermophilic *Campylobacter* species in cats and dogs in North-Western Nigeria," *Veterinaria Italiana*, vol. 46, p. 6, 2010.
- [85] N. S. Karshima, J. A. Benschak, S. I. Bata, I. J. Barde, and S. Aaron, "Molecular characterization of thermophilic *Campylobacter* species isolated from companion dogs presented to a veterinary facility in Jos, Nigeria," *Tropical Veterinarian*, vol. 32, no. 1-2, pp. 17–27, 2014.
- [86] H. E. Nonga and A. P. Muhairwa, "Prevalence and antibiotic susceptibility of thermophilic *Campylobacter* isolates from free range domestic duck (*Cairina moschata*) in Morogoro municipality, Tanzania," *Tropical Animal Health and Production*, vol. 42, no. 2, pp. 165–172, 2010.
- [87] E. Moré, T. Ayats, P. G. Ryan et al., "Seabirds (Laridae) as a source of *Campylobacter* spp., *Salmonella* spp. and antimicrobial resistance in South Africa," *Environmental Microbiology*, vol. 19, no. 10, pp. 4164–4176, 2017.
- [88] S. S. Ngulukun, S. I. Oboegbulem, I. O. Fagbamila et al., "Isolation of thermophilic *Campylobacter* species from Japanese quails (*Coturnix coturnix*) in Vom, Nigeria," *Veterinary Record*, vol. 166, no. 5, pp. 147–148, 2010.
- [89] I. P. Kashoma, A. Kumar, Y. M. Sanad et al., "Phenotypic and genotypic diversity of thermophilic *Campylobacter* spp. in commercial Turkey flocks: a longitudinal study," *Foodborne Pathogens and Disease*, vol. 11, no. 11, pp. 850–860, 2014.
- [90] B. Szczepanska, M. Andrzejewska, D. Spica, and J. J. Klawe, "Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from children and environmental sources in urban and suburban areas," *BMC Microbiology*, vol. 17, no. 1, 2017.
- [91] Y.-S. Kang, Y.-S. Cho, S.-K. Yoon et al., "Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from raw chicken meat and human stools in Korea," *Journal of Food Protection*, vol. 69, no. 12, pp. 2915–2923, 2006.
- [92] M. E. Kendall, S. Crim, K. Fullerton et al., "Travel-associated enteric infections diagnosed after return to the United States, foodborne diseases active surveillance network (FoodNet), 2004–2009," *Clinical Infectious Diseases*, vol. 54, no. 5, pp. S480–S487, 2012.
- [93] M. R. Evans, C. D. Ribeiro, and R. L. Salmon, "Hazards of healthy living: bottled water and salad vegetables as risk factors for *Campylobacter* infection," *Emerging Infectious Diseases*, vol. 9, no. 10, pp. 1219–1225, 2003.
- [94] P. Padungtod and J. B. Kaneene, "Campylobacter in food animals and humans in Northern Thailand," *Journal of Food Protection*, vol. 68, no. 12, pp. 2519–2526, 2005.
- [95] K. Kottawatta, M. Van Bergen, P. Abeynayake, J. Wagenaar, K. Veldman, and R. Kalupahana, "Campylobacter in broiler chicken and broiler meat in Sri Lanka: influence of semi-automated vs. wet market processing on *Campylobacter* contamination of broiler neck skin samples," *Foods*, vol. 6, no. 12, p. 105, 2017.
- [96] J. J. Carrique-Mas, J. E. Bryant, N. V. Cuong et al., "An epidemiological investigation of *Campylobacter* in pig and poultry farms in the Mekong Delta of Vietnam," *Epidemiology and Infection*, vol. 142, no. 7, pp. 1425–1436, 2014.
- [97] O. Rosef, B. Gondrosen, G. Kapperud, and B. Underdal, "Isolation and characterization of *Campylobacter jejuni* and *Campylobacter coli* from domestic and wild mammals in Norway," *Applied and Environmental Microbiology*, vol. 46, no. 4, pp. 855–859, 1983.
- [98] J. Oosterom, R. Dekker, G. J. A. de Wilde, F. van Kempde Troye, and G. B. Engels, "Prevalence of *Campylobacter jejuni* and *Salmonella* during pig slaughtering," *Veterinary Quarterly*, vol. 7, no. 1, pp. 31–34, 1985.
- [99] A. A. Adesiyun, J. S. Kaminjolo, R. Loregnard, and W. Kitson-Piggott, "Campylobacter infections in calves, piglets, lambs and kids in Trinidad," *British Veterinary Journal*, vol. 148, no. 6, pp. 547–556, 1992.
- [100] A.-K. Schilling, H. Hotzel, U. Methner et al., "Zoonotic agents in small ruminants kept on city farms in Southern Germany," *Applied and Environmental Microbiology*, vol. 78, no. 11, pp. 3785–3793, 2012.
- [101] Y. M. Sanad, I. I. Kassem, M. Abley, W. Gebreyes, J. T. LeJeune, and G. Rajashekhara, "Genotypic and phenotypic properties of cattle-associated *Campylobacter* and their implications to public health in the USA," *PLoS One*, vol. 6, no. 10, Article ID e25778, 2011.
- [102] R. Khoshbakht, M. Tabatabaei, S. Hoseinzadeh, M. Raeisi, H. S. Aski, and E. Berizi, "Prevalence and antibiotic resistance profile of thermophilic *Campylobacter* spp. of slaughtered cattle and sheep in Shiraz, Iran," *Veterinary Research Forum*, vol. 7, no. 3, pp. 241–246, 2016.
- [103] B. R. Hoar, E. R. Atwill, C. Elmi, and T. B. Farver, "An examination of risk factors associated with beef cattle shedding pathogens of potential zoonotic concern," *Epidemiology and Infection*, vol. 127, no. 1, pp. 147–155, 2001.
- [104] K. Wiczorek, E. Denis, O. Lynch, and J. Osek, "Molecular characterization and antibiotic resistance profiling of *Campylobacter* isolated from cattle in polish slaughterhouses," *Food Microbiology*, vol. 34, no. 1, pp. 130–136, 2013.
- [105] J. F. Prescott and C. W. Bruin-Mosch, "Carriage of *Campylobacter jejuni* in healthy and diarrheic animals," *American Journal of Veterinary Research*, vol. 42, no. 1, pp. 164–165, 1981.
- [106] P. Whyte, K. McGill, and J. D. Collins, "An assessment of steam pasteurization and hot water immersion treatments

- for the microbiological decontamination of broiler carcasses," *Food Microbiology*, vol. 20, no. 1, pp. 111–117, 2003.
- [107] G. Pezzotti, A. Serafin, I. Luzzi, R. Mioni, M. Milan, and R. Perin, "Occurrence and resistance to antibiotics of *Campylobacter jejuni* and *Campylobacter coli* in animals and meat in northeastern Italy," *International Journal of Food Microbiology*, vol. 82, no. 3, pp. 281–287, 2003.
- [108] S. Mayrhofer, P. Paulsen, F. J. M. Smulders, and F. Hilbert, "Antimicrobial resistance profile of five major food-borne pathogens isolated from beef, pork and poultry," *International Journal of Food Microbiology*, vol. 97, no. 1, pp. 23–29, 2004.
- [109] I. Hussain, M. Shahid Mahmood, M. Akhtar, and A. Khan, "Prevalence of *Campylobacter* species in meat, milk and other food commodities in Pakistan," *Food Microbiology*, vol. 24, no. 3, pp. 219–222, 2007.
- [110] J.-R. Yang, H.-S. Wu, C.-S. Chiang, and J.-J. Mu, "Pediatric campylobacteriosis in Northern Taiwan from 2003 to 2005," *BMC Infectious Diseases*, vol. 8, p. 151, 2008.
- [111] S. F. Altekruse, N. J. Stern, P. I. Fields, and D. L. Swerdlow, "*Campylobacter jejuni*-an emerging foodborne pathogen," *Emerging Infectious Diseases*, vol. 5, no. 1, pp. 28–35, 1999.
- [112] A. J. Lastovica, "Emerging *Campylobacter* spp.: the tip of the iceberg," *Clinical Microbiology Newsletter*, vol. 28, no. 7, pp. 49–56, 2006.
- [113] E. Bessede, A. Delcamp, E. Sifre, A. Buissonniere, and F. Megraud, "New methods for detection of *Campylobacter* in stool samples in comparison to culture," *Journal of Clinical Microbiology*, vol. 49, no. 3, pp. 941–944, 2011.
- [114] N. P. Varela, R. M. Friendship, and C. E. Dewey, "Prevalence of *Campylobacter* spp isolated from grower-finisher pigs in Ontario," *The Canadian Veterinary Journal=La Revue Veterinaire Canadienne*, vol. 48, no. 48, pp. 515–517, 2007.
- [115] D. A. Tadesse, P. B. Bahnson, J. A. Funk et al., "Prevalence and antimicrobial resistance profile of *Campylobacter* spp. isolated from conventional and antimicrobial-free swine production systems from different U.S. regions," *Foodborne Pathogens and Disease*, vol. 8, no. 3, pp. 367–374, 2011.
- [116] A. N. Jensen, M. T. Andersen, A. Dalsgaard, D. L. Baggesen, and E. M. Nielsen, "Development of real-time PCR and hybridization methods for detection and identification of thermophilic *Campylobacter* spp. in pig faecal samples," *Journal of Applied Microbiology*, vol. 99, no. 2, pp. 292–300, 2005.
- [117] R. H. Madden, L. Moran, and P. Scates, "Optimising recovery of *Campylobacter* spp. from the lower porcine gastrointestinal tract," *Journal of Microbiological Methods*, vol. 42, pp. 115–119, 2000.
- [118] P. Lubber, J. Wagner, H. Hahn, and E. Bartelt, "Antimicrobial resistance in *Campylobacter jejuni* and *Campylobacter coli* strains isolated in 1991 and 2001-2002 from poultry and humans in Berlin, Germany," *Antimicrobial Agents and Chemotherapy*, vol. 47, no. 12, pp. 3825–3830, 2003.
- [119] S. Lévesque, E. Frost, and S. Michaud, "Comparison of antimicrobial resistance of *Campylobacter jejuni* isolated from humans, chickens, raw milk, and environmental water in Québec," *Journal of Food Protection*, vol. 70, no. 3, pp. 729–735, 2007.
- [120] M. L. Signorini, E. Rossler, D. C. Díaz David et al., "Antimicrobial resistance of thermotolerant *Campylobacter* species isolated from humans, food-producing animals, and products of animal origin: a worldwide meta-analysis," *Microbial Drug Resistance*, vol. 24, no. 8, pp. 1174–1190, 2018.
- [121] K. Wiczorek, R. Szewczyk, and J. Osek, "Prevalence, antimicrobial resistance, and molecular characterization of *Campylobacter jejuni* and *C. coli* isolated from retail raw meat in Poland," *Veterinárni Medicína*, vol. 57, no. 6, pp. 293–299, 2012.
- [122] A. Noormohamed and M. Fakhr, "A higher prevalence rate of *Campylobacter* in retail beef livers compared to other beef and pork meat cuts," *International Journal of Environmental Research and Public Health*, vol. 10, no. 5, pp. 2058–2068, 2013.
- [123] S. Payot, J. M. Bolla, D. Corcoran, S. Fanning, F. Mégraud, and Q. Zhang, "Mechanisms of fluoroquinolone and macrolide resistance in *Campylobacter* spp.," *Microbes and Infection*, vol. 8, no. 7, pp. 1967–1971, 2006.
- [124] X. Chen, G.-W. Naren, C.-M. Wu et al., "Prevalence and antimicrobial resistance of *Campylobacter* isolates in broilers from China," *Veterinary Microbiology*, vol. 144, no. 1-2, pp. 133–139, 2010.
- [125] E. Shin and Y. Lee, "Characterization of erythromycin-resistant porcine isolates of *Campylobacter coli*," *Microbial Drug Resistance*, vol. 16, no. 3, pp. 231–239, 2010.
- [126] A. Caprioli, "Monitoring of antibiotic resistance in bacteria of animal origin: epidemiological and microbiological methodologies," *International Journal of Antimicrobial Agents*, vol. 14, no. 4, pp. 295–301, 2000.
- [127] M. Lehtopolku, P. Kotilainen, P. Puukka et al., "Inaccuracy of the disk diffusion method compared with the agar dilution method for susceptibility testing of *Campylobacter* spp.," *Journal of Clinical Microbiology*, vol. 50, no. 1, pp. 52–56, 2012.



Hindawi

Submit your manuscripts at
www.hindawi.com

