

Research

Prevention of spina bifida: folic acid intake during pregnancy in Gulu district, northern Uganda

Femke Bannink^{1,8}, Rita Larok², Peter Kirabira², Lieven Bauwens³, Geert van Hove¹

¹Ghent University, Faculty of Psychology and Educational Sciences, Ghent University Henry Dunantlaan 2, B 9000 Ghent, Belgium, ²International Health Sciences University, Kampala, Uganda, ³International Federation for Spina Bifida and Hydrocephalus, Celebroersstraat 16 - 1000 Brussels, Belgium

⁸Corresponding author: Femke Bannink, Ghent University, based in Kampala, Uganda

Key words: Folic acid, spina bifida, pregnancy, antenatal care, Uganda

Received: 04/09/2014 - Accepted: 15/01/2015 - Published: 30/01/2015

Abstract

Introduction: The intake of folic acid before conception and during the first trimester of pregnancy can prevent spina bifida. This paper describes folic acid intake in women in Gulu district in northern Uganda. **Methods:** Structured interviews were held with 394 women attending antenatal care (ANC), 15 mothers of children with spina bifida, and 35 health workers in 2012 and 2013. SPSS16 was used for data analysis. **Results:** 1/4 mothers of children with spina bifida took folic acid during late pregnancy, none preconception. None had knowledge about folic acid and spina bifida prevention. 33.5% of women attending ANC had ever heard about spina bifida, 1% knew folic acid intake can prevent spina bifida. 42.4% took folic acid supplements in late pregnancy, 8.1% during the first trimester, none preconception. All women said to have eaten food rich in folic acid. None were aware about fortified foods. 7% of health workers understood the importance of early folic acid intake. All health workers recommended folic acid intake to women attending ANC. 20% of the health workers and 25% of the women said folic acid supplements are not always available. **Conclusion:** Folic acid intake is limited in northern Uganda. This is attributed to limited education and understanding of women and health workers about the importance of early folic acid intake, late presentation of women at ANC, poor supply chain and dilapidated health services caused by war and poverty. A combination of food fortification, sensitization of health workers, women, and improving folic acid supply is recommended.

Pan African Medical Journal. 2015; 20:90 doi:10.11604/pamj.2015.20.90.5338

This article is available online at: <http://www.panafrican-med-journal.com/content/article/20/90/full/>

© Femke Bannink et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Spina Bifida is a neural tube defect (NTD) caused by a fault in the development of the central nervous system in the first 25 days of the pregnancy [1]. The worldwide incidence varies between 0.17 and 6.39 per 1000 live births [2-5]. An estimated 1,400 children are born with spina bifida in Uganda annually [6]. Incidence and prevalence rates in Uganda are probably higher due to absence of folate consumption by pregnant women, lack of pre-natal care [7], absence of secondary prevention [8], higher exposure to environmental risk factors such as dioxins [9] and fumonisins intake [10-12], and high birth rates [6].

Folic acid (also referred to as folate, folacin, Vitamin B9, pteroylglutamic acid), delivered through supplementation or fortification (or other strategies) prevents the first occurrence of NTDs [13, 14] as well as the recurrence of NTDs in families with previous NTD-affected pregnancy [15]. The World Health Organization (WHO) recommends that all women of childbearing age consume 400 µg of folic acid daily and that women with pregnancies previously affected by NTDs consume 5000 µg of FA daily [16]. Women should consume these amounts in the peri-conceptual period as it takes 8 weeks to reach the optimal level of serum folate [17]. Supplements are an effective way to prevent spina bifida, but there is low compliance [18, 19]. Countries fortifying the flour with at least folic acid see an average 46% reduction of NTD prevalence [20].

In Uganda only 17% of pregnant women attend antenatal care (ANC) before the fourth month of pregnancy [21]. Of those who do attend ANC women are 5.5 months pregnant (median) when they come for their first visit. The Ugandan Government advises intake of folic acid following WHO recommendations [22]. In a study in rural Uganda 13.2% of women attending ANC took folic acid during pregnancy [23]. The health indicators for the northern region are worse compared to the rest of the country [21], partly attributed to conflict between the Government of Uganda and the Lord's Resistance Army (1986 - 2006), which displaced an estimated 2 million people into internally displaced people (IDP) camps [24]. Lack of human resources, limited knowledge, shortages of medical supplies and inadequate infrastructure hamper implementation of quality health services [25, 26]. The study area, Gulu district had 410,673 inhabitants with 15,992 deliveries, 33% of pregnant women attending 4 ANC visits, and 72% of staff positions filled in

health facilities in 2012 [27]. Uganda passed a national legislation to require fortification of wheat and maize flours with iron, zinc, folic acid and other B vitamins in 2011. The legislation requires fortification of white and brown wheat flours and maize products produced in Uganda as well as those imported to Uganda. Whilst some flour producers have started fortifying their products, not all are compliant [28].

Methods

Quantitative and qualitative descriptive cross-sectional survey methods were employed to collect data on folic acid intake and understanding of its effect on the prevention of spina bifida in Gulu district. In total 15 mothers of children with spina bifida were asked about folic acid intake during their pregnancy. In addition 394 women in reproductive age group (15-49 years) attending ANC were surveyed. As well 35 health workers, working at ANCat government health facilities were interviewed. The 394 women and 35 health workers were selected by simple random sampling at antenatal clinics. The mothers of children with spina bifida were purposefully selected from a mobile specialist review clinic for children with neurological conditions in Gulu. The data collection tools included semi structured questionnaires and interviews. Data was entered and analyzed using SPSS16.

Results

Study participants had a median age of 29 years, the majority was married, Christian, completed primary school, and had 3 or 4 children. Table 1 shows the demographic characteristics of 15 mothers of children with spina bifida, 394 women in reproductive age attending antenatal clinics, and 35 health workers. Of the 15 mothers of children with spina bifida interviewed, only a fourth took folic acid supplements during pregnancy, and none of them took these in the first semester (table 2). All mothers of children with spina bifida had never heard about spina bifida before they gave birth to their child, and had not known about the preventative effect of folic acid. All said they ate vegetables rich in folic acid such as spinach and avocado before and during pregnancy, but not on a daily basis. Intake varied during the year depending on whether they were dependent on food handouts during their time in the IDP

camps, and/or the season and availability. None had heard about food fortification before having a child with spina bifida.

Of the women attending ANC, 65.8% lived within 2 kilometers of the health facility (259), 23.4% (92) within 5kms, and 11% (43) beyond 5kms. 50.5% (197) took folic acid, none before pregnancy, and only 8.1% (32) during the first trimester of pregnancy (table 2). In total 33.5% (132/394) of the women attending ANC had ever heard about spina bifida. Only 1% (2) knew about the preventative effect of folic acid intake. Women who had knowledge about spina bifida were more likely to take folic acid than those who did not (table 3, $p < 0.001$). All women said they ate vegetables rich in folic acid such as spinach, okra, beans, lentils and avocado regularly. Intake varied during the year depending on the season and availability. None of the women had heard about food fortification or recognized the signs on available food products. Of the women taking folic acid, 75.5% (149/197) said folic acid (in 5mg tablets) was available at the health facility during their visits. The other 24.5% (46) said sometimes stock outs occurred. Those who had access to folic acid at the health facility were more likely to take it ($p < 0.001$). Women attending health education at the health facilities were more likely to take folic acid compared to those who did not receive health education ($p < 0.001$).

All health workers interviewed had ever heard of spina bifida. The majority of the health workers, 80% (28/35), believed folic acid intake and good antenatal care could prevent spina bifida, 9% (3/35) believed there is no prevention, while 11% (4/35) were not aware of any. Of those who believed folic acid intake could prevent spina bifida (28/35), 7% (2/35) said it should be taken before pregnancy and early weeks of pregnancy, 25% (7/35) said preferably before birth or as soon as the woman realizes she is pregnant for a period of 90 days, 39% (11/35) recommended folic acid use (5mg tablets once a day) throughout pregnancy, and 29% (8/35) health workers recommended folic acid use in only the second and third trimester. All health workers felt that intake of folic acid was important for the babies general development, and mother's health. A fifth (7/35) of the health workers said folic acid is not always available at the health facility they work at. None of the health workers had heard about food fortification.

Discussion

Our findings show that knowledge about the preventative effect of folic acid on spina bifida, and folate intake before and during the first weeks of pregnancy is very limited in northern Uganda. Awareness of spina bifida (33.5%) in women attending ANC was in line with earlier studies, e.g. 25.5% in Nigeria [29], and 53% in Congo [30]. With 17.3% of the Ugandan women in our study taking folic acid supplements during the first trimester of pregnancy, the percentage is slightly lower than the worldwide estimate of only 20% of pregnant women being able to follow the recommendation of folic acid intake for prevention [31]. Whilst peri-conceptual intake of folic acid supplements was 17.2% in Tanzania [32], our study population did not report any early intake of supplements. Women did report to regularly consume foods rich in folic acid. One of the limitations of our study is that it could not define the amounts consumed, as no food samples or biomedical investigations were conducted. The lack of folic acid intake in northern Uganda could partially be explained by the lack of knowledge on the importance of intake prior to and in the first 28 days of pregnancy amongst the health workers interviewed. Health workers in our study had knowledge about the importance of folic acid intake during pregnancy but not about the crucial pre-conceptual and early conceptual period, they were also not aware of the recommended dosage too.

Sensitization of health workers and women in reproductive age is key to encourage early ANC attendance and folic acid intake. Alongside sensitization at the antenatal clinic, women should receive sexual and reproductive health education from an early age. Following the conflict, educational attainment has been lacking in the northern region, and girls secondary school completion remains low [33]. Counseling on folic acid use in the media and marriage counseling may be considered to inform young couples as many will not have received formal sexual and reproductive health education. In Congo patient education through video media increased awareness and knowledge of spina bifida and folic acid [30]. In northern Uganda mobile video media could be explored. Radio messages have proven powerful in transferring health messages in the region and could also be used [34-36].

To improve access to folic acid supplements, there is need to strengthen the supply chain and address stock outs, so that folic acid supplements are provided at the first antenatal visit, or ideally

to all women in child bearing age. Gulu district health facilities had a 57% basic drug stock out in 2012 [27]. Aside assuring folic acid stocks in health facilities, we recommend sensitization on the correct dosage. In a Uganda a huge difference was found between and within districts in the dosage of folic acid provided to pregnant women [37]. The differences were not only caused by lack of supplies but also by health worker's lack of knowledge on updated guidelines. During the war health structures in northern Uganda dilapidated and health workers in the area may not have received the necessary training, resulting in knowledge gaps. With targeted recovery plans of the Ugandan Government, supported by donors, and national guidelines from the Ministry of Health, the health system is recovering and targets are set to improve maternal and child health [22, 38, 39]. Whilst educating health workers on the correct dosage, supplies of 400mcg rather than 5mg tablets should be availed by the National Medical Stores to enable health workers provide the correct preventative dosage, as 1mg and not 5mg is the upper tolerable daily limit [40].

Together with improving folic acid supplementation, we argue for mandatory food fortification and implementation of this process at national level to increase levels of folic acid intake for all women in childbearing age. Participants in our study were not familiar with food fortification. Globally 25% of folic acid-preventable spina bifida is being prevented, and mandatory food fortification is recommended to achieve total prevention [41]. Sensitization about food fortification and recognition of the label can increase use of fortified foods. In South Africa folic acid fortification of staple foods reduced the prevalence of spina bifida with 41.6%, benefits outweighed the costs of food fortification [42]. It should be noted that not all women in reproductive age will benefit from food fortification, as many are subsistent farmers and are more likely to feed from their own produce than goods sold on the markets. We recommend further studies involving women prior to conception, during pregnancy and after birth with frequent measurements of folic acid intake to understand the actual levels of folic acid intake in the population, and evaluation studies to measure possible effects of sensitization campaigns and training of health workers in future.

Conclusion

Folic acid intake is limited in northern Uganda. This is attributed to limited education and understanding of women and health workers

about the importance of early folic acid intake, late presentation of women at ANC, poor supply chain and dilapidated health services caused by war and poverty. A combination of food fortification, sensitization of health workers, women, and improving folic acid supply is recommended to reduce the number of children born with neural tube defects such as spina bifida.

Competing interests

The authors declare no competing interests.

Authors' contributions

Femke Bannink participated in formulation of the research question, designed and collected data from mothers of children with spina bifida, and health workers, analyzed the data, and drafted the manuscript. Rita Larok participated in formulating the research question, design of the study, carried out data collection in women attending antenatal care, and carried out data analysis. She contributed to the writing of the manuscript. Peter Kirabira participated in formulating the research question, design of the study, and supervised data collection and analysis. He contributed to the writing of the manuscript too. Lieven Bauwens participated in formulating the research question, design of the study, data analysis and writing the manuscript. Geert van Hove supervised and participated in formulating the research question, study design, data analysis and writing the manuscript. This study complies with the Forum for African Medical Editors ethical principles. Informed consents of all subjects were obtained and the study was approved by the Science and Ethics Committee Ghent University, the International Health Sciences University Research and Ethics Committee and the Uganda National Council for Science and Technology. All authors read and approved the final version of the manuscript.

Acknowledgments

The authors thank all participants, parents of children with spina bifida, staff of the Gulu District Health Offices, the Gulu Regional Orthopaedic Workshop and Rehabilitation Center, AVSI Foundation,

and CURE Children's Hospital Uganda for their participation and help in this study. Our gratitude goes to the International Federation for Spina Bifida and Hydrocephalus, and the IF Uganda office staff for their logistic support.

Tables

Table 1: Demographic characteristics respondents northern Uganda

Table 2: Folic acid intake in mothers of children with spina bifida and women attending ANC in northern Uganda

Table 3: Factors associated with folic acid intake in women attending ANC in northern Uganda

References

1. Northrup H, Volcik KA. Spina bifida and other neural tube defects. *Curr Probl Pediatr.* 2000 Nov-Dec;30(10):313-32. **PubMed | Google Scholar**
2. Bowman RM, Boshnjaku V, McLone DG. The changing incidence of myelomeningocele and its impact on pediatric neurosurgery: a review from the Children's Memorial Hospital; *Child's nervous system: ChNS: official journal of the International Society for Pediatric Neurosurgery.* 2009 Jul;25(7):801-6. **PubMed | Google Scholar**
3. Shaer CM, Chescheir N, Schulkin J. Myelomeningocele: a review of the epidemiology, genetics, risk factors for conception, prenatal diagnosis, and prognosis for affected individuals. *Obstet Gynecol Surv.* 2007 Jul;62(7):471-9. **PubMed | Google Scholar**
4. Msamati BC, Igbigbi PS, Chisi JE. The incidence of cleft lip, cleft palate, hydrocephalus and spina bifida at Queen Elizabeth Central Hospital, Blantyre, Malawi. *The Central African journal of medicine.* 2000 Nov;46(11):292-6. **PubMed | Google Scholar**
5. Kinasha AD, Manji K. The incidence and pattern of neural tube defects in Dar es Salaam, Tanzania. *Eur J Pediatr Surg.* 2002 Dec; vol 12, Suppl 1: S38-9. **PubMed | Google Scholar**
6. Warf BC, Wright EJ, Kulkarni AV. Factors affecting survival of infants with myelomeningocele in southeastern Uganda. *J Neurosurg Pediatr.* 2011 Feb; 7(2): 127-33. **PubMed | Google Scholar**
7. Miles M. Children with spina bifida and hydrocephalus in Africa: can medical, family and community resources improve the life chances?. The Independent Living Institute. 2006. **Google Scholar**
8. Frey L, Hauser WA. Epidemiology of neural tube defects. *Epilepsia.* 2003; vol 44, Suppl 3: 4-13. **PubMed | Google Scholar**
9. Safi J, Joyeux L, Chalouhi GE. Periconceptional folate deficiency and implications in neural tube defects. *Journal of pregnancy.* 2012; 2012: 295083. **PubMed | Google Scholar**
10. Hendricks K. Fumonisin and neural tube defects in South Texas. *Epidemiology.* 1999 Mar; 10(2): 198-200. **PubMed | Google Scholar**
11. Marasas WF, Riley RT, Hendricks KA, Stevens VL, Sadler TW, Gelineau-van Waes J, et al. Fumonisin disrupt sphingolipid metabolism, folate transport, and neural tube development in embryo culture and in vivo: a potential risk factor for human neural tube defects among populations consuming fumonisin-contaminated maize. *The Journal of nutrition.* 2004 Apr; 134(4): 711-6. **PubMed | Google Scholar**
12. Wild CP, Gong YY. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis.* 2010 Jan; 31(1): 71-82. **PubMed | Google Scholar**
13. Czeizel AE. Primary prevention of neural-tube defects and some other major congenital abnormalities: recommendations for the appropriate use of folic acid during pregnancy. *Paediatric drugs.* 2000 Nov-Dec; 2(6): 437-49. **PubMed | Google Scholar**

14. Wilson RD, Davies G, Desilets V, Reid GJ, Summers A, Wyatt P, et al. The use of folic acid for the prevention of neural tube defects and other congenital anomalies. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC.* 2003 Nov; 25(11): 959-73. **PubMed | Google Scholar**
15. MRC VitaminStudy Research Group. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. *Lancet.* 1991 Jul 20; 338(8760): 131-7. **PubMed | Google Scholar**
16. Pena-Rosas JP, De-Regil LM, Dowswell T, Viteri FE. Daily oral iron supplementation during pregnancy. *The Cochrane database of systematic reviews.* 2012; 12: CD004736. **PubMed | Google Scholar**
17. World Health Organization. Daily iron and folic acid supplementation in pregnant women http://www.who.int/nutrition/publications/micronutrients/guidelines/daily_ifa_supp_pregnant_women/en/2012. Accessed on September 04 2014. **Google Scholar**
18. Nilsen RM, Vollset SE, Gjessing HK, Magnus P, Meltzer HM, Haugen M, et al. Patterns and predictors of folic acid supplement use among pregnant women: the Norwegian Mother and Child Cohort Study. *The American journal of clinical nutrition.* 2006 Nov; 84(5): 1134-41. **PubMed | Google Scholar**
19. Brough L, Rees GA, Crawford MA, Dorman EK. Social and ethnic differences in folic acid use preconception and during early pregnancy in the UK: effect on maternal folate status. *Journal of human nutrition and dietetics : the official journal of the British Dietetic Association.* 2009 Apr; 22(2): 100-7. **PubMed | Google Scholar**
20. Pachón H, Kancherla V, Handforth B, Tyler V, Bauwens L. Folic acid fortification of wheat flour: A cost-effective public health intervention to prevent birth defects in Europe. *Nutrition Bulletin.* 2013; 38(2): 201-9. **PubMed | Google Scholar**
21. Uganda Government Ministry of Health- Uganda Demographic Health Survey www.ubos.org/onlinefiles/uploads/ubos/UDHS/UDHS2011.pdf2011 accessed on September 04 2014. **PubMed | Google Scholar**
22. Uganda Government Ministry of Health- Guidelines for maternal nutrition in Uganda www.health.go.ug/docs/GI_MN.pdf: Government of Uganda; 2010 accessed on September 04 2014. **PubMed | Google Scholar**
23. Mbule MA, Byaruhanga YB, Kabahenda M, Lubowa A. Determinants of anaemia among pregnant women in rural Uganda. *Rural and remote health.* 2013 Apr-Jun; 13(2): 2259. **PubMed | Google Scholar**
24. Annan JB, Carlson C, Mazurana D. The State of Female Youth and in Northern Uganda: Findings from the Survey of War Affected Youth, Phase II. 2008. **Google Scholar**
25. AVSI Foundation, UNICEF Uganda. Health Facilities Survey of 191 Health facilities in the Acholi sub-region in northern Uganda. Kampala: AVSI Uganda, 2011. **Google Scholar**
26. Ahoua L, Ayikoru H, Gnauck K, Odaru G, Odar E, Ondo-Onama C, et al. Evaluation of a 5-year programme to prevent mother-to-child transmission of HIV infection in Northern Uganda. *Journal of tropical pediatrics.* 2010 Feb; 56(1): 43-52. **PubMed | Google Scholar**
27. AVSI Foundation, Gulu District Health Office. Monthly HMIS reports Gulu district. 2012. **Google Scholar**
28. FoodFortificationInitiative. Uganda Requires Wheat and Maize Flour Fortification. Periodic Update: FFI Newsletter March 20 12 2012. **Google Scholar**
29. Rabi TB, Tihamiyu LO, Awoyinka BS. Awareness of spina bifida and periconceptional use of folic acid among pregnant women in a developing economy; *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery.* 2012 Dec; 28(12): 2115-9. **PubMed | Google Scholar**

30. Claude KM, Juvenal KL, Hawkes M. Applying a knowledge-to-action framework for primary prevention of spina bifida in tropical Africa. *Maternal & child nutrition*. 2012 Apr; 8(2): 174-84. **PubMed | Google Scholar**
31. Scott J. Folic acid consumption throughout pregnancy: differentiation between trimesters. *Annals of nutrition & metabolism*. 2011; 59(1): 46-9. **PubMed | Google Scholar**
32. Ogundipe O, Hoyo C, Ostbye T, Oneko O, Manongi R, Lie RT, et al. Factors associated with prenatal folic acid and iron supplementation among 21,889 pregnant women in Northern Tanzania: a cross-sectional hospital-based study. *BMC public health*. 2012; 12: 481. **PubMed | Google Scholar**
33. Uganda Government, Ministry of Education and Sports. Uganda Education Statistical Abstract. Kampala, Uganda 2011. **PubMed | Google Scholar**
34. Kitara DL, Ocerro A, Lanyero J, Ocom F. Roll-out of medical male circumcision (MMC) for HIV prevention in non-circumcising communities of Northern Uganda. *The Pan African medical journal*. 2013; 15: 100. **PubMed | Google Scholar**
35. Bannink-Mbazzi F, Lowicki-Zucca M, Ojom L, Kabasomi SV, Esiru G, Homsy J. High PMTCT program uptake and coverage of mothers, their partners, and babies in northern Uganda: achievements and lessons learned over 10 years of implementation (2002-2011). *J Acquir Immune Defic Syndr*. 2013 Apr 15; 62(5): e138-45. **PubMed | Google Scholar**
36. Mertens P, Bannink F. Interdisciplinary care for children with spina bifida in east and southern Africa. *Journal of Medicine and the Person*. 2012; Volume 10, Issue 3: pp 129-135. **PubMed | Google Scholar**
37. World Health Organization. Action - The MOST Project - Iron and folic acid supplementation - Pregnant women (PW) <https://extranet.who.int/nutrition/gina/en/node/114862001> [cited 2014 30-6] accessed on September 04 2014. **PubMed | Google Scholar**
38. Uganda Government. Peace Recovery and Development Plan for Northern Uganda. Kampala, Uganda 2007. **Google Scholar**
39. Uganda Government. National Development Plan 2010/11 - 2014/15. Kampala, Uganda. 2010. **PubMed | Google Scholar**
40. Health National Institute of Health. Folate: Dietary Supplement Fact Sheet 2012 [cited 2014 02-07]. <http://ods.od.nih.gov/factsheets/FolateHealthProfessional/#en2>. **Google Scholar**
41. Youngblood ME, Williamson R, Bell KN, Johnson Q, Kancherla V, Oakley GP, Jr. 2012 Update on global prevention of folic acid-preventable spina bifida and anencephaly. *Birth defects research Part A, Clinical and molecular teratology*. 2013 Oct; 97(10): 658-63. **PubMed | Google Scholar**
42. Sayed AR, Bourne D, Pattinson R, Nixon J, Henderson B. Decline in the prevalence of neural tube defects following folic acid fortification and its cost-benefit in South Africa. *Birth defects research Part A, Clinical and molecular teratology*. 2008 Apr; 82(4): 211-6. **PubMed | Google Scholar**

Characteristics	Mothers of children with spina bifida (N=15)	Women attending antenatal clinics (N=394)	Health workers in antenatal clinics (N=35)
Age (mean)	31 (range 22 – 36)	29 years (range 18 – 40, SD 7.8)	28 years (range 22 – 42)
Marital status	6.7% (1) single, 60% (9) married, 13.3% (2) separated, 20% (3) widowed	19% (75) single, 72.2% (284) married, 5% (20) separated, 3.8% (15) widowed	11.4% (4) single, 88.6% (31) married
Number of children	4.2 mean (range 1 – 8)	3.8 children (range 0 - 9)	3.1 mean (0 – 6)
Education	26.6 % (4) never went to school, 60.0% completed primary (9), 6.7% (1) completed secondary, 6.7% (1) higher education	15.8% (63) never went to school, 55.6% (219) completed primary, 22.4% (88) completed secondary, 6.2% (24) higher education	54% (19) nursing - enrolled, 11% (4) registered nursing - registered, 35% (12) midwifery
Religion	86.7% (13) Christian 13.3% (2) Muslim	97% (382) Christian 3% (12) Muslim	98% (34) Christian 2% (1) Muslim
Occupation	73.2% (11) peasant farmers, 13.4% (2) petty traders, 13.4% (2) others	44.7% (176) peasant farmers, 39.9% (157) petty traders, 15.5% (61) others	54% (19) enrolled nurse, 11% (4) registered nurse, 35% (12) midwife

Started taking FA	Mother of children with spina bifida	Women attending antenatal care
Before conception	0% (0)	0% (0)
During pregnancy (first trimester)	0% (0)	8.1% (32)
During pregnancy (2nd and 3rd trimester)	26.7% (4)	42.4% (167)
Did not take FA	(11)	49.5% (195)

Table 3: Factors associated with folic acid intake in women attending ANC in northern Uganda					
Factors		Folic Acid Intake		N (total)	χ²
		Yes	No		
Knowledge of spina bifida	Yes	91 (45.5)	41 (21.1)	132 (33.5)	26.24*
	No	109 (54.5)	153 (78.9)	262 (66.5)	
Availability of drugs	Yes	168 (87.0)	121 (63.7)	289 (75.5)	28.22*
	No	25 (13.0)	69 (36.3)	94 (24.5)	
Health education	Yes	52 (25.7)	13 (6.6)	65 (16.3)	26.59*
	No	150 (74.3)	183 (93.4)	333 (83.7)	
p<0.001					