

# PRINCIPAL DIFFERENTIAL IDEALS AND A GENERIC INVERSE DIFFERENTIAL GALOIS PROBLEM FOR $GL_n$

LOURDES JUAN

ABSTRACT. We characterize the principal differential ideals of a polynomial ring in  $n^2$  indeterminates with coefficients in the ring of differential polynomials in  $n^2$  indeterminates and derivation given by a “general” element of  $\text{Lie}(GL_n)$  and use this characterization to construct a generic Picard-Vessiot extension for  $GL_n$ . In the case when the differential base field has finite transcendence degree over its field of constants we provide necessary and sufficient conditions for solving the inverse differential Galois problem for this group via specialization from our generic extension.

## INTRODUCTION

Given a differential field  $F$  and differential indeterminates  $Y_{ij}$ ,  $i, j = 1, \dots, n$  over  $F$  one writes  $F\{Y_{ij}\}$  for the ring of differential polynomials in the  $Y_{ij}$ , i.e., the polynomial ring  $F[Y_{1,1,0}, Y_{1,1,1}, \dots, Y_{1,1,s}, \dots, Y_{n,n,0}, \dots]$  with derivation extending the derivation on  $F$  by  $D(Y_{i,j,k}) = Y_{i,j,k+1}$ . For convenience, denote  $Y_{i,j,k}$  by  $Y_{i,j}^{(k)}$  and  $Y_{i,j,0}$  by  $Y_{ij}$ . Then one can extend this derivation to the ring  $R = F\{Y_{ij}\}[X_{ij}]$  where the  $X_{ij}$  are algebraically independent over the differential quotient field  $F\langle Y_{ij} \rangle$  of  $F\{Y_{ij}\}$  using the formula  $D(X_{ij}) = \sum_{\ell=1}^n Y_{i\ell} X_{\ell j}$ . If we pass to the above quotient field  $F\langle Y_{ij} \rangle$  and then localize  $F\langle Y_{ij} \rangle[X_{ij}]$  at  $\det[X_{ij}]$ , we obtain the coordinate ring of  $GL_n$  over  $F\langle Y_{ij} \rangle$  and  $D$  becomes a “general” element of  $\text{Lie}(GL_n)$ .

In this paper we show that the principal differential ideals of  $R$  (i.e., the ideals  $\mathcal{I} = (p)$  with  $p$  dividing  $D(p)$ ) are the differential ideals generated by elements of the form  $\det^a[X_{ij}]$ , with  $a \in \mathbb{N}$ . A polynomial  $p$  that divides its derivative is called a *Darboux polynomial*. Our result can then be stated as follows:

---

*Date:* August 22, 2001.

*1991 Mathematics Subject Classification.* Primary 12H05; Secondary 12F12, 12Y05, 20G15i.

*Key words and phrases.* Constants, Darboux polynomials, differential field, differential Galois group, differential specialization, inverse differential Galois problem, generic, multivariable division algorithm, new constant, Picard-Vessiot extension, principal differential ideals, wronskian.

This paper was written while the author held an NSF-funded postdoctoral fellowship at the MSRI. The author wishes to thank the MSRI for its hospitality.

**Theorem 1.** *Let  $R = F\{Y_{ij}\}[X_{ij}]$  and let  $p$  be a Darboux polynomial in  $R$ . Then there are  $\ell \in F$  and  $a \in \mathbb{N}$  such that  $p = \ell \det^a[X_{ij}]$ . Therefore, the only principal differential ideals in  $R$  are those of the form  $\mathcal{I} = (\det^a[X_{ij}])$ .*

The proof of Theorem 1 involves some long and delicate computations that make use of Gröbner bases machinery.

Now, suppose that the field  $C$  of constants of  $F$  is algebraically closed. We use Theorem 1 to show that the quotient field  $F\langle Y_{ij}\rangle(X_{ij})$  of  $R$  is a no-new-constant extension of  $F\langle Y_{ij}\rangle$ . Similar to the above,  $F\langle Y_{ij}\rangle(X_{ij})$  is the function field of  $\mathrm{GL}_n$  over  $F\langle Y_{ij}\rangle$ . This allows us to give an affirmative answer, for the group  $\mathrm{GL}_n(C)$ , to the following

**Generic Inverse Differential Galois Problem:** *For a connected algebraic group  $G$  over  $C$  find a generic Picard-Vessiot extension of  $F$  with differential Galois group  $G$ .*

By *generic extension* we mean a Picard-Vessiot extension of a *generic field* that contains  $F$  and such that every Picard-Vessiot extension of  $F$  for  $G$  in the usual sense can be obtained from the generic one by specialization. Conversely, any such specialization will provide a solution to the inverse differential Galois problem in the usual sense, namely, to determine, given  $F$  and  $C$  as above, and a linear algebraic group  $G$  over  $C$ , what differential field extensions  $E \supset F$  are Picard-Vessiot extensions with differential Galois group  $G$  and, in particular, whether there are any.

This result is the content of:

**Theorem 2.** *The differential field extension  $F\langle Y_{ij}\rangle(X_{ij}) \supset F\langle Y_{ij}\rangle$  is a generic Picard-Vessiot extension of  $F$  with differential Galois group  $\mathrm{GL}_n(C)$ .*

We point out that Theorem 2 is a consequence of Theorem 1 but not equivalent to it: the fact that  $F\langle Y_{ij}\rangle(X_{ij}) \supset F\langle Y_{ij}\rangle$  is a no-new-constant extension does not automatically give information about what the Darboux polynomials in  $R$  are. Darboux polynomials are also interesting in other related applications such as studying the integrability of differential equations [2, 9, 18, 16, 17, 35].

A more direct proof for Theorem 2 was pointed out to us by Michael Singer. Singer proves that  $F\langle Y_{ij}\rangle(X_{ij}) \supset F\langle Y_{ij}\rangle$  is a no-new-constant extension by showing that  $F\langle Y_{ij}\rangle(X_{ij})$  is isomorphic to  $F\langle X_{ij}\rangle$ . Singer's proof and our generalization of it to all connected linear algebraic groups will appear in a subsequent publication [12].

Now, suppose that  $F$  has finite transcendence degree over  $C$  say,  $F = C(t_1, \dots, t_m)[z_1, \dots, z_k]$ , where the  $t_i$  are algebraically independent over  $C$  and the  $z_i$  are algebraic over  $C(t_1, \dots, t_m)$ . Consider the differential field  $F\langle X_{ij}\rangle$  with derivation given by  $D(X_{ij}) = \sum_{\ell=1}^n f_{i\ell} X_{\ell j}$ . Let  $\mathcal{C}$  denote its field of constants. Let  $R = F\{Y_{ij}\}[X_{ij}]$  be the differential ring defined above. For  $k \geq 1$  let  $\mathbb{T}_k$  denote the set of monomials in  $R$  which have total degree less than or equal to  $k$  and which involve both the  $t_i$  and the  $X_{ij}$ . Fix a term order on the set  $\mathbb{T}$  of monomials in the  $t_i$  and the  $X_{ij}$  and let  $W_k(Y_{ij})$  denote the wronskian of  $\mathbb{T}_k$  relative to that order (the order will

only affect the wronskian by a sign). The following theorem summarizes our specialization results:

**Theorem 3.**  *$F(X_{ij}) \supset F$  is a Picard-Vessiot extension for  $\mathrm{GL}_n(C)$  if and only if all the wronskians  $W_k(Y_{ij})$  map to nonzero elements in  $F(X_{ij})$  via the specialization  $Y_{ij} \mapsto f_{ij} \in F$ .*

The above condition on the wronskians means that all the sets  $\mathbb{T}_k$ , for  $k \geq 1$ , are linearly independent over  $\mathcal{C}$ . This is in turn equivalent to the fact that the set of all the  $t_i$  and all the  $X_{ij}$  are algebraically independent over  $\mathcal{C}$ . Unfortunately, Theorem 3 gives infinitely many conditions. We do not know at present how to use these conditions to effectively construct solutions to the inverse problem, and this constitutes an interesting open problem.

A specialization as in Theorem 3, however, is known to exist by a result of C. Mitschi and M. Singer [23]. They give a constructive algebraic solution to the inverse problem for all connected linear algebraic groups (and, in particular, for  $\mathrm{GL}_n(C)$ ) when  $F$  has finite transcendence degree over  $C$ . An interesting direction of research in connection with the previous open problem is to give a complete description of the solutions (isomorphic and non-isomorphic) that may arise in this situation.

The work of Mitschi and Singer in [23] makes use of the logarithmic derivative and an inductive technique developed by Kovacic [14, 15] to lift a solution to the inverse problem from  $G/R_u$ , where  $R_u$  is the unipotent radical of  $G$ , to the full group  $G$ . Using this machinery Kovacic proved that it is enough to find a solution to the inverse problem for reductive groups (observe that  $G/R_u$  is reductive). In [25], van der Put explains and partly proves the results in [23].

In the introduction of [23] the authors briefly review previous work on the inverse problem such as results of Bialynicki-Birula in [4], Kovacic [14, 15], Ramis [26, 27], Singer [30], Tretkoff and Tretkoff [32], Beukers and Heckman [3], Katz [13], Duval and Mitschi [8], Mitschi [21, 22], Duval and Loday-Richaud [7], Ulmer and Weil [33] and Singer and Ulmer [31]. A more extensive survey on the inverse problem can be found in M. Singer's [29].

The constructive algebraic solutions to the inverse differential Galois problem for connected linear algebraic groups that are currently available are based on Kolchin's Main Structure Theorem for Picard-Vessiot extensions (see Theorem 2.1.1 below). In particular, a corollary to this theorem (see Theorem 2.1.2) establishes that if  $E \supset F$  is Picard-Vessiot and  $G$  is, for example, unipotent or solvable or  $G = \mathrm{GL}_n$  or  $G = \mathrm{SL}_n$ , then  $E$  is isomorphic as an  $F$ -module and as a  $G$ -module to the function field of the group  $G_F$  obtained from  $G$  by extension of scalars from  $C$  to  $F$ . Therefore, to get a Picard-Vessiot extension  $E \supset F$  with group  $G$  (if it exists) one can begin by taking  $E$  to be the function field of  $G_F$  and then the problem reduces to extending the derivation from  $F$  to  $E$  in such a way that  $E \supset F$  is Picard-Vessiot for that derivation. In this paper we use this approach for our construction.

The idea of tackling the inverse problem by constructing generic extensions is inspired by the works of E. Noether [24] for the Galois theory of algebraic equations. Following her approach, L. Goldman in [10] introduced the notion of a *generic differential equation with group  $G$* . Goldman explicitly constructed a generic equation with group  $G$  for some groups. However, after specializing Goldman's equation, the group of the new equation obtained is a subgroup of the original group. In order to solve the inverse problem by this means, we need to keep the original group as the group of the equation after specialization. Goldman's generic equation for  $\mathrm{GL}_n$  is equivalent to Magid's *general equation of order  $n$*  (Example 5.26 in [19]).

More work in the spirit of Goldman's generic equation came some years later in J. Miller's dissertation [20]. He defined the notion of hilbertian differential field and gave a sufficient condition for the generic equation with group  $G$  to specialize to an equation over such a field with group  $G$  as well. However, as pointed out by Mitschi and Singer in [23], his condition was stronger than the analogous one for algebraic equations and this made the theory especially difficult to apply for those groups that were not already known to be Galois groups.

We use the terminology of A. Magid's book [19]. In [19] the reader may also find definitions and proofs of some results from differential Galois theory that will be recalled here.

This paper contains the results of the author's Ph.D. dissertation [11]. I wish to thank my Ph.D. advisor Andy Magid for the many valuable research meetings that we had. I am also grateful to Michael Singer for many enlightening conversations on the inverse problem.

**Notation.** Throughout this paper  $F$  denotes a differential field with algebraically closed field of constants  $C$ .

## 1. PRINCIPAL DIFFERENTIAL IDEALS IN $F\{Y_{ij}\}[X_{ij}]$

### 1.1. Darboux polynomials in $\mathbf{F}\{\mathbf{Y}_{ij}\}[\mathbf{X}_{ij}]$ .

**Definition 1.1.1.** Let  $D$  be a derivation on the polynomial ring  $A = k[Z_1, \dots, Z_s]$ . A polynomial  $p \in A$  is called a Darboux polynomial if there is a polynomial  $q \in A$  such that  $D(p) = qp$ . That is,  $p$  divides  $D(p)$ .

An ideal  $\mathcal{I}$  of  $A$  is a differential ideal if  $D(\mathcal{I}) \subset \mathcal{I}$ . In particular,  $\mathcal{I} = (p)$  is a principal differential ideal if  $p$  divides  $D(p)$ . Hence, Darboux polynomials in  $A$  correspond to principal differential ideals.

Let  $F\{Y_{ij}\}$  be the ring of differential polynomials in the  $Y_{ij}$  and  $F\langle Y_{ij} \rangle$  its differential quotient field. By that we mean the usual quotient field endowed with the natural derivation:

$$\mathcal{D}\left(\frac{p}{q}\right) = \frac{D(p)q - pD(q)}{q^2}.$$

for  $p, q \in F\{Y_{ij}\}$ , where  $D$  is the derivation on  $F\{Y_{ij}\}$ .

Consider the differential ring  $R = F\{Y_{ij}\}[X_{ij}]$  where the  $X_{ij}$ ,  $1 \leq i, j \leq n$ , are algebraically independent over  $F\langle Y_{ij} \rangle$  and derivation extending the derivation on  $F\{Y_{ij}\}$  by a formula

$$D(X_{ij}) = \sum_{\ell=1}^n Y_{i\ell} X_{\ell j}.$$

An elementary computation shows that an element of the form  $p = \ell \det^a[X_{ij}]$  with  $\ell \in F$  and  $a \in \mathbb{N}$  is a Darboux polynomial in  $R$  with  $D(p) = (\frac{\ell'}{\ell} + a \sum_{i=1}^n Y_{ii})p$ . The rest of this section is devoted to showing that all the Darboux polynomials in  $R$  are of this form.

The multinomial notation  $a_\alpha \mathbf{Z}^\alpha$  will be used to denote a term of the form  $a_{\alpha_1 \dots \alpha_s} Z_1^{\alpha_1} \dots Z_s^{\alpha_s}$ .

First, we show that there are no non-trivial Darboux polynomials in the  $Y_{ij}$ . For simplicity, if  $h(Y) \in F\{Y_{ij}\}$ , we write  $h'(Y)$  for  $D(h(Y))$ . Notice that this is not the usual meaning  $h'(Y) = \sum h'_\alpha \mathbf{Y}^\alpha$ .

**Proposition 1.1.2.** *If  $h(Y) \in F\{Y_{ij}\}$  satisfies  $h'(Y) = g(Y)h(Y)$  for some  $g(Y) \in F\{Y_{ij}\}$  then  $h(Y) \in F$ . That is, there are no non-trivial Darboux polynomials in  $F\{Y_{ij}\}$ .*

*Proof.* Write  $Y_{ij,k}$  for  $Y_{ij}^{(k)}$  and order the set of subindices  $\{ij, k\}$ ,  $i, j, k \in \mathbb{N}$ , with the lexicographical ordering. That is,  $\{i_1 j_1, k_1\} > \{i_2 j_2, k_2\}$  if and only if the first coordinates  $s_1$  and  $s_2$  from the left, for  $s = i, j, k$  above, which are different satisfy  $s_1 > s_2$ .

Let  $h(Y_{ij})$  and  $g(Y_{ij})$  be as in the hypothesis. Denote by  $\{mn, t\}$  the largest subindex such that  $Y_{mn,t}$  occurs in  $h(Y)$  and put

$$h(Y) = \sum_{\alpha} a_{\alpha} Y_{11}^{\alpha_{11}} \dots Y_{mn,t}^{\alpha_{mn,t}}.$$

Then

$$\begin{aligned} h'(Y) &= \sum_{\alpha} a'_{\alpha} Y_{11}^{\alpha_{11}} \dots Y_{mn,t}^{\alpha_{mn,t}} + \sum_{\alpha} a_{\alpha} \alpha_{11} Y_{11}^{\alpha_{11}-1} Y_{11,1}^{\alpha_{11,1}+1} \dots Y_{mn,t}^{\alpha_{mn,t}} \\ &\quad + \dots + \sum_{\alpha} a_{\alpha} \alpha_{mn,t} Y_{11}^{\alpha_{11}} \dots Y_{mn,t}^{\alpha_{mn,t}-1} Y_{mn,t+1} \\ &= h_1(Y_{11}, \dots, Y_{mn,t}) + \left( \sum_{\alpha} a_{\alpha} \alpha_{mn,t} Y_{11}^{\alpha_{11}} \dots Y_{mn,t}^{\alpha_{mn,t}-1} \right) Y_{mn,t+1} \\ &= g(Y)h(Y). \end{aligned}$$

Now, for  $Y_{mn,t+1} = Y'_{mn,t}$  we have  $\{mn, t+1\} > \{mn, t\}$ . Thus it may not occur in  $h(Y)$  by the choice of  $\{mn, t\}$ . Also, it does not occur in  $h_1(Y_{11}, \dots, Y_{mn,t})$ . Thus, the above equation implies that  $Y_{mn,t+1}$  must occur in  $g(Y)$ . Let  $g_{t+1}(Y)$  be its coefficient in  $g(Y)$  and write

$$h_2(Y) = \sum_{\alpha} a_{\alpha} \alpha_{mn,t} Y_{11}^{\alpha_{11}} \dots Y_{mn,t}^{\alpha_{mn,t}-1}.$$

We have

$$h(Y)g_{t+1}(Y)Y_{mn,t+1} = h_2(Y)Y_{mn,t+1}$$

or

$$h(Y)g_{t+1}(Y) = h_2(Y).$$

But the total degree of  $h_2(Y)$  is strictly less than the total degree of  $h(Y)$ . This forces  $h(Y) \in F$ .  $\square$

Next, we proceed to the computations in  $R$ . The ring  $F[X_{ij}]$  is assumed to be ordered with the degree reverse lexicographical order (*degrevlex*). That is, the set

$$\mathbb{T}^{n^2} = \{\mathbf{X}^\beta \mid \mathbf{X} = (X_{ij}), \beta = (\beta_{ij}) \in \mathbb{N}^{n^2}\}$$

of the power products in the  $X_{ij}$  is ordered by  $X_{11} > \cdots > X_{1n} > \cdots > X_{n1} > \cdots > X_{nn}$ , and

$$\mathbf{X}^\alpha < \mathbf{X}^\beta \iff \begin{cases} \sum_{j=1}^n \sum_{i=1}^n \alpha_{ij} < \sum_{j=1}^n \sum_{i=1}^n \beta_{ij} \\ \text{or} \\ \sum_{j=1}^n \sum_{i=1}^n \alpha_{ij} = \sum_{j=1}^n \sum_{i=1}^n \beta_{ij}, \text{ and the first coordinates} \\ \alpha_{ij}, \beta_{ij} \text{ from the right which are different satisfy } \alpha_{ij} > \beta_{ij}. \end{cases}$$

We will refer to the leading term of a polynomial with respect to this order as its leading power product.

**Remarks. 1.1.3** (Derivative of a power product in the  $X_{ij}$ ). *Let*

$$\mathbf{X}^\alpha = X_{11}^{\alpha_{11}} \cdots X_{1n}^{\alpha_{1n}} \cdots X_{n1}^{\alpha_{n1}} \cdots X_{nn}^{\alpha_{nn}},$$

then

$$\begin{aligned} D(\mathbf{X}^\alpha) &= \left( \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ij} \right) \mathbf{X}^\alpha \\ &+ \sum_{i=1}^n \sum_{j=1}^n \left( \sum_{\ell > i} \alpha_{ij} Y_{i\ell} X_{11}^{\alpha_{11}} \cdots X_{ij}^{\alpha_{ij}-1} \cdots X_{\ell j}^{\alpha_{\ell j}+1} \cdots X_{nn}^{\alpha_{nn}} \right. \\ &\quad \left. + \sum_{\ell < i} \alpha_{ij} Y_{i\ell} X_{11}^{\alpha_{11}} \cdots X_{\ell j}^{\alpha_{\ell j}+1} \cdots X_{ij}^{\alpha_{ij}-1} \cdots X_{nn}^{\alpha_{nn}} \right). \end{aligned}$$

**1.1.4.** *For a given  $\alpha$  and  $\mathbf{X}^\alpha$  as before, we want find all the power products  $\mathbf{X}^\beta$  such that  $\mathbf{X}^\alpha$  occurs in  $D(\mathbf{X}^\beta)$ . If that is the case,  $\mathbf{X}^\alpha$  will appear in  $D(\mathbf{X}^\beta)$  in a product of the form  $Y_{rt}\mathbf{X}^\alpha$ . By Remark 1.1.3 all such power products are of the form*

$$\mathbf{X}^{\alpha_{rs,t}} = \begin{cases} X_{11}^{\alpha_{11}} \cdots X_{rs}^{\alpha_{rs}+1} \cdots X_{ts}^{\alpha_{ts}-1} \cdots X_{nn}^{\alpha_{nn}} & \text{if } r < t \\ X_{11}^{\alpha_{11}} \cdots X_{ts}^{\alpha_{ts}-1} \cdots X_{rs}^{\alpha_{rs}+1} \cdots X_{nn}^{\alpha_{nn}} & \text{if } r > t \end{cases}$$

for  $1 \leq r, s \leq n$ ,  $t \neq r$ , and  $\mathbf{X}^\alpha$  itself.

**1.1.5.** *Let  $p \in R$ . Since  $D(X_{ij}) = \sum_{\ell=1}^n Y_{i\ell} X_{\ell j}$ , the total degree of  $p$  with respect to the  $X_{ij}$  does not change after differentiation. Therefore, if  $D(p) = qp$  then  $q \in F\{Y_{ij}\}$ .*

**Proposition 1.1.6.** *Let  $p \in R$ . Write it as  $p = \sum_{\alpha} p_{\alpha}(Y) \mathbf{X}^{\alpha}$ , with  $p_{\alpha}(Y) \in F\{Y_{ij}\}$ . Then for any  $\alpha$  with  $p_{\alpha}(Y) \neq 0$ , the coefficient of  $\mathbf{X}^{\alpha}$  in  $D(p)$  is*

$$p'_{\alpha}(Y) + p_{\alpha}(Y) \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii} + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y) Y_{i\ell},$$

where  $\alpha_{ij,\ell}$  is the exponent vector of the power product

$$\mathbf{X}^{\alpha_{ij,\ell}} = \begin{cases} X_{11}^{\alpha_{11}} \dots X_{ij}^{\alpha_{ij}+1} \dots X_{\ell j}^{\alpha_{\ell j}-1} \dots X_{nn}^{\alpha_{nn}} & \text{if } i < \ell \\ X_{11}^{\alpha_{11}} \dots X_{\ell j}^{\alpha_{\ell j}-1} \dots X_{ij}^{\alpha_{ij}+1} \dots X_{nn}^{\alpha_{nn}} & \text{if } \ell > i \end{cases}$$

as in Remark 1.1.4.

*Proof.* This is a direct consequence of Remarks 1.1.3 and 1.1.4.  $\square$

**Proposition 1.1.7.** *Let  $p \in R$  and suppose that  $D(p) = qp$ , for some  $q \in F\{Y_{ij}\}$ . Then  $p \in F[X_{ij}]$ .*

*Proof.* Let  $p = \sum_{\alpha} p_{\alpha}(Y) \mathbf{X}^{\alpha}$ . Then

$$\begin{aligned} D(p) &= \sum_{\alpha} p'_{\alpha}(Y) \mathbf{X}^{\alpha} + p_{\alpha}(Y) D(\mathbf{X}^{\alpha}) \\ &= qp \\ &= \sum_{\alpha} q(Y) p_{\alpha}(Y) \mathbf{X}^{\alpha}. \end{aligned}$$

By Proposition 1.1.6, for each  $\alpha$  with  $p_{\alpha}(Y) \neq 0$  the corresponding coefficient of  $\mathbf{X}^{\alpha}$  in  $D(p)$  is

$$\begin{aligned} D(p)_{\alpha} &= p'_{\alpha}(Y) + p_{\alpha}(Y) \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii} \\ &\quad + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y) Y_{i\ell}. \end{aligned}$$

Since  $D(p) = qp$ , it must be  $D(p)_{\alpha} = q(Y) p_{\alpha}(Y)$  or, equivalently,

$$\begin{aligned} q(Y) p_{\alpha}(Y) &= p'_{\alpha}(Y) + p_{\alpha}(Y) \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii} \\ &\quad + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y) Y_{i\ell}. \end{aligned}$$

This means that for each  $\alpha$ , the coefficient  $p_{\alpha}(Y)$  of  $\mathbf{X}^{\alpha}$  in  $p$  divides the expression

$$p'_{\alpha}(Y) + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y) Y_{i\ell}.$$

Thus, for each  $\alpha$ , there is  $u_\alpha(Y)$  such that

$$p_\alpha(Y)u_\alpha(Y) = p'_\alpha(Y) + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y)Y_{i\ell}.$$

As in the proof of Proposition 1.1.2, order the triples  $\{ij, k\}$ ,  $i, j, k \in \mathbb{N}$ , with the lexicographical order. Let  $\{mn, t\}$  be the largest subindex such that  $Y_{mn,t}$  occurs in  $p$ . We have  $D(Y_{mn,t}) = Y_{mn,t+1}$  and  $\{mn, t+1\} > \{mn, t\}$ .

Now, for each  $\alpha$  such that  $Y_{mn,t}$  occurs in  $p_\alpha(Y)$  we have that  $Y_{mn,t+1}$  will occur in  $p'_\alpha(Y)$  but not in  $p_\alpha(Y)$  or in

$$\sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}}(Y)Y_{i\ell}$$

by the choice of  $\{mn, t\}$ . Therefore, it must occur in  $p_\alpha(Y)u_\alpha(Y)$ . Let

$$p_\alpha(Y) = \sum a_\beta Y_{11}^{\beta_{11}} Y_{12}^{\beta_{12}} \cdots Y_{mn,t}^{\beta_{mn,t}}$$

then

$$\begin{aligned} p'_\alpha(Y) &= \sum a'_\beta Y_{11}^{\beta_{11}} \cdots Y_{mn,t}^{\beta_{mn,t}} \\ &\quad + \sum a_\beta \beta_{11} Y_{11}^{\beta_{11}-1} Y_{11,1}^{\beta_{11,1}+1} \cdots Y_{mn,t}^{\beta_{mn,t}} + \cdots \\ &\quad + \sum a_\beta \beta_{mn,t} Y_{11}^{\beta_{11}} \cdots Y_{mn,t}^{\beta_{mn,t}-1} Y_{mn,t+1}. \end{aligned}$$

So  $Y_{mn,t+1}$  occurs in  $p'_\alpha(Y)$  only in

$$\begin{aligned} &\sum a_\beta \beta_{mn,t} Y_{11}^{\beta_{11}} \cdots Y_{mn,t}^{\beta_{mn,t}-1} Y_{mn,t+1} \\ &= \left( \sum a_\beta \beta_{mn,t} Y_{11}^{\beta_{11}} \cdots Y_{mn,t}^{\beta_{mn,t}-1} \right) Y_{mn,t+1} \\ &= v(Y)Y_{mn,t+1}. \end{aligned}$$

Since  $Y_{mn,t+1}$  occurs in  $p_\alpha(Y)u_\alpha(Y)$  and not in  $p_\alpha(Y)$  it must occur in  $u_\alpha(Y)$ . Let  $u_{\alpha,t+1}(Y)$  be the coefficient of  $Y_{mn,t+1}$  in  $u_\alpha(Y)$ . Then it has to be

$$p_\alpha(Y)u_{\alpha,t+1}(Y)Y_{mn,t+1} = v(Y)Y_{mn,t+1}.$$

The above equation implies that  $p_\alpha(Y)$  divides  $v(Y)$ . But this is impossible since the total degree of  $v(Y)$  is strictly less than the total degree of  $p_\alpha(Y)$ . This contradiction yields the result.  $\square$

**Lemma 1.1.8.** *Let  $p \in F[X_{ij}]$  and suppose that there is  $q \in F\{Y_{ij}\}$  such that  $D(p) = qp$ . Then  $q$  is a linear polynomial in the  $Y_{ij}$ . If  $\beta = (\beta_{ij})$  is such that  $\mathbf{X}^\beta$  occurs in  $p$ , then for  $1 \leq i \leq n$  the coefficient of  $Y_{ii}$  in  $q$  is  $\sum_{j=1}^n \beta_{ij}$ . In particular, the sums  $\sum_{j=1}^n \beta_{ij}$ , for  $1 \leq i \leq n$ , are independent of the choice of  $\mathbf{X}^\beta$ .*

*Proof.* We have  $p = \sum a_\beta \mathbf{X}^\beta$ , with  $a_\beta \in F$ .

Thus,

$$\begin{aligned} D(p) &= \sum a'_\beta \mathbf{X}^\beta + a_\beta D(\mathbf{X}^\beta) \\ &= qp \\ &= \sum q(Y) a_\beta \mathbf{X}^\beta. \end{aligned}$$

By Proposition 1.1.6, the coefficient of  $\mathbf{X}^\beta$  in  $D(p)$  is

$$a'_\beta + a_\beta \sum_{i=1}^n \sum_{j=1}^n \beta_{ij} Y_{ii} + \sum_{i=1}^n \sum_{j=1}^n (\beta_{ij} + 1) \sum_{\ell \neq i} a_{\beta_{ij,\ell}} Y_{i\ell}.$$

Hence, it must be

$$q(Y) a_\beta = a'_\beta + a_\beta \left( \sum_{i=1}^n \sum_{j=1}^n \beta_{ij} Y_{ii} + \sum_{i=1}^n \sum_{j=1}^n (\beta_{ij} + 1) \sum_{\ell \neq i} a_{\beta_{ij,\ell}} Y_{i\ell} \right).$$

From this,

$$q(Y) = \frac{a'_\beta}{a_\beta} + \sum_{i=1}^n \sum_{j=1}^n \beta_{ij} Y_{ii} + \sum_{i=1}^n \sum_{j=1}^n (\beta_{ij} + 1) \sum_{\ell \neq i} \frac{a_{\beta_{ij,\ell}}}{a_\beta} Y_{i\ell}.$$

The coefficient of  $Y_{ii}$  in the above expression is  $\sum_{j=1}^n \beta_{ij}$ , for  $1 \leq i \leq n$ . Since this expression for  $q$  is valid for any index  $\beta$ , the “in particular” part follows immediately.  $\square$

**Corollary 1.1.9.** *Let  $p$  be as in Lemma 1.1.8. Let  $\mathbf{X}^\alpha$  be the leading power product of  $p$ . Let  $\mathbf{X}^\beta$  be any power product with non-zero coefficient in  $p$ . Then  $\sum_{j=1}^n \beta_{ij} = \sum_{j=1}^n \alpha_{ij}$ , for  $1 \leq i \leq n$ . Thus  $p$  is homogeneous of degree  $\sum_{j=1}^n \sum_{i=1}^n \alpha_{ij}$ .*

*Proof.* This is an immediate consequence of the “in particular” part in Lemma 1.1.8.  $\square$

**Corollary 1.1.10.** *Let  $p \in F[X_{ij}]$  and suppose that  $D(p) = qp$ , for some  $q \in F\{Y_{ij}\}$ . Let  $\mathbf{X}^\alpha$  be the leading power product of  $p$ , and let  $\ell \in F$  be its coefficient. Then*

$$q = \frac{\ell'}{\ell} + \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii}.$$

*Proof.* By Proposition 1.1.6 and since  $D(p) = qp$ , the coefficient of  $\mathbf{X}^\alpha$  in  $D(p)$  is

$$(1) \quad \ell q = \ell' + \ell \left( \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii} + \sum_{i=1}^n \sum_{j=1}^n (\alpha_{ij} + 1) \sum_{\ell \neq i} p_{\alpha_{ij,\ell}} Y_{i\ell} \right).$$

The  $p_{\alpha_{ij,k}}$  are the coefficients of the power products  $\mathbf{X}^{\alpha_{ij,k}}$  in  $p$ , with  $\alpha_{ij,k} \neq \alpha$ , such that  $D(\mathbf{X}^{\alpha_{ij,k}})$  contains an expression of the form  $Y_{st} \mathbf{X}^\alpha$ . By

Remark 1.1.4, these power products are

$$\mathbf{X}^{\alpha_{rs,t}} = \begin{cases} X_{11}^{\alpha_{11}} \cdots X_{rs}^{\alpha_{rs}+1} \cdots X_{ts}^{\alpha_{ts}-1} \cdots X_{nn}^{\alpha_{nn}} & \text{if } r < t \\ X_{11}^{\alpha_{11}} \cdots X_{ts}^{\alpha_{ts}-1} \cdots X_{rs}^{\alpha_{rs}+1} \cdots X_{nn}^{\alpha_{nn}} & \text{if } r > t \end{cases},$$

all of which violate Corollary 1.1.9 for  $i = r$  and  $i = t$ . Therefore it must be  $p_{\alpha_{ij,k}} = 0$ , for all  $1 \leq i, j \leq n$ ;  $k \neq i$ . But now, substituting back in (1), we see that

$$\ell q = \ell' + \ell \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii}.$$

Hence,

$$q = \frac{\ell'}{\ell} + \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij} Y_{ii}.$$

□

Our next step in order to show that the Darboux polynomials  $p \in R$  have the desired form will be to show that such a  $p$  is not reduced with respect to  $\det[X_{ij}]$ . For that we will show that the leading power product of  $p$  is a power of the leading power product of  $\det[X_{ij}]$ . First, we have

**Lemma 1.1.11.** *Let  $p \in F[X_{ij}]$  be such that  $D(p) = qp$ ,  $q \in F\{Y_{ij}\}$ . Let  $\mathbf{X}^\alpha$  be its leading power product. Then  $\alpha_{ij} = 0$  for  $j \neq n - i + 1$  and  $\alpha_{i,n-i+1} > 0$ ,  $1 \leq i \leq n$ . That is,  $\mathbf{X}^\alpha = X_{1n}^{\alpha_{1n}} X_{2,n-1}^{\alpha_{2,n-1}} \cdots X_{n1}^{\alpha_{n1}}$ .*

*Proof.* To prove that  $\alpha_{ij} = 0$  for  $j \neq n - i + 1$  we first show that  $\alpha_{ij} = 0$  for  $j > n - k + 1$ ,  $i \geq k$ ,  $2 \leq k \leq n$ . Indeed, for  $k = 2$  we have  $j > n - 1$ , so  $j = n$  and

$$D(\mathbf{X}^\alpha) = \alpha_{nn} \sum_{k=1}^{n-1} Y_{nk} X_{11}^{\alpha_{11}} \cdots X_{kn}^{\alpha_{kn}+1} \cdots X_{nn}^{\alpha_{nn}-1} + \cdots$$

Since  $q$  has no  $Y_{ij}$  with  $i \neq j$ , each term in  $D(\mathbf{X}^\alpha)$  containing such a  $Y_{ij}$  must be cancelled. In particular we need to cancel the terms containing

$$Y_{nj} X_{11}^{\alpha_{11}} \cdots X_{jn}^{\alpha_{jn}+1} \cdots X_{nn}^{\alpha_{nn}-1}$$

for  $1 \leq j \leq n - 1$  above. For that we can only use the derivatives of power products of the form

$$\mathbf{X}^{\alpha_{nl,j}} = X_{11}^{\alpha_{11}} \cdots X_{j\ell}^{\alpha_{j\ell}-1} \cdots X_{jn}^{\alpha_{jn}+1} \cdots X_{n1}^{\alpha_{n1}} \cdots X_{n\ell}^{\alpha_{n\ell}+1} \cdots X_{nn}^{\alpha_{nn}-1}, \quad \ell < n.$$

But these are all strictly greater than  $\mathbf{X}^\alpha$  (the leading power product of  $p$ ), and they may not occur in  $p$ . As a consequence, it has to be  $\alpha_{nn} = 0$ . Now let  $k > 2$  be such that  $\alpha_{in} = 0$  for  $i \geq k$ . Then

$$\mathbf{X}^\alpha = X_{11}^{\alpha_{11}} \cdots X_{k-1,n}^{\alpha_{k-1,n}} \cdots X_{k,n-1}^{\alpha_{k,n-1}} X_{k+1,1}^{\alpha_{k+1,1}} \cdots X_{k+1,n-1}^{\alpha_{k+1,n-1}} \cdots X_{n,n-1}^{\alpha_{n,n-1}}$$

and

$$D(\mathbf{X}^\alpha) = \alpha_{k-1,n} \left( \sum_{i < k-1} Y_{k-1,i} X_{11}^{\alpha_{11}} \cdots X_{in}^{\alpha_{in}+1} \cdots X_{k-1,n}^{\alpha_{k-1,n}-1} \cdots X_{n,n-1}^{\alpha_{n,n-1}} \right. \\ \left. + \sum_{i > k-1} Y_{k-1,i} X_{11}^{\alpha_{11}} \cdots X_{k-1,n}^{\alpha_{k-1,n}-1} \cdots X_{in}^{\alpha_{in}+1} \cdots X_{n,n-1}^{\alpha_{n,n-1}} \right) + \dots$$

Likewise, we need to cancel all the terms in  $D(\mathbf{X}^\alpha)$  that contain  $Y_{k-1,i}$ , for  $i \neq k-1$ . In particular, we need to cancel

$$Y_{k-1,i} X_{11}^{\alpha_{11}} \cdots X_{in}^{\alpha_{in}+1} \cdots X_{k-1,n}^{\alpha_{k-1,n}-1} \cdots X_{n,n-1}^{\alpha_{n,n-1}},$$

for  $i < k-1$ . For that we can only use the power products of the form

$$\mathbf{X}^{\alpha_{k-1,\ell,i}} = \\ X_{11}^{\alpha_{11}} \cdots X_{i\ell}^{\alpha_{i\ell}-1} \cdots X_{in}^{\alpha_{in}+1} \cdots X_{k-1,\ell}^{\alpha_{k-1,\ell}+1} \cdots X_{k-1,n}^{\alpha_{k-1,n}-1} \cdots X_{n,n-1}^{\alpha_{n,n-1}},$$

for  $i < k-1$ .

But all of them are strictly greater than  $\mathbf{X}^\alpha$  and cannot occur in  $p$ . Thus, it has to be  $\alpha_{k-1,n} = 0$ . Since this argument is valid for any  $k > 2$ , it follows that  $\alpha_{kn} = 0$ , for  $2 \leq k \leq n$ . This makes the statement that  $\alpha_{ij} = 0$  for  $j > n - k + 1$ ,  $i \geq k$ , true for  $k = 2$ .

Now assume that  $k$  is such that  $\alpha_{ij} = 0$  for  $j > n - k + 1$ ,  $i \geq k$ . So

$$\mathbf{X}^\alpha = \\ X_{11}^{\alpha_{11}} \cdots X_{1n}^{\alpha_{1n}} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}} X_{k+1,1}^{\alpha_{k+1,1}} \cdots X_{k+1,n-k+1}^{\alpha_{k+1,n-k+1}} \cdots X_{n,n-k+1}^{\alpha_{n,n-k+1}}$$

and for  $i > k$

$$\alpha_{i,n-k+1} Y_{ij} X_{11}^{\alpha_{11}} \cdots X_{1n}^{\alpha_{1n}} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}+1} \cdots X_{i,n-k+1}^{\alpha_{i,n-k+1}-1} \cdots X_{n,n-k+1}^{\alpha_{n,n-k+1}}$$

occurs in  $D(\mathbf{X}^\alpha)$ . Thus we need to cancel it. For that we can only use the derivatives of power products of the form

$$\mathbf{X}^{\alpha_{i,j,k}} = \\ X_{11}^{\alpha_{11}} \cdots X_{kj}^{\alpha_{kj}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}+1} \cdots X_{ij}^{\alpha_{ij}+1} \cdots X_{i,n-k+1}^{\alpha_{i,n-k+1}-1} \cdots X_{n,n-k+1}^{\alpha_{n,n-k+1}}$$

with  $j < n - k + 1$  since  $\alpha_{kj} = 0$  for all  $j > n - k + 1$  by hypothesis. But all such power products are strictly greater than  $\mathbf{X}^\alpha$  and therefore they cannot occur in  $p$ . This forces  $\alpha_{i,n-k+1} = 0$  for  $i > k$ . We can repeat this process until  $k = n$  and get  $\alpha_{ij} = 0$  for all  $j > n - k + 1$ ,  $i \geq k$ ,  $2 \leq k \leq n$ , that is,

$$\mathbf{X}^\alpha = \\ X_{11}^{\alpha_{11}} \cdots X_{1n}^{\alpha_{1n}} X_{21}^{\alpha_{21}} \cdots X_{2,n-1}^{\alpha_{2,n-1}} X_{31}^{\alpha_{31}} \cdots X_{n-1,2}^{\alpha_{n-1,2}} X_{n1}^{\alpha_{n1}}.$$

Now we show that  $\alpha_{ij} = 0$  for  $j < n - k + 1$ ,  $1 \leq k \leq n - 1$ ,  $i \leq k$ . The process is analogous to what we just did. First we show that  $\alpha_{i1} = 0$  for  $i < n$ . Indeed, for each  $i$  we have for  $\ell > i$  that

$$\alpha_{i1} Y_{i\ell} X_{11}^{\alpha_{11}} \cdots X_{i1}^{\alpha_{i1}-1} \cdots X_{\ell 1}^{\alpha_{\ell 1}+1} \cdots X_{n1}^{\alpha_{n1}}$$

occurs in  $D(\mathbf{X}^\alpha)$ . So, in order to cancel it, we need to use the derivatives of power products of the form

$$\mathbf{X}^{\alpha_{ij,\ell}} = X_{11}^{\alpha_{11}} \dots X_{i1}^{\alpha_{i1}-1} \dots X_{ij}^{\alpha_{ij}+1} \dots X_{\ell 1}^{\alpha_{\ell 1}+1} \dots X_{\ell j}^{\alpha_{\ell j}-1} \dots X_{n1}^{\alpha_{n1}}$$

with  $j > 1$ , all of which are strictly greater than  $\mathbf{X}^\alpha$  if  $\ell < n$ , and for  $\ell = n$  we cannot simply have one of those since  $\alpha_{nj} = 0$  for  $j \neq 1$ . Thus such power products cannot occur in  $p$  and it has to be  $\alpha_{i1} = 0$  for  $i < n$ .

Let  $k \leq n-1$  be such that  $\alpha_{ij} = 0$  for  $j < n-k+1$ ,  $i \leq k$ . We have

$$\mathbf{X}^\alpha = X_{1,n-k+1}^{\alpha_{1,n-k+1}} \dots X_{1n}^{\alpha_{1n}} \dots X_{k,n-k+1}^{\alpha_{k,n-k+1}} \dots X_{n1}^{\alpha_{n1}}$$

and for all  $i < k$ ,  $\ell > i$ , we have that

$$\alpha_{i,n-k+1} Y_{i\ell} X_{1,n-k+1}^{\alpha_{1,n-k+1}} \dots X_{i,n-k+1}^{\alpha_{i,n-k+1}-1} \dots X_{\ell,n-k+1}^{\alpha_{\ell,n-k+1}+1} \dots X_{n1}^{\alpha_{n1}}$$

occurs in  $D(\mathbf{X}^\alpha)$  and in order to cancel it we only have the derivatives of power products of the form

$$\mathbf{X}^{\alpha_{ij,\ell}} = X_{1,n-k+1}^{\alpha_{1,n-k+1}} \dots X_{i,n-k+1}^{\alpha_{i,n-k+1}-1} \dots X_{ij}^{\alpha_{ij}+1} \dots X_{\ell,n-k+1}^{\alpha_{\ell,n-k+1}+1} \dots X_{\ell j}^{\alpha_{\ell j}-1} \dots X_{n1}^{\alpha_{n1}}$$

with  $j > n-k+1$  since  $\alpha_{ij} = 0$  for  $i \leq k$ ,  $j < n-k+1$ .

For  $\ell < k$ , all these power products are strictly greater than  $\mathbf{X}^\alpha$  and therefore they cannot occur in  $p$ . For  $\ell \geq k$  we cannot simply have such power products since for  $\ell \geq k$ ,  $\alpha_{\ell j} = 0$  if  $j > n-k+1$ . Thus it has to be  $\alpha_{i,n-k+1} = 0$  for  $i \leq k-1$ .

We can repeat this process until  $k = n-1$  and get  $\alpha_{ij} = 0$ ,  $j < n-k+1$ ,  $i \leq k$ ,  $1 \leq k \leq n-1$ . This completes the proof of the first part of the lemma.

To prove that  $\alpha_{i,n-i+1} \neq 0$ , for all  $1 \leq i \leq n$ , suppose that there is  $i$  such that  $\alpha_{i,n-i+1} = 0$  and let  $j \neq i$  be such that  $\alpha_{j,n-j+1} \neq 0$ . Then  $D(\mathbf{X}^\alpha)$  will contain

$$\alpha_{j,n-j+1} Y_{ji} X_{1n}^{\alpha_{1n}} \dots X_{j,n-j+1}^{\alpha_{j,n-j+1}-1} \dots X_{i,n-j+1} \dots X_{n1}^{\alpha_{n1}} + \dots \quad \text{if } i > j$$

or

$$\alpha_{j,n-j+1} Y_{ji} X_{1n}^{\alpha_{1n}} \dots X_{i,n-j+1} \dots X_{j,n-j+1}^{\alpha_{j,n-j+1}-1} \dots X_{n1}^{\alpha_{n1}} + \dots \quad \text{if } i < j.$$

As noted above, since  $q$  does not contain any  $Y_{ij}$  with  $i \neq j$ , we need to cancel the terms in  $D(p)$  involving either of the above. But that is impossible since  $\alpha_{ij} = 0$  for all  $j$  and by Corollary 1.1.9 all the power products

$$X_{11}^{\beta_{11}} \dots X_{ij}^{\beta_{ij}} \dots X_{nn}^{\beta_{nn}}$$

in  $p$  must have  $\beta_{ij} = 0$  for  $j = 1, \dots, n$ . In particular, we cannot have in  $p$  power products of the form  $\mathbf{X}^{\alpha_{j,n-j+1,i}}$  as in Remark 1.1.4  $\square$

Next we show that the exponents  $\alpha_{st}$  of the  $X_{st}$  in  $\mathbf{X}^\alpha$ , the leading power product of  $p$ , are all equal:

**Lemma 1.1.12.** *Let  $p \in F[X_{ij}]$  be such that  $D(p) = qp$ ,  $q \in F\{Y_{ij}\}$ . Let*

$$\mathbf{X}^\alpha = X_{1n}^{\alpha_{1n}} X_{2,n-1}^{\alpha_{2,n-1}} \cdots X_{n1}^{\alpha_{n1}}$$

*be its leading power product. Then  $\alpha_{i,n-i+1} = \alpha_{1n}$ , for  $i > 1$ , that is, if  $a = \alpha_{1n}$ , then*

$$\mathbf{X}^\alpha = (X_{1n} X_{2,n-1} \cdots X_{n1})^a.$$

*Proof.* Let  $\ell$  be the coefficient of  $\mathbf{X}^\alpha$  in  $p$ . We have

$$\begin{aligned} D(\ell X_{1n}^{\alpha_{1n}} X_{2,n-1}^{\alpha_{2,n-1}} \cdots X_{n1}^{\alpha_{n1}}) &= \\ & \left( \sum_{i=1}^n \alpha_{i,n-i+1} \ell Y_{ii} \right) X_{1n}^{\alpha_{1n}} X_{2,n-1}^{\alpha_{2,n-1}} \cdots X_{n1}^{\alpha_{n1}} \\ & + \alpha_{1n} \ell \sum_{k \neq 1} Y_{1k} X_{1n}^{\alpha_{1n}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}} \cdots X_{kn} \cdots X_{n1}^{\alpha_{n1}} \\ & + \ell \sum_{1 < i} \alpha_{i,n-i+1} \sum_{k > i} Y_{ij} X_{1n}^{\alpha_{1n}} \cdots X_{i,n-i+1}^{\alpha_{i,n-i+1}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}} \cdots X_{k,n-i+1} \cdots X_{n1}^{\alpha_{n1}} \\ & + \ell \sum_{1 < i} \alpha_{i,n-i+1} \sum_{k > i} Y_{ij} X_{1n}^{\alpha_{1n}} \cdots X_{k,n-i+1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}} \cdots X_{i,n-i+1}^{\alpha_{i,n-i+1}-1} \cdots X_{n1}^{\alpha_{n1}} \\ & + \ell' X_{1n}^{\alpha_{1n}} X_{2,n-1}^{\alpha_{2,n-1}} \cdots X_{n1}^{\alpha_{n1}}. \end{aligned}$$

In order to cancel

$$\alpha_{1n} \ell Y_{1k} X_{1n}^{\alpha_{1n}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}} \cdots X_{kn} \cdots X_{n1}^{\alpha_{n1}}, \quad k \neq 1,$$

above, we can only use the derivatives of the power product

$$\begin{aligned} \mathbf{X}^{\alpha_{1,n-k+1,k}} &= \\ X_{1,n-k+1} \cdots X_{1n}^{\alpha_{1n}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}-1} \cdots X_{kn} \cdots X_{n1}^{\alpha_{n1}}, \end{aligned}$$

since for  $j \neq n-k+1$  we have  $\alpha_{kj} = 0$ .

Let  $a_{\alpha_{1,n-k+1,k}}$  be the coefficient of  $X^{\alpha_{1,n-k+1,k}}$  in  $p$ . Then

$$(2) \quad a_{\alpha_{1,n-k+1,k}} = -\ell \alpha_{1n}$$

On the other hand, in order to cancel

$$\alpha_{k,n-k+1} \ell Y_{k1} X_{1,n-k+1} \cdots X_{1n}^{\alpha_{1n}} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}-1} \cdots X_{n1}^{\alpha_{n1}}, \quad k \neq 1$$

above, the only power product that we can use is, again,

$$\begin{aligned} \mathbf{X}^{\alpha_{k,n-k+1,k}} &= X_{1,n-k+1} \cdots X_{1n}^{\alpha_{1n}-1} \cdots X_{k,n-k+1}^{\alpha_{k,n-k+1}-1} \cdots X_{kn} \cdots X_{n1}^{\alpha_{n1}} \\ &= X^{\alpha_{1,n-k+1,k}}, \end{aligned}$$

since  $\alpha_{1j} = 0$  for  $j \neq n$ . Thus it must be

$$(3) \quad a_{\alpha_{1,n-k+1,k}} = -\ell \alpha_{k,n-k+1}$$

as well.

From (2) and (3) it follows that, for  $k \neq 1$ ,  $\alpha_{1n} = \alpha_{k,n-k+1}$ .  $\square$

As a consequence of the above results we obtain the following expression for  $q$ :

**Corollary 1.1.13.** *Let  $p \in F[X_{ij}]$  and suppose that  $D(p) = qp$ ,  $q \in F\{Y_{ij}\}$ . Let  $\mathbf{X}^\alpha$  be the leading power product of  $p$ . Let  $a \in \mathbb{N}$  be such that*

$$\mathbf{X}^\alpha = (X_{1n}X_{2,n-1} \cdots X_{n1})^a$$

and let  $\ell \in F$  be the coefficient of  $\mathbf{X}^\alpha$  in  $p$ . Then

$$q = \frac{\ell'}{\ell} + a \sum_{i=1}^n Y_{ii}.$$

*Proof.* This is a consequence of Corollary 1.1.10 and Lemma 1.1.12.  $\square$

**Corollary 1.1.14.** *Let  $p$  be as in Corollary 1.1.13. Then  $p$  is homogeneous of degree  $na$ .*

*Proof.* This is a consequence of Corollary 1.1.8 and Lemma 1.1.12  $\square$

Lemma 1.1.12 implies that  $p$  is not reduced with respect to  $\det[X_{ij}]$ . Since this is a key point in the proof of our main result we restate it as the following

**Theorem 1.1.15.** *Let  $p \in F[X_{ij}]$  be such that  $D(p) = qp$ ,  $q \in F\{Y_{ij}\}$ . Let  $\mathbf{X}^\alpha$  be its leading power product. Then*

$$\mathbf{X}^\alpha = (X_{1n}X_{2,n-1} \cdots X_{n1})^a = \text{lp}(\det[X_{ij}])^a.$$

Thus  $p$  is not reduced with respect to  $\det[X_{ij}]$ .

**Note.** If  $f$  is a polynomial,  $\text{lp}(f)$  denotes its leading power product with respect to a given order.

*Proof.* This is just a restatement of Lemma 1.1.12.  $\square$

**Remark 1.1.16.** *Let  $p_1, p_2 \in F[X_{ij}]$  be two polynomials such that  $\text{lp}(p_1) = X^\alpha = \text{lp}(p_2)$ . Then we can write  $p_1 = f p_2 + r$  where  $f \in F$  and  $r$  is reduced with respect to  $p_2$ . Indeed, since  $\text{lp}(p_1) = \text{lp}(p_2)$ , we have that  $\text{lp}(p_2)$  divides  $\text{lp}(p_1)$ . So  $p_1$  is not reduced with respect to  $p_2$ . We may apply the Multivariable Division Algorithm (see [1]) to  $p_1$  and  $p_2$ , to get  $f, r \in F[X_{ij}]$ , such that  $p_1 = f p_2 + r$ , with  $r$  reduced with respect to  $p_2$  and  $\text{lp}(p_1) = \text{lp}(f)\text{lp}(p_2)$ . The last equation implies that  $\text{lp}(f) = 1$ . Hence,  $f \in F$ .*

We are now ready to prove our main result on the form of the Darboux polynomials in  $R$ :

**Theorem 1.1.17.** *Let  $p \in F[X_{ij}]$  and  $q \in F\{Y_{ij}\}$  be polynomials in  $R$  that satisfy the Darboux condition  $D(p) = qp$ . Then there is  $a \in \mathbb{N}$  and  $\ell \in F$  such that*

$$p = \ell \det[X_{ij}]^a$$

and

$$q = \frac{\ell'}{\ell} + a \sum_{i=1}^n Y_{ii}.$$

*Proof.* Let  $q_1 = \sum_{i=1}^n a Y_{ii}$ , so that,

$$D(\det[X_{ij}]^a) = q_1 \det[X_{ij}]^a = \left(q - \frac{\ell'}{\ell}\right) \det[X_{ij}]^a.$$

By Remark 1.1.16 we can write  $p = \ell \det[X_{ij}]^a + r$ , with  $r$  reduced with respect to  $\det[X_{ij}]^a$ . Now,

$$\begin{aligned} D(p) &= D(\ell \det[X_{ij}]^a) + D(r) \\ &= \ell' \det[X_{ij}]^a + \ell \left(q - \frac{\ell'}{\ell}\right) \det[X_{ij}]^a + D(r) \\ &= \ell' \det[X_{ij}]^a + q\ell \det[X_{ij}]^a - \ell' \det[X_{ij}]^a + D(r) \\ &= q\ell \det[X_{ij}]^a + D(r). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} D(p) &= qp \\ &= q\ell \det[X_{ij}]^a + qr. \end{aligned}$$

Therefore, it has to be  $D(r) = qr$ . But  $r$  is reduced with respect to  $\det[X_{ij}]^a$ . It follows, by Theorem 1.1.15, that  $r = 0$ . The statement about the form of  $q$  is just the content of Corollary 1.1.13.  $\square$

**1.2. Principal differential ideals in  $\mathbf{F}\{\mathbf{Y}_{ij}\}[\mathbf{X}_{ij}]$ .** As mentioned in the introduction, if we pass to the quotient field  $F\langle Y_{ij} \rangle$  of  $F\{Y_{ij}\}$  and localize  $F\langle Y_{ij} \rangle[X_{ij}]$  at  $\det[X_{ij}]$ , we get the coordinate ring of  $\mathrm{GL}_n$  over  $F\langle Y_{ij} \rangle$ . The derivation  $D$  on  $F\langle Y_{ij} \rangle[X_{ij}]$  defined above can then be seen as a ‘‘general’’ element of  $\mathrm{Lie}(\mathrm{GL}_n)$ . In particular,  $D$  is a linear combination of the basis of  $\mathrm{Lie}(\mathrm{GL}_n)$  consisting of the derivations  $D_{E(ij)}$  given by multiplication by the matrix  $E(ij)$ , with 1 in position  $(i,j)$  and zero elsewhere and the coefficient of  $D_{E(ij)}$  in  $D$  is  $Y_{ij}$ .

We will show next that the result in Theorem 1.1.17 is true for any other such element of  $\mathrm{Lie}(\mathrm{GL}_n)$ . That is, the result does not depend on the particular basis of  $\mathrm{Lie}(\mathrm{GL}_n)$  used.

**Theorem 1.2.1.** *Let  $\mathcal{D}_{st}$ ,  $1 \leq s, t \leq n$ , be any basis of  $\mathrm{Lie}(\mathrm{GL}_n)$ . Define a derivation in the ring  $R = F\{Y_{ij}\}[X_{ij}]$  by  $\mathcal{D} = \sum Y_{st} \mathcal{D}_{st}$ . Let  $p$  and  $q$  be polynomials in  $R$  that satisfy the Darboux condition  $\mathcal{D}(p) = qp$ . Then there is  $a \in \mathbb{N}$  and  $\ell \in F$  such that  $p = \ell \det[X_{ij}]^a$  and  $q = \frac{\ell'}{\ell} + a \sum_{i=1}^n Y_{ii}$ .*

*Proof.* Since  $\{D_{E(ij)} \mid 1 \leq i, j \leq n\}$  is a basis of  $\mathrm{Lie}(\mathrm{GL}_n(C))$  we have

$$\mathcal{D}_{st} = \sum c_{st,ij} D_{E(ij)},$$

with  $c_{st,ij} \in C$ . Thus,

$$\begin{aligned}
\mathcal{D} &= \sum_{s,t} Y_{st} \mathcal{D}_{st} \\
&= \sum_{s,t} Y_{st} \sum_{i,j} c_{st,ij} D_{E(ij)} \\
&= \sum_{i,j} \sum_{s,t} c_{st,ij} Y_{st} D_{E(ij)} \\
&= \sum_{i,j} Z_{ij} D_{E(ij)},
\end{aligned}$$

where  $Z_{ij} = \sum_{s,t} c_{st,ij} Y_{st}$ . Now,  $[c_{st,ij}]$  is a matrix of change of basis so it is invertible. Also the  $c_{st,ij}$  are constants for  $D$ , thus the map  $Z_{ij,k} \rightarrow Y_{ij,k}$  is a differential bijection. In other words, the differential rings

$$R = F\{Y_{ij}\}[X_{ij}], D$$

and

$$R' = F\{Z_{ij}\}[X_{ij}], \mathcal{D}$$

are isomorphic and therefore we can apply Theorem 1.1.17 to  $R'$ .  $\square$

**Theorem 1.2.2.** *Let  $R = F\{Y_{ij}\}[X_{ij}]$  be a differential ring with derivation obtained by restriction of a general element of  $\text{Lie}(\text{GL}_n)$  in the sense described above. Then the principal differential ideals in  $R$  are those of the form  $\mathcal{I} = (\det^a[X_{ij}])$  for  $a \in \mathbb{N}$ .*

*Proof.* This is a consequence of Theorems 1.1.17, 1.2.1 and of the observation that Darboux polynomials correspond to principal differential ideals in  $R$ .  $\square$

## 2. A GENERIC PICARD-VESSIOT EXTENSION FOR $\text{GL}_n(C)$

**2.1. Preliminaries on Differential Galois Theory.** As before,  $F$  is a differential field with algebraically closed field of constants  $C$ . If  $E \supseteq F$  is a differential field extension then the group of differential automorphisms of  $E$  over  $F$  is denoted by  $G(E/F)$ .

If  $G$  is a linear algebraic group over  $C$  and  $K$  is an overfield of  $C$  we denote by  $G_K$  the group obtained from  $G$  by extending scalars from  $C$  to  $K$ .

We will show that  $F\langle Y_{ij} \rangle(X_{ij})$  is a generic Picard-Vessiot extension of  $F$  for the group  $\text{GL}_n(C)$ . Notice that  $F\langle Y_{ij} \rangle(X_{ij})$  is the function field of  $G_K$  with  $G = \text{GL}_n(C)$  and  $K = F\langle Y_{ij} \rangle$ . The following two results ([19], Theorem 5.12 and Corollary 5.29) will be used:

**Theorem 2.1.1** (Kolchin Structure Theorem). *Let  $E \supseteq F$  be a Picard-Vessiot extension, let  $G \leq G(E/F)$  be a Zariski closed subgroup and let  $T$  be the set of all  $f$  in  $E$  that satisfy a linear homogeneous differential equation over  $K = E^G$ . Then  $T$  is a finitely generated  $G$ -stable differential*

$K$ -algebra with quotient field  $E$ , and if  $\overline{K}$  denotes the algebraic closure of  $K$ , then there is a  $G$ -algebra isomorphism

$$\overline{K} \otimes_K T \rightarrow \overline{K} \otimes_C C[G].$$

Note that  $C[G]$  denotes the affine coordinate ring of  $G$  and that the target of the above isomorphism is the affine coordinate ring of the group  $G_{\overline{K}}$  obtained from  $G$  by extension of scalars from  $C$  to  $\overline{K}$ .

**Theorem 2.1.2.** *Let  $E \supseteq F$  be a Picard-Vessiot extension, let  $G \leq G(E/F)$  be a Zariski closed subgroup with  $E^G = F$ . Let  $\overline{F}$  be an algebraic closure of  $F$ , and suppose the Galois cohomology  $H^1(\overline{F}/F, G(\overline{F}))$  is a singleton. Let  $T(E/F)$  be the set of all  $f$  in  $E$  that satisfy a linear homogeneous differential equation over  $F$ . Then there are  $F$ - and  $G$ -isomorphisms  $T(E/F) \rightarrow F[G_F]$  and  $E \rightarrow F(G_F)$ . In particular, this holds if  $G$  is unipotent or solvable, or if  $G = \mathrm{GL}_n(C)$  or if  $G = \mathrm{SL}_n$ .*

The following characterization of Picard-Vessiot extension (see [19], Proposition 3.9) will be employed:

**Theorem 2.1.3.** *Let  $E \supseteq F$  be a differential field extension. Then  $E$  is a Picard-Vessiot extension if and only if:*

1.  $E = F\langle V \rangle$ , where  $V \subset E$  is a finite-dimensional vector space over  $C$ ;
2. There is a group  $G$  of differential automorphisms of  $E$  with  $G(V) \supseteq V$  and  $E^G = F$ ;
3.  $E \supset F$  has no new constants.

In particular, if the above conditions hold and if  $\{y_1, \dots, y_n\}$  is a  $C$ -basis of  $V$ , then  $E$  is a Picard-Vessiot extension of  $F$  for the linear homogeneous differential operator

$$L(Y) = \frac{w(Y, y_1, \dots, y_n)}{w(y_1, \dots, y_n)}$$

where  $w(-)$  denotes the wronskian determinant and  $L^{-1}(0) = V$ .

For the base field  $F\langle Y_{ij} \rangle$  and group  $G = \mathrm{GL}_n(C)$  we first show that  $F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  is a Picard-Vessiot extension with differential Galois group  $\mathrm{GL}_n(C)$ . To that end, we only need to show that  $F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  is a no-new-constant extension. Conditions 1. and 2. in Theorem 2.1.3 are then easily verified with  $V$  the  $C$ -span of the  $X_{ij}$  and  $G = \mathrm{GL}_n(C)$ .

**2.2. Darboux polynomials and the constants of  $F\langle \mathbf{Y}_{ij} \rangle(X_{ij})$ .** We will show that the field of constants  $\mathcal{C}$  of  $F\langle Y_{ij} \rangle(X_{ij})$  coincides with the field of constants  $C$  of  $F$ . We first show (Corollary 2.2.2) that this can be reduced to proving that the only *Darboux polynomials* in  $R$  are, up to a scalar multiple in  $F$ , powers of  $\det[X_{ij}]$ .

The following basic proposition (proven in [34] for  $A$  as in Definition 1.1.1) characterizes new constants for the extension  $F\langle Y_{ij} \rangle(X_{ij}) \supset F$  in terms of Darboux polynomials:

**Proposition 2.2.1.** *Let  $p_1, p_2 \in R = F\langle Y_{ij} \rangle[X_{ij}]$ ,  $p_1, p_2 \neq 0$ , be relatively prime. Then  $D(\frac{p_1}{p_2}) = 0$ , if and only if  $p_1$  and  $p_2$  are Darboux polynomials. Moreover, if  $q_1, q_2 \in R$  are such that  $D(p_1) = q_1 p_1$  and  $D(p_2) = q_2 p_2$ , then  $q_1 = q_2$ .*

*Proof.* For the necessity of the condition we have

$$D\left(\frac{p_1}{p_2}\right) = \frac{D(p_1)p_2 - p_1 D(p_2)}{p_2^2} = 0,$$

thus  $D(p_1)p_2 - p_1 D(p_2) = 0$ , that is

$$(1) \quad D(p_1)p_2 = p_1 D(p_2).$$

Since  $p_1$  and  $p_2$  are relatively prime, the last equation implies that  $p_1$  divides  $D(p_1)$  and  $p_2$  divides  $D(p_2)$ .

Now, let  $q_1, q_2 \in R$  be such that  $D(p_1) = q_1 p_1$  and  $D(p_2) = q_2 p_2$ , respectively. Then it follows from (1) that

$$q_1 p_1 p_2 = q_2 p_1 p_2.$$

Hence,  $q_1 = q_2$ .

The proof of the converse is obvious.  $\square$

**Corollary 2.2.2.** *Let  $f \in F\langle Y_{ij} \rangle(X_{ij})$  be such that  $D(f) = 0$  and assume that  $f \notin F$  then there are relatively prime Darboux polynomials  $p_1, p_2 \in R$  which satisfy the Darboux condition with respect to the same  $q \in R$  (i.e.,  $D(p_i) = q p_i$ ,  $i = 1, 2$ ) and such that  $f = \frac{p_1}{p_2}$ . Therefore, if such relatively prime Darboux polynomials in  $R$  do not exist, the constants of  $F\langle Y_{ij} \rangle(X_{ij})$  coincide with the constants of  $F$ .*

*Proof.*  $F\langle Y_{ij} \rangle(X_{ij})$  is the fraction field of  $R$ .  $\square$

### 2.3. The generic extension.

**Theorem 2.3.1.**  *$F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  is a generic Picard-Vessiot extension with differential Galois group  $\mathrm{GL}_n(C)$ .*

*Proof.* First we need to show that  $F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  is a Picard-Vessiot extension with differential Galois group  $\mathrm{GL}_n(C)$ . We will use the characterization of Theorem 2.1.3. We have

1.  $F\langle Y_{ij} \rangle(X_{ij}) = F\langle Y_{ij} \rangle\langle V \rangle$ , where  $V \subset F\langle Y_{ij} \rangle(X_{ij})$  is the finite dimensional vector space over  $C$  spanned by the  $X_{ij}$ .
2. The group  $G = \mathrm{GL}_n(C)$  acts as a group of differential automorphisms of  $F\langle Y_{ij} \rangle(X_{ij})$  with  $G(V) \subseteq V$  and  $F\langle Y_{ij} \rangle(X_{ij})^G = F\langle Y_{ij} \rangle$ . This follows from the fact that  $F\langle Y_{ij} \rangle(X_{ij})$  is the function field of  $\mathrm{GL}_n(C)_{F\langle Y_{ij} \rangle}$ .
3.  $F\langle Y_{ij} \rangle(X_{ij}) \supseteq F\langle Y_{ij} \rangle$  has no new constants. This is a consequence of Proposition 2.2.1, Corollary 2.2.2 and Theorem 1.1.17.

Now, suppose that  $E \supseteq F$  is a Picard-Vessiot extension of  $F$  with differential Galois group  $\mathrm{GL}_n(C)$ . By Theorems 2.1.1 and 2.1.2, we have that in this situation  $E$  is isomorphic to  $F(X_{ij})$  (the function field of  $\mathrm{GL}_n(C)_F$ ) as

a  $\mathrm{GL}_n(C)$ -module and as an  $F$ -module. Any  $\mathrm{GL}_n(C)$  equivariant derivation  $D_E$  on  $F(X_{ij})$  extends the derivation on  $F$  in such a way that

$$D_E(X_{ij}) = \sum_{\ell=1}^n f_{i\ell} X_{\ell j}$$

with  $f_{ij} \in F$ . Since  $E \supset F$  is a Picard-Vessiot extension for  $\mathrm{GL}_n(C)$ , then so is  $C\langle f_{ij} \rangle(X_{ij}) \supset C\langle f_{ij} \rangle$ , the derivation on  $C\langle f_{ij} \rangle(X_{ij})$  being the corresponding restriction of  $D_E$ . From this Picard-Vessiot extension one can retrieve  $F(X_{ij}) \supset F$  by extension of scalars from  $C$  to  $F$ . In this way, any Picard-Vessiot extension  $E \supset F$  with differential Galois group  $\mathrm{GL}_n(C)$  can be obtained from  $F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  via the specialization  $Y_{ij} \mapsto f_{ij}$ . This means that  $F\langle Y_{ij} \rangle(X_{ij}) \supset F\langle Y_{ij} \rangle$  is a generic Picard-Vessiot extension of  $F$  for  $\mathrm{GL}_n(C)$ .  $\square$

**2.4. Specializing to a Picard-Vessiot extension of  $F$ .** In this section we give necessary and sufficient conditions for a specialization  $Y_{ij} \mapsto f_{ij}$ ,  $f_{ij} \in F$ , with  $C\langle f_{ij} \rangle(X_{ij}) \supset C\langle f_{ij} \rangle$  a Picard-Vessiot extension, to exist. We restrict ourselves to the case when  $F$  has finite transcendence degree over  $C$ .

Our goal is to find  $f_{ij} \in F$  such that the specialization (homomorphism) from  $C\{Y_{ij}\}$  to  $F$  given by  $Y_{ij} \mapsto f_{ij}$  is such that  $C\langle f_{ij} \rangle(X_{ij}) \supset C\langle f_{ij} \rangle$ , with derivation given by  $D(X_{ij}) = \sum_{\ell=1}^n f_{i\ell} X_{\ell j}$ , has no new constants. We have:

**Theorem 2.4.1.** *Let  $F = C(t_1, \dots, t_m)[z_1, \dots, z_k]$  where the  $t_i$  are algebraically independent over  $C$  and the  $z_i$  are algebraic over  $C(t_1, \dots, t_m)$ . Assume that the derivation on  $F$  has field of constants  $C$  and that it extends to  $F(X_{ij})$  so that  $D(f \otimes X_{ij}) = D(f) \otimes X_{ij} + f \otimes \sum_{\ell=1}^n f_{i\ell} X_{\ell j}$  on  $F \otimes C[X_{ij}]$ . Let  $\mathcal{C}$  be the field of constants of  $F(X_{ij})$ . Then  $\mathcal{C} = C$  if and only if the set of all the  $t_i$  and all the  $X_{ij}$  are algebraically independent over  $C$ .*

*Proof.* (Sufficiency) Suppose that  $\mathcal{C}$  properly contains  $C$ . Let  $r$  be the transcendence degree of  $\mathcal{C}$  over  $C$ . Since  $C$  is algebraically closed,  $r$  has to be at least one.

We have the tower of fields

$$C \subset \mathcal{C} \subset \mathcal{C}(X_{ij}) \subset F(X_{ij})$$

where the transcendence degree of  $\mathcal{C} \subset \mathcal{C}(X_{ij})$  is  $n^2$  and the transcendence degree of  $C \subset F(X_{ij})$  is  $n^2 + m$ . Since  $r \geq 1$  the transcendence degree  $\ell$  of  $\mathcal{C}(X_{ij}) \subset F(X_{ij})$  has to be  $\ell < m$  and therefore there is an algebraic relation among the  $t_i$  over  $\mathcal{C}(X_{ij})$ . Let  $g(X_{ij}), f_i(X_{ij}) \in \mathcal{C}[X_{ij}]$ ,  $g(X_{ij}) \neq 0$ , be such that

$$t^{\delta_s} + \frac{f_{s-1}(X_{ij})}{g(X_{ij})} t^{\delta_{s-1}} + \dots + \frac{f_0(X_{ij})}{g(X_{ij})} = 0.$$

Then

$$g(X_{ij})t^{\delta_s} + f_{s-1}(X_{ij})t^{\delta_{s-1}} + \dots + f_0(X_{ij}) = 0.$$

Since the  $f_i(X_{ij})$  and  $g(X_{ij})$  are polynomials in the  $X_{ij}$  with coefficients in  $\mathcal{C}$ , the last equation gives an algebraic relation among the  $t_i$  and the  $X_{ij}$  over  $\mathcal{C}$ .

For the necessity we only need to point out that by construction the set of all the  $t_i$  and all the  $X_{ij}$  are algebraically independent over  $\mathcal{C}$ .  $\square$

Now to check whether the set of all the  $t_i$  and all the  $X_{ij}$  are algebraically independent over  $\mathcal{C}$ , we let  $\mathbb{T}_k$ ,  $k \geq 1$ , denote the set of monomials in both the  $t_i$  and the  $X_{ij}$  of total degree less than or equal to  $k$ . Then the set of all the  $t_i$  and all the  $X_{ij}$  are algebraically independent over  $\mathcal{C}$  if and only if, for each  $k$ , the set  $\mathbb{T}_k$  is linearly independent over  $\mathcal{C}$ .

Fix a term order on the set  $\mathbb{T}$  of all monomials in both the  $t_i$  and the  $X_{ij}$  and let  $W_k$  denote the wronskian of the set  $\mathbb{T}_k$  relative to that order. Then the above condition is equivalent to the fact that  $W_k \neq 0$  for  $k \geq 1$ . Now go back to  $C\{Y_{ij}\}[X_{ij}]$  and extend scalars from  $C$  to  $F$ . Let  $W_k(Y_{ij})$  be the Wronskian of  $\mathbb{T}_k$  in  $F \otimes C\{Y_{ij}\}[X_{ij}]$ .

Then, the condition of Theorem 2.4.1 for finding a specialization  $Y_{ij} \mapsto f_{ij}$  so that  $C\langle f_{ij} \rangle(X_{ij}) \supset C\langle f_{ij} \rangle$  has no new constants can be expressed as follows:

**Theorem 2.4.2.** *There is a specialization of the  $Y_{ij}$  with no new constants if and only if there are  $f_{ij} \in F$  such that all the wronskians  $W_k(Y_{ij})$ ,  $k \geq 1$ , map to non-zero elements under  $Y_{ij} \mapsto f_{ij}$ .*

## 2.5. Specialization results for connected linear algebraic groups.

The proofs of the specialization theorems in 2.4 do not make any special use of the fact that the group under consideration is  $\mathrm{GL}_n(C)$  and can be applied to arbitrary connected linear algebraic groups as follows:

As in the previous section,  $F = C(t_1, \dots, t_m)[z_1, \dots, z_k]$  where the  $t_i$  are algebraically independent over  $C$  and the  $z_i$  are algebraic over  $C(t_1, \dots, t_m)$ . We let  $Y_1, \dots, Y_n$  denote differential indeterminates over  $F$  and  $X_1, \dots, X_n$  algebraically independent elements over  $F\langle Y_i \rangle$ .

In this section  $G$  is assumed to be a connected linear algebraic group with function field  $C(G) = C(X_i)$ .

If  $\{D_1, \dots, D_n\}$  is a basis for  $\mathrm{Lie}(G)$ ,  $D_Y = \sum_{i=1}^n Y_i D_i$  is a  $G$ -equivariant derivation on  $F\langle Y_i \rangle(X_i)$ . Let  $D = \sum_{i=1}^n f_i D_i$ ,  $f_i \in F$ , be a specialization of  $D_Y$  to a  $G$ -equivariant derivation on  $F(X_i)$  with field of constants  $\mathcal{C}$ . We have,

**Theorem 2.5.1.** *The field of constant  $\mathcal{C}$  of  $F(X_i)$  coincides with  $C$  if and only if the set of all the  $t_i$  and the  $X_i$  are algebraically independent over  $\mathcal{C}$ .*

Now, fix an order in the set  $\mathbb{T}$  of monomials in both the  $t_i$  and the  $X_i$  and let  $W_k(Y_i)$  be the wronskian (with respect to this order) of the monomials in both the  $t_i$  and the  $X_i$  of degree less than or equal to  $k$  computed in  $F \otimes C\{Y_i\}[X_i]$ . Then,

**Theorem 2.5.2.** *There is a specialization of the  $Y_i$  with no new constants if and only if there are  $f_i \in F$  such that all the wronskians  $W_k(Y_i)$ ,  $k \geq 1$ , map to non-zero elements under  $Y_i \mapsto f_i$ .*

For the proofs of Theorems 2.5.1 and 2.5.2 we only need to replace the  $X_{ij}$  with  $X_i$ , the  $Y_{ij}$  with  $Y_i$  and  $n^2$  with  $n$  in the proofs of Theorems 2.4.1 and 2.4.2.  $\square$

Observe that the proofs of Theorems 2.5.1 and 2.5.2 do not use the fact that  $C(X_i)$  is the function field of  $G$ . However, this hypothesis is used in the following theorem to show that  $F(X_i) \supset F$  is a Picard-Vessiot extension with group  $G$ .

Under the hypothesis and notation of Theorems 2.5.1 and 2.5.2 we have:

**Theorem 2.5.3.**  *$F(X_i) \supset F$  is a Picard-Vessiot extension with Galois group  $G$  if and only if the set of all the  $t_i$  and all the  $X_i$  is algebraically independent over the field of constants  $\mathcal{C}$  of  $F(X_i)$ .*

*Proof.* First assume that  $F(X_i) \supset F$  is a Picard-Vessiot extension. Then the field of constants  $\mathcal{C}$  of  $F(X_i)$  coincides with  $C$ . So we can apply Theorem 2.5.1 and get the result.

Conversely, if the set of all the  $t_i$  and all the  $X_i$  are algebraically independent over  $\mathcal{C}$ , by Theorem 2.5.1,  $F(X_i) \supset F$  is a no-new-constant extension. On the other hand,  $F(X_i)$  is obtained from  $C(X_i)$  by the extension of scalars:

$$\begin{aligned} F(X_i) &= q.f.(F \otimes_C C(X_i)) \\ &= q.f.(F \otimes_C C[G]) \end{aligned}$$

where  $C[G]$  is the coordinate ring of  $G$  and  $G$  acts on  $F \otimes_C C[G]$  fixing  $F$ . So,  $G \subseteq G(F(X_i)/F)$ . Counting dimensions we get that  $G = G(F(X_i)/F)$  since  $C(X_i) = C(G)$ , the function field of  $G$ . Finally,  $F(X_i) = F\langle V \rangle$ , where  $V$  is the finite-dimensional vector space over  $C$  spanned by the  $X_i$ . By Theorem 2.1.3,  $F(X_i) \supset F$  is a Picard-Vessiot extension.  $\square$

Applying Theorems 2.5.2 and 2.5.3 we also obtain:

**Theorem 2.5.4.** *There is a specialization of the  $Y_i$  such that  $F(X_i) \supset F$  is a Picard-Vessiot extension if and only if there are  $f_i \in F$  such that all the  $W_k(Y_i)$ ,  $k \geq 1$ , map to non-zero elements via  $Y_i \mapsto f_i$ .*

**2.6. An example.** The previous Theorem 2.4.1 says that if there is an algebraic relation among the set of all the  $t_i$  and all the  $X_{ij}$  over the field of constants  $\mathcal{C}$  of  $F(X_{ij})$  then  $\mathcal{C}$  properly contains  $C$ .

In this section we give an example in which a new constant is produced from such an algebraic relation. We assume  $F = C$ . So, in particular, the coefficients  $f_{ij}$  in the derivation of  $F$  are constant. In this situation, since the transcendence degree of  $F$  over  $C$  is zero, if  $\mathcal{C} \not\supseteq C$ , the condition of Theorem 2.4.1 means that the  $X_{ij}$  are algebraically dependent over  $\mathcal{C}$ .

We restrict ourselves to the case  $n = 2$  and consider the following particular dependence relation.

Let

$$D(X_{ij}) = \sum_{\ell=1}^2 f_{i\ell} X_{\ell j},$$

where the  $f_{ij}$  are such that the wronskian  $W_1 = w(X_{11}, X_{12}, X_{21}, X_{22}) = 0$ . That is, the  $X_{ij}$  are linearly dependent over  $\mathcal{C}$ . Furthermore, assume that the linear relation among the  $X_{ij}$  is such that there are  $\beta_{12}, \beta_{21}, \beta_{22} \in \mathcal{C}$  with

$$(1) \quad X_{11} = \beta_{12}X_{12} + \beta_{21}X_{21} + \beta_{22}X_{22}$$

and that  $X_{12}, X_{21}$  and  $X_{22}$  are linearly independent. In order to simplify the computations we will also assume that  $\det[f_{ij}] = 0$ .

We want to find  $a, b, c \in F$  such that  $p = aX_{12} + bX_{21} + cX_{22}$  is a Darboux polynomial in  $F[X_{ij}]$ , that is  $D(aX_{12} + bX_{21} + cX_{22}) = q(aX_{12} + bX_{21} + cX_{22})$  for certain  $q \in F$ .

We have,

$$\begin{aligned} D(aX_{12} + bX_{21} + cX_{22}) &= a(f_{11}X_{12} + f_{12}X_{22}) + b(f_{21}X_{11} + f_{22}X_{21}) + c(f_{21}X_{12} + f_{22}X_{22}) \\ &= bf_{21}X_{11} + (af_{11} + cf_{21})X_{12} + bf_{22}X_{21} + (af_{12} + cf_{22})X_{22} \\ &= bf_{21}(\beta_{12}X_{12} + \beta_{21}X_{21} + \beta_{22}X_{22}) + (af_{11} + cf_{21})X_{12} + bf_{22}X_{21} \\ &\quad + (af_{12} + cf_{22})X_{22} \\ &= (af_{11} + bf_{21}\beta_{12} + cf_{21})X_{12} + b(f_{22} + f_{21}\beta_{12})X_{21} \\ &\quad + (af_{12} + bf_{21}\beta_{22} + cf_{22})X_{22} \\ &= qaX_{12} + qbX_{21} + qcX_{22}. \end{aligned}$$

Therefore,

$$(2) \quad \begin{aligned} [a(f_{11} - q) + bf_{21}\beta_{12} + cf_{21}]X_{12} + b(f_{22} + f_{21}\beta_{12} - q)X_{21} \\ + (af_{12} + bf_{21}\beta_{22} + c(f_{22} - q))X_{22} = 0. \end{aligned}$$

Since we are assuming that  $X_{12}, X_{21}$  and  $X_{22}$  are linearly independent their coefficients in (2) must be equal to zero. So we have the following homogeneous linear system in  $a, b, c$ :

$$\begin{aligned} (f_{11} - q)a + f_{21}\beta_{12}b + f_{21}c &= 0 \\ (f_{22} + f_{21}\beta_{12} - q)b &= 0 \\ f_{12}a + f_{21}\beta_{22}b + (f_{22} - q)c &= 0 \end{aligned}$$

In order for the above system to have non-trivial solutions we need that

$$\det \begin{bmatrix} f_{11} - q & f_{21}\beta_{12} & f_{21} \\ 0 & f_{22} + f_{21}\beta_{12} - q & 0 \\ f_{12} & f_{21}\beta_{22} & f_{22} - q \end{bmatrix} = 0.$$

But,

$$\begin{aligned}
 & \det \begin{bmatrix} f_{11} - q & f_{21}\beta_{12} & f_{21} \\ 0 & f_{22} + f_{21}\beta_{12} - q & 0 \\ f_{12} & f_{21}\beta_{22} & f_{22} - q \end{bmatrix} \\
 &= (f_{22} + f_{21}\beta_{12} - q) \det \begin{bmatrix} f_{11} - q & f_{21} \\ f_{12} & f_{22} - q \end{bmatrix} \\
 &= (f_{22} + f_{21}\beta_{12} - q)(\det[f_{ij}] - \left(\sum_{i=1}^2 f_{ii}\right)q + q^2) \\
 &= 0.
 \end{aligned}$$

This gives either

$$(3) \quad f_{22} + f_{21}\beta_{12} - q = 0$$

or

$$(4) \quad \det[f_{ij}] - \left(\sum_{i=1}^2 f_{ii}\right)q + q^2 = 0.$$

From (3)-(4) we get

$$(5) \quad q = f_{22} + f_{21}\beta_{12}$$

or

$$(6) \quad q = \frac{\sum_{i=1}^2 f_{ii} \pm \sqrt{(\sum_{i=1}^2 f_{ii})^2 - 4 \det[f_{ij}]}}{2}$$

Since we are assuming that  $\det[f_{ij}] = 0$ , (6) becomes:

$$(7) \quad q = \begin{cases} \sum_{i=1}^2 f_{ii}, & \text{or} \\ 0 \end{cases}$$

Choose  $q = \sum_{i=1}^2 f_{ii}$  and assume that  $q \neq 0$ ,  $q \neq f_{22} + f_{21}\beta_{12}$ . Then the second equation in the system implies that  $b = 0$  and the system becomes:

$$\begin{aligned}
 -f_{22}a + f_{21}c &= 0 \\
 f_{12}a - f_{11}c &= 0
 \end{aligned}$$

If  $f_{22} \neq 0$  then the above system has the general solution

$$a = \frac{f_{21}}{f_{22}}c, \text{ where } c \in \mathcal{C}.$$

In particular, if we take  $c = 1$  then  $p = \frac{f_{21}}{f_{22}}X_{12} + X_{22}$  satisfies

$$D\left(\frac{f_{21}}{f_{22}}X_{12} + X_{22}\right) = \left(\sum_{i=1}^2 f_{ii}\right)\left(\frac{f_{21}}{f_{22}}X_{12} + X_{22}\right).$$

On the other hand we also have that

$$D(\det[X_{ij}]) = \left( \sum_{i=1}^2 f_{ii} \right) \det[X_{ij}].$$

Let

$$\theta = \frac{\frac{f_{21}}{f_{22}} X_{12} + X_{22}}{\det[X_{ij}]}.$$

We have,

$$\begin{aligned} D(\theta) &= D\left(\frac{\frac{f_{21}}{f_{22}} X_{12} + X_{22}}{\det[X_{ij}]}\right) \\ &= \frac{D\left(\frac{f_{21}}{f_{22}} X_{12} + X_{22}\right) \det[X_{ij}] - \left(\frac{f_{21}}{f_{22}} X_{12} + X_{22}\right) D(\det[X_{ij}])}{\det[X_{ij}]^2} \\ &= \frac{\left(\sum_{i=1}^2 f_{ii}\right) \left(\frac{f_{21}}{f_{22}} X_{12} + X_{22}\right) \det[X_{ij}] - \left(\frac{f_{21}}{f_{22}} X_{12} + X_{22}\right) \left(\sum_{i=1}^2 f_{ii}\right) \det[X_{ij}]}{\det[X_{ij}]^2} \\ &= 0. \end{aligned}$$

That is,  $\theta$  is a new constant in  $F(X_{ij})$ .

Now we show that under the restrictions that we imposed on the  $f_{ij}$  it is possible to find a non-zero  $f_{22}$ .

Since we have a linear dependence relation among the  $X_{ij}$ , the wronskian  $W_1$  must be equal to zero. This Wronskian can be expressed, up to a sign, as the following product of determinants:

$$W_1 = \begin{vmatrix} 1 & 0 & 0 & 1 \\ f_{11} & f_{12} & f_{21} & f_{22} \\ A & B & E & F \\ C & D & G & H \end{vmatrix} \begin{vmatrix} X_{11} & X_{12} & 0 & 0 \\ X_{21} & X_{22} & 0 & 0 \\ 0 & 0 & X_{11} & X_{12} \\ 0 & 0 & X_{21} & X_{22} \end{vmatrix} = M(f_{ij}) \det[X_{ij}]^2,$$

where

$$\begin{aligned} A &= f'_{11} + f_{11}^2 + f_{12}f_{21}, \\ B &= f'_{12} + f_{11}f_{12} + f_{12}f_{22} \\ C &= f_{11}A + f_{21}B + A' \\ &= 3f_{11}f'_{11} + 2f_{11}f_{12}f_{21} + 2f'_{12}f_{21} + f''_{11} + f_{12}f'_{21} + f_{11}^3, \end{aligned}$$

$$\begin{aligned}
 D &= f_{12}A + f_{22}B + B' \\
 &= 2f'_{11}f_{12} + f_{11}^2f_{12} + f_{12}^2f_{21} + f_{21}f_{22}^2 + 2f'_{12}f_{22} + f_{11}f'_{12} \\
 &\quad + f_{12}f'_{22} + f_{12}'' + f_{11}f_{12}f_{22}, \\
 E &= f'_{21} + f_{21}f_{11} + f_{22}f_{21}, \\
 F &= f'_{22} + f_{12}f_{21} + f_{22}^2, \\
 G &= f_{11}E + f_{21}F + E' \\
 &= 2f'_{21}f_{11} + f_{21}f_{11}^2 + f_{22}f_{21}f_{11} + 2f'_{22}f_{21} + f_{12}f_{21}^2 \\
 &\quad + f_{22}^2f_{21} + f_{21}'' + f_{21}f'_{11} + f_{22}f'_{21}, \\
 H &= f_{22}F + f_{12}E + F' \\
 &= f_{21}f_{11}f_{12} + 2f_{22}f_{21}f_{12} + 3f_{22}f'_{22} + 2f_{12}f'_{21} + f'_{12}f_{21} \\
 &\quad + f_{22}^3 + f_{22}'' .
 \end{aligned}$$

and

$$M(f_{ij}) = \begin{vmatrix} 1 & 0 & 0 & 1 \\ f_{11} & f_{12} & f_{21} & f_{22} \\ A & B & E & F \\ C & D & G & H \end{vmatrix} .$$

We have after simplifying using the hypothesis that  $\det[f_{ij}] = 0$ ,

$$\begin{aligned}
 M(f_{ij}) &= (f_{22} - f_{11})(f'_{12}f'_{21} - f'_{21}f'_{12}) + (f'_{22} - f'_{11})(f''_{12}f_{21} - f_{12}f''_{21}) \\
 &\quad - f'_{12}f'_{21}(f_{11} - f_{22})^2 - f_{12}f_{21}(f'_{11} - f'_{22}) \\
 &\quad + f_{12}f'_{21}(f_{11}f'_{11} + f_{22}f'_{22} - f'_{11}f_{22} - f_{11}f'_{22} + f''_{22} - f''_{11} + f_{12}f'_{21} - f'_{12}f_{21}) \\
 &\quad + f'_{12}f_{21}(f_{11}f'_{11} + f_{22}f'_{22} - f'_{11}f_{22} - f_{11}f'_{22} + f''_{11} - f''_{22} + f'_{12}f_{21} - f_{12}f'_{21}) .
 \end{aligned}$$

Getting the above expression for  $M(f_{ij})$  took long and involved computations. We first computed the determinant directly and then we checked the result using Dogson's method [6, 28].

The wronskian  $W_1 = 0$  if and only if  $M(f_{ij}) = 0$ . Now, observe that if  $f_{12} = 0$  then  $f'_{12} = 0$  which implies that  $B = 0$  and  $D = 0$  as well. Therefore  $M(f_{ij}) = 0$ . So, if we let  $M(Y_{ij})$  be the differential polynomial in the  $Y_{ij}$  whose specialization to the  $f_{ij}$  is  $M(f_{ij})$  then  $M(Y_{ij})$  is in the differential ideal

$$\begin{aligned}
 \mathcal{I} &= \{\det[Y_{ij}], Y_{12}\} \\
 &= \{Y_{11}Y_{22} - Y_{12}Y_{21}, Y_{12}\} \\
 &= \{Y_{11}Y_{22}, Y_{12}\}
 \end{aligned}$$

of  $C\{Y_{11}, Y_{12}, Y_{21}, Y_{22}\}$ . It is easy to see that  $Y_{22}$  is not in  $\mathcal{I}$ . Indeed, suppose that

$$(8) \quad Y_{22} = pY_{11}Y_{22} + qY_{12} + r,$$

where  $p, q \in C\{Y_{11}, Y_{12}, Y_{21}, Y_{22}\}$ ,

$$r = \sum_{i,j} [p_i (Y_{11}Y_{22})^{(i)} + q_j Y_{12}^{(j)}]$$

with  $p_i, q_j \in C\{Y_{11}, Y_{12}, Y_{21}, Y_{22}\}$ .

Now, consider the map

$$\psi : C\{Y_{11}, Y_{21}, Y_{22}\} \longrightarrow C[Y_{11}, Y_{21}, Y_{22}]$$

given by  $\psi(Y_{22}) = Y_{22}$  and  $\psi(Y_{ij}) = 0$  for  $i, j \neq 2$ . Let  $\bar{p} = \psi(p)$ ,  $\bar{q} = \psi(q)$ ,  $\bar{r} = \psi(r)$ . We have that  $\bar{r} = 0$  and (8) becomes

$$Y_{22} = 0.$$

which is impossible. □

#### REFERENCES

1. W. W. Adams and Ph. Loustau, *An Introduction to Gröbner Bases*, Graduate Studies in Mathematics, American Mathematical Society (1994).
2. D. W. Albrecht, E. L. Mansfield, A. E. Milne, Algorithms for special integrals of ordinary differential equations. *J. Phys. A* **29** (1996), 5, 973–991.
3. F. Beukers, G. Heckmann, Monodromy for the hypergeometric function  ${}_nF_{n-1}$ , *Invent. Math.* **95** (1989), 325–354.
4. A. Bialynicki-Birula, On the inverse problem of Galois theory of differential fields, *Bull. Amer. Math. Soc.* **16** (1963), 960–964.
5. K. Bousset, *Groupes de Galois des équations hypergéométriques*, C. R. Acad. Sci. Paris. **309**, I (1989), 587–589.
6. C. L. Dodgson, Condensation of determinants, *Proc. Royal Soc. London* **15** (1866), 150–155.
7. A. Duval, M. Loday-Richaud, Kovacic’s algorithm and its applications to some families of special functions, *AAECC Journal* **3** (1992).
8. A. Duval, C. Mitschi, Matrices de Stokes et groupe de Galois des équations hypergéométriques confluentes généralisées, *Pacific J. Math.* **138** (1989), 25–56.
9. A. Figueiredo, T. M. Rocha-Filho, L. Brenig, Necessary conditions for the existence of quasi-polynomial invariants: the quasi-polynomial and Lotka-Volterra systems. *Phys. A* **262** (1999), 1-2, 158–180.
10. L. Goldman, Specializations and the Picard–Vessiot theory, *Trans. Amer. Math. Soc.* **85** (1957), 327–356.
11. L. Juan, *A Generic Picard–Vessiot extension with group  $GL_n$  and the inverse differential Galois problem*, Ph. D. Thesis, University of Oklahoma (2000).
12. L. Juan, Generic Picard–Vessiot extensions for connected linear algebraic groups. (*In preparation.*)
13. N. Katz, On the calculation of some differential Galois groups, *Invent. Math.* **87** (1987), 13–61.
14. J. Kovacic, The inverse problem in the Galois theory of differential fields, *Ann. of Math.* **89** (1969), 583–608.
15. J. Kovacic, On the inverse problem in the Galois theory of differential fields, *Ann. of Math.* **93** (1971), 269–284.
16. S. Labrunie, On the non-integrability of the  $SU(2)$  invariant Kähler-Einstein metrics. *Phys. Lett. A* **236** (1997), 5-6, 439–444.
17. S. Labrunie, Simon On the polynomial first integrals of the  $(a, b, c)$  Lotka-Volterra system. *J. Math. Phys.* **37** (1996), 11, 5539–5550.

18. A. J. Maciejewski, J. -M. Strelcyn, Non-integrability of the generalized Halphen system. *Regul. Khaoticheskaya Din.* **1** (1996), 2, 3–12.
19. A. Magid, *Lectures in Differential Galois Theory*, University Lecture Series, American Mathematical Society (1994).
20. J. Miller *On Differentially Hilbertian Differential Fields*, Ph.D. thesis, Columbia University (1970).
21. C. Mitschi, Groupe de Galois des équations hypergéométriques confluentes généralisées, *C. R. Acad. Sci. Paris.* **309**, I (1989), 217–220.
22. C. Mitschi, Differential Galois Groups of Generalized Hypergeometric Equations: An Approach Using Stokes Multipliers, *Pacific J. Math.* **176** 2, (1996), 365–405.
23. C. Mitschi and M. F. Singer, Connected Linear Groups as Differential Galois Groups, *Journal of Algebra.* **184** (1996), 333–361.
24. E. Noether, Gleichungen mit vorgeschriebener Gruppen, *Math. Ann.* **78** (1918), 221–229.
25. M. van der Put, Recent Work on Differential Galois Theory, *Séminaire BOURBAKI*, 50ème année, (1997-1998), n° 849.
26. J. -P. Ramis, About the solution of some inverse problems in differential Galois theory by Hamburger equations, in *Differential Equations, Dynamical Systems, and Control Science*, Elworthy, Everett, and Lee, eds., Lecture Notes in Pure and Applied Mathematics. **157** (1994), 277–300.
27. J. -P. Ramis, About the Inverse Problem in Differential Galois Theory: The Differential Abhyankar Conjecture, *The Stokes Phenomenon and Hilbert's 16th Problem*; B. I. J. Braaksma, et al., eds. World Scientific, Singapore (1996)
28. D. P. Robins and H. Rumsey, Jr., Determinants and Alternating Sign Matrices, *Advances in Mathematics.* **62** (1986), 169–184.
29. M. F. Singer, Direct and Inverse Problems in Differential Galois Theory, *Selected Works of Ellis Kolchin with Commentary*, Bass, Buium, Cassidy eds., American Mathematical Society (1999), 527–554.
30. M. F. Singer, Moduli of linear differential equations on the Riemann sphere with fixed Galois groups, *Pacific J. Math.* **106**, 2 (1993), 343–395.
31. M. F. Singer, F. Ulmer, Liouvillian and algebraic solutions of second and third order linear differential equations, *Journal of Symbolic Computation.* **16** (1993), pp. 37–74.
32. C. Tretkoff and M. Tretkoff, Solution of the inverse problem of differential Galois theory in the classical case, *Amer. J. Math.* **101** (1979), 1327–1332.
33. F. Ulmer, J. -A. Weil, Note on Kovacic's algorithm, *J. Symbolic Comput.* **28**, 2, (1996) 179–200.
34. J. -A. Weil, *Constantes et polynômes de Darboux en algèbre différentielle: applications aux systèmes différentiels linéaires*, Ph.D. Thesis, École Polytechnique de France (1995).
35. J. -A. Weil, First integrals and Darboux polynomials of homogeneous linear differential systems. *Applied algebra, algebraic algorithms and error-correcting codes* (Paris, 1995), 469–484, Lecture Notes in Comput. Sci., 948, Springer, Berlin, 1995.

DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, BOX 41042, LUBBOCK, TX 79409-1042, USA

*E-mail address:* ljuan@math.ttu.edu

*URL:* www.math.ttu.edu/~ljuan