

# PRO CDH-DESCENT FOR CYCLIC HOMOLOGY AND $K$ -THEORY

MATTHEW MORROW

*Mathematisches Institut, Universität Bonn, Endenicher Allee 60,  
53115 Bonn, Germany* ([morrow@math.uni-bonn.de](mailto:morrow@math.uni-bonn.de))  
<http://www.math.uni-bonn.de/people/morrow/>

(Received 25 May 2014; revised 6 November 2014; accepted 6 November 2014;  
first published online 27 November 2014)

*Abstract* In this paper, we prove that cyclic homology, topological cyclic homology, and algebraic  $K$ -theory satisfy a pro Mayer–Vietoris property with respect to abstract blow-up squares of varieties, in both zero and finite characteristic. This may be interpreted as the well-definedness of  $K$ -theory with compact support.

*Keywords:*  $K$ -theory; cyclic homology; cdh-topology

2010 *Mathematics subject classification:* Primary 19D55  
Secondary 13D03

## Contents

<b>Introduction</b>	<b>539</b>
<b>1 Review of pro modules and Artin–Rees properties</b>	<b>542</b>
1.1 Pro abelian groups and pro modules . . . . .	542
1.2 Artin–Rees properties . . . . .	543
<b>2 Formal function properties and pro descent for André–Quillen homology</b>	<b>544</b>
2.1 André–Quillen homology for schemes . . . . .	544
2.2 Formal function properties for AQ homology of schemes . . . . .	547
2.3 Pro descent for AQ homology with respect to blow-up squares . . . . .	550
<b>3 Formal function properties and pro descent for <math>HH</math>, <math>HC</math>, <math>THH</math>, <math>TC</math>, and <math>K</math>-theory</b>	<b>553</b>
3.1 Hochschild and cyclic homology . . . . .	554
3.2 Topological Hochschild and cyclic homology . . . . .	557
3.3 Pro cdh-descent for $K$ -theory . . . . .	559
<b>4 Applications of Theorem 3.7</b>	<b>562</b>
4.1 $K$ -theory with compact support . . . . .	562
4.2 Negative $K$ -groups . . . . .	564
<b>References</b>	<b>566</b>

Introduction

The primary goal of this paper is to prove that algebraic  $K$ -theory satisfies a pro Mayer–Vietoris property with respect to abstract blow-up squares of varieties, in both zero and finite characteristic, at least assuming strong resolution of singularities. This may also be interpreted as the well-definedness of a theory of  $K$ -groups with compact support.

To state the main result, we first recall that an *abstract blow-up square* of schemes is a pull-back diagram

$$\begin{array}{ccc}
 Y' & \longrightarrow & X' \\
 \downarrow & & \downarrow \\
 Y & \longrightarrow & X
 \end{array}$$

where  $X' \rightarrow X$  is proper,  $Y \rightarrow X$  is a closed embedding, and the induced map on the open subschemes  $X' \setminus Y' \rightarrow X \setminus Y$  is an isomorphism. We denote by  $rY$  the  $r$ th infinitesimal thickening of  $Y$  inside  $X$ , i.e.,  $rY := \text{Spec } \mathcal{O}_X/\mathcal{I}^r$ , where  $\mathcal{I} \subseteq \mathcal{O}_X$  is the sheaf of ideals defining  $Y$ , and similarly for other closed embeddings.

Our pro descent theorem is the following.

**Theorem 0.1** (Pro cdh-descent for  $K$ -theory; see Theorem 3.7). *Let  $k$  be an infinite perfect field which has strong resolution of singularities. For any abstract blow-up square, as above, consisting of finite type  $k$ -schemes, the resulting square of pro spectra*

$$\begin{array}{ccc}
 K(X) & \longrightarrow & K(X') \\
 \downarrow & & \downarrow \\
 \{K(rY)\}_r & \longrightarrow & \{K(rY')\}_r
 \end{array}$$

is homotopy Cartesian. In other words, the canonical map of pro abelian relative  $K$ -groups

$$\{K_n(X, rY)\}_r \longrightarrow \{K_n(X', rY')\}_r$$

is an isomorphism for all  $n \in \mathbb{Z}$ .

The main applications of Theorem 0.1 are to zero cycles; since the theorem may be accepted as a black box for these applications, they are presented separately in an accompanying paper [25]. There, we solve cases of an outstanding conjecture of Srinivas and Bloch [30, p. 6] concerning the Levine–Weibel Chow group of zero cycles on singular varieties, and relate Chow groups with modulus, which play a prominent role in Kerz and Saito’s higher-dimensional class field theory [19], to both Levine–Weibel Chow groups and  $K$ -theory via cycle class maps.

Theorem 0.1 may be interpreted as a definition of  $K$ -theory with compact support as follows. Given a separated  $k$ -variety  $X$ , we may choose a proper compactification  $\bar{X}$ , set  $Y := \bar{X} \setminus X$ , and then define  $K^c(X) := \text{holim}_r K(\bar{X}, rY)$ . Theorem 0.1 implies that this definition does not depend on the chosen compactification  $\bar{X}$  of  $X$ , and we prove in Proposition 4.4 that there is a pushforward localization sequence relating  $K^c(U)$

and  $K^c(X)$  for any open subvariety  $U \subseteq X$ , thereby justifying the nomenclature of  $K^c$  as a theory with compact support.

In characteristic zero, we actually establish Theorem 0.1 in much greater generality, namely for any abstract blow-up square of Noetherian, quasi-excellent,  $\mathbb{Q}$ -schemes of finite Krull dimension. This is obtained by first extending Haesemeyer’s argument [15, §§ 5, 6], for checking whether a presheaf of spectra satisfies cdh-descent, to this generality; see Proposition 3.6. This also allows us to prove the vanishing part of Weibel’s  $K$ -dimension conjecture for such schemes; see Theorem 4.7.

To contextualize Theorem 0.1, we should make a few comments about cdh-descent. Algebraic  $K$ -theory (as well as André–Quillen, Hochschild, topological Hochschild, cyclic, and topological cyclic homologies) does not satisfy descent in Voevodsky’s cdh-topology [33], in contrast to, for example, Weibel’s homotopy-invariant  $K$ -theory [15], or periodic cyclic homology [4]. Since it is known that  $K$ -theory does satisfy Nisnevich descent, this precisely means that taking the  $K$ -theory of an abstract blow-up square does not necessarily yield a homotopy Cartesian square of spectra. In fact, the failure of  $K$ -theory to satisfy cdh-descent in characteristic zero (respectively,  $p > 0$ ) is precisely equal to the analogous obstruction in cyclic homology [3, 4, 15] (respectively, topological cyclic homology [12]); this has led to new calculations of the  $K$ -theory of singular algebraic varieties, especially by Cortiñas, Haesemeyer, Schlichting, Walker, and Weibel [4–8] in characteristic zero. The moral of Theorem 0.1 is that the failure of  $K$ -theory to satisfy cdh-descent can be remedied by taking all infinitesimal thickenings of the exceptional subschemes  $Y$  and  $Y'$  into account.

We now briefly discuss how Theorem 0.1 is proved and simultaneously state our other main results. Using the aforementioned cdh comparison between  $K$ -theory and cyclic homology, Theorem 0.1 in characteristic zero may be reduced to an analogous pro Mayer–Vietoris assertion about cyclic homology, or even Hochschild or André–Quillen homology. We establish this pro Mayer–Vietoris property in considerable generality (see Theorem 2.10 for the André–Quillen case).

**Theorem 0.2** (Pro descent for  $HH$  and  $HC$  with respect to abstract blow-ups; see Theorem 3.3). *Let  $k$  be a Noetherian ring, and let*

$$\begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

*be an abstract blow-up square of Noetherian, finite Krull-dimensional  $k$ -schemes. Then the canonical maps*

$$\{HH_n^k(X, rY)\}_r \longrightarrow \{HH_n^k(X', rY')\}_r, \quad \{HC_n^k(X, rY)\}_r \longrightarrow \{HC_n^k(X', rY')\}_r$$

*are isomorphisms of pro abelian groups for all  $n \in \mathbb{Z}$ .*

We should mention at this point that our Hochschild and cyclic homology groups are always computed in the derived sense, that is, after replacing rings by free simplicial resolutions (see § 3.1 for details).

The key input to proving Theorem 0.2 is new formal function theorems for André–Quillen, Hochschild, and cyclic homology, in the style of Grothendieck’s formal functions theorem for coherent cohomology. In particular, in the case of Hochschild homology, we prove the following (see Corollary 2.5 for the André–Quillen case).

**Theorem 0.3** (Formal functions theorem for *HH*; see Theorem 3.2(v)). *Let  $A$  be a Noetherian ring,  $I \subseteq A$  an ideal, and  $X$  a proper scheme over  $A$  of finite Krull dimension. Then the canonical map*

$$\{HH_n^A(X) \otimes_A A/I^r\}_r \longrightarrow \{HH_n^{A/I^r}(X \times_A A/I^r)\}_r$$

is an isomorphism of pro  $A$ -modules for all  $n \in \mathbb{Z}$ .

If  $A$ , as in the previous theorem, is moreover  $I$ -adically complete, then it follows that the canonical map  $HH_n^A(X) \longrightarrow \varprojlim_r HH_n^{A/I^r}(X \times_A A/I^r)$  is an isomorphism for all  $n \in \mathbb{Z}$ , which is the statement closest to the usual formulation of Grothendieck’s formal functions theorem for coherent cohomology.

The proof of Theorem 0.1 in finite characteristic is similar in outline, except that the results concerning André–Quillen, Hochschild, and cyclic homology in §§ 2 and 3.1 must be replaced by similar results for topological Hochschild and cyclic homology. These results are not contained in the current paper, but may rather be found in recent joint work with Dundas [10], which includes in particular the *THH* analogue of Theorem 0.3. The few details missing from [10], where for example the analogue of Theorem 0.2 for *THH* and *TC* was not explicitly stated, are given in § 3.2.

**Notation, etc.** All rings are commutative, associative, and unital. If  $A_\bullet$  is a simplicial abelian group, then we often abuse notation by identifying  $A_\bullet$  with its associated complex  $NA_\bullet$ ; in particular, we speak of the homology groups  $H_n(A_\bullet)$  of  $A_\bullet$ , rather than more correctly, but equivalently, the simplicial homotopy groups  $\pi_n(A_\bullet)$  or the homology groups  $H_n(NA_\bullet)$ .

## 1. Review of pro modules and Artin–Rees properties

### 1.1. Pro abelian groups and pro modules

Everything we need about categories of pro objects may be found in one of the standard references, such as the appendix to [2], or [17]. We will often use  $\text{Pro}(A\text{-mod})$ , the category of pro  $A$ -modules for some ring  $A$ , and  $\text{Pro } Ab$ , the category of pro abelian groups.

Let  $\mathcal{C}$  be a category. In this paper, an object of  $\text{Pro } \mathcal{C}$  is simply an inverse system  $\cdots \rightarrow A_2 \rightarrow A_1$  of objects and morphisms in  $\mathcal{C}$ , which is denoted  $\{A_r\}_r$  or very occasionally  $A_\infty$ . Morphisms in  $\text{Pro } \mathcal{C}$  are given by the rule

$$\text{Hom}_{\text{Pro } \mathcal{C}}(\{A_r\}_r, \{B_s\}_s) := \varprojlim_s \varinjlim_r \text{Hom}_{\mathcal{C}}(A_r, B_s),$$

where the right-hand side is a genuine pro-ind limit in the category of sets, and composition is defined in the obvious way. For example, a pro object  $\{A_r\}_r$  is isomorphic to zero (assuming that a zero object exists in  $\mathcal{C}$ , and hence also in  $\text{Pro } \mathcal{C}$ ) if and only if for each  $r \geq 1$  there exists  $s \geq r$  such that the transition map  $A_s \rightarrow A_r$  is zero.

There is a fully faithful embedding  $\mathcal{C} \rightarrow \text{Pro } \mathcal{C}$ . Assuming that  $\mathcal{C}$  has countable inverse limits, this has a right adjoint  $\text{Pro } \mathcal{C} \rightarrow \mathcal{C}$ ,  $\{A_r\}_r \mapsto \varprojlim_r A_r$ , which is left exact but not right exact. Moreover, if  $\mathcal{C}$  is an abelian category then so is  $\text{Pro } \mathcal{C}$ : given a inverse system of exact sequences

$$\cdots \longrightarrow A_{n-1}(r) \longrightarrow A_n(r) \longrightarrow A_{n+1}(r) \longrightarrow \cdots,$$

the ‘limit as  $r \rightarrow \infty$ ’, namely

$$\cdots \longrightarrow \{A_{n-1}(r)\}_r \longrightarrow \{A_n(r)\}_r \longrightarrow \{A_{n+1}(r)\}_r \longrightarrow \cdots,$$

is an exact sequence in  $\text{Pro } \mathcal{C}$ .

Pro spectral sequences play an important role in the paper, which we will discuss for concreteness only in the case of abelian groups. Suppose that

$$E_{pq}^1(r) \implies H_{p+q}(r),$$

for  $r \geq 1$ , are spectral sequences of abelian groups, which are functorial in that we have morphisms of spectral sequences  $\cdots \rightarrow E_{pq}^\bullet(2) \rightarrow E_{pq}^\bullet(1)$ . To avoid convergence issues, suppose that each spectral sequence is bounded, by a bound independent of  $r$ ; for example, each spectral sequence might be zero outside the first quadrant. Then we will often ‘let  $r \rightarrow \infty$ ’ to obtain a spectral sequence of pro abelian groups

$$E_{pq}^1(\infty) := \{E_{pq}^1(r)\}_r \implies \{H_{p+q}(r)\}_r.$$

For further discussion and a dummy example, see [24, Appendix A].

### 1.2. Artin–Rees properties

For the sake of reference, we now formally state a fundamental Artin–Rees result which will be used several times; this result appears to have been first noticed and exploited by André [1, Proposition 10 & Lemma 11] and Quillen [27, Lemma 9.9].

**Theorem 1.1** (André–Quillen). *Let  $A$  be a Noetherian ring, and  $I \subseteq A$  any ideal.*

- (i) *If  $M$  is a finitely generated  $A$ -module, then the pro  $A$ -module  $\{\text{Tor}_n^A(A/I^r, M)\}_r$  vanishes for all  $n > 0$ .*
- (ii) *The ‘completion’ functor*

$$\begin{aligned} - \otimes_A A/I^\infty : A\text{-mod} &\longrightarrow \text{Pro}(A\text{-mod}) \\ M &\mapsto \{M \otimes_A A/I^r\}_r \end{aligned}$$

*is exact on the subcategory of finitely generated  $A$ -modules.*

**Sketch of proof.** By picking a resolution  $P_\bullet$  of  $M$  by finitely generated projective  $A$ -modules and applying the classical Artin–Rees property to the pair  $d(P_n) \subseteq P_{n-1}$ , one sees that for each  $r \geq 1$  there exists  $s \geq r$  such that the map

$$\text{Tor}_n^A(A/I^s, M) \rightarrow \text{Tor}_n^A(A/I^r, M)$$

is zero. This proves (i). (ii) is just a restatement of (i). □

Next, we quote a similar Artin–Rees result for André–Quillen homology from [24], a companion to this paper which treats pro excision problems; the basics of André–Quillen homology will be recalled in § 2.1:

**Theorem 1.2.** *Let  $k \rightarrow A$  be a homomorphism of Noetherian rings,  $I \subseteq A$  an ideal, and  $M$  an  $A$ -module. Then the following hold.*

- (i)  $\{D_n^i(A/I^r|A, M/I^r M)\}_r = 0$  for all  $n \geq 0, i \geq 1$ .
- (ii) The canonical map  $\{D_n^i(A|k, M/I^r M)\}_r \rightarrow \{D_n^i(A/I^r|k, M/I^r M)\}_r$  is an isomorphism for all  $n \geq 0, i \geq 0$ .

**Proof.** This is [24, Theorem 4.4], where the pro H-unitality hypothesis of the cited theorem is satisfied thanks to [24, Theorem 0.3]. □

### 2. Formal function properties and pro descent for André–Quillen homology

The aim of this section is to prove that André–Quillen homology of proper schemes satisfies certain formal function properties. These will be established in § 2.2, after preliminary material on André–Quillen homology is presented in § 2.1. From these formal function properties we show in § 2.3 that André–Quillen homology satisfies the desired pro Mayer–Vietoris property with respect to abstract blow-up squares; this is a key step in deducing the same for  $K$ -theory in characteristic zero.

#### 2.1. André–Quillen homology for schemes

In this preliminary section, we recall the fundamentals of André–Quillen homology [1, 26, 28], though we expect that the reader already has some familiarity with it, before extending it from algebras to schemes in the standard way using hypercohomology, as was done for Hochschild and cyclic homology in [34, 38].

Let  $k \rightarrow A$  be a homomorphism of rings; let  $P_\bullet \rightarrow A$  be a simplicial resolution of  $A$  by free  $k$ -algebras, and set

$$\mathbb{L}_{A|k} := \Omega_{P_\bullet|k}^1 \otimes_{P_\bullet} A.$$

Thus  $\mathbb{L}_{A|k}$  is a simplicial  $A$ -module which is free in each degree; it is called the *cotangent complex* of the  $k$ -algebra  $A$ . The cotangent complex up to homotopy depends only on  $A$ , since the free simplicial resolution  $P_\bullet \rightarrow A$  is unique up to homotopy.

Given simplicial  $A$ -modules  $M_\bullet, N_\bullet$ , the tensor product and alternating powers are new simplicial  $A$ -modules defined degreewise:  $(M_\bullet \otimes_A N_\bullet)_n = M_n \otimes_A N_n$  and  $(\bigwedge_A^i M_\bullet)_n = \bigwedge_A^i M_n$ .

In particular, we set  $\mathbb{L}_{A|k}^i := \bigwedge_A^i \mathbb{L}_{A|k}$  for each  $i \geq 1$ . The *André–Quillen homology* of the  $k$ -algebra  $A$ , with coefficients in any  $A$ -module  $M$ , is defined by

$$D_n^i(A|k, M) := H_n(\mathbb{L}_{A|k}^i \otimes_A M),$$

for  $n \geq 0, i \geq 1$ . When  $M = A$ , the notation is simplified to

$$D_n^i(A|k) := D_n^i(A|k, A) = H_n(\mathbb{L}_{A|k}^i).$$

When  $i = 1$ , the superscript is often omitted, writing  $D_n(A|k, M) = H_n(\mathbb{L}_{A|k} \otimes_A M)$  and  $D_n(A|k) = H_n(\mathbb{L}_{A|k})$  instead. If  $k \rightarrow A$  is essentially of finite type and  $k$  is Noetherian,

then  $D_n^i(A|k, M)$  is a finitely generated  $A$ -module for all  $n, i$ , and for all finitely generated  $A$ -modules  $M$ .

To avoid any ambiguity, we also remark that the notation  $D_n^i(A|k, M)$  is defined in the same way if  $i \leq 0$  or  $n < 0$ . However,  $D_n^i(A|k, M) = 0$  if  $n < 0$  and

$$D_n^0(A|k, M) = \begin{cases} M & n = 0, \\ 0 & \text{else,} \end{cases}$$

since  $\mathbb{L}_{A|k}^0 \otimes_A M \simeq M$ .

Now, we extend the definition of André–Quillen homology to schemes; since we restrict consideration to Noetherian schemes of finite Krull dimension, we may follow a classical approach, as we explain in the next remark.

**Remark 2.1** (Hypercohomology of non-bounded-below complexes). Our main results concerning André–Quillen (and later Hochschild and cyclic) homology of schemes concern Noetherian schemes or even varieties; therefore, we will restrict consideration throughout to Noetherian schemes of finite Krull dimension to simplify the hypercohomology of complexes which are not necessarily bounded below.

The Zariski hypercohomology of a cochain complex of sheaves  $M^\bullet$  on a Noetherian scheme  $X$  of finite Krull dimension is defined as follows [23, Appendix C]: there exists a quasi-isomorphism  $M^\bullet \xrightarrow{\sim} \mathcal{F}^\bullet$ , where  $\mathcal{F}^\bullet$  is a cochain complex of acyclic sheaves on  $X$ , and one shows that

$$\mathbb{H}^*(X, M^\bullet) := H^*(\mathcal{F}^\bullet(X))$$

is independent of  $\mathcal{F}$ . We may use the Cartan–Eilenberg approach to construct  $\mathcal{F}^\bullet$ : first, pick functorial acyclic resolutions  $M^p \xrightarrow{\sim} \mathcal{F}^{p\bullet}$  of length  $\leq \dim X$  for each  $p \in \mathbb{Z}$  (e.g., the Godement resolution), and then set  $\mathcal{F}^\bullet = \text{Tot } \mathcal{F}^{\bullet\bullet}$ ; note that at most  $1 + \dim X$  terms appear in each direct sum  $\mathcal{F}^n = \bigoplus_{p+q=n} \mathcal{F}^{p,q}$ .

The familiar spectral sequences

$$\begin{aligned} E_1^{p,q} = H^q(X, M^p) &\implies \mathbb{H}^{p+q}(X, M^\bullet) \\ {}'E_2^{p,q} = H^p(X, H^q(M^\bullet)) &\implies \mathbb{H}^{p+q}(X, M^\bullet) \end{aligned}$$

remain valid and bounded. We will also require two variations on Deligne’s spectral sequence for the hypercohomology of a bounded-below filtered complex [9, 1.4.5], which are not hard to prove given the boundedness assumptions we impose.

- (i) Let  $M^{\bullet\bullet}$  be a bounded (i.e.,  $M^{p,q}$  is non-zero for only finitely many  $p, q$  in each bounded region of the plane) double cochain complex of sheaves. Then there is a bounded spectral sequence

$$E_2^{p,q} = \mathbb{H}^p(X, H_v^q(M^{\bullet\bullet})) \implies \mathbb{H}^{p+q}(X, \text{Tot } M^{\bullet\bullet}),$$

where  $H_v$  indicates taking cohomology in the second index.

- (ii) Let  $M^\bullet$  be a cochain complex of sheaves equipped with a descending filtration  $FM^\bullet$  which satisfies  $F^0M^\bullet = M^\bullet$  and  $F^pM^\bullet = 0$  for  $p \gg 0$ . Then there is a bounded spectral sequence

$$E_1^{p,q} = \mathbb{H}^{p+q}(X, \text{gr}^q M^\bullet) \implies \mathbb{H}^{p+q}(X, M^\bullet).$$

Now, we are prepared to define André–Quillen homology of schemes. Let  $k$  be a ring, and fix  $i \geq 0$ . First, notice that if  $A$  is a  $k$ -algebra then it is possible to choose a simplicial resolution  $P_\bullet(A) \rightarrow A$  by free  $k$ -algebras in a way which is functorial in  $A$ , for example, via the comonad associating to  $A$  the free  $k$ -algebra generated by the set  $A$ ; see [37, e.g., 8.6.16]. So, given a scheme  $X$  over  $k$ , let  $\tilde{P}_\bullet(X)$  denote the simplicial sheaf of  $k$ -algebras obtained by degreewise sheafifying  $U \mapsto P_\bullet(\mathcal{O}_X(U))$ ; also, put  $\mathbb{L}_{X|k}^i := \Omega_{\tilde{P}_\bullet(X)/k}^i \otimes_{\tilde{P}_\bullet(X)} \mathcal{O}_X$ . Given a quasi-coherent  $\mathcal{O}_X$ -module  $M$ , and assuming that  $X$  is Noetherian of finite Krull dimension, define the André–Quillen homology of  $X$ , relative to  $k$ , with coefficients in  $M$  to be

$$D_n^i(X|k, M) := \mathbb{H}^{-n}(X, \mathbb{L}_{X|k, \text{neg}}^i \otimes_{\mathcal{O}_X} M),$$

where the subscript *neg* indicates replacing the simplicial sheaf  $\mathbb{L}_{X|k}^i$  by its associated chain complex and then negating the indexing to get a cochain complex of sheaves.

We continue the standard notational abuses of omitting  $i$  when it equals 1 and omitting  $M$  when it equals  $\mathcal{O}_X$ . Note, from the first two properties below, that  $D_n^i(X|k)$  can be non-zero for  $n$  both positive and negative.

The following consequences are either automatic from the construction, or follow from the previous remark on hypercohomology and mimicking the arguments for Hochschild homology given in [34, 38], bearing in mind that André–Quillen homology of rings behaves well under localization.

**$i = 0$ :** Since  $\mathbb{L}_{X|k, \text{neg}}^0 \otimes_{\mathcal{O}_X} M \simeq M$ , we have  $D_n^0(X|k, M) \cong H^{-n}(X, M)$ .

**Agreement on affines:** If  $X = \text{Spec } A$  is an affine  $k$ -scheme, then  $D_n^i(X|k, M) = D_n^i(A|k, M)$ .

**Mayer–Vietoris property:** If  $U, V$  are an open cover of  $X$ , then there is a long exact Mayer–Vietoris sequence

$$\dots \rightarrow D_n^i(X|k, M) \rightarrow D_n^i(U|k, M|_U) \oplus D_n^i(V|k, M|_V) \rightarrow D_n^i(U \cap V|k, M|_{U \cap V}) \rightarrow \dots$$

**Hypercohomology spectral sequence:** There is a bounded fourth quadrant spectral sequence

$$E_2^{pq} = H^p(X, \mathcal{D}_{-q}^i(X|k, M)) \implies D_{-p-q}^i(X|k, M),$$

where  $\mathcal{D}_{-q}^i(X|k, M)$  denotes the Zariski sheafification of  $U \mapsto D_{-q}^i(U|k, M|_U)$ . This shows that  $D_n^i(X|k, M) = 0$  for  $n < -\dim X$ .

Moreover,

**Homology long exact sequence:** If  $0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$  is a short exact sequence of quasi-coherent  $\mathcal{O}_X$ -modules, then there is a long exact sequence of André–Quillen homology

$$\dots \rightarrow D_n^i(X|k, M) \rightarrow D_n^i(X|k, N) \rightarrow D_n^i(X|k, P) \rightarrow \dots$$

(Proof: This is the long exact hypercohomology sequence associated to a short exact sequence of complexes of sheaves.)



**Higher Jacobi–Zariski spectral sequence:** Let  $k \rightarrow A$  be a homomorphism of rings, let  $X$  be a Noetherian scheme over  $A$  of finite Krull dimension, let  $M$  be a quasi-coherent  $\mathcal{O}_X$ -module, and fix  $i \geq 0$ . Then there is a natural bounded spectral sequence of  $A$ -modules  $E_{pq}^1 \implies D_{p+q}^i(X|k, M)$  whose columns may be described as follows.

- (i) Suppose that  $p < -i$  or  $p > 0$ . Then  $E_{pq}^1 = 0$ .
- (ii) Suppose that  $-i \leq p \leq 0$ . Then the  $p$ th column of the  $E^1$ -page is given by a bounded spectral sequence

$$\mathcal{E}_{\alpha\beta}^2 = D_{\alpha}^{-p}(A|k, D_{\beta}^{i+p}(X|A, M)) \implies E_{p,\alpha+\beta-p}^1.$$

(Note that  $\mathcal{E}_{\alpha\beta}^2 = 0$  if  $\alpha$  or  $\beta$  is  $< -\dim X$ .)

(Proof: In the case that  $X$  is affine, this is due to Kassel and Sletsjøe [18]. Examining the proof in the affine case reveals that  $\mathbb{L}_{X|k}^i$  has a filtration with graded pieces  $\mathrm{gr}^p \mathbb{L}_{X|k}^i \simeq \mathbb{L}_{A|k}^p \otimes_A \mathbb{L}_{X/A}^{i-p}$  for  $p = 0, \dots, i$ . Tensoring by  $M$ , and applying the variations on Deligne’s spectral sequence for a filtered complex given in the Remark 2.1, one easily obtains the desired spectral sequence.)

**Example 2.2.** Suppose that  $X$  is smooth over  $k$ , or more generally that  $X$  has an affine open cover by the spectra of geometrically regular  $k$ -algebras. Then  $\mathbb{L}_{X|k}^i \simeq \Omega_{X|k}^i$ , and so  $D_n^i(X|k) \cong H^{-n}(X, \Omega_{X|k}^i)$  for all  $n \in \mathbb{Z}, i \geq 0$ .

This completes our discussion of André–Quillen homology for schemes.

## 2.2. Formal function properties for AQ homology of schemes

Now that the basic properties of André–Quillen homology for schemes have been established, we show that it satisfies formal function properties akin to Grothendieck’s formal functions theorem. We begin with a lemma in the affine case.

**Lemma 2.3.** *Let  $A \rightarrow B$  be an essentially finite type morphism of Noetherian rings,  $I \subseteq B$  an ideal, and  $M$  a finitely generated  $B$ -module. Then the canonical map*

$$\{D_n^i(B|A, M) \otimes_B B/I^r\} \longrightarrow \{D_n^i(B|A, M/I^r M)\}_r$$

*is an isomorphism for all  $n, i \geq 0$ .*

**Proof.** Given another  $B$ -module  $N$ , there is a natural first quadrant spectral sequence

$$E_{pq}^2 = \mathrm{Tor}_p^B(D_q^i(B|A, M), N) \implies D_{p+q}^i(B|A, M \otimes_B N).$$

This is proved in the usual way by choosing a resolution of  $N$  by flat  $B$ -modules and applying the Kunnetth spectral sequence for a tensor product of complexes.

Applying this with  $N = B/I^r$  gives spectral sequences

$$E_{pq}^2(r) = \mathrm{Tor}_p^B(D_q^i(B|A, M), B/I^r) \implies D_{p+q}^i(B|A, M/I^r M),$$

and taking the limit over  $r$  yields a first quadrant spectral sequence of pro  $B$ -modules:

$$E_{pq}^2(\infty) = \{\mathrm{Tor}_p^B(D_q^i(B|A, M), B/I^r)\}_r \implies \{D_{p+q}^i(B|A, M/I^r M)\}_r.$$

Our hypotheses on  $A, B$  and  $M$  ensure that the  $B$ -modules  $D_q^i(B|A, M)$  are finitely generated, whence Theorem 1.1(i) implies that  $E_{pq}^2(\infty) = 0$  for  $p > 0$ . We thus obtain edge map isomorphisms  $\{D_n^i(B|A, M) \otimes_B B/I^r\} \xrightarrow{\cong} \{D_n^i(B|A, M/I^r M)\}$ , as desired.  $\square$

In the following results, a sheaf of ideals  $\mathcal{I} \subseteq \mathcal{O}_X$  plays a prominent role; to keep the notation clear, we write  $rY$  for the  $r$ th infinitesimal thickening of the closed subscheme defined by  $\mathcal{I}$ , i.e.,

$$rY := \mathrm{Spec} \mathcal{O}_X/\mathcal{I}^r.$$

**Theorem 2.4.** *Let  $A$  be a Noetherian ring and  $\pi : X \rightarrow \mathrm{Spec} A$  a Noetherian scheme over  $A$  of finite Krull dimension; let  $\mathcal{I} \subseteq \mathcal{O}_X$  be an ideal sheaf and  $M$  a coherent  $\mathcal{O}_X$ -module. Then the following hold.*

- (i) *The canonical map  $\{D_n^i(X|A, M/\mathcal{I}^r M)\}_r \longrightarrow \{D_n^i(rY|A, M/\mathcal{I}^r M)\}_r$  is an isomorphism for all  $n \in \mathbb{Z}, i \geq 0$ .*
- (ii) *Assuming that  $\pi$  is essentially of finite type, the canonical map of pro sheaves  $\{D_n^i(X|A, M) \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r\}_r \longrightarrow \{D_n^i(X|A, M/\mathcal{I}^r M)\}_r$  is an isomorphism for all  $n \in \mathbb{Z}, i \geq 0$ .*

Now, suppose further that  $\pi$  is proper and that  $\mathcal{I} = I\mathcal{O}_X$  for some ideal  $I \subseteq A$ . Then the following hold.

- (iii)  *$D_n^i(X|A, M)$  is a finitely generated  $A$ -module for all  $n \in \mathbb{Z}, i \geq 0$ .*
- (iv) *The canonical map  $\{D_n^i(X|A, M) \otimes_A A/I^r\}_r \longrightarrow \{D_n^i(X|A, M/\mathcal{I}^r M)\}_r$  is an isomorphism for all  $n \in \mathbb{Z}, i \geq 0$ .*

**Proof.** (i) By the Mayer–Vietoris long exact descent sequence, and the usual two inductions starting with an affine open cover of  $X$  (note that  $X$  is quasi-separated, since it is Noetherian), this claim reduces to the case that  $X$  is affine, in which case it is Theorem 1.2(ii).

(ii) Assume that  $\pi$  is essentially of finite type. This claim also reduces to the affine case, which is Lemma 2.3, but since this is the only occurrence of pro sheaves, we provide a few more details. Fixing  $n \in \mathbb{Z}$  and  $i \geq 0$ , let  $K_r$  and  $C_r$  respectively denote the kernel and cokernel of the canonical map

$$\mathcal{D}_n^i(X|A, M) \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r \longrightarrow \mathcal{D}_n^i(X|A, M/\mathcal{I}^r M).$$

Let  $\{U_i\}$  be a finite cover of  $X$  by affine opens. Then, given  $r \geq 1$  and applying the previous lemma, there exists  $s \geq r$  such that the maps  $K_s(U_i) \rightarrow K_r(U_i)$  and  $C_s(U_i) \rightarrow C_r(U_i)$  are zero; since these are quasi-coherent sheaves, it follows that the maps  $K_s \rightarrow K_r$  and  $C_s \rightarrow C_r$  are also zero, whence  $\{K_r\}_r$  and  $\{C_r\}_r$  vanish, completing the proof.

For the rest of the proof, we assume that  $\pi$  is proper and that  $\mathcal{I} = I\mathcal{O}_X$  for some ideal  $I \subseteq A$ .

(iii) Each  $\mathcal{O}_X$ -module  $\mathcal{D}_n^i(X|A, M)$  is coherent, since  $X$  is of finite type over  $A$ . Since  $X$  is proper over  $A$ , this implies that each of the cohomology groups of  $\mathcal{D}_n^i(X|A, M)$  is

a finitely generated  $A$ -module; so the claim follows from the hypercohomology spectral sequence.

(iv) As explained in the proof of (iii), the hypercohomology spectral sequence

$$E_2^{pq} = H^p(X, \mathcal{D}_{-q}^i(X|A, M)) \implies D_{-p-q}^i(X|A, M)$$

consists of finitely generated  $A$ -modules. Since  $\{- \otimes_A A/I^r\}_r$  is an exact functor on the category of finitely generated  $A$ -modules, by Theorem 1.1(ii), we may apply it to the spectral sequence to obtain a spectral sequence of pro  $A$ -modules

$$E_2^{pq}(\infty) = \{H^p(X, \mathcal{D}_{-q}^i(X|A, M)) \otimes_A A/I^r\}_r \implies \{D_{-p-q}^i(X|A, M) \otimes_A A/I^r\}_r.$$

This spectral sequence maps to the limit of the hypercohomology spectral sequences for the  $\mathcal{O}_X$ -modules  $M/\mathcal{I}^r M$ , namely

$$'E_2^{pq}(\infty) = \{H^p(X, \mathcal{D}_{-q}^i(X|A, M/\mathcal{I}^r M))\}_r \implies \{D_{-p-q}^i(X|A, M/\mathcal{I}^r M)\}_r,$$

and so to complete the proof it is enough to show that the canonical maps  $E_2^{pq}(\infty) \rightarrow 'E_2^{pq}(\infty)$  are all isomorphisms.

First, since  $\mathcal{D}_{-q}^i(X|A, M)$  is a coherent sheaf and  $\pi$  is proper, Grothendieck's formal functions theorem<sup>1</sup>[13, Corollary 4.1.7] states that the canonical map

$$E_2^{pq}(\infty) = \{H^p(X, \mathcal{D}_{-q}^i(X|A, M)) \otimes_A A/I^r\}_r \longrightarrow \{H^p(X, \mathcal{D}_{-q}^i(X|A, M) \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r)\}_r$$

is an isomorphism. Second, part (ii) implies that the canonical map

$$\{H^p(X, \mathcal{D}_{-q}^i(X|A, M) \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r)\}_r \longrightarrow \{H^p(X, \mathcal{D}_{-q}^i(X|A, M/\mathcal{I}^r M))\}_r = 'E_2^{pq}(\infty)$$

is also an isomorphism, completing the proof. □

It is part (iv) of the previous theorem which will be so essential in § 2.3; a secondary application of it is the following corollaries, which will not be needed but which most closely mimic Grothendieck's formal functions theorem for coherent cohomology.

**Corollary 2.5.** *Let  $A$  be a Noetherian ring,  $X$  a proper scheme over  $A$  of finite Krull dimension,  $I \subseteq A$  an ideal, and  $M$  a coherent  $\mathcal{O}_X$ -module. Then the canonical map*

$$\{D_n^i(X|A, M) \otimes_A A/I^r\}_r \longrightarrow \{D_n^i(rY|A/I^r, M/I^r M)\}_r$$

*is an isomorphism for all  $n \in \mathbb{Z}$ ,  $i \geq 0$ , where  $rY = X \times_A A/I^r$ .*

**Proof.** Using parts (iv) and (i) of Theorem 2.4, it remains only to prove that the canonical map  $\{D_n^i(rY|A, M/I^r M)\}_r \rightarrow \{D_n^i(rY|A/I^r, M/I^r M)\}_r$  is an isomorphism. From the higher Jacobi–Zariski spectral sequence, this follows from the vanishing of  $\{D_n^i(A/I^r|A)\}_r$ ; the details are as follows.

Applying the higher Jacobi–Zariski spectral sequences to the homomorphism  $A \rightarrow A/I^r$  and  $A/I^r$ -scheme  $rY$ , we obtain a bounded spectral sequence of pro  $A$ -modules

<sup>1</sup>This theorem is normally stated as the isomorphism  $\varprojlim_r H^p(X, N) \otimes_A A/I^r \xrightarrow{\sim} \varprojlim_r H^p(X, N/\mathcal{I}^r N)$  for any coherent  $\mathcal{O}_X$ -module  $N$ . But a quick examination of the cited proof in EGA shows that the stronger isomorphism of pro  $A$ -modules  $\{H^p(X, N) \otimes_A A/I^r\}_r \xrightarrow{\sim} \{H^p(X, N/\mathcal{I}^r N)\}_r$  also holds.

$E_{pq}^1(\infty) \implies \{D_{p+q}^i(rY|A, M/I^r M)\}_r$  which is supported in the range  $-i \leq p \leq 0$ , where it is described by bounded right-half-plane spectral sequences

$$\mathcal{E}_{\alpha\beta}^2(\infty) = \{D_{\alpha}^{-p}(A/I^r|A, D_{\beta}^{i+p}(rY|A/I^r, M/I^r M))\}_r \implies E_{p,\alpha+\beta-p}^1(\infty).$$

For any  $r, \alpha > 0$ , Theorem 1.2(i) may be applied to the ideal  $I^r$  and module  $D_{\beta}^{i+p}(rY|A/I^r, M/I^r M)$  to deduce that there exists  $s > 1$  such that the second of the following maps, and hence the composition, is zero:

$$\begin{aligned} D_{\alpha}^{-p}(A/I^{sr}|A, D_{\beta}^{i+p}(srY|A/I^{sr}, M/I^{sr} M)) &\longrightarrow D_{\alpha}^{-p}(A/I^s|A, D_{\beta}^{i+p}(rY|A/I^r, M/I^r M)) \\ &\longrightarrow D_{\alpha}^{-p}(A/I^r|A, D_{\beta}^{i+p}(rY|A/I^r, M/I^r M)). \end{aligned}$$

That is,  $\mathcal{E}_{\alpha\beta}^2(\infty) = 0$  for  $\alpha \neq 0$ . Also,  $\mathcal{E}_{0\beta}^2(\infty) = 0$  for  $p \neq 0$ . So the  $\mathcal{E}$ -spectral sequences degenerate to edge map isomorphisms

$$E_{p,n-p}^1(\infty) \cong \begin{cases} 0 & p \neq 0, \\ D_n^i(rY|A/I^r, M/I^r M) & p = 0. \end{cases}$$

Thus the  $E$ -spectral sequence degenerates to the desired edge map isomorphisms. □

**Corollary 2.6** (Formal functions theorem for André–Quillen homology). *Under the hypotheses of the previous corollary, the canonical map*

$$D_n^i(X|A, M)^\wedge \longrightarrow \varprojlim_r D_n^i(rY|A/I^r, M/I^r M)$$

is an isomorphism for all  $n \in \mathbb{Z}, i \geq 0$ , where  $D_n^i(X|A, M)^\wedge$  denotes the  $I$ -adic completion of  $D_n^i(X|A, M)$ .

**Proof.** Take  $\varprojlim_r$  of the pro isomorphism of the previous corollary. □

**Remark 2.7.** Let  $A$  be a Noetherian ring,  $X$  a proper scheme over  $A$  of finite Krull dimension, and  $I \subseteq A$  an ideal with respect to which  $A$  is assumed to be adically complete. Applying Corollary 2.6 to  $M = \mathcal{O}_X$ , and noting that  $D_n^i(X|A)$  is a finitely generated  $A$ -module, and hence already  $I$ -adically complete, we obtain

$$D_n^i(X|A) \xrightarrow{\cong} \varprojlim_r D_n^i(rY|A/I^r).$$

### 2.3. Pro descent for AQ homology with respect to blow-up squares

With the formal function theorems for André–Quillen homology established, we may now prove that André–Quillen homology satisfies the desired pro Mayer–Vietoris property with respect to abstract blow-up squares. We begin by developing further the results of Theorem 2.4 in the case that  $\pi$  is an isomorphism away from  $V(I)$ .

**Lemma 2.8.** *Let  $k \rightarrow A$  be a homomorphism of Noetherian rings,  $I \subseteq A$  an ideal, and  $X$  a proper scheme over  $A$  of finite Krull dimension such that the induced morphism*

$X \setminus V(I\mathcal{O}_X) \rightarrow \text{Spec } A \setminus V(I)$  is an isomorphism. Then the following canonical maps are isomorphisms for all  $n \in \mathbb{Z}, i \geq 0$ .

- (i)  $\{D_n^i(X|A, I^r\mathcal{O}_X)\}_r \longrightarrow \{I^r D_n^i(X|A)\}_r \stackrel{(*)}{\cong} 0$   
 (vanishing  $(*)$  not necessarily valid if  $i = n = 0$ ).
- (ii)  $\{D_n^i(A|k, I^r)\} \longrightarrow \{D_n^i(X|k, I^r\mathcal{O}_X)\}_r$ .

**Proof.** (i) The short exact sequences  $0 \rightarrow I^r\mathcal{O}_X \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X/I^r\mathcal{O}_X \rightarrow 0$  induce a long exact sequence of pro  $A$ -modules

$$\dots \longrightarrow \{D_n^i(X|A, I^r\mathcal{O}_X)\}_r \longrightarrow D_n^i(X|A) \longrightarrow \{D_n^i(X|A, \mathcal{O}_X/I^r\mathcal{O}_X)\}_r \longrightarrow \dots$$

According to Theorem 2.4(iv),  $\{D_n^i(X|A, \mathcal{O}_X/I^r\mathcal{O}_X)\}_r \cong \{D_n^i(X|A) \otimes_A A/I^r\}_r$ , from which it follows that the long exact sequences breaks into short sequences and induce the desired isomorphisms  $\{D_n^i(X|A, I^r\mathcal{O}_X)\}_r \xrightarrow{\cong} \{I^r D_n^i(X|A)\}_r$ .

Regarding the vanishing claim, suppose first that  $i = 0$ ; then  $D_n^0(X|A) = H^{-n}(X, \mathcal{O}_X)$ , which is a finitely generated  $A$ -module (since  $X$  is proper over  $A$ ) which is supported on  $V(I)$  when  $n \neq 0$  (since  $X \setminus V(I\mathcal{O}_X) \cong \text{Spec } A \setminus V(I)$ ), and hence is killed by a power of  $I$ . Next, suppose that  $i > 0$ ; in this case, the vanishing claim merely requires that  $X$  be essentially of finite type over  $A$ ; induction on the size of an affine open cover of  $X$  reduces us to proving that if  $B$  is an essentially finite type  $A$ -algebra then  $D_n^i(B|A)$  is killed by a power of  $I$ . But  $D_n^i(B|A)$  is a finitely generated  $B$ -module supported on  $V(IB)$ , since André–Quillen homology behaves well under localization; thus it is killed by a power of  $I$ , completing the proof of all the vanishing claims.

(ii) The higher Jacobi–Zariski spectral sequence states that for each  $r \geq 1$  there is a natural, third quadrant, bounded spectral sequence

$$E_{pq}^1(r) \implies D_{p+q}^i(X|k, I^r\mathcal{O}_X)$$

which vanishes outside  $-i \leq p \leq 0$ , and whose columns in the range  $-i \leq p \leq 0$  are described by natural, first quadrant spectral sequences

$$\mathcal{E}_{\alpha\beta}^2(r) = D_{\alpha}^{-p}(A|k, D_{\beta}^{i+p}(X|A, I^r\mathcal{O}_X)) \implies E_{p, \alpha+\beta-p}^1(r).$$

According to part (i),  $\{\mathcal{E}_{\alpha\beta}^2(r)\}_r = 0$  unless  $\beta = 0$  and  $p = -i$ , whence the limit of the spectral sequences collapse to edge map isomorphisms

$$\{D_n^i(A|k, D_0^0(X|A, I^r\mathcal{O}_X))\}_r \xrightarrow{\cong} \{D_n^i(X|k, I^r\mathcal{O}_X)\}_r.$$

Since  $\{D_0^0(X|A, I^r\mathcal{O}_X)\}_r \cong \{I^r D_0^0(X|A)\}_r = \{I^r H^0(X, \mathcal{O}_X)\}_r$  by part (i) and the usual description of André–Quillen homology when  $i = 0$ , we will have completed the proof as soon as we show that the canonical map

$$\{I^r\}_r \longrightarrow \{I^r H^0(X, \mathcal{O}_X)\}_r$$

is an isomorphism of pro  $A$ -modules. Write  $B := H^0(X, \mathcal{O}_X)$ , which is a finite  $A$ -algebra with the property that  $A_{\mathfrak{p}} \xrightarrow{\cong} B \otimes_A A_{\mathfrak{p}}$  for each prime ideal  $\mathfrak{p} \subseteq A$  not containing  $I$ ; hence

the kernel  $K$  and cokernel  $C$  of the structure map  $A \rightarrow B$  are killed by a power of  $I$ . Now, consider the following commutative diagram of pro  $A$ -modules:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & K & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \\
 & & \downarrow (1) & & \downarrow (2) & & \downarrow (3) & & \downarrow (4) & & \\
 0 & \longrightarrow & \{K \otimes_A A/I^r\}_r & \longrightarrow & \{A/I^r\}_r & \longrightarrow & \{B \otimes_A A/I^r\}_r & \longrightarrow & \{C \otimes_A A/I^r\}_r & \longrightarrow & 0
 \end{array}$$

The top row is an exact sequence of finitely generated  $A$ -modules, whence the bottom row is an exact sequence of pro  $A$ -modules by Theorem 1.1(ii). Since  $K$  and  $C$  are killed by a power of  $I$ , maps (1) and (4) are isomorphisms, and so the central square in the diagram is bicartesian. It follows that maps (2) and (3) have isomorphic kernels; i.e.,  $\{I^r\}_r \xrightarrow{\sim} \{I^r B\}_r$ , as required. □

**Corollary 2.9.** *Let  $k$  be a Noetherian ring, let  $X' \rightarrow X$  be a proper morphism between Noetherian  $k$ -schemes of finite Krull dimension, and suppose that  $\mathcal{I} \subseteq \mathcal{O}_X$  is an ideal sheaf such that the induced morphism  $X' \setminus V(\mathcal{I}\mathcal{O}_{X'}) \rightarrow X \setminus V(\mathcal{I})$  is an isomorphism. Then the canonical map*

$$\{D_n^i(X|k, \mathcal{I}^r)\}_r \longrightarrow \{D_n^i(X'|k, \mathcal{I}^r\mathcal{O}_{X'})\}_r$$

is an isomorphism for all  $n \in \mathbb{Z}$ ,  $i \geq 0$ .

**Proof.** By the usual two inductions starting with an affine open cover of  $X$  (since  $X$  is Noetherian, and hence quasi-separated) we may assume that  $X = \text{Spec } A$  is affine, in which case the statement is precisely Lemma 2.8(ii) (with  $X$  rechristened  $X'$ ). □

We are now equipped to prove our main theorem about André–Quillen homology of schemes, namely that it satisfies the pro Mayer–Vietoris property with respect to abstract blow-up squares. Recall that an *abstract blow-up square* of schemes is a pull-back diagram

$$\begin{array}{ccc}
 Y' & \longrightarrow & X' \\
 \downarrow & & \downarrow \\
 Y & \longrightarrow & X
 \end{array}$$

where  $X' \rightarrow X$  is proper,  $Y \rightarrow X$  is a closed embedding, and the induced map on the open subschemes  $X' \setminus Y' \rightarrow X \setminus Y$  is an isomorphism.

To state our descent result, we need the following piece of notation which we did not introduce in Remark 2.1 when discussing hypercohomology: if  $M^\bullet$  is a cochain complex of sheaves, and  $M^\bullet \xrightarrow{\sim} \mathcal{F}^\bullet$  is a quasi-isomorphism to a cochain complex of acyclic sheaves, then set

$$\mathbb{H}(X, M^\bullet) := \mathcal{F}^\bullet(X),$$

which is well defined up to quasi-isomorphism and satisfies  $H^*(\mathbb{H}(X, M^\bullet)) = \mathbb{H}^*(X, M^\bullet)$  by definition.

The following theorem is true more generally with a coherent module  $M$  in place of  $\mathcal{O}_X$ , but we omit it to simplify the notation.

**Theorem 2.10** (Pro descent for AQ homology with respect to abstract blow-ups). *Let  $k$  be a Noetherian ring, and let*

$$\begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

*be an abstract blow-up square of Noetherian, finite Krull-dimensional  $k$ -schemes. Then the following square of cochain complexes of pro  $k$ -modules is homotopy Cartesian*

$$\begin{array}{ccc} \mathbb{H}(X, \mathbb{L}_{X|k, \text{neg}}^i) & \longrightarrow & \mathbb{H}(X', \mathbb{L}_{X'|k, \text{neg}}^i) \\ \downarrow & & \downarrow \\ \{\mathbb{H}(rY, \mathbb{L}_{rY|k, \text{neg}}^i)\}_r & \longrightarrow & \{\mathbb{H}(rY', \mathbb{L}_{rY'|k, \text{neg}}^i)\}_r \end{array}$$

*resulting in a long exact Mayer–Vietoris sequence of pro  $k$ -modules*

$$\dots \longrightarrow D_n^i(X|k) \longrightarrow \{D_n^i(rY|k)\}_r \oplus \{D_n^i(X'|k)\}_r \longrightarrow D_n^i(rY'|k)_r \longrightarrow \dots$$

**Proof.** According to Theorem 2.4(i), the cohomologies of the complexes in the bottom row of the diagram are unchanged if we replace them by

$$\{\mathbb{H}(X, \mathbb{L}_{X|k, \text{neg}}^i \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r)\}_r \longrightarrow \{\mathbb{H}(X', \mathbb{L}_{X'|k, \text{neg}}^i \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{X'}/\mathcal{I}^r \mathcal{O}_{X'})\}_r,$$

where  $\mathcal{I}$  is the sheaf of ideals defining  $Y$ . Having made this replacement, we must show that the homotopy fibres of the two resulting vertical arrows

$$\begin{array}{ccc} \mathbb{H}(X, \mathbb{L}_{X|k, \text{neg}}^i) & & \mathbb{H}(X', \mathbb{L}_{X'|k, \text{neg}}^i) \\ \downarrow & & \downarrow \\ \{\mathbb{H}(X, \mathbb{L}_{X|k, \text{neg}}^i \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r)\}_r & & \{\mathbb{H}(X', \mathbb{L}_{X'|k, \text{neg}}^i \otimes_{\mathcal{O}_{X'}} \mathcal{O}_{X'}/\mathcal{I}^r \mathcal{O}_{X'})\}_r \end{array}$$

are quasi-isomorphic. These homotopy fibres are respectively

$$\{\mathbb{H}(X, \mathbb{L}_{X|k, \text{neg}}^i \otimes_{\mathcal{O}_X} \mathcal{I}^r)\}_r \quad \text{and} \quad \{\mathbb{H}(X', \mathbb{L}_{X'|k, \text{neg}}^i \otimes_{\mathcal{O}_{X'}} \mathcal{I}^r \mathcal{O}_{X'})\}_r,$$

between which the canonical map is indeed a quasi-isomorphism, since  $\{D_n^i(X|k, \mathcal{I}^r)\}_r \xrightarrow{\cong} \{D_n^i(X'|k, \mathcal{I}^r \mathcal{O}_{X'})\}_r$  for all  $n \in \mathbb{Z}, i \geq 0$ , by Corollary 2.9.  $\square$

### 3. Formal function properties and pro descent for $HH, HC, THH, TC$ , and $K$ -theory

Now, we extend the main results of § 2 to Hochschild and cyclic homology of schemes, and their topological counterparts, and then deduce that  $K$ -theory also satisfies the pro Mayer–Vietoris property with respect to abstract blow-up squares of varieties.

### 3.1. Hochschild and cyclic homology

Here, we discuss the theories of derived Hochschild and cyclic homology for schemes which will concern us.

Let  $k \rightarrow A$  be a homomorphism of commutative rings. Given an  $A$ -module  $M$ , we let  $H_*^{\text{naive},k}(A, M)$  denote the ‘usual’ Hochschild homology of  $A$  as a  $k$ -algebra with coefficients in  $M$ ; it is the homology of the Hochschild complex  $C_{\bullet}^k(A, M)$ . In particular,  $HH_*^{\text{naive},k}(A) = H_*^{\text{naive},k}(A, A)$  denotes the usual Hochschild homology of  $A$  as a  $k$ -algebra. However, we will work throughout with the derived version of Hochschild homology, for which we use the notation  $H_*^k(A, M)$ . That is, letting  $P_{\bullet} \rightarrow A$  be a simplicial resolution of  $A$  by free  $k$ -algebras,  $H_*^k(A, M)$  is defined to be the homology (of the diagonal) of the bisimplicial  $A$ -module

$$C_q^k(P_p, M) = M \otimes_k P_p^{\otimes kq}. \tag{p, q \ge 0}$$

In the special case when  $A = M$ , we write  $HH_*^k(A) = H_*^k(A, A)$ .

Next, we discuss cyclic homology. First,  $HC_*^{\text{naive},k}(A)$  denotes the usual cyclic homology of the  $k$ -algebra  $A$ , which is the homology of the cyclic bicomplex  $CC_{\bullet\bullet}^k(A)$ . Just as for Hochschild homology, we prefer to denote by  $HC_*^k(A)$  the derived version, defined as the homology of (the diagonal of) the simplicial tricocomplex  $CC_{\bullet\bullet}^k(P_{\bullet})$ , where  $P_{\bullet} \rightarrow A$  is a simplicial resolution of  $A$  by free  $k$ -algebras. The usual SBI sequence remains valid in the derived setting:

$$\dots \longrightarrow HH_n^k(A) \xrightarrow{I} HC_n^k(A) \xrightarrow{S} HC_{n-2}^k(A) \xrightarrow{B} \dots$$

Next, consider a Noetherian scheme  $X$  of finite Krull dimension over  $k$ . As in § 2.1, let  $\tilde{P}_{\bullet}^k(X)$  denote the simplicial sheaf of  $k$ -algebras obtained by degreewise sheafifying  $U \mapsto P_{\bullet}(\mathcal{O}_X(U))$ , where  $P_{\bullet}(\mathcal{O}_X(U)) \rightarrow \mathcal{O}_X(U)$  is a functorially chosen simplicial resolution by free  $k$ -algebras. Given a quasi-coherent  $\mathcal{O}_X$ -module  $M$ , set

$$C_{\bullet}^k(X, M) := M \otimes_k \tilde{P}_{\bullet}^k(X)^{\otimes k\bullet},$$

and let  $C_{\text{neg}}^k(X, M)$  be the cochain complex of quasi-coherent  $\mathcal{O}_X$ -modules, supported in negative degrees, obtained by negating the numbering of the chain complex of  $C_{\bullet}^k(X, M)$ . Define the derived Hochschild homology of  $X$ , relative to  $k$ , with coefficients in  $M$  to be its hypercohomology

$$H_*^k(X, M) := \mathbb{H}^{-*}(X, C_{\text{neg}}^k(X, M)).$$

In the special case when  $M = \mathcal{O}_X$ , we write  $C_{\bullet}^k(X) = C_{\bullet}^k(X, \mathcal{O}_X)$  and  $HH_*^k(X) = H_*^k(X, \mathcal{O}_X)$ . The obvious analogues of the ‘Agreement on affines’, ‘Mayer–Vietoris property’, ‘Hypercohomology spectral sequence’, and ‘Homology long exact sequence’ stated for André–Quillen homology in § 2.1 are true for this derived Hochschild homology of schemes.

We omit the analogous construction of derived cyclic homology for schemes, since the procedure should now be clear. As in the affine case, the usual SBI sequence remains valid:

$$\dots \longrightarrow HH_n^k(X) \xrightarrow{I} HC_n^k(X) \xrightarrow{S} HC_{n-2}^k(X) \xrightarrow{B} \dots$$

The relationships between André–Quillen homology, derived Hochschild/cyclic homology, and usual Hochschild/cyclic homology are summarized by the next lemma.



**Lemma 3.1.** *Let  $k$  be a ring,  $X$  a Noetherian scheme of finite Krull dimension over  $k$ , and  $M$  a quasi-coherent  $\mathcal{O}_X$ -module. Then the following hold.*

(i) *If  $X$  is flat over  $k$ , then the canonical maps*

$$H_n^k(X, M) \rightarrow H_n^{\text{naive},k}(X, M) \quad \text{and} \quad HC_n^k(X) \rightarrow HC_n^{\text{naive},k}(X)$$

*are isomorphisms for all  $n \in \mathbb{Z}$ , where  $H^{\text{naive},k}$  and  $HC^{\text{naive},k}$  denote the ‘usual’ Hochschild and cyclic homology of a scheme, as constructed in, for example, [34].*

(ii) *There is a bounded spectral sequence of  $k$ -modules*

$$E_{pq}^2 = D_p^q(X|k, M) \implies H_{p+q}^k(X, M).$$

**Proof.** We briefly explain the proofs of these standard results in the affine case when  $X = \text{Spec } A$ , after which the case of a scheme should be clear. (i) If  $A$  is flat over  $k$  then the map  $M \otimes_k P_{\bullet}^{\otimes kq} \rightarrow M \otimes_k A^{\otimes kq}$  is a quasi-isomorphism for all  $q \geq 0$ , and so the diagonal of the bisimplicial  $A$ -module  $C_{\bullet}^k(P_{\bullet}, M)$  is quasi-isomorphic to  $C_{\bullet}^k(A, M)$ . (ii) This is the spectral sequence associated to the bisimplicial  $A$ -module  $C_{\bullet}^k(P_{\bullet}, M)$ , since  $C_{\bullet}^k(P_p, M)$  has homology  $\Omega_{P_p/k}^* \otimes_{P_p} M$  for each  $p \geq 0$ . □

The following formal function results for Hochschild homology are the analogues of those given for André–Quillen homology in § 2.

**Theorem 3.2.** *Let  $k \rightarrow A$  be a homomorphism of Noetherian rings and  $\pi : X \rightarrow \text{Spec } A$  a Noetherian scheme over  $A$  of finite Krull dimension; let  $\mathcal{I} \subseteq \mathcal{O}_X$  be an ideal sheaf and  $M$  a coherent  $\mathcal{O}_X$ -module. Then the following hold.*

(i) *The canonical map  $\{H_n^A(X, M/\mathcal{I}^r M)\}_r \rightarrow \{H_n^A(rY, M/\mathcal{I}^r M)\}_r$  is an isomorphism for all  $n \in \mathbb{Z}$ .*

(ii) *Assuming that  $\pi$  is essentially of finite type, the canonical map of pro sheaves  $\{\mathcal{H}_n^A(X, M) \otimes_{\mathcal{O}_X} \mathcal{O}_X/\mathcal{I}^r\}_r \rightarrow \{\mathcal{H}_n^A(X, M/\mathcal{I}^r M)\}_r$  is an isomorphism for all  $n \in \mathbb{Z}$ .*

*Now, suppose further that  $\pi$  is proper and that  $\mathcal{I} = I\mathcal{O}_X$  for some ideal  $I \subseteq A$ . Then the following hold.*

(iii)  *$H_n^A(X, M)$  is a finitely generated  $A$ -module for all  $n \in \mathbb{Z}$ ,*

*and the following canonical maps are isomorphisms for all  $n \in \mathbb{Z}$ .*

(iv)  $\{H_n^A(X, M) \otimes_A A/I^r\}_r \rightarrow \{H_n^A(X, M/\mathcal{I}^r M)\}_r$ .

(v)  $\{H_n^A(X, M) \otimes_A A/I^r\}_r \rightarrow \{H_n^{A/I^r}(rY, M/\mathcal{I}^r M)\}_r$ .

*Now suppose still further that the induced morphism  $X \setminus V(I\mathcal{O}_X) \rightarrow \text{Spec } A \setminus V(I)$  is an isomorphism. Then the following canonical maps are isomorphisms for all  $n \in \mathbb{Z}$ .*

(vi)  $\{H_n^A(X, I^r \mathcal{O}_X)\}_r \rightarrow \{I^r HH_n^A(X)\}_r \stackrel{(*)}{\cong} 0$   
*(vanishing  $(*)$  not necessarily valid if  $n = 0$ ).*

(vii)  $\{H_n^k(A, I^r)\}_r \rightarrow \{H_n^k(X, I^r \mathcal{O}_X)\}_r$ .

**Proof.** Thanks to the André–Quillen to Hochschild homology spectral sequence for schemes, i.e., Lemma 3.1(ii), these claims immediately reduce to the analogous assertions for André–Quillen homology, which we have already proved. For the sake of reference, (i)–(iv) correspond to Theorem 2.4, (v) to Corollary 2.5, and (vi) and (vii) to Lemma 2.8. □

Given a closed embedding  $Y \rightarrow X$  of Noetherian schemes of finite Krull dimension, we will write  $HH_*^k(X, Y)$ ,  $HC_*^k(X, Y)$  for the relative derived Hochschild and cyclic homology groups, fitting into long exact sequences

$$\begin{aligned} \dots &\longrightarrow HH_n^k(X, Y) \longrightarrow HH_n^k(X) \longrightarrow HH_n^k(Y) \longrightarrow \dots \\ \dots &\longrightarrow HC_n^k(X, Y) \longrightarrow HC_n^k(X) \longrightarrow HC_n^k(Y) \longrightarrow \dots \end{aligned}$$

Now, we prove the analogue of Theorem 2.10 for  $HH$  and  $HC$ , namely that they also satisfy the pro Mayer–Vietoris property for abstract blow-up squares.

**Theorem 3.3** (Pro descent for  $HH$  and  $HC$  with respect to abstract blow-ups). *Let  $k$  be a Noetherian ring, and let*

$$\begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

*be an abstract blow-up square of Noetherian, finite Krull-dimensional  $k$ -schemes. Then the canonical maps*

$$\{HH_n^k(X, rY)\}_r \longrightarrow \{HH_n^k(X', rY')\}_r, \quad \{HC_n^k(X, rY)\}_r \longrightarrow \{HC_n^k(X', rY')\}_r$$

*are isomorphisms of pro abelian groups for all  $n \in \mathbb{Z}$ .*

**Proof.** The claim for  $HH$  is equivalent to the statement that the following square of cochain complexes of pro  $k$ -modules is homotopy Cartesian, where  $\mathbb{H}$  is the hypercohomology replacement functor introduced immediately preceding Theorem 2.10:

$$\begin{array}{ccc} \mathbb{H}(X, C_{\text{neg}}^k(X)) & \longrightarrow & \mathbb{H}(X', C_{\text{neg}}^k(X')) \\ \downarrow & & \downarrow \\ \{\mathbb{H}(rY, C_{\text{neg}}^k(rY))\}_r & \longrightarrow & \{\mathbb{H}(rY', C_{\text{neg}}^k(rY'))\}_r \end{array}$$

This is proved exactly as it was for André–Quillen homology in Theorem 2.10, using the appropriate parts of Theorem 3.2. More precisely, by Theorem 3.2(i), we may replace the bottom of the diagram by

$$\{\mathbb{H}(X, C_{\text{neg}}^k(X, \mathcal{O}_X/\mathcal{I}^r))\}_r \longrightarrow \{\mathbb{H}(X, C_{\text{neg}}^k(X', \mathcal{O}_{X'}/\mathcal{I}^r \mathcal{O}_{X'}))\}_r$$

without changing the cohomologies of the complexes, where  $\mathcal{I}$  is the sheaf of ideals defining  $Y$ . Then the map between the homotopy fibres of the vertical arrows becomes

$$\{\mathbb{H}(X, C_{\text{neg}}^k(X, \mathcal{I}^r))\}_r \longrightarrow \{\mathbb{H}(X, C_{\text{neg}}^k(X', \mathcal{I}^r \mathcal{O}_{X'}))\}_r,$$

which is a quasi-isomorphism by part Theorem 3.2(vii) (to be precise, by the  $HH$  analogue of Corollary 2.9, which follows from Theorem 3.2(vii) by induction on the size of an affine open cover of the base). This proves the claim for  $HH$ .

To pass to  $HC$ , use the five lemma and induction up the limit of the SBI sequences

$$\begin{array}{ccccccc} \cdots & \longrightarrow & HH_n^k(X, rY) & \longrightarrow & HC_n^k(X, rY) & \longrightarrow & HC_{n-2}^k(X, rY) & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & HH_n^k(X', rY') & \longrightarrow & HC_n^k(X', rY') & \longrightarrow & HC_{n-2}^k(X', rY') & \longrightarrow & \cdots \end{array}$$

(note that it is possible to start the induction since the SBI sequences vanish in degrees  $< -\max\{\dim X, \dim X'\}$ ). □

### 3.2. Topological Hochschild and cyclic homology

The analogues for topological Hochschild and cyclic homology of the formal function properties given in Theorem 3.2 were recently proved in joint work with Dundas [10]; from this we will quickly prove that these theories also satisfy the pro Mayer–Vietoris property with respect to abstract blow-up squares.

We will assume that the reader is familiar with topological Hochschild and cyclic homology, and so only offer a brief summary. For a quasi-compact quasi-separated scheme  $X$ , the spectra  $THH(X)$ ,  $TR^m(X; p)$ , and  $TC^m(X; p)$  were defined by Geisser and Hesselholt in [11] in such a way that all these presheaves of non-connective spectra satisfy Zariski descent (see the proof of [11, Corollary 3.3.3]). They were shown in [10] to have reasonable finiteness and continuity properties under the assumption of F-finiteness, where we fix a prime number  $p$ .

**Definition 3.4.** A  $\mathbb{Z}_{(p)}$ -algebra is said to be *F-finite* if and only if the  $\mathbb{F}_p$ -algebra  $A/pA$  is finitely generated over its subring of  $p$ th-powers. A  $\mathbb{Z}_{(p)}$ -scheme is said to be F-finite if and only if it admits a finite open cover by the spectra of F-finite  $\mathbb{Z}_{(p)}$ -algebras.

In particular, given a Noetherian F-finite  $\mathbb{Z}_{(p)}$ -algebra  $A$ , the ring of truncated  $p$ -typical Witt vectors  $W_m(A)$  is again a Noetherian F-finite  $\mathbb{Z}_{(p)}$ -algebra; see [10, § 2] for a detailed discussion of such issues. Recall that  $TR_n^m(A; p)$  is naturally a  $W_m(A)$ -algebra.

The following establishes that topological Hochschild homology and the related theories all have the pro Mayer–Vietoris property for abstract blow-up squares.

**Theorem 3.5** (Pro descent for  $THH$ ,  $TR^m$ ,  $TC^m$  with respect to abstract blow-ups). *Let*

$$\begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

be an abstract blow-up square of Noetherian,  $F$ -finite, finite Krull-dimensional  $\mathbb{Z}_{(p)}$ -schemes. Then the canonical maps

$$\begin{aligned} \{THH_n(X, rY; \mathbb{Z}/p^v)\}_r &\longrightarrow \{THH_n(X', rY'; \mathbb{Z}/p^v)\}_r \\ \{TR_n^m(X, rY; p, \mathbb{Z}/p^v)\}_r &\longrightarrow \{TR_n^m(X', rY'; p, \mathbb{Z}/p^v)\}_r \\ \{TC_n^m(X, rY; p, \mathbb{Z}/p^v)\}_r &\longrightarrow \{TC_n^m(X', rY'; p, \mathbb{Z}/p^v)\}_r \end{aligned}$$

are isomorphisms of pro abelian groups for all  $n \in \mathbb{Z}$  and  $m, v \geq 1$ .

**Proof.** Since  $THH = TR^1$ , and since  $TC^m = \text{holim}(TR^m \xrightarrow{1-F} TR^{m-1})$ , it is sufficient to prove the claim for  $TR^m$ . By Zariski descent, we may assume that  $X = \text{Spec } A$  is affine.

That is,  $A$  is a Noetherian  $F$ -finite  $\mathbb{Z}_{(p)}$ -algebra,  $I \subseteq A$  is an ideal,  $\pi : X' \rightarrow \text{Spec } A$  is a proper morphism which induces an isomorphism  $X' \setminus V(I\mathcal{O}_X) \rightarrow \text{Spec } A \setminus V(I)$ , and we must prove that the square of pro spectra

$$\begin{array}{ccc} TR^m(A; p, \mathbb{Z}/p^v) & \longrightarrow & TR^m(X'; p, \mathbb{Z}/p^v) \\ \downarrow & & \downarrow \\ \{TR^m(A/I^r; p, \mathbb{Z}/p^v)\}_r & \longrightarrow & \{TR^m(X' \times_A A/I^r; p, \mathbb{Z}/p^v)\}_r \end{array} \tag{†}$$

is homotopy Cartesian. By [10, Theorem 5.6], the hypotheses imply that the canonical maps

$$\begin{aligned} \{TR_n^m(A; p, \mathbb{Z}/p^v) \otimes_{W_m(A)} W_m(A/I^r)\}_r &\longrightarrow \{TR_n^m(A/I^r; p, \mathbb{Z}/p^v)\}_r \\ \{TR_n^m(X'; p, \mathbb{Z}/p^v) \otimes_{W_m(A)} W_m(A/I^r)\}_r &\longrightarrow \{TR_n^m(X' \times_A A/I^r; p, \mathbb{Z}/p^v)\}_r \end{aligned}$$

are isomorphisms of pro  $W_m(A)$ -modules for all  $n \in \mathbb{Z}$ . Hence the vertical arrows in (†) induce surjections at the level of the homotopy groups, and so it is sufficient to show that the induced horizontal map on the kernels, namely

$$\{W_m(I^r) TR_n^m(A; p, \mathbb{Z}/p^v)\}_r \longrightarrow \{W_m(I^r) TR_n^m(X'; p, \mathbb{Z}/p^v)\}_r,$$

is an isomorphism for all  $n \in \mathbb{Z}$ . The rest of the proof is devoted to proving this isomorphism.

First, since  $TR_n^m$  behaves well under localization (see Lemma 5.1 of [10] and its proof), the canonical map

$$TR_n^m(X'; p, \mathbb{Z}/p^v) \otimes_{W_m(A)} W_m(A_f) \longrightarrow TR_n^m(X' \times_A A_f; p, \mathbb{Z}/p^v)$$

is an isomorphism for any  $f \in A$ . In particular, if  $f \in I$  whence  $X' \times_A A_f \cong \text{Spec } A_f$ , then the canonical map

$$TR_n^m(A; p, \mathbb{Z}/p^v) \otimes_{W_m(A)} W_m(A_f) \longrightarrow TR_n^m(X'; p, \mathbb{Z}/p^v) \otimes_{W_m(A)} W_m(A_f)$$

is an isomorphism. Since the rest of the proof is commutative algebra, write  $M := TR_n^m(A; p, \mathbb{Z}/p^v)$  and  $N := TR_n^m(X'; p, \mathbb{Z}/p^v)$ ; these are finitely generated as  $W_m(A)$ -modules, by [10, Theorem 5.3], and we have just proved that the map  $M \rightarrow N$  becomes an isomorphism after applying  $-\otimes_{W_m(A)} W_m(A_f)$  for any  $f \in I$ .

Second, we claim that the spectra  $\text{Spec } W_m(A_f)$ ,  $f \in I$ , form an open affine cover of  $\text{Spec } W_m(A) \setminus V(W_m(I))$ . This follows from the following two facts:

$$W_m(A_f) = W_m(A)_{[f]},$$

$$W_m(I)^M \subseteq \langle \text{ideal of } W_m(A) \text{ generated by } [f] : f \in I \rangle \text{ for } M \gg 0,$$

where  $[f] \in W_m(A)$  denotes the Teichmüller lift of an element  $f \in A$ . Both these facts are relatively standard results about Witt rings; for example, see [29, Lemma A.6(i)] and [10, Lemma 2.1 & Remark 2.2], respectively.

It now follows from commutative algebra that the kernel and cokernel of the map  $M \rightarrow N$  are killed by a power of  $W_m(I)$ . So, by the same argument as used at the end of the proof of Lemma 2.8, the induced map  $\{W_m(I)^r M\}_r \rightarrow \{W_m(I)^r N\}_r$  is an isomorphism. Finally, to complete the proof, note that the chains of ideals  $W_m(I)^r$  and  $W_m(I')^r$  are intertwined, by [10, Lemma 2.1]. □

### 3.3. Pro cdh-descent for K-theory

We now prove the main theorem of the paper, namely that  $K$ -theory satisfies the pro Mayer–Vietoris property with respect to abstract blow-up squares of varieties.

In the interest of not being restricted to varieties over a field, we prefer to work in the generality of Noetherian quasi-excellent schemes whenever possible. Recall that a Noetherian scheme  $X$  is called quasi-excellent if and only if the formal fibre of each point of  $X$  is geometrically regular and the regular locus of any finite type  $X$ -scheme is open. All ‘naturally occurring’ schemes in algebraic geometry are quasi-excellent, and the notion is intimately related to resolution of singularities: according to Grothendieck [14, Proposition 7.9.5], if  $X$  is a Noetherian scheme with the property that every integral finite type  $X$ -scheme admits a desingularization, then  $X$  is quasi-excellent. If  $X$  is furthermore assumed to be a  $\mathbb{Q}$ -scheme, then the converse is true by Hironaka [16] and Temkin [31, Theorem 2.3.6].

The following refinement of the Haesemeyer argument is the key to understanding Noetherian quasi-excellent  $\mathbb{Q}$ -schemes; I am grateful to Haesemeyer for discussions about this.

**Proposition 3.6.** *Let  $\mathcal{E}$  be a presheaf of spectra on the category of Noetherian quasi-excellent  $\mathbb{Q}$ -schemes with the following properties:  $\mathcal{E}$  is invariant under nilpotent extensions, satisfies Nisnevich descent, and satisfies the Mayer–Vietoris property for blow-ups along regularly embedded centres. Then  $\mathcal{E}$  satisfies the Mayer–Vietoris property for all abstract blow-up squares.*

**Proof.** To simplify the explanation, we begin by indicating an abstract blow-up square of Noetherian quasi-excellent  $\mathbb{Q}$ -schemes and its value under  $\mathcal{E}$ :

$$(1) \quad \begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array} \qquad (2) \quad \begin{array}{ccc} \mathcal{E}(X) & \longrightarrow & \mathcal{E}(X') \\ \downarrow & & \downarrow \\ \mathcal{E}(Y) & \longrightarrow & \mathcal{E}(Y') \end{array}$$

According to the Haesemeyer argument [15, §§ 5, 6], axiomized into the form in which we are using it in [4, Theorem 3.12], our assumptions apply that square (2) is homotopy Cartesian as soon as square (1) consists of finite type  $k$ -schemes for some characteristic-zero field  $k$ . However, examination of Haesemeyer’s original argument reveals that the only required properties of  $k$  are that it be infinite and have strong resolution of singularities in the sense specified in [15, Theorem 2.4]. But Hironaka [16] proved strong resolution of singularities over any Noetherian quasi-excellent local ring  $A$  with characteristic-zero residue field; so Haesemeyer’s argument works in this generality, and we deduce that square (2) is homotopy Cartesian whenever square (1) consists of finite type  $A$ -schemes, for such a ring  $A$ .

Finally, let square (1) consist of arbitrary Noetherian quasi-excellent  $\mathbb{Q}$ -schemes. Since  $\mathcal{E}$  satisfies Zariski descent, square (2) is homotopy Cartesian as soon as the squares

$$\begin{array}{ccc}
 \mathcal{E}(\mathrm{Spec} \mathcal{O}_{X,x}) & \longrightarrow & \mathcal{E}(X' \times_X \mathrm{Spec} \mathcal{O}_{X,x}) \\
 \downarrow & & \downarrow \\
 \mathcal{E}(Y \times_X \mathrm{Spec} \mathcal{O}_{X,x}) & \longrightarrow & \mathcal{E}(Y' \times_X \mathrm{Spec} \mathcal{O}_{X,x})
 \end{array}$$

are homotopy Cartesian for all points  $x \in X$ . But, since  $A = \mathcal{O}_{X,x}$  is a Noetherian quasi-excellent local ring with characteristic-zero residue field, these squares are indeed homotopy Cartesian by the previous paragraph. □

We have reached the main theorem of the paper.

**Theorem 3.7** (Pro descent for  $K$ -theory with respect to abstract blow-ups). *Let*

$$\begin{array}{ccc}
 Y' & \longrightarrow & X' \\
 \downarrow & & \downarrow \\
 Y & \longrightarrow & X
 \end{array}$$

be an abstract blow-up square of either

- (i) Noetherian quasi-excellent  $\mathbb{Q}$ -schemes of finite Krull dimension; or
- (ii) finite type schemes over an infinite perfect field  $k$  which has strong resolution of singularities.

Then the canonical map

$$\{K_n(X, rY)\}_r \longrightarrow \{K_n(X', rY')\}_r$$

of pro abelian relative  $K$ -groups is an isomorphism for all  $n \in \mathbb{Z}$ .

**Proof.** (i) For any quasi-compact quasi-separated  $\mathbb{Q}$ -scheme  $Z$ , let  $HN^{\mathbb{Q}}(Z)$  and  $HP^{\mathbb{Q}}(Z)$  denote its negative and periodic cyclic homologies over  $\mathbb{Q}$ , as formulated in [4]. The infinitesimal  $K$ -theory of  $Z$  is defined to be the homotopy fibre of the Chern character from  $K$ -theory to negative cyclic homology, i.e.,

$$K^{\mathrm{inf}}(Z) := \mathrm{hofib}(K(Z) \xrightarrow{ch} NH^{\mathbb{Q}}(Z)).$$

According to [4, Theorem 4.6] and [4, Corollary 3.13], or rather their improvements using Proposition 3.6, the presheaves of spectra  $K^{\text{inf}}$  and  $HP$  carry any abstract blow-up square of Noetherian quasi-excellent  $\mathbb{Q}$ -schemes to a homotopy Cartesian square of spectra; so the canonical maps of relative spectra

$$K^{\text{inf}}(X, rY) \longrightarrow K^{\text{inf}}(X', rY'), \quad HP^{\mathbb{Q}}(X, rY) \longrightarrow HP^{\mathbb{Q}}(X', rY')$$

are weak equivalences for all  $r \geq 1$ .

From the long exact sequence  $\cdots \rightarrow HN_n^{\mathbb{Q}} \rightarrow HP_n^{\mathbb{Q}} \rightarrow HC_{n-2}^{\mathbb{Q}} \rightarrow \cdots$ , Theorem 3.3 with  $k = \mathbb{Q}$ , and the usual five lemma argument, we may now deduce that the canonical map  $\{HN_n^{\mathbb{Q}}(X, rY)\}_r \rightarrow \{HN_n^{\mathbb{Q}}(X', rY')\}_r$  is an isomorphism for all  $n \in \mathbb{Z}$ . Applying the same argument to the long exact sequence  $\cdots \rightarrow K_n^{\text{inf}} \rightarrow K_n \rightarrow NH_n \rightarrow \cdots$ , we deduce that  $\{K_n(X, rY)\}_r \rightarrow \{K_n(X', rY')\}_r$  is an isomorphism for all  $n \in \mathbb{Z}$ , as desired.

(ii) We may evidently assume that  $p := \text{char } k$  is non-zero, since otherwise the claim is covered by case (i). The infinitesimal  $K$ -theory  $\{K^{\text{inf},m}(Z)\}_m$  of a finite type  $k$ -scheme  $Z$  is defined to be the pro spectrum arising as the homotopy fibres of the trace map from  $K$ -theory to topological cyclic homology, i.e.,

$$K^{\text{inf},m}(Z) := \text{hofib}(K(Z) \xrightarrow{tr} TC^m(Z; p)).$$

According to [12, Theorem B] (see also [20]), the square of pro spectra

$$\begin{array}{ccc} \{K^{\text{inf},m}(X)\}_m & \longrightarrow & \{K^{\text{inf},m}(X')\}_m \\ \downarrow & & \downarrow \\ \{K^{\text{inf},m}(Y)\}_m & \longrightarrow & \{K^{\text{inf},m}(Y')\}_m \end{array}$$

is homotopy Cartesian. Therefore our claim for  $K$ -theory follows, in a similar way to characteristic zero, from the analogous assertion for topological cyclic homology, namely Theorem 3.3 (note that the hypotheses of Theorem 3.3 are satisfied for finite type  $k$ -schemes, and that for such schemes the finite coefficients can be eliminated everywhere; see [10, Corollary 5.9] for related discussion).  $\square$

**Remark 3.8.** There are three special cases in which Theorem 3.7 is already known or can be otherwise deduced.

- (i) If  $Y \rightarrow X$  is a regular immersion and  $X'$  is the blow-up of  $X$  along  $Y$ , then it follows from Thomason’s blow-up formula [32] that there are short exact sequences of pro abelian groups

$$0 \longrightarrow K_n(X) \longrightarrow \{K_n(rY)\}_r \oplus K_n(X') \longrightarrow \{K_n(rY')\}_r \longrightarrow 0,$$

which is a stronger statement than the conclusion of Theorem 3.7.

- (ii) If  $X' \rightarrow X$  is a finite morphism, then Theorem 0.1 reduces to pro excision, which was established for arbitrary commutative Noetherian rings in [24].
- (iii) If  $X$  is a Cohen–Macaulay variety over an infinite field with only isolated singularities,  $Y = X_{\text{sing}}$ , and  $X' \rightarrow X$  is a resolution of singularities, then a technique of Weibel [35] using minimal reduction ideals implies that  $X'$  may be

obtained by first blowing up  $X$  along a regular immersion and then normalizing. So, in this case, Theorem 3.7 reduces to combining cases (i) and (ii). This case of desingularizing a Cohen–Macaulay variety with isolated singularities was first proved by Krishna [21, Proof of Theorem 1.2] in characteristic zero.

The validity of these special cases of Theorem 3.7 was our motivation for establishing it in full generality. It is a commonly encountered issue in cdh-descent problems that typical resolutions of singularities cannot be factored into regularly embedded blow-ups and normalizations; see, for example, [15, e.g., 6.3]. Since the Haesemeyer argument of Proposition 3.6 is not valid here (the pro spectra depend not only on the given schemes, but also on the embeddings), there appears to be no way to assemble the aforementioned cases (i)–(iii) into a proof of Theorem 3.7.

#### 4. Applications of Theorem 3.7

As mentioned in the introduction, the main applications of Theorem 3.7, which are to zero cycles on singular varieties, are presented in an accompanying paper [25]. Here, we offer some more straightforward applications to demonstrate its use, first by reformulating it as a definition of  $K$ -theory with compact support, and then using it to give a quick proof of parts of Weibel’s  $K$ -dimension conjecture on negative  $K$ -theory.

For simplicity, we adopt the following conventions in this section.

A field  $k$  will be called *good* if and only if it is infinite, perfect, and has strong resolution of singularities; for example,  $\text{char } k = 0$  suffices.

A  $k$ -variety means simply a finite type  $k$ -scheme; further assumptions will be specified when required, and the reference to  $k$  will occasionally be omitted.

##### 4.1. $K$ -theory with compact support

Here, we explain how Theorem 3.7 may be interpreted as the well-definedness of ‘ $K$ -theory with compact support’; I am grateful to H. Gillet for suggesting this interpretation to me. In this section,  $k$  is a good field.

By Nagata, any separated  $k$ -variety  $X$  has a compactification  $\bar{X}$ , i.e.,  $\bar{X}$  is proper over  $k$  and  $X$  is a dense open subset of  $\bar{X}$ ; we refer the reader to [22] for the proof.

**Definition 4.1.** Let  $X$  be a separated  $k$ -variety. Let  $\bar{X}$  be any compactification of  $X$ , and set  $Y = \bar{X} \setminus X$ , equipped with any structure as a closed subscheme. The  $K$ -theory of  $X$  with compact support is defined to be the spectrum

$$K^c(X) := \text{holim}_r K(\bar{X}, rY),$$

and the  $K$ -groups of  $X$  with compact support are defined to be its homotopy groups.

The content of the following proposition is equivalent to taking homotopy limits in Theorem 3.7 in the case of separated  $k$ -varieties.

**Proposition 4.2.** *The previous definition does not depend on the chosen compactification  $\bar{X}$  of  $X$ .*



**Proof.** Suppose that  $\bar{X}_i$ ,  $i = 1, 2$ , are different compactifications of  $X$ ; then, by [22, Corollary 2.4], there exists a third compactification  $\bar{X}_3$  which is a blow-up of both  $\bar{X}_1$  and  $\bar{X}_2$ . Let  $Y_1, Y_2, Y_3$  be the complements of  $X$  in the compactifications. Then, the diagram

$$\begin{array}{ccc} Y_3 & \longrightarrow & X_3 \\ \downarrow & & \downarrow \\ Y_i & \longrightarrow & X_i \end{array}$$

is an abstract blow-up square for  $i = 1, 2$ , and so Theorem 3.7 implies that the maps

$$\{K_n(\bar{X}_1, rY_1)\}_r \longrightarrow \{K_n(\bar{X}_3, rY_3)\}_r \longleftarrow \{K_n(\bar{X}_2, rY_2)\}_r$$

are isomorphisms for all  $n \in \mathbb{Z}$ . Hence the maps

$$\text{holim}_r K(\bar{X}_1, rY_1) \longrightarrow \text{holim}_r K(\bar{X}_3, rY_3) \longleftarrow \text{holim}_r K(\bar{X}_2, rY_2)$$

are weak equivalences, as required. □

**Remark 4.3.** If  $\bar{X}$  is a proper variety containing  $X$  as an open, but not necessarily dense, subscheme, then

$$\begin{array}{ccc} Y' := X' \setminus X & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y := \bar{X} \setminus X & \longrightarrow & \bar{X} \end{array}$$

is an abstract blow-up square, where  $X'$  denotes the closure of  $X$  inside  $\bar{X}$ . So Theorem 3.7 implies that  $K^c(X) \simeq \text{holim}_r K(\bar{X}, rY)$ . In other words,  $K^c(X)$  may be defined with respect to any proper variety containing  $X$  as an open subset: it is not necessary that  $X$  be dense. This will be implicitly used in the next result.

To justify its definition as a theory with compact support, we show that  $K^c$  fits into a localization sequence.

**Proposition 4.4.** *Let  $Y \rightarrow X$  be a closed embedding of separated  $k$ -varieties, with open complement  $U \subseteq X$ . Then there is a functorial homotopy fibre sequence*

$$K^c(U) \longrightarrow K^c(X) \longrightarrow \text{holim}_r K^c(rY).$$

**Proof.** The proof is a chase of the definitions in which some care is required regarding infinitesimal thickenings.

First, we may suppose that  $Y$  is reduced since that does not affect its system of thickenings in  $X$ . Let  $\bar{X}$  be a compactification of  $X$ , and set  $Z := \bar{X} \setminus X$  and  $W := \bar{X} \setminus U$ , with their reduced subscheme structure to obtain closed embeddings  $Z \rightarrow W \rightarrow \bar{X}$ . Then there is a resulting commutative diagram of homotopy fibre sequences

$$\begin{array}{ccccc} K(\bar{X}, r_{\bar{X}}Z) & \longrightarrow & K(\bar{X}) & \longrightarrow & K(r_{\bar{X}}Z) \\ \downarrow & & \parallel & & \downarrow \\ K(\bar{X}, r_{\bar{X}}W) & \longrightarrow & K(\bar{X}) & \longrightarrow & K(r_{\bar{X}}W) \end{array}$$

in which the notations  $r_{\bar{X}}Z = rZ$  and  $r_{\bar{X}}W = rW$  are to remind the reader and author that the infinitesimal thickenings are taken inside  $\bar{X}$ . Taking  $\text{holim}_r$  and homotopy cofibres of the vertical maps yields a homotopy fibre sequence

$$K^c(U) \longrightarrow K^c(X) \longrightarrow \text{holim}_r K(r_{\bar{X}}W, r_{\bar{X}}Z).$$

Next, we claim that the open complement of the closed embedding  $r_{\bar{X}}Z \rightarrow r_{\bar{X}}W$  is precisely  $r_X Y$ , where this latter infinitesimal thickening is taken inside  $X$ ; this follows from the identifications

$$r_{\bar{X}}Z \times_{\bar{X}} X = r_X(Z \times_{\bar{X}} X) = r_X Y.$$

Hence  $K^c(r_X Y) = \text{holim}_s K^c(r_{\bar{X}}W, s(r_{\bar{X}}Z))$ , where we are taking the  $s$ th infinitesimal thickening of  $r_{\bar{X}}Z$  inside  $r_{\bar{X}}W$ . To complete the proof, it remains to check that the canonical map

$$\text{holim}_{r,s} K(r_{\bar{X}}W, s(r_{\bar{X}}Z)) \longrightarrow \text{holim}_r K(r_{\bar{X}}W, r_{\bar{X}}Z)$$

is a weak equivalence. This is true, simply because the inverse systems on each side are intertwined. (This is easiest to see in the affine case, when  $\bar{X} = \text{Spec } A$ ,  $W = \text{Spec } A/J$ , and  $Z = \text{Spec } A/I$  with  $I \supseteq J$ ; then the canonical map of interest can be written as  $\text{holim}_{r,s} K(A/J^r, (J^r + I^{sr})/J^r) \rightarrow \text{holim}_r K(A/J^r, I^r/J^r)$ , in which the two systems are intertwined because the transition map  $K(A/J^{sr}, I^{sr}/J^{sr}) \rightarrow K(A/J^r, I^r/J^r)$  factors through  $K(A/J^r, (J^r + I^{sr})/J^r)$ .) □

**Remark 4.5.** We finish this section with two remarks about this  $K$ -theory with compact support.

- (i) More generally, suppose that  $\pi : X \rightarrow S$  is a separated morphism of  $k$ -varieties. Picking a proper morphism  $\bar{X} \rightarrow S$  of  $k$ -varieties compactifying  $\pi$ , and setting  $Y := \bar{X} \setminus X$ , we may define the  $K$ -theory with compact support of the family  $\pi$  as

$$K^c(X/S) := \text{holim}_r K(\bar{X}, rY).$$

Repeating the proofs of Propositions 4.2 and 4.4 verbatim, we see that this is well defined and satisfies localization.

- (ii) The construction in (i) remains valid for any separated morphism of Noetherian quasi-excellent  $\mathbb{Q}$ -schemes of finite Krull dimension. For example, if  $\text{char } k = 0$  and  $X$  is a smooth proper variety over  $k((t))$ , then

$$K^c(X/k[[t]]) := \text{holim}_r K(\bar{X}, rY)$$

is a well-defined invariant of  $X$ , where  $\bar{X} \rightarrow \text{Spec } k[[t]]$  is any proper model of  $X$ , with special fibre  $Y := \bar{X} \times_{k[[t]]} k$ . It seems likely that  $K_0^c(X/k[[t]])$  is related to nearby cycles.

### 4.2. Negative $K$ -groups

The following conjecture concerning negative  $K$ -theory was raised by Weibel in 1980 [36, Qu. 2.9].

**$K$ -dimension conjecture:** If  $X$  is a Noetherian scheme of dimension  $d$ , then  $K_n(X) = 0$  for  $n < d$ , and  $X$  is  $K_{-d}$ -regular.

The conjecture was proved in [4] for varieties over characteristic-zero fields, where it was also shown that  $K_{-d}$  of any such scheme is a finitely generated group (this is not explicitly stated, but is a straightforward consequence of [4, Theorem 0.2(1)]), which had been proved earlier for mild types of singularities in [15, § 8].

The vanishing part of the conjecture was proved in [12] and [20] for varieties over good fields of finite characteristic.

Our aim in this section is to show that Theorem 3.7 offers a very quick proof of the vanishing part of the  $K$ -dimension conjecture, and of the finite generation of  $K_{-d}$ , for the following types of scheme:

- (i) Noetherian quasi-excellent  $\mathbb{Q}$ -schemes; and
- (ii) varieties over good fields of finite characteristic.

We do not consider the  $K_{-d}$ -regularity part of the conjecture.

The following standard lemma reduces the problems to reduced schemes.

**Lemma 4.6.** *If  $X$  is a Noetherian scheme of dimension  $d$ , then the canonical map  $K_n(X) \rightarrow K_n(X_{\text{red}})$  is an isomorphism for  $n \leq -d$ .*

**Proof.** Given a nilpotent ideal  $I$  of a ring  $R$ , the canonical map  $K_n(R) \rightarrow K_n(R/I)$  is an isomorphism for  $n = 0$ , and hence for  $n < 0$ . So the canonical map from the descent spectral sequence for  $X$ , namely

$$E_2^{pq} = H^p(X, \mathcal{K}_{-q, X}) \implies K_{-p-q}(X),$$

to that for  $X_{\text{red}}$ ,

$${}_{\text{red}}E_2^{pq} = H^p(X_{\text{red}}, \mathcal{K}_{-q, X_{\text{red}}}) \implies K_{-p-q}(X_{\text{red}}),$$

is an isomorphism on the  $E_2$  page whenever  $q \geq 0$ . The claim easily follows from this.  $\square$

**Theorem 4.7.** *Let  $X$  be a  $d$ -dimensional scheme which is either*

- (i) *a Noetherian quasi-excellent  $\mathbb{Q}$ -scheme; or*
- (ii) *a variety over a good field.*

*Then  $K_n(X) = 0$  for  $n < -d$ , and the group  $K_{-d}(X)$  is finitely generated.*

**Proof.** The proof is by induction on  $d$ ; for any fixed  $d$ , it is enough to treat the case that  $X$  is reduced, thanks to Lemma 4.6.

First, if  $X$  is zero dimensional and reduced, then it is a disjoint union of the spectra of fields. So all the negative  $K$ -groups vanish, and  $K_0(X)$  is the free abelian group generated by the finitely many points of  $X$ .

The inductive step proceeds by picking a desingularization  $X' \rightarrow X$ ; in case (ii) this exists by assumption, while in case (i) it exists by Temkin’s extension [31, Theorem 2.3.6] of Hironaka’s resolution theorem. That is, there exists an abstract blow-up square

$$\begin{array}{ccc} Y' & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

in which  $X'$  is regular, and  $Y'$  and  $Y$  have dimension  $< d$ . By Theorem 3.7, there is a resulting long exact Mayer–Vietoris sequence of pro abelian groups

$$\cdots \longrightarrow K_n(X) \longrightarrow \{K_n(rY)\}_r \oplus K_n(X') \longrightarrow \{K_n(rY')\}_r \longrightarrow \cdots$$

But  $K_n(X') = 0$  for all  $n < 0$ , since  $X'$  is regular, and  $K_n(rY) = K_n(Y)$  and  $K_n(rY') = K_n(Y')$  for all  $n \leq -(d - 1)$  by Lemma 4.6; hence the Mayer–Vietoris sequence simplifies in degrees  $\leq -d$  to an exact sequence

$$K_{-(d-1)}(Y') \rightarrow K_{-d}(X) \rightarrow K_{-d}(Y) \rightarrow K_{-d}(Y') \rightarrow K_{-d-1}(X) \rightarrow K_{-d-1}(Y) \rightarrow \cdots .$$

But, by the inductive hypothesis,  $K_n(Y)$  and  $K_n(Y')$  vanish for  $n < -(d - 1)$ , and  $K_{-(d-1)}(Y')$  is finitely generated. This evidently completes the proof.  $\square$

**Acknowledgements.** I thank B. Dundas for the collaboration [10], without which Theorem 0.1 would be restricted to characteristic zero. The majority of this work was done while I was supported by the Simons foundation as a postdoctoral fellow at the University of Chicago, and I am grateful to the foundation for their support.

**References**

1. M. ANDRÉ, *Homologie des algèbres commutatives (Die Grundlehren der mathematischen Wissenschaften, Band 206)* (Springer, Berlin, 1974).
2. M. ARTIN AND B. MAZUR, *Étale Homotopy (Lecture Notes in Mathematics, Volume 100)* (Springer, Berlin, 1986). Reprint of the 1969 original.
3. G. CORTIÑAS, The obstruction to excision in  $K$ -theory and in cyclic homology, *Invent. Math.* **164**(1) (2006), 143–173.
4. G. CORTIÑAS, C. HAESEMEYER, M. SCHLICHTING AND C. WEIBEL, Cyclic homology,  $\text{cdh}$ -cohomology and negative  $K$ -theory, *Ann. of Math. (2)* **167**(2) (2008), 549–573.
5. G. CORTIÑAS, C. HAESEMEYER, M. E. WALKER AND C. WEIBEL, Bass’  $NK$  groups and  $\text{cdh}$ -fibrant Hochschild homology, *Invent. Math.* **181**(2) (2010), 421–448.
6. G. CORTIÑAS, C. HAESEMEYER, M. E. WALKER AND C. WEIBEL,  $K$ -theory of cones of smooth varieties, *J. Algebraic Geom.* **22**(1) (2013), 13–34.
7. G. CORTIÑAS, C. HAESEMEYER AND C. WEIBEL,  $K$ -regularity,  $\text{cdh}$ -fibrant Hochschild homology, and a conjecture of Vorst, *J. Amer. Math. Soc.* **21**(2) (2008), 547–561.
8. G. CORTIÑAS, C. HAESEMEYER AND C. WEIBEL, Infinitesimal cohomology and the Chern character to negative cyclic homology, *Math. Ann.* **344**(4) (2009), 891–922.
9. P. DELIGNE, Théorie de Hodge II, *Publ. Math. Inst. Hautes Études Sci.* **40** (1971), 5–57.
10. B. DUNDAS AND M. MORROW, Finite generation and continuity of topological Hochschild and cyclic homology (2014), [arXiv:1403.0534](https://arxiv.org/abs/1403.0534).
11. T. GEISSER AND L. HESSELHOLT, Topological cyclic homology of schemes, in *Algebraic K-Theory, Proceedings of an AMS-IMS-SIAM Summer Research Conference, Seattle, WA, USA, July 13–24, 1997*, pp. 41–87 (American Mathematical Society, Providence, RI, 1999).
12. T. GEISSER AND L. HESSELHOLT, On the vanishing of negative  $K$ -groups, *Math. Ann.* **348**(3) (2010), 707–736.
13. A. GROTHENDIECK, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I, *Publ. Math. Inst. Hautes Études Sci.* **11** (1961), 5–167.
14. A. GROTHENDIECK, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II, *Publ. Math. Inst. Hautes Études Sci.* **24** (1965), 5–231.
15. C. HAESEMEYER, Descent properties of homotopy  $K$ -theory, *Duke Math. J.* **125**(3) (2004), 589–620.

16. H. HIRONAKA, Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II, *Ann. of Math.* (2) **79** (1964), 109–203. *ibid.* (2) **79** (1964), 205–326.
17. D. C. ISAKSEN, Calculating limits and colimits in pro-categories, *Fund. Math.* **175**(2) (2002), 175–194.
18. C. KASSEL AND A. B. SLETSJØE, Base change, transitivity and Künneth formulas for the Quillen decomposition of Hochschild homology, *Math. Scand.* **70**(2) (1992), 186–192.
19. M. KERZ AND S. SAITO, Chow group of 0-cycles with modulus and higher dimensional class field theory (2013), [arXiv:1304.4400](https://arxiv.org/abs/1304.4400).
20. A. KRISHNA, On the negative *K*-theory of schemes in finite characteristic, *J. Algebra* **322**(6) (2009), 2118–2130.
21. A. KRISHNA, An Artin–Rees theorem in *K*-theory and applications to zero cycles, *J. Algebraic Geom.* **19**(3) (2010), 555–598.
22. W. LÜTKEBOHMERT, On compactification of schemes, *Manuscripta Math.* **80**(1) (1993), 95–111.
23. J. S. MILNE, *Étale Cohomology (Princeton Mathematical Series, Volume 33)* (Princeton University Press, Princeton, NJ, 1980).
24. M. MORROW, Pro unitality and pro excision in algebraic *K*-theory and cyclic homology (2014), [arXiv:1404.4179](https://arxiv.org/abs/1404.4179).
25. M. MORROW, Zero cycles on singular varieties and their desingularisations (2014), [arXiv:1404.4649](https://arxiv.org/abs/1404.4649).
26. D. QUILLEN, On the (co-) homology of commutative rings, in *Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968)*, pp. 65–87 (American Mathematical Society, Providence, RI, 1970).
27. D. G. QUILLEN, Homology of commutative rings, *Unpublished MIT Notes* (1968).
28. M. RONCO, On the Hochschild homology decompositions, *Comm. Algebra* **21**(12) (1993), 4699–4712.
29. K. RÜLLING, The generalized de Rham–Witt complex over a field is a complex of zero-cycles, *J. Algebraic Geom.* **16**(1) (2007), 109–169.
30. V. SRINIVAS, Zero cycles on a singular surface. II, *J. Reine Angew. Math.* **362** (1985), 4–27.
31. M. TEMKIN, Desingularization of quasi-excellent schemes in characteristic zero, *Adv. Math.* **219**(2) (2008), 488–522.
32. R. W. THOMASON, Les *K*-groupes d’un schéma éclaté et une formule d’intersection excédentaire, *Invent. Math.* **112**(1) (1993), 195–215.
33. V. VOEVODSKY, Homotopy theory of simplicial sheaves in completely decomposable topologies, *J. Pure Appl. Algebra* **214**(8) (2010), 1384–1398.
34. C. WEIBEL, Cyclic homology for schemes, *Proc. Amer. Math. Soc.* **124**(6) (1996), 1655–1662.
35. C. WEIBEL, The negative *K*-theory of normal surfaces, *Duke Math. J.* **108**(1) (2001), 1–35.
36. C. WEIBEL, *K*-theory and analytic isomorphisms, *Invent. Math.* **61**(2) (1980), 177–197.
37. C. WEIBEL, *An Introduction to Homological Algebra (Cambridge Studies in Advanced Mathematics, Volume 38)* (Cambridge University Press, Cambridge, 1994).
38. C. WEIBEL AND S. C. GELLER, Étale descent for Hochschild and cyclic homology, *Comment. Math. Helv.* **66**(3) (1991), 368–388.