

Probiotic therapy for the prevention and treatment of *Clostridium difficile*-associated diarrhea: a systematic review

Nandini Dendukuri, Vania Costa, Maurice McGregor, James M. Brophy

Abstract

Background: The recent increase in the number and severity of cases of nosocomial *Clostridium difficile*-associated diarrhea (CDAD) has prompted interest in the use of probiotics for the prevention and treatment of this disease. We performed a systematic review of randomized controlled trials to assess the effectiveness of probiotic therapy.

Methods: We searched the PubMed, EMBASE, INAHTA, HEN and Cochrane Collaboration databases to identify trials in which the prevention or treatment of CDAD with probiotic therapy was the primary or secondary outcome. We extracted data on the number of patients randomly assigned to receive probiotic or placebo, the number of patients with CDAD, the type of probiotic, criteria for diagnosing CDAD, persistence of infection after treatment, compliance and adverse effects.

Results: We identified 4 eligible studies in which prevention ($n = 1$) or treatment ($n = 3$) of CDAD was the primary outcome. The benefit of probiotic therapy seen in 2 of the studies was restricted to subgroups characterized by severe CDAD and increased use of vancomycin. The remaining 2 studies were too methodologically flawed for us to draw meaningful conclusions. We also identified 4 trials in which prevention of antibiotic-associated diarrhea with probiotics was the primary outcome and prevention of CDAD a secondary outcome. These studies were limited primarily by too few CDAD cases and provided no evidence of effective prophylaxis. Overall, heterogeneity in choice and dose of probiotic and in criteria for diagnosing CDAD makes it difficult to synthesize information from the 8 studies.

Interpretation: Studies conducted to date provide insufficient evidence for the routine clinical use of probiotics to prevent or treat CDAD. Better designed and larger studies are needed.

CMAJ 2005;173(2):167-70

Both the incidence and severity of *Clostridium difficile*-associated diarrhea (CDAD) have increased in hospitals across North America.¹⁻⁴ Recent outbreaks in Montreal and Sherbrooke resulted in a 4-fold increase in the number of CDAD cases as well as an increase in the number of attributable deaths.^{1,5-7} CDAD is a debilitating and costly illness, particularly among patients with recurring episodes. Probiotics, or naturally occurring “good bacteria,” have been suggested as a means of both preventing and treating the disease.⁸⁻¹¹ The disturbance of normal gastrointestinal flora, particularly after antibiotic use, is be-

lieved to predispose patients to colonization by *C. difficile*;² probiotics, by delivering bacteria to the gastrointestinal tract, are believed to restore equilibrium in the altered gastrointestinal flora and thus protect against colonization.⁸ Probiotics that have been proposed for prevention and treatment of antibiotic-associated diarrhea and CDAD include various bacteria (*Bifidobacterium*, *Lactobacillus* GG, *L. rhamnosus*, *L. casei*, *L. plantarum* 299v, *Enterococcus faecium* [SF68]) and yeasts (*Saccharomyces boulardii*, *S. cerevisiae*). They are commonly available as lyophilized capsules or in the form of a fermented drink. We performed a systematic review of randomized controlled trials of probiotic therapy for the prevention and treatment of CDAD.

Methods

We followed the guidelines for systematic reviews described in the QUORUM statement.¹² We searched the databases of PubMed, EMBASE, INAHTA, HEN and the Cochrane Collaboration up to Mar. 19, 2005, using the following search terms: (Probiotic, Probiotics, Lactobacillus, lactic-acid, acidophilus, casei, bulgaricus, plantarum, rhamnosus, yeast, Saccharomyces, boulardii, cervisiae, Bifidobacterium, bifidum, SF68) and (Clostridium, difficile, diarrhea, antibiotic-associated) and (patients, subjects). We selected randomized controlled trials where prevention or treatment of CDAD was either a primary or secondary outcome and where the study subjects were adults and recruited from a hospital setting. Further details of the database search strategy are given in online Appendix 1 (www.cmaj.ca/cgi/content/full/173/2/167/DC1). We also reviewed the reference lists from articles fulfilling the inclusion criteria.

Two of the authors independently extracted the following data from all studies meeting the inclusion criteria: the number of patients randomly assigned to probiotic and placebo groups, the number of patients with CDAD in each group, the type of probiotic, criteria for diagnosing CDAD, persistence of *C. difficile* infection after treatment, compliance and adverse effects. We calculated a risk difference and 95% confidence interval (CI) for comparing probiotic and placebo groups. CIs were calculated using the hybrid score method with continuity correction.¹³

The quality of each trial was assessed using the scale described by Jadad and associates,¹⁴ which rates studies on randomization, blinding and attrition. The score ranges from 0 to 5, with 5 indicating best quality. A more detailed summary of each study is given in a recent Health Technology Assessment report prepared by the Technology Assessment Unit of the McGill University Health Centre (“Probiotics,” available at www.mcgill.ca/tau/publications/2005/).

Results

The results of our search strategy are summarized in Fig. 1. We identified 4 studies in which prevention ($n = 1$)¹⁵ or treatment ($n = 3$)¹⁶⁻¹⁹ of CDAD was the primary outcome. No meta-analysis was performed because there was marked heterogeneity between the studies. We also identified 4 randomized, placebo-controlled trials of probiotic therapy for the prevention of antibiotic-associated diarrhea where prevention of CDAD was measured as a secondary outcome¹⁹⁻²² (Table 1; a longer version of this table is available online at www.cmaj.ca/cgi/content/full/1723/2/167/DC1). Study quality ranged from 3 to 5, with all the randomized studies being double-blinded.

As seen in Table 1, there was considerable protocol variation in the choice of probiotic formulation and duration of treatment. There was also variability in the criteria for defining a case of CDAD: various definitions of diarrhea and different combinations of tests for *C. difficile* and its associated toxins A and B were used. In some studies patients were considered to have CDAD if they had diarrhea and were positive for *C. difficile* as determined by at least 1 standard culture or toxin A or B test. This definition, although probably the most inclusive, has a low specificity because it classifies patients with nontoxigenic *C. difficile* as having CDAD.² In other studies, patients were tested for toxin presence only if a standard culture was positive for the bacillus. However, the poor sensitivity of standard culture means that subjects who had a false-negative result may not have been tested.




Fig. 1: Flow of articles in the search strategy. CDAD = *Clostridium difficile*-associated diarrhea, ADD = antibiotic-associated diarrhea. No new studies were found when the reference lists of eligible studies were reviewed.

In the single study on the effectiveness of probiotic therapy in preventing CDAD,¹⁵ no difference was found between probiotic and placebo groups in the percentage of patients who tested positive for *C. difficile* infection after treatment (Table 1). The authors also reported a 32% (statistically nonsignificant) decrease in the incidence of diarrhea among patients who were positive for infection after culture and toxin test, but this comparison is not appropriate since it is not between all subjects randomly assigned to the 2 groups.

Among the 3 trials of probiotic treatment of CDAD,¹⁶⁻¹⁹ only 1, by McFarland and associates,¹⁶ reported a significant beneficial effect overall. However, in a post hoc analysis, the authors found that this effect was almost entirely limited to a subgroup with recurrent CDAD (risk difference 30, 95% CI 2.3 to 50.6). These subjects were characterized by a greater use of vancomycin, although vancomycin use was not significantly associated with the outcome in a multivariate analysis. In the study by Surawicz and colleagues,¹⁷ a subgroup analysis of subjects receiving high-dose vancomycin, who were more likely to have severe CDAD, demonstrated a beneficial effect of the probiotic (risk difference 33, 95% CI -0.3 to 62.0). However, the CIs in the subgroup analyses from both studies were very wide, which made it difficult to ascertain the magnitude of the effect. The study by Wullt and associates¹⁸ was chiefly limited by the very small sample ($n = 21$).

In the 4 studies in which prevention of antibiotic-associated diarrhea was the primary outcome, probiotic therapy had no significant effect on the prevention of CDAD.¹⁹⁻²² In 1 study,²⁰ the only available data on CDAD involved a comparison that was not between randomly assigned subjects. Probiotic therapy was found to be beneficial in preventing antibiotic-associated diarrhea in only 2 of the studies.^{19,20} Within each study there was variability in the antibiotic therapy (type, dose and duration) used concomitantly with the probiotic.

No serious adverse effects or problems with compliance were found in the probiotic group in any of the studies.

Interpretation

We identified 8 eligible trials of probiotic therapy in which CDAD prevention or treatment was either a primary or secondary outcome. The scant literature does not provide convincing proof of the clinical benefit of probiotics. Interestingly, in 2 treatment studies^{16,17} a beneficial effect (with a wide CI) was observed in a subgroup of patients characterized by more severe disease. In the first study these subjects had had at least 1 previous episode of CDAD in the last year and were more likely to be taking vancomycin than metronidazole. In the second study the patients were taking high doses of vancomycin and were more likely to have colitis. In both studies the apparent beneficial effect of the probiotic was seen only in patients with severe CDAD. The third treatment study¹⁸ was too small for us to draw any meaningful conclusion. In all 4 studies where CDAD prevention or treatment was the pri-

mary outcome, there was no difference between probiotic and placebo groups in the persistence of *C. difficile* after the intervention.^{15–18} Studies in which prevention of CDAD was measured as a secondary outcome had small samples and varied definitions of CDAD. In 3 studies^{19,21,22} the duration of follow-up after antibiotic use was less than 6 weeks, which may be too short for CDAD to develop.²³

Some authors have suggested that probiotic therapy has not been found to be beneficial because the doses used were

too low.²² This may be supported by the fact that in some of the studies involving adults in which no benefit was found, the dose used was the same as that used in pediatric studies that reported positive results.²⁴ Another concern is that the probiotic preparations may have become nonviable after manufacture. The effectiveness of the probiotic depends on the number of viable bacteria at the time of use.²⁵ Several articles have reviewed commercially available probiotic products such as food supplements, yoghurt and fermented

Table 1: Summary of randomized controlled trials where prevention or treatment of *Clostridium difficile*-associated diarrhea (CDAD) was the primary or secondary outcome (abridged)*

Trial	Probiotic and duration	Outcome measure (length of follow-up)	Risk difference (95% CI) (placebo-probiotic)	Quality score†	Comments
Primary outcome					
Plummer et al, 2004; ¹⁵ prevention	<i>Lactobacillus acidophilus</i> and <i>Bifidobacterium bifidum</i> for 20 d	Presence of <i>C. difficile</i> (20 d)	-2.8 (-8.7 to 15.3)	3	No difference in toxin positivity between treatment arms among patients with positive culture; lower incidence of diarrhea reported among culture- and toxin-positive subjects in probiotic group; short follow-up
McFarland et al, 1994; ¹⁶ treatment	<i>Saccharomyces boulardii</i> for 4 wk	Diarrhea and at least 1 positive assay for <i>C. difficile</i> by culture or toxin A or B (8 wk)	20.5 (2.2 to 37.0) First-time CDAD: 4.9 (-17.6 to 26.4) Recurrent CDAD: 30.0 (2.3 to 50.6)	5	Concomitant use of different antibiotics for differing lengths of time; no difference in culture positivity between treatment arms, but significantly fewer toxin B positive cases in probiotic group
Surawicz et al, 2000; ¹⁷ treatment	<i>S. boulardii</i> for 28 d	Diarrhea and at least 1 positive assay for <i>C. difficile</i> by culture or toxin A or B (8 wk)	9.8 (-6.7 to 25.6) High-dose vancomycin: 33.0 (-0.3 to 62.0) Low-dose vancomycin: -6.4 (-16.2 to 28.1) Metronidazole: -1.9 (-25.8 to 29.2)	3	Concomitant antibiotic assigned to patients after randomization; no difference in culture or toxin positivity in subgroup taking high-dose vancomycin; results not reported for other groups
Wullt et al, 2003; ¹⁸ treatment	<i>L. plantarum</i> 299v for 38 d	Diarrhea (5–10 d) Positive assay for <i>C. difficile</i> toxins A or B (11–13 d)	-8.3 (-40.3 to 29.6) -19.4 (-64.7 to 15.5)	4	Small sample; no difference in culture positivity after treatment among 20 patients who had no diarrhea after 5–10 d
Secondary outcome					
Surawicz et al, 1989 ¹⁹	<i>S. boulardii</i> starting within 48 h of beginning antibiotic until 2 wk post-antibiotic	Acquired <i>C. difficile</i> after enrolment (duration of probiotic treatment; minimum of 8 d)	-13.3 (-25.7 to 4.7)	5	Probiotic resulted in a significant reduction in risk of AAD; <i>C. difficile</i> tested only in those with ≥ 3 stool samples; short follow-up
McFarland et al, 1995 ²⁰	<i>S. boulardii</i> starting within 72 h of beginning antibiotic until 3 d after last dose of antibiotic	Diarrhea among <i>C. difficile</i> - or toxin-positive subjects (7 wk after probiotic was discontinued)	-1.4 (-3.5 to 4.1)	4	Probiotic resulted in a significant reduction in risk of AAD; the only available result for CDAD was not based on a comparison between randomized subjects
Lewis et al, 1998 ²¹	<i>S. boulardii</i> for duration of antibiotic treatment	Positive assay for <i>C. difficile</i> toxin (duration of antibiotic treatment)	-6.8 (-22.6 to 8.0)	4	Probiotic had no effect on AAD; <i>C. difficile</i> tested regularly every 4 d and whenever subjects had diarrhea; small number of CDAD cases; short follow-up
Thomas et al, 2001 ²²	<i>L. GG</i> starting within 24 h of beginning antibiotic treatment for 14 d	Diarrhea and positive assay for <i>C. difficile</i> toxin (21 d)	0.7 (-3.0 to 4.5)	4	Probiotic had no effect on AAD; <i>C. difficile</i> toxin results obtained from medical chart; small number of CDAD cases; short follow-up

Note: CI = confidence interval, AAD = antibiotic-associated diarrhea.

*See the unabridged table at www.cmaj.ca/cgi/content/full/173/2/167/DC1.

†The quality of the studies was assessed using the scale described by Jadad et al.¹⁴ A score of 5 indicates best quality.

milk^{25,26} and found that the number of colony-forming units can be much lower than advertised and that the bacterial species listed on product labels are often incorrect. It is also possible that, in prevention trials, the probiotic does not survive to colonize the colon because it is eliminated by the acidity of the stomach or because of concomitant use of antibiotics. Only 1 of the studies we reviewed evaluated the presence of the probiotic in fecal samples.¹⁸ Studies are needed to establish the minimum effective adult dose. Although side effects appear to be rare, there has been a small number of case reports of fungemia and bacteremia attributed to probiotic use, particularly among immunocompromised patients.²⁷ These specific side effects were either not observed or not reported in the studies we reviewed. The varied definitions of CDAD also complicate the interpretation of these studies.

The studies conducted to date provide insufficient evidence for the routine clinical use of probiotics to prevent or treat CDAD in adults. Better designed trials are therefore needed. To demonstrate a prevention effect, studies would need to randomize a sufficiently large number of patients within categories of antibiotic regimens and have a sufficiently long follow-up after antibiotic use. Therapeutic studies would need to randomize patients within strata defined by the concomitant antibiotic regimen. In both types of studies a clearer rationale is needed for the type and dose of probiotic and for how CDAD is defined. Measuring the levels of probiotic bacteria in the stool of patients given different doses may help to determine acceptable dosing. The need for more rigorous proof of the efficacy of probiotics is particularly important given the increasing problem of nosocomial CDAD in Canada and the simultaneous growth in sales of over-the-counter probiotic products.

This article has been peer reviewed.

From the Technology Assessment Unit, McGill University Health Centre (Dendukuri, Costa, McGregor, Brophy); the Department of Medicine (McGregor, Brophy); and Department of Epidemiology and Biostatistics, McGill University, Montréal, Que. (Dendukuri)

Competing interests: None declared.

Contributors: Nandini Dendukuri, Vania Costa and Maurice McGregor were responsible for the literature searches and data extraction. Nandini Dendukuri, Maurice McGregor and James Brophy were responsible for interpreting the data and drafting the paper. Vania Costa was responsible for revising the paper. All of the authors approved the final version.

References

- Eggertson L. *C. difficile* hits Sherbrooke, Que., hospital: 100 deaths. *CMAJ* 2004;171(5):436.
- Poutanen SM, Simor AE. *Clostridium difficile*-associated diarrhea in adults. *CMAJ* 2004;171(1):51-8.
- Pepin J, Valiquette L, Alary ME, Villemure P, Pelletier A, Forget K, et al. *Clostridium difficile*-associated diarrhea in a region of Quebec from 1991 to 2003: a changing pattern of disease severity. *CMAJ* 2004;171(5):466-72.
- Archibald LK, Banerjee SN, Jarvis WR. Secular trends in hospital-acquired *Clostridium difficile* disease in the United States, 1987-2001. *J Infect Dis* 2004;189:1585-9.
- Valiquette L, Low DE, Pepin J, McGeer A. *Clostridium difficile* infection in hospitals: a brewing storm. *CMAJ* 2004;171(1):27-9.
- Loo VG, Libman MD, Miller MA, Bourgault AM, Frenette CH, Kelly M, et al. *Clostridium difficile*: a formidable foe. *CMAJ* 2004;171(1):47-8.
- Eggertson L, Sibbald B. Hospitals battling outbreaks of *C. difficile*. *CMAJ* 2004;171(1):19-21.
- Sullivan A, Nord CE. Probiotics in human infections. *J Antimicrob Chemother* 2002;50:625-7.
- Gill HS. Probiotics to enhance anti-infective defences in the gastrointestinal tract. *Best Pract Res Clin Gastroenterol* 2003;17:755-73.
- Elmer GW. Probiotics: living drugs. *Am J Health Syst Pharm* 2001;58:1101-9.
- Johnston BC, Vohra S. Treating *C. difficile*. *CMAJ* 2005;172(4):447-8.
- Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of reporting of meta-analyses. *Lancet* 1999;354:1896-900.
- Newcombe RG. Interval estimation for the difference between independent proportions: comparison of eleven methods. *Stat Med* 1998;17:873-90.
- Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1-12.
- Plummer S, Weaver MA, Harris JC, Dee P, Hunter J. *Clostridium difficile* pilot study: effects of probiotic supplementation on the incidence of *C. difficile* diarrhoea. *Int Microbiol* 2004;7:59-62.
- McFarland LV, Surawicz CM, Greenberg RN, Fekety R, Elmer GW, Moyer KA, et al. A randomized placebo-controlled trial of *Saccharomyces boulardii* in combination with standard antibiotics for *Clostridium difficile* disease. *JAMA* 1994;271:1913-8.
- Surawicz CM, McFarland LV, Greenberg RN, Rubin M, Fekety R, Mulligan ME, et al. The search for a better treatment for recurrent *Clostridium difficile* disease: use of high-dose vancomycin combined with *Saccharomyces boulardii*. *Clin Infect Dis* 2000;31:1012-7.
- Wullt M, Hagslatt ML, Odenholt I. *Lactobacillus plantarum* 299v for the treatment of recurrent *Clostridium difficile*-associated diarrhoea: a double-blind, placebo-controlled trial. *Scand J Infect Dis* 2003;35:365-7.
- Surawicz CM, Elmer GW, Speelman P, McFarland LV, Chinn J, van Belle G. Prevention of antibiotic-associated diarrhea by *Saccharomyces boulardii*: a prospective study. *Gastroenterology* 1989;96:981-8.
- McFarland LV, Surawicz CM, Greenberg RN, Elmer GW, Moyer KA, Melcher SA, et al. Prevention of beta-lactam-associated diarrhea by *Saccharomyces boulardii* compared with placebo. *Am J Gastroenterol* 1995;90:439-48.
- Lewis SJ, Potts LF, Barry RE. The lack of therapeutic effect of *Saccharomyces boulardii* in the prevention of antibiotic-related diarrhoea in elderly patients. *J Infect* 1998;36:171-4.
- Thomas MR, Litin SC, Osmon DR, Corr AP, Weaver AL, Lohse CM. Lack of effect of *Lactobacillus GG* on antibiotic-associated diarrhea: a randomized, placebo-controlled trial. *Mayo Clin Proc* 2001;76:883-9.
- Mylonakis E, Ryan ET, Calderwood SB. *Clostridium difficile*-associated diarrhea: A review. *Arch Intern Med* 2001;161:525-33.
- Vanderhoof JA, Whitney DB, Antonson DL, Hanner TL, Lupo JV, Young RJ. *Lactobacillus GG* in the prevention of antibiotic-associated diarrhea in children. *J Pediatr* 1999;135:564-8.
- Coeuret V, Gueguen M, Vernoux JP. Numbers and strains of lactobacilli in some probiotic products. *Int J Food Microbiol* 2004;97:147-56.
- Huff BA. Caveat emptor: "Probiotics" might not be what they seem. *Can Fam Physician* 2004;50:583-7.
- Marteau P, Seksik P. Tolerance of probiotics and prebiotics. *J Clin Gastroenterol* 2004;38(6 Suppl):S67-9.

Correspondence to: Dr. Nandini Dendukuri, Technology Assessment Unit, R4.09 Ross Pavilion, Royal Victoria Hospital, Montréal QC H3A 1A1; fax 514 843-1493; nandini.dendukuri@mcgill.ca