

Process Optimization for the Extraction of Polyphenols from Okara

Anshu Singh, Arindam Kuila, Geetanjali Yadav and Rintu Banerjee*

Microbial Biotechnology and Downstream Processing Laboratory, Agricultural and Food Engineering Department, Indian Institute of Technology, 721302 Kharagpur, India

Received: November 23, 2010

Accepted: February 23, 2011

Summary

The objective of the present investigation is to examine okara, a suitable substrate for polyphenol extraction, and to develop a feasible eco-friendly process to maximize the yield of antioxidant phenolics. Box-Behnken design (BBD) based on response surface methodology (RSM) was employed to investigate the effect of temperature (°C), solvent fraction (%) and incubation time (min) on polyphenol extraction by using MINITAB 15 software. Acetone was used as solvent to extract the phenolic compounds possessing the antioxidant properties (DPPH radical scavenging activity, reducing power, and metal chelating activity). Extraction under the optimum conditions yielded total polyphenolic content of 1.16 mg/mL, DPPH radical scavenging activity of 61.07 %, metal chelating activity of 61.20 % and better reducing power. The effective model developed for antioxidant mining from okara under mild operational conditions can be a valuable technique for soybean-based food industry.

Key words: natural antioxidants, polyphenolics, soybean, okara, response surface methodology (RSM)

Introduction

Health consciousness of consumers has led to a dynamic increase in the demand for natural antioxidants, which contribute to nutritional quality of products. The food industries are meeting the demand for antioxidants by the extraction from natural sources. Plant phenolics play a major antioxidative role in the diet. They are aromatic compounds responsible for the protection against degenerative diseases (1). Dietary antioxidants quench the free radicals generated during metabolism which are responsible for disruption of membrane fluidity, lipid peroxidation, oxidative DNA damage and alteration of platelet functions because they cause quality deterioration and nutritional loss (2). Antioxidants not only have health-related effects, but they also increase the shelf life of food.

Food industries are driven to use antioxidants in the form of food additives to maintain the market value of the product by enhancing its colour, flavour and nutri-

tional profile. Although synthetic antioxidants like butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) are common food additives, due to their toxicity and carcinogenicity (3) their application is questioned. Therefore, the need for natural antioxidants is rapidly increasing not only in food industry but also in medicine.

Agricultural and food waste generated during the processing is becoming the ideal substrate for extraction of antioxidant polyphenolics. Several food and agro-residues such as apple, potato and onion peels, carob pods and olive tree leaves (4), raspberry waste (5), *etc.* have been assessed for extraction of phenolics. Among these food processing residues, okara, a waste by-product of soybean industry, can be a potential feedstock for efficient recovery of bioactive antioxidant phytochemicals. Mateos-Aparicio *et al.* (6) measured the carbohydrate digestion (pectin) of okara by sequential extraction and correlated it with the antioxidant activity. Okara is

*Corresponding author; Phone: ++91 3222 283 104; Fax: ++91 3222 282 244; E-mail: rb@agfe.iitkgp.ernet.in

rich in indigestible complex polysaccharides and it has not been explored yet for the extraction of polyphenolics. However, total phenolic content of okara is almost half of the soybean seeds (7), but until now okara has been exploited only as nitrogen source during fermentation, animal feed or for dietary fibre extraction.

Natural antioxidants can be extracted from plant and animal sources under effective process conditions. Extraction process differs with the raw material and is also affected by several physicochemical factors. The economic viability and feasibility of the technological process of extraction is highly dependent on the availability and cost of raw material. Another important aspect apart from the cost-effective extraction and selection of substrate is the safety concern associated with the use of organic solvents. In past decades, methanol/hexane/benzene extractions were performed, but their safety issue due to potential toxic effects from the residual solvent was a point of concern. Optimization process for the reduction of use of organic solvents would be advantageous in their application in food and pharmaceutical sector. There are several reports on the extraction of polyphenols from food waste by using several solvents (2,8,9). Among them, acetone/water is the most efficient solvent for polyphenol extraction, because water in combination with acetone contributes to the creation of a moderately polar medium that ensures the efficient extraction of polyphenols and their antioxidant activities (10). The advantage of using food grade acetone is that it reduces the persistent problem of waste solvent disposal and hazards of environmental pollution. Therefore, the present article deals with the assessment of okara, a potential low-cost feedstock for extraction of antioxidant-rich phenolics with radical scavenging, reducing power and metal chelating activities, using food grade solvent by adopting Box-Behnken design based on the response surface methodology (RSM).

Materials and Methods

Preparation of okara

Soybeans (*Glycine max*) were purchased from a local market at Kharagpur, India. Accurately weighed clean soybeans were soaked for 2 h in a ratio of soybean to water of 1:2.48 and incubated at 40 °C in a thermostatically controlled water bath (11). Soaked soybeans were subsequently dehulled and weighed to add twice the volume of water. It was blended and extracted through cheese cloth to obtain the soy milk (12). The leftover residue after cheese extraction, okara, was dried overnight at 60 °C and grinded to powder for subsequent extraction.

Extraction of antioxidant phytochemicals from okara

A mass of 1 g of powdered okara was mixed with various fractions of solvent (acetone) and incubated under different time and temperature conditions. After the incubation, the sample was centrifuged at 2000×g for 10 min to separate the insoluble fractions, and antioxidant potential was estimated in the supernatant.

Determination of the total phenolic content in the extracts

Solvent extraction of polyphenols was performed in 50-mL stoppered conical flask containing 1 g of dried

okara sample. Samples were taken from the reaction mixture at specific time interval according to the experimental design. Each sample taken from the reaction mixture was centrifuged at 2000×g for 5 min. Then polyphenolic content (mg/mL) in the supernatant was estimated by Folin-Ciocalteu method (13).

Antioxidant activity

Antioxidant activity of the phytochemicals extracted from okara was assessed by measuring their radical scavenging activity, reducing power, and metal chelating activity.

Radical scavenging activity

Hydrogen atom or electron-donating ability of the corresponding extracts was measured by the bleaching of the purple-coloured methanol solution of 2,2-diphenyl-1-picrylhydrazyl (DPPH). This spectrophotometric assay uses stable DPPH radical as a reagent (14). To 1 mL of sample, 0.5 mL of DPPH (1 mM in methanol) was added. The control sample was prepared in a similar way by adding 1 mL of acetone instead of sample. The mixtures were shaken vigorously and left to settle for 15 min at room temperature. After the incubation period, the absorbance was read against a blank at 517 nm. Inhibition of DPPH free radical (in %) was calculated as follows:

$$\text{Inhibition} = [(A_0 - A_s) / A_0] \cdot 100 \quad /1/$$

where A_0 is the absorbance of the control, and A_s is the absorbance of the sample at 517 nm.

Reducing power

Reducing power of samples was determined according to the method of Benzie and Strain (15) by measuring the coloured ferrous-tripyridyltriazine complex formed due to ferric to ferrous ion reduction at low pH. To 3 mL of freshly prepared ferric reducing ability of plasma (FRAP) reagent (300 mM acetate buffer, pH=3.6, 10 mM 2,4,6-tripyridyl-s-triazine (TPTZ) solution and 20 mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in a 10:1:1 ratio), 0.3 mL of distilled water and 0.1 mL of extract were added. The absorbance was measured after 8 min of incubation at 593 nm. Higher the absorbance, higher the reducing power of sample.

Metal chelating activity

Ferrous ion chelating activity of the extract was determined by the method of Dinis *et al.* (16). Into test tubes containing 1.7 mL of distilled water and 50 μL of 0.2 mM $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$, 50 μL of sample solution were added and the mixture was left at room temperature for 1 min. To this mixture 0.2 mL of 5 mM ferrozine were added and final colour was measured at 562 nm after 10 min of incubation. The metal chelating efficiency of samples was determined by comparing with the chelating activity of ethylenediaminetetraacetic acid (EDTA). The inhibition percentage of Fe^{2+} -ferrozine complex formation against blanks containing FeCl_2 and ferrozine was calculated by the formula:

$$\text{Inhibition} = [(A_0 - A_1) / A_0] \cdot 100 \quad /2/$$

where A_0 is the absorbance of the control, and A_1 is the absorbance of the sample at 562 nm.

Box-Behnken design of experiment

A three-level three-factor fractional factorial design was adopted in this study. Initially the influencing parameters were selected by preliminary experiments on the basis of one-factor-at-a-time approach. The input variables considered to be important during the extraction process were temperature (40–60 °C), solvent fraction (25–75 %) and incubation time (15–45 min). In coded terms the lowest, central and the highest levels of five variables were –1, 0 and +1, respectively. Table 1 shows the coded and actual values of the experimental variables.

rical model to examine the interaction of different associated parameters responsible for the extraction of phenolic constituents present in okara using RSM, and also to identify the optimum conditions for a multivariable system of extraction. The predicted values were compared with the experimentally observed values to check the performance of the model as shown in Table 1. Analysis of variance for experimental set up was done to evaluate the fitness of response function. The linearity and quadratic effect of the independent variables, their interaction and regression coefficients on the response variables (Table 2) were analyzed. The goodness of fit of the mod-

Table 1. Experimental design for total polyphenolic content and antioxidant activity (DPPH radical scavenging activity, metal chelation and reducing power)

Run order	A ₁	A ₂	A ₃	Experimental			
				γ(total polyphenols)/(mg/mL)	DPPH radical scavenging activity/%	Metal chelation/%	Reducing power (absorbance)
1	0	–1	–1	0.855	60.47	54.70	0.643
2	0	–1	+1	0.825	49.00	38.08	0.607
3	0	+1	–1	0.825	48.67	38.00	0.608
4	0	+1	+1	0.792	48.13	27.35	0.574
5	–1	0	–1	0.873	57.89	37.12	0.589
6	–1	0	+1	0.264	45.81	46.20	0.656
7	+1	0	–1	0.338	55.43	45.85	0.728
8	+1	0	+1	0.230	54.81	43.19	0.726
9	–1	–1	0	0.227	40.36	42.87	0.631
10	–1	+1	0	1.148	58.01	31.74	0.607
11	+1	–1	0	0.844	48.69	46.47	0.634
12	+1	+1	0	0.639	45.93	59.12	0.662
13	0	0	0	0.610	53.18	16.70	0.546
14	0	0	0	0.230	53.18	10.70	0.704
15	0	0	0	0.230	53.18	10.70	0.704
16	0	0	0	0.230	53.17	10.70	0.704

The experimental data were analyzed by the response surface regression (RSREG) procedure to fit the following second-order polynomial equation:

$$Y = \beta_{k0} + \sum_{i=1}^5 \beta_{ki} X_i + \sum_{i=1}^5 \beta_{kii} X_i^2 + \sum_{i=1}^4 \sum_{j=i+1}^5 \beta_{kij} X_i X_j \quad /3/$$

where Y is the response (total polyphenolic content, DPPH radical scavenging activity, metal chelation and reducing power); β_{k0} , β_{ki} , β_{kii} and β_{kij} are constant coefficients, and x_i and x_j are the coded independent variables, which influence the response variable Y. This response was preferred because a relatively few experimental combinations of the variables were adequate to estimate a potentially complex response function. Data were analyzed using MINITAB 15 software (Minitab Inc., State College, PA, USA) to find the interaction between the variables and the responses.

Results and Discussion

In the present investigation, the experimental design has been formulated in such a way to develop an empi-

el was checked by the coefficient of determination (R^2) for stating a good statistical model, and also to justify its robustness. The coefficient of determination (R^2) was calculated to be 0.9608, 0.9958, 0.9982 and 0.9999 for total polyphenolic content, DPPH radical scavenging activity, metal chelation and reducing power, respectively, which are all close to 1.

Quadratic equation for solvent extraction

Second-order polynomial equations were used to correlate the input process variables with the responses. The second-order polynomial coefficient for each term of the equation was determined through multiple regression analysis using MINITAB 15.

The mathematical expression of relationship for total polyphenolic content, DPPH radical scavenging, metal chelation and reducing power with variables A₁, A₂ and A₃ (temperature (°C), solvent fraction (%) and incubation time (min), respectively) are given in Eqs. 4–7:

$$\begin{aligned}
 &\text{Total polyphenolic content}/(\text{mg}/\text{mL})= \\
 &=4.049-0.028A_1-0.110A_2-0.004A_3+0.00007A_1^2+ \\
 &\quad 0.00092A_2^2+0.00012A_3^2+0.00027A_1A_2- \\
 &\quad -0.00013A_1A_3+0.00009A_2A_3 \quad /4/ \\
 &\text{DPPH activity}/\%=-11.60+3.47A_1-1.53A_2+ \\
 &\quad +1.47A_3-0.033A_1^2+0.00252A_2^2-0.0036A_3^2+ \\
 &\quad +0.0166A_1A_2-0.0401A_1A_3+0.014A_2A_3 \quad /5/ \\
 &\text{Metal chelation}/\%=+444.02-13.82A_1+0.25A_2- \\
 &\quad -5.57A_3+0.17A_1^2+0.02A_2^2+0.077A_3^2- \\
 &\quad -0.06A_1A_2+ \\
 &\quad +0.018A_2A_3 \quad /6/ \\
 &\text{Reducing power}=-0.454+0.02A_1+0.02A_2+ \\
 &\quad +0.016A_3-0.000049A_1^2+0.00014A_2^2-0.00006A_3^2- \\
 &\quad -0.00014A_1A_2-0.00028A_1A_3+0.00003A_2A_3 \quad /7/
 \end{aligned}$$

Table 2. ANOVA analysis of response surface quadratic model for extraction of total polyphenols and antioxidant activity (DPPH radical scavenging activity, metal chelation and reducing power)

Source	DF ^a	Seq SS ^b	Adj SS ^b	Adj MS ^c	F	p
γ(TPC)/(mg/mL)						
regression	9	1.436	1.436	0.160	16.34	0.001
linear	3	0.150	0.656	0.219	22.39	0.001
square	3	1.260	1.265	0.422	43.17	<0.001
interaction	3	0.026	0.026	0.009	0.88	0.502
residual error	6	0.059	0.059	0.010		
lack-of-fit	3	0.059	0.059	0.020		
pure error	3					
total	16	1.494				
R ² =96.08 %				R ² =90.20 %		
DPPH						
regression	9	420.340	420.340	46.704	157.91	<0.001
linear	3	34.838	242.514	80.838	273.32	<0.001
square	3	42.201	55.765	18.588	62.85	<0.001
interaction	3	343.300	343.300	114.433	386.90	<0.001
residual error	6	1.775	1.775	0.296		
lack-of-fit	3	1.720	1.720	0.573	31.55	0.009
pure error	3	0.055	0.055	0.018		
total	16	422.115				
R ² =99.58 %				R ² =98.95 %		
Metal chelation						
regression	9	3665.290	3665.290	407.254	360.12	<0.001
linear	3	411.590	1150.670	383.557	339.17	<0.001
square	3	2227.170	2266.080	755.360	667.94	<0.001
interaction	3	1026.530	1026.530	342.177	302.57	<0.001
residual error	6	6.790	6.790	1.131		
lack-of-fit	3	6.780	6.780	2.261	2119.41	<0.001
pure error	3			0.001		
total	16	3672.080				
R ² =99.82 %				R ² =99.54 %		
Reducing power						
regression	9	0.047	0.047	0.005	5144.57	<0.001
linear	3	0.004	0.022	0.007	7361.69	<0.001
square	3	0.030	0.029	0.010	9645.74	<0.001
interaction	3	0.013	0.013	0.004	4225.11	<0.001
residual error	6					
lack-of-fit	3				11.14	0.039
pure error	3					
total	16	0.047				
R ² =99.99 %				R ² =99.97 %		

^adegrees of freedom, ^bsum of squares, ^cmean squares

From the regression equations mentioned above, it can be seen that A_2^2 , A_3^2 and A_1A_2 were significant model terms for total polyphenolic content. The surface plot in Fig. 1 shows the effect of temperature ($^{\circ}\text{C}$) and solvent fraction (%) on the extraction of total polyphenols (mg/mL). By increasing the temperature and solvent fraction, total polyphenolic content decreased significantly. This can be due to the degradation of polyphenols and decrease in the polarity of solvent at higher temperatures (17,18). At specific temperature (40°C) and solvent fraction (25 %), a maximum yield of polyphenols (1.16 mg/mL) was obtained.




Fig. 1. RSM plot showing the effect of temperature and solvent fraction on total polyphenolic content

For DPPH radical scavenging activity, A_1 , A_3 and A_1A_2 were significant model terms. From a 3D surface plot (Fig. 2), it was observed that radical scavenging ability increased with the increase in temperature and solvent fraction up to 50°C and 33 % respectively, but with further increase in these parameters, the percentage of inhibition of DPPH radicals started declining. Liu *et al.* (19) also observed the inhibition of DPPH radical scavenging activity at higher temperature.

The influencing model terms for metal chelation property were A_2 , A_1^2 and A_2A_3 . The surface plot (Fig. 3) indicates that at specific solvent fraction (25 %) and incubation time (15 min), maximum metal chelating activity can be achieved.

For reducing power, A_1 , A_2 and A_2A_3 were significant model terms. The surface plot (Fig. 4) shows the effect of interaction between solvent fraction and incubation time on the reducing power activity of okara. It demonstrated maximum reducing power activity at 42 % solvent fraction and 20 min of incubation time.

Validation of the model

Validation of the optimal conditions for solvent extraction of polyphenols was done by MINITAB 15 software. The results in Table 3 present the optimal conditions for each individual response with the predicted and experimental values. The optimum conditions for radical scavenging properties were obtained at 50°C with incubation time of 15 min using solvent fraction of 33 %. The radical scavenging activity present at 50°C was probably due to the increasing diffusivity of the solvent in




Fig. 2. RSM plot showing the effect of temperature and solvent fraction on DPPH radical scavenging activity




Fig. 3. RSM plot showing the effect of solvent fraction and incubation time on metal chelating activity




Fig. 4. RSM plot showing the effect of solvent fraction and incubation time on reducing power activity

the solid matrix, which favours the extraction. The reducing power activity was obtained at 60°C , solvent fraction of 42 % and incubation time of 20 min. Optimum condition for reducing power was noticed at higher temperature (60°C) than that for radical scavenging proper-

Table 3. Predicted and experimental values under optimum conditions for maximum total polyphenolic content and antioxidant activity (DPPH radical scavenging activity, metal chelation and reducing power)

Responses	Temperature/°C	Solvent fraction/%	Incubation time/min	Predicted value	Experimental value
γ (total polyphenols)/(mg/mL)	40	25	15	1.156	1.160
DPPH radical scavenging activity/%	50	33	15	61.230	61.070
Metal chelation/%	41	25	15	60.120	61.200
Reducing power (A)	60	42	20	0.730	0.732

ty. The stability of reducing power activity could be partly due to the formation of products of Maillard reaction. It was reported that there was an alteration in the phenolic compounds after heating, which contributed to the increase in reducing power (20).

Chelating agents are effective as secondary antioxidants because they reduce redox potential, thereby stabilizing the oxidized form of the metal ion (21). The extraction of chelating agent under optimized conditions (41 °C, 15 min, 25 % solvent fraction) showed chelation activity up to 61.20 %. Total polyphenolic content of okara (1.16 mg/mL) is similar to that reported by Gan and Latiff (8). Report by Tabart *et al.* (22) also agreed that aqueous acetone gave better yield of total polyphenolics than methanol and ethanol. Validation of the model suggests that okara may serve as a good substrate for extraction of a significant amount of polyphenols and other antioxidants.

Conclusion

The obtained results provide innovativeness and eco-friendly approach for recovery of the antioxidant phenolics from a low-cost feedstock okara, which can retain the maximum bioactivity of extracted biomolecule. In the present study, maximum amount of extracted polyphenols was achieved by using a food grade solvent. It can be concluded that the optimum conditions for maximum polyphenol extraction from okara (1.16 mg/mL) can be achieved by using acetone as solvent under 40 °C, 15 min and 25 % of solvent fraction, DPPH radical scavenging activity of 61.07 % at 50 °C for 15 min and 33 % of solvent fraction, metal chelating activity of 61.20 % at 41 °C for 15 min and 25 % of solvent fraction, and better reducing power at 60 °C for 20 min and 42 % of solvent fraction.

References

- C.A. Rice-Evans, N.J. Miller, G. Paganga, Antioxidant properties of phenolic compounds, *Trends Plant Sci.* 2 (1997) 152–159.
- F. Sharififar, G. Dehghn-Nudeh, M. Mirtajaldini, Major flavonoids with antioxidant activity from *Teucrium polium* L., *Food Chem.* 112 (2009) 885–888.
- H. Zhang, F. Chen, X. Wang, H.Y. Yao, Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents, *Food Res. Int.* 39 (2006) 833–839.
- C. Kaur, H.C. Kapoor, Antioxidants in fruits and vegetables – The millennium's health, *Int. J. Food Sci. Technol.* 36 (2001) 703–725.
- L. Laroze, C. Soto, M.E. Zúñiga, Phenolic antioxidants extraction from raspberry wastes assisted by-enzymes, *Electron. J. Biotechnol.* 13 (2010) 1–11.
- I. Mateos-Aparicio, C. Mateos-Peinado, A. Jiménez-Escrig, P. Rupérez, Multifunctional antioxidant activity of polysaccharide fractions from the soybean byproduct okara, *Carbohydr. Polym.* 82 (2010) 245–250.
- G. Préstamo, P. Rupérez, I. Espinosa-Martos, M.J. Villanueva, M.A. Lasunción, The effects of okara on rat growth, cecal fermentation, and serum lipids, *Eur. Food. Res. Technol.* 225 (2007) 925–928.
- C.Y. Gan, A.A. Latiff, Optimisation of the solvent extraction of bioactive compounds from *Parkia speciosa* pod using response surface methodology, *Food Chem.* 124 (2011) 1277–1283.
- Z. Khiari, D.P. Makris, P. Kefalas, Recovery of bioactive flavonols from onion solid wastes employing water/ethanol-based solvent system, *Food Sci. Technol. Int.* 14 (2008) 497–502.
- R. Chirinos, H. Rogez, D. Camposa, R. Pedreschi, Y. Laron-delle, Optimization of extraction conditions of antioxidant phenolic compounds from mashua (*Tropaeolum tuberosum* Ruiz & Pavón) tubers, *Sep. Purif. Technol.* 55 (2007) 217–225.
- T. Gulati, M. Chakrabarti, A. Singh, M. Duvuuri, R. Banerjee, Comparative study of response surface methodology, artificial neural network and genetic algorithms for optimization of soybean hydration, *Food Technol. Biotechnol.* 48 (2010) 11–18.
- M. Rivas, R.P. Garay, J.F. Escanero, P. Cia Jr., P. Cia, J.O. Alda, Soy milk lowers blood pressure in men and women with mild to moderate essential hypertension, *J. Nutr.* 132 (2002) 1900–1902.
- A. Scalbert, B. Monties, G. Janin, Tannins in wood: Comparison of different estimation methods, *J. Agric. Food Chem.* 37 (1989) 1324–1329.
- T. Yamaguchi, H. Takamura, T. Matoba, J. Terao, HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl, *Biosci. Biotech. Biochem.* 62 (1998) 1201–1204.
- I.F.F. Benzie, J.J. Strain, The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay, *Anal. Biochem.* 239 (1996) 70–76.
- T.C.P. Dinis, V.M.C. Madeira, L.M. Almeida, Action of phenolic derivatives (acetoaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxyl radical scavengers, *Arch. Biochem. Biophys.* 315 (1994) 161–169.
- C.F. Yap, C.W. Ho, W.M. Wan Aida, S.W. Chan, C.Y. Lee, Y.S. Leong, Optimization of extraction conditions of total phenolic compounds from star fruit (*Averrhoa carambola* L.) residues, *Sains Malaysiana*, 38 (2009) 511–520.
- M.A. Al-Farsi, C.Y. Lee, Optimization of phenolics and dietary fibre extraction from date seeds, *Food Chem.* 108 (2008) 977–985.

19. Q.M. Liu, X.M. Yang, L. Zhang, G. Majetich, Optimization of ultrasonic-assisted extraction of chlorogenic acid from *Folium eucommiae* and evaluation of its antioxidant activity, *J. Med. Plant. Res.* 4 (2010) 2503–2511.
20. P.J. Tsai, C.H. She, Significance of phenol-protein interactions in modifying the antioxidant capacity of peas, *J. Agric. Food Chem.* 54 (2006) 8491–8494.
21. M.H. Gordon: The Mechanism of Antioxidant Action *in vitro*. In: *Food Antioxidants*, B.J.F. Hudson (Ed.), Springer, London, UK (1990) pp. 1–18.
22. J. Tabart, C. Kevers, A. Sipel, J. Pincemail, J.O. Defraigne, J. Dommes, Optimisation of extraction of phenolics and antioxidants from black currant leaves and buds and of stability during storage, *Food Chem.* 105 (2007) 1268–1275.