



Production of Reactive Oxygen Species by Photosystem II as a Response to Light and Temperature Stress

Pavel Pospíšil*

Department of Biophysics, Centre of the Region Haná for Biotechnological and Agricultural Research, Faculty of Science, Palacký University, Olomouc, Czechia

OPEN ACCESS

Edited by:

Maya Velitchkova,
Institute of Biophysics and Biomedical
Engineering, Bulgarian Academy
of Science, Bulgaria

Reviewed by:

Christine Helen Foyer,
University of Leeds, UK
Anjana Jajoo,
Devi Ahilya University, India

*Correspondence:

Pavel Pospíšil
pavel.pospisil@upol.cz

Specialty section:

This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

Received: 23 September 2016

Accepted: 07 December 2016

Published: 26 December 2016

Citation:

Pospíšil P (2016) Production
of Reactive Oxygen Species by
Photosystem II as a Response
to Light and Temperature Stress.
Front. Plant Sci. 7:1950.
doi: 10.3389/fpls.2016.01950

The effect of various abiotic stresses on photosynthetic apparatus is inevitably associated with formation of harmful reactive oxygen species (ROS). In this review, recent progress on ROS production by photosystem II (PSII) as a response to high light and high temperature is overviewed. Under high light, ROS production is unavoidably associated with energy transfer and electron transport in PSII. Singlet oxygen is produced by the energy transfer from triplet chlorophyll to molecular oxygen formed by the intersystem crossing from singlet chlorophyll in the PSII antennae complex or the recombination of the charge separated radical pair in the PSII reaction center. Apart to triplet chlorophyll, triplet carbonyl formed by lipid peroxidation transfers energy to molecular oxygen forming singlet oxygen. On the PSII electron acceptor side, electron leakage to molecular oxygen forms superoxide anion radical which dismutates to hydrogen peroxide which is reduced by the non-heme iron to hydroxyl radical. On the PSII electron donor side, incomplete water oxidation forms hydrogen peroxide which is reduced by manganese to hydroxyl radical. Under high temperature, dark production of singlet oxygen results from lipid peroxidation initiated by lipoxygenase, whereas incomplete water oxidation forms hydrogen peroxide which is reduced by manganese to hydroxyl radical. The understanding of molecular basis for ROS production by PSII provides new insight into how plants survive under adverse environmental conditions.

Keywords: photoinhibition, heat inactivation, singlet oxygen, free oxygen radicals, lipid peroxidation

INTRODUCTION

Photosystem II (PSII) is water-plastoquinone oxidoreductase embedded in the thylakoid membrane that catalyzes light-driven H₂O oxidation to O₂ and plastoquinone (PQ) reduction to plastoquinol (PQH₂; Dau et al., 2012; Vinyard et al., 2013; Nelson and Junge, 2015; Suga et al., 2015; Najafpour et al., 2016). In this reaction, primary charge separation between the chlorophyll monomer (Chl_{D1}) and pheophytin (Pheo_{D1}) of D1 protein forms ¹[Chl_{D1}^{•+}Pheo_{D1}^{•-}] radical pair which is fast stabilized by the oxidation of the weakly coupled chlorophyll dimer P_{D1} and P_{D2} (P680) forming ¹[P680^{•+}Pheo_{D1}^{•-}] radical pair (Cardona et al., 2012). ¹[P680^{•+}Pheo_{D1}^{•-}] radical pair is stabilized by the electron transport from Pheo_{D1} to the tightly bound plastoquinone Q_A forming Q_A^{•-} and from the redox active tyrosine residue D1:161Y (Y_Z) to P680^{•+} forming Y_Z[•]. Electron transport

form $Q_A^{\bullet-}$ to loosely bound plastoquinone Q_B and the reduction of Y_Z^{\bullet} by the proton-coupled electron transport from the Mn_4O_5Ca cluster forms reducing and oxidizing equivalent at Q_B and Mn_4O_5Ca cluster, respectively. When two reducing equivalents are formed at Q_B site, its protonation forms plastoquinol (PQH_2) which is liberated to PQ pool via channels (Lambrev et al., 2014). Formation of four oxidizing equivalents in the Mn_4O_5Ca cluster causes four-electron oxidation of two H_2O to O_2 which is released via channels into the lumen (Vogt et al., 2015).

Light-driven processes comprising both energy transfer and electron transport are accompanied by formation of reactive oxygen species (ROS). In the energy transfer, singlet oxygen (1O_2) is formed by the energy transfer from triplet chlorophyll to O_2 (Triantaphylides and Havaux, 2009; Pospíšil, 2012; Fischer et al., 2013). In electron transport, ROS are formed by the consecutive one-electron reduction of O_2 and by the concerted two-electron oxidation of H_2O on the PSII electron acceptor and donor sides, respectively (Pospíšil, 2009). The one-electron reduction of O_2 forms superoxide anion radical ($O_2^{\bullet-}$) which dismutates spontaneously or enzymatically to hydrogen peroxide (H_2O_2) and subsequently is reduced to hydroxyl radical (HO^{\bullet}) via Fenton reaction. The two-electron oxidation of water forms H_2O_2 which is oxidized and reduced to $O_2^{\bullet-}$ and HO^{\bullet} , respectively. Non-enzymatic and enzymatic scavenging systems have been engaged to eliminate ROS and thus control level of ROS formed under various types of abiotic (adverse environmental conditions such as high light, high and low temperatures, UV-radiation, and drought) and biotic (herbivores and pathogens such as viruses, bacteria, and fungi) stresses.

Under moderate stress, when scavenging system maintains ROS level low, ROS serves as signaling molecules which activate an acclimation response and programmed cell death (Apel and Hirt, 2004; Dietz et al., 2016). Several lines of evidence have been provided that ROS play a crucial role in intracellular signaling from the chloroplast to the nucleus under high light (Gollan et al., 2015; Laloi and Havaux, 2015) and high temperature (Sun and Guo, 2016). However, due high reactivity of ROS toward proteins and lipids, ROS diffusion is limited. It seems to be unlikely that ROS might transmit signal from the chloroplast to the nucleus. It is considered that products of protein oxidation and lipid peroxidation might serve as signaling molecules (Fischer et al., 2012). As ROS formed by energy transfer (1O_2) and electron transport (H_2O_2) are produced simultaneously, it seems to be likely that their action in signaling pathways interferes. It was demonstrated that H_2O_2 antagonizes the 1O_2 signaling pathways in the *flu* Arabidopsis mutant (Laloi et al., 2007).

Under severe stress, when scavenging system is unable to sufficiently eliminate undesirable ROS formation, PSII proteins and lipids might be oxidized by ROS. Several lines of evidence were provided in the last three decades on the oxidative damage of PSII proteins by ROS under high light (Aro et al., 1993) and high temperature (Yamamoto et al., 2008). It is widely accepted that 1O_2 is major ROS responsible for oxidative modification of PSII proteins. Contrary, H_2O_2 has low capability to oxidize PSII protein; however, when free or protein-bound metals are available, HO^{\bullet} formed by Fenton reaction oxidizes nearby

proteins. It has to be pointed that experimental evidence on PSII protein oxidation was obtained *in vitro* and thus it remains to be clarified whether oxidative modification of PSII proteins by ROS occurs *in vivo*. Apart to involvement of ROS in PSII protein damage, the inhibition of *de novo* protein synthesis by ROS was proposed under high light (Nishiyama et al., 2006) and high temperature (Allakhverdiev et al., 2008). Whereas PSII protein oxidation is widely described, limited evidence has been provided on lipid peroxidation near PSII. It was shown that 1O_2 formed in PSII initiates lipid peroxidation in the thylakoid membrane (Triantaphylides et al., 2008).

In this review, an update on the latest findings on molecular mechanism of ROS formation at high light and high temperature is presented. In spite of the fact that molecular mechanism of ROS formation is substantially different at high light and high temperature, high light regularly combined with high temperature might bring about more serious impact on ROS formation.

HIGH LIGHT

When light energy which is driving force for photosynthetic reactions exceeds the photosynthetic capacity, a light-induced decline in photochemical activity in PSII denoted as photoinhibition occurs. Limitations in the energy transfer and electron transport result in the generation of ROS. Limitation in energy transfer occurs, when the excess energy absorbed by chlorophyll in the PSII antennae complex is not fully utilized in the PSII reaction center by charge separation. Under these conditions, singlet chlorophyll might be converted to deleterious triplet chlorophyll. To prevent formation of triplet chlorophyll, quenching of singlet chlorophyll to heat is maintained directly by xanthophylls or indirectly by the rearrangement of Lhcb protein by PsbS (Ruban et al., 2012). However, when quenching of singlet chlorophyll is not sufficient, singlet chlorophyll is converted to triplet chlorophyll which transfers energy to O_2 forming 1O_2 . Limitation in electron transport on the PSII electron acceptor side is accompanied by full reduction of PQ pool. As the Q_B site becomes unoccupied by PQ due to the full reduction of PQ pool, forward electron from Q_A to Q_B is blocked. Under these conditions, back electron transport from $Q_A^{\bullet-}$ to Pheo and consequent recombination of Pheo $^{\bullet-}$ with $P680^{\bullet+}$ forms deleterious triplet chlorophyll which transfer to O_2 forming 1O_2 . Under highly reducing conditions, double reduction and protonation of Q_A might result in the release of Q_AH_2 from its binding site. To prevent double reduction of Q_A , electron from $Q_A^{\bullet-}$ leaks to O_2 forming $O_2^{\bullet-}$. Superoxide anion radical is eliminated by its spontaneous and enzymatic dismutation to H_2O_2 . In the interior of the thylakoid membrane, $O_2^{\bullet-}$ is eliminated by the intrinsic SOD activity of *cyt b₅₅₉*, whereas $O_2^{\bullet-}$ which diffuse out the thylakoid membrane is eliminated by FeSOD attached to the stromal side of the thylakoid membrane at the vicinity of PSII. Limitation in electron transport on the PSII electron donor side is associated with incomplete H_2O oxidation catalyzed by the Mn_4O_5Ca cluster. Incomplete H_2O oxidation results in the formation of H_2O_2 which serves as precursor for

HO^\bullet . Under conditions, when H_2O_2 is not properly eliminated by catalase, HO^\bullet is formed by Fenton reactions catalyzed by iron and manganese on the PSII electron acceptor and donor sides, respectively.

Singlet Oxygen


Singlet oxygen is formed by the triplet-triplet energy transfer from triplet chlorophyll or triple carbonyl to O_2 . Triplet-triplet energy transfer from triplet chlorophyll to O_2 occurs in both the PSII antennae complex and the PSII reaction center. In the PSII antennae complex, triplet chlorophyll is formed by the photosensitization reaction, whereas in PSII reaction center triplet chlorophyll is formed by the charge recombination of triplet radical pair $^3[\text{P680}^{\bullet+}\text{Pheo}^{\bullet-}]$. Triplet-triplet energy transfer from triplet carbonyl to O_2 proceeds during lipid peroxidation initiated by ROS formed by light. Whereas $^1\text{O}_2$ formation by the energy transfer from triplet chlorophyll is well documented and represents the main source of $^1\text{O}_2$ at high light, $^1\text{O}_2$ formation by the energy transfer from triplet carbonyls is rarely evidenced and has marginal contribution to the overall $^1\text{O}_2$ formation.

Triplet Chlorophyll

Light energy absorbed by chlorophylls is transferred from the PSII antennae complex toward the PSII reaction center (van Amerongen and Croce, 2013). However, when energy transfer is limited, chlorophylls might serve as photosensitizers which form $^1\text{O}_2$ by the energy transfer from their triplet state to O_2 (Figure 1A). To prevent this, chlorophylls are coupled with carotenoids which have capability to quench triplet chlorophylls. Carotenoids consist of carotenes (β -carotene) and their oxygenated derivatives xanthophylls (lutein, zeaxanthin; Domonkos et al., 2013). In the PSII antennae complex, lutein and zeaxanthin play a crucial role in triplet chlorophyll quenching (Dall'Osto et al., 2006, 2012). Whereas lutein is permanently

coordinated to Lhcb proteins, zeaxanthin is accumulated under high light by the reversible de-epoxidation of violaxanthin and is either free in the thylakoid membrane or bound to Lhcb protein (Havaux and Niyogi, 1999; Pinnola et al., 2013). Four xanthophyll binding sites were documented in the monomeric (Lhcb4-6) and the trimeric (LHCII) antenna proteins of PSII (Liu et al., 2004). Xanthophylls bound in both L1 (lutein) and L2 (lutein in LHCII and lutein or zeaxanthin in monomeric Lhcb4-6 proteins) sites can efficiently quench the neighboring triplet chlorophylls. Lutein in L1 (Lut620) and L2 (Lut621) are coupled with chlorophylls Chl610-Chl614 and Chl602-Chl604, respectively. The quenching of triplet chlorophylls 602 and 603 by lutein in L2 is highly efficient, whereas lutein in L1 site had no effect on quenching of triplet chlorophyll 612 (Ballottari et al., 2013). To maintain effective quenching of triplet chlorophyll by carotenoids, carotenoids has to be properly distanced and oriented from chlorophylls. Triplet-triplet energy transfer from chlorophylls to carotenoids is mediated by Dexter mechanism (Dexter, 1953), which needs overlap between the electron clouds of the donor and acceptor. When distance or orientation of carotenoid and chlorophyll is changed, the capability of carotenoids to quench excitation energy of triplet chlorophylls is diminished (Cupellini et al., 2016). Under such conditions, when O_2 is in the proximity of triplet chlorophyll, the transfer of excitation energy from triplet chlorophyll to O_2 forms $^1\text{O}_2$. Comparison of the monomeric and the trimeric antenna proteins of PSII showed that the monomeric antenna proteins (Lhcb6 > Lhcb5 > Lhcb4) produced more $^1\text{O}_2$ as compared to trimeric antenna proteins (LHCII; Ballottari et al., 2013).

When electron transport on the PSII electron acceptor side is limited due to the slow electron transport to the Q_A and Q_B , several types of charge recombination of $[\text{P680}^{\bullet+}\text{Q}_\text{A}^{\bullet-}]$ and $^1[\text{P680}^{\bullet+}\text{Pheo}_{\text{D1}}^{\bullet-}]$ radical pairs occur. Whereas $[\text{P680}^{\bullet+}\text{Q}_\text{A}^{\bullet-}]$ radical pair recombines solely to the ground state P680, primary radical pair $^1[\text{P680}^{\bullet+}\text{Pheo}_{\text{D1}}^{\bullet-}]$ formed by the reverse



electron transport from $Q_A^{\bullet-}$ to Pheo_{D1} either recombines to the ground state P680 or converts to the triplet radical pair $^3[P680^{\bullet+}Pheo_{D1}^{\bullet-}]$ by change in the spin orientation. Recombination of triplet radical pair $^3[P680^{\bullet+}Pheo_{D1}^{\bullet-}]$ forms triplet chlorophyll $^3P680^*$ delocalized on the weakly coupled chlorophyll dimer P_{D1} and P_{D2} (Fischer et al., 2013; Telfer, 2014). Evidence has been provided that triplet state is localized on the Chl_{D1} at low temperature (Noguchi et al., 2001). The formation of $^3Chl_{D1}$ was proposed to occur either directly by the charge recombination of the triplet radical pair $^3[P680^{\bullet+}Pheo_{D1}^{\bullet-}]$ or by the triplet energy transfer from $^3P680^*$ to Chl_{D1}. As two β -carotenes (Car_{D1} and Car_{D2}) are distanced from chlorophyll dimer P_{D1} and P_{D2}, β -carotenes are not able to quench triplet chlorophyll $^3P680^*$ (Figure 1B).

Triplet Carbonyl

Lipid peroxidation initiated by radical ROS ($O_2^{\bullet-}$, HO^{\bullet}) forms the primary and the secondary lipid peroxidation products. The primary lipid peroxidation product are lipid hydroperoxides (lipid hydroperoxy fatty acids, LOOH) which decompose to the secondary lipid peroxidation products lipid hydroxides (hydroxy fatty acids, LOH), reactive carbonyl species (RCS), and electronically excited species. Hydrogen abstraction from polyunsaturated fatty acid by HO^{\bullet} forms lipid alkyl radical (L^{\bullet}) which interacts with O_2 forming lipid peroxy radical (LOO^{\bullet}). Lipid peroxy radical abstracts hydrogen from the adjacent polyunsaturated fatty acid forming LOOH. Lipid hydroperoxide is stable; however, under oxidizing or reducing condition it is oxidized or reduced to LOO^{\bullet} or alkoxy radical (LO^{\bullet}). Cyclization or recombination of LOO^{\bullet} forms high energy intermediates, dioxetane, or tetroxide. High energy intermediates are highly unstable and decompose to triplet excited carbonyls ($^3L^*$) which might transfer triplet energy to O_2 forming 1O_2 . Alternatively, tetroxide might directly decompose to 1O_2 via the Russell mechanism. Evidence has been provided that 1O_2 is formed through lipid peroxidation under light stress in spinach PSII membranes deprived by the Mn_4O_5Ca cluster (Yadav and Pospíšil, 2012a). The authors demonstrated that the oxidation of lipids by highly oxidizing $P680^{\bullet+}$ and $TyrZ^{\bullet}$ caused 1O_2 formation via the Russell mechanism. It has to be noted that amount of 1O_2 formed by the triplet-triplet energy transfer from triplet chlorophyll is considerably higher than from triplet carbonyl.

Superoxide Anion Radical


Superoxide anion radical is formed by the one-electron reduction of O_2 on the PSII electron acceptor side (Figure 2). Pheophytin (Pheo_{D1}^{•-}), tightly bound plastoquinone ($Q_A^{\bullet-}$), loosely bound plastoquinones ($Q_B^{\bullet-}$ or $Q_C^{\bullet-}$), free PQ ($PQ^{\bullet-}$), and ferrous iron of LP form of cyt *b*₅₅₉ were proposed to serve as electron donors to O_2 (Ananyev et al., 1994; Cleland and Grace, 1999; Pospíšil et al., 2004, 2006; Yadav et al., 2014). As Pheo_{D1}^{•-} has highly negative redox potential, the reduction of O_2 by Pheo_{D1}^{•-} is thermodynamically feasible; however, its short lifetime makes the diffusion limited reduction of O_2 less reasonable. Contrary, plastoquinones ($Q_A^{\bullet-}$, $Q_B^{\bullet-}$) does not fulfill thermodynamic criteria due to their more positive

redox potential, whereas they accomplish the kinetic criteria due their long lifetime. However, due to the different concentration of O_2 and $O_2^{\bullet-}$, the standard redox potential of $O_2/O_2^{\bullet-}$ redox couple is shifted according Nernst equation to more positive and thus the reduction of O_2 by plastoquinones becomes feasible (Pospíšil, 2009). The observation that exposure of isolated D1/D2/cyt *b*₅₅₉ complexes which lacks Q_A to high light causes a significant rate of cytochrome (III) reduction revealed that Pheo_{D1}^{•-} has capability to reduce O_2 . The detection of $O_2^{\bullet-}$ in isolated thylakoids by a voltammetric method showed $O_2^{\bullet-}$ production by the tightly bound plastoquinone $Q_A^{\bullet-}$ (Cleland and Grace, 1999). Experimental evidence has been recently provided on the reduction of O_2 by the loosely bound plastoquinones (Yadav et al., 2014). The authors demonstrated that plastoquinone is formed by the one-electron reduction of plastoquinone at the Q_B site and the one-electron oxidation of plastoquinol by cyt *b*₅₅₉ at the Q_C site. Apart to cofactors involved in the linear transport, the ferrous heme iron of LP form of cyt *b*₅₅₉ was shown to reduce O_2 forming $O_2^{\bullet-}$ (Pospíšil et al., 2006).

It has been demonstrated that PsbS knock-out rice mutants produced more $O_2^{\bullet-}$ compared to WT under high light (Zulfugarov et al., 2014). The authors proposed that the lack of PsbS may cause shift in the midpoint redox potential of $Q_A/Q_A^{\bullet-}$ redox couple to more negative value and thus enhance $O_2^{\bullet-}$ production by $Q_A^{\bullet-}$. The D1 protein phosphorylation which is associated with the migration of damaged PSII complexes from the grana to the stroma lamellae during D1 protein repair cycle was shown to decrease $O_2^{\bullet-}$ production (Chen et al., 2012). The author proposed that the D1 protein phosphorylation causes conformation change of D1 protein and thus modifies the binding of loosely bound plastoquinone to Q_B site. Consequently, the alternation of Q_B site brings about the decrease in $O_2^{\bullet-}$ formed by the loosely bound plastoquinone $Q_B^{\bullet-}$. In agreement with this proposal, it has been recently demonstrated that $O_2^{\bullet-}$ production is enhanced in STN8 kinase knock-out rice mutants under high light (Poudyal et al., 2016). It has been proposed that enhancement in $O_2^{\bullet-}$ production is due to the absence of conformational changes caused by STN8 kinase-induced phosphorylation. Using PsbY knock-out Arabidopsis plants, it has been shown that redox potential property of cyt *b*₅₅₉ is controlled by PsbY protein (von Sydow et al., 2016). It has to be explored whether PsbY protein controls $O_2^{\bullet-}$ production.

Hydrogen Peroxide


Hydrogen peroxide is formed by the one-electron reduction of $O_2^{\bullet-}$ and the two-electron oxidation of H_2O on the PSII electron acceptor and donor sides, respectively (Figure 2). Hydrogen peroxide formation by the one-electron reduction of $O_2^{\bullet-}$ occurs as dismutation or is maintained by plastoquinone. In the dismutation, two $O_2^{\bullet-}$ are simultaneously reduced and oxidized forming H_2O_2 and O_2 , respectively. In the spontaneous dismutation, the interaction of two $O_2^{\bullet-}$ is restricted due to repulsion of the negative charge on the molecule, whereas the interaction of the protonated form of superoxide, hydroperoxyl radical (HO_2^{\bullet}), either with $O_2^{\bullet-}$ or HO_2^{\bullet} is feasible. Spontaneous dismutation has been recently monitored



by real-time detection of H_2O_2 in PSII membrane under high light using highly sensitive and selective osmium-horseradish modified electrode (Prasad et al., 2015). In the enzymatic dismutation, reduction and oxidation of $O_2^{\bullet-}$ is associated with the redox change of the redox active metal center which serves as a superoxide oxidase (SOO) and superoxide reductase (SOR), respectively. It was demonstrated that the interaction of $O_2^{\bullet-}$ with the non-heme iron results in the oxidation of the ferrous iron and the formation of ferric-peroxo species which is protonated to ferric-hydroperoxo species (bound peroxide; Pospíšil et al., 2004) (Figure 3A). Evidence has been provided that the ferric and ferrous heme irons of *cyt b₅₅₉* exhibit the SOO and the SOR activities, respectively (Tiwari and Pospíšil, 2009; Pospíšil, 2011). Apart to dismutation, free $PQ^{\bullet-}$ in PQ pool was proposed to participate in H_2O_2 formation. Hydrogen peroxide was shown to be formed by reduction of $O_2^{\bullet-}$ by free $PQ^{\bullet-}$ (Borisova-Mubarakshina et al., 2015). The authors showed that H_2O_2 formed in PQ pool regulates the size of PSII antenna complex at high light. Furthermore, evidence has been provided that H_2O_2 might be formed by reduction 1O_2 of by PQH_2 (Khorobrykh et al., 2015). It was demonstrated that 1O_2 generated by photosensitizer Rose Bengal interacts with PQH_2

forming H_2O_2 . The authors proposed that H_2O_2 formed by reduction of 1O_2 by PQH_2 in the thylakoid membrane might cause dimerization of the protein kinase STN7 and thus activates the enzyme.

Hydrogen peroxide formation by the two-electron oxidation of H_2O is maintained by the Mn_4O_5Ca cluster when the complete four-electron oxidation of H_2O to O_2 is limited. Whereas all four manganese are redox active in four-electron oxidation of H_2O to O_2 , the incomplete oxidation of H_2O to H_2O_2 involves two redox active manganese. The two-electron oxidation of H_2O has been proposed to involve the transition from either S_2 to S_0 state or S_1 to S_{-1} state. Evidence has been provided that release of chloride from its binding site near to the Mn_4O_5Ca cluster enhanced H_2O_2 formation (Bradley et al., 1991; Fine and Frasc, 1992; Arato et al., 2004). A nucleophilic attack of hydroxo group on oxo group was proposed as an attractive model for formation of hydroperoxo species. It is proposed that nucleophilic attack of hydroxo group coordinated to Mn(4) and Cl(2) and oxo group coordinated to Ca forms hydroperoxo intermediate (Figure 3B). The hydroxo group is formed by deprotonation of the H_2O substrate coordinated to Mn(4) and Cl(2), whereas the oxo group is formed by



double deprotonation of H₂O substrate coordinated to Ca. A nucleophilic attack of manganese-coordinated hydroxo group on the calcium-coordinated electrophilic oxo group forms a peroxide intermediate that substitutes Cl(2) in coordination to Mn(4). Chloride controls accessibility of H₂O substrate to Mn(4) and the nucleophilicity of hydroxo group and thus interaction of hydroxo and oxo groups. Water substrate, which serves as a precursor for the hydroxo group, enters into the catalytic site, when the Cl(2) binding site becomes opened to the solvent H₂O due to its release.

Hydroxyl Radical

Hydroxyl radical is formed by the one-electron reduction of H₂O₂ formed on the both PSII electron acceptor and donor sides (Figure 2). Hydroxyl radical formation by the one-electron reduction of free H₂O₂ and bound peroxide on the PSII electron acceptor side was shown to be maintained by free iron and the non-heme iron, respectively (Pospíšil et al., 2004). The authors demonstrated that the reduction of bound peroxide (ferric iron-hydroperoxo intermediate) formed by the interaction of O₂^{•-} with the ferrous non-heme iron forms HO[•] via ferric iron-oxo intermediate (Figure 3A).

Hydroxyl radical formation by the one-electron reduction of H₂O₂ on the PSII electron donor side is likely to be maintained by manganese. From thermodynamic point of view, the reduction of H₂O₂ by manganese is not feasible. It was proposed that the reduction of H₂O₂ by manganese becomes thermodynamically more favorable by (1) the coordination of manganese to the protein due to the decrease in the redox potential of manganese and (2) the pH decrease in the lumen due to the increase in the

standard redox potential of H₂O₂/HO[•] redox couple (Pospíšil, 2012). It was demonstrated that PSII membranes depleted by chloride shows higher HO[•] formation compared to control PSII membranes (Arato et al., 2004). Based on the observation that HO[•] formation was not completely suppressed by exogenous SOD, the authors proposed that HO[•] is formed by reduction of H₂O₂ produced by the incomplete water oxidation on the PSII electron donor side.

HIGH TEMPERATURE

When PSII is exposed to high temperature, decline in the PSII activity denoted as heat inactivation occurs (Mathur et al., 2014). Heat inactivation occurs on the both PSII electron acceptor and donor sides. On the PSII electron donor side, heat inactivation is associated with the inhibition of water oxidation accompanied with release of PsbO, PsbP, and PsbQ proteins, calcium, chloride, and manganese from their binding sites (Coleman et al., 1988; Enami et al., 1994; Pospíšil et al., 2003; Barra et al., 2005). On the PSII electron acceptor side, heat inactivation is linked to the inhibition of electron transport from Q_A to Q_B (Pospíšil and Tyystjärvi, 1999). The authors demonstrated that increase in the midpoint redox potential of Q_A/Q_A^{•-} redox couple is responsible for the inhibition of Q_A to Q_B electron transport. Contrary to high light, ROS formation at high temperature is not driven by energy absorbed by chlorophylls; however, it is associated with heat-induced structural and functional changes in the thylakoid membrane. On the PSII electron acceptor side, ¹O₂ is formed decomposition of high energy intermediates formed by

lipid peroxidation. On the PSII electron donor side, incomplete H_2O oxidation forms H_2O_2 which is reduced by manganese to HO^\bullet via Fenton reaction.

Singlet Oxygen

Singlet oxygen is formed by the triplet-triplet energy transfer from $^3\text{L}^*$ to O_2 produced by the decomposition of high energy intermediates, dioxetane, or tetroxide, formed during lipid peroxidation (Havaux et al., 2006; Pospíšil and Prasad, 2014). The observation that elimination of HO^\bullet formation by mannitol did not suppress $^1\text{O}_2$ formation revealed that lipid peroxidation is unlikely initiated by HO^\bullet (Pospíšil et al., 2007). More recently, it has been demonstrated that inhibition of lipoxygenase by catechol and caffeic acid in *Chlamydomonas* cells prevented $^1\text{O}_2$ formation (Prasad et al., 2016). Singlet oxygen was proposed to be generated at the lipid phase near the Q_B site (Yamashita et al., 2008). It was pointed that PQH_2 formed by reduction of PQ by stromal reducing compound might cause ROS production which can damage D1 protein (Marutani et al., 2012).


Hydrogen Peroxide

Hydrogen peroxide is formed by the two-electron oxidation of H_2O on the PSII electron donor side (Figure 4). It was proposed that the release of extrinsic proteins (PsbO, PsbP, and PsbQ) leads to the inadequate accessibility of water to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster and consequently to the formation of H_2O_2 (Thompson et al., 1989). Indeed, it was demonstrated using the amplex red fluorescent assay that exposure of PSII membranes to high temperature (40°C) results in H_2O_2 formation (Yadav and Pospíšil, 2012b). The authors demonstrated that the binding of acetate to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster in the competition with chloride and blockage of water channel prevented H_2O_2 formation. Based on these observations, it was suggested that the release of chloride from its binding site near to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster leads to uncontrolled accessibility of H_2O to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster. To maintain controlled four-electron oxidation of H_2O to O_2 , the accessibility of H_2O to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster has to be regulated. Chloride coordinated to amino acids nearby the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster controls the accessibility of H_2O to the metal center and thus maintain proper four-electron oxidation of H_2O to O_2 .

However, when chloride is released from its binding site, the delivery of H_2O to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster is unrestricted and incomplete oxidation of O_2 to H_2O_2 occurs. Crystal structure of PSII from cyanobacteria *Thermosynechococcus vulcanus* reveals that two chlorides are located at distances of 6.67 and 7.40 Å from the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster (Umena et al., 2011). To avoid oxidation of nearby amino acid, diffusion of H_2O_2 into the lumen has to be restricted to the channels. As H_2O_2 is larger polar molecule similar to H_2O , it seems to be likely that H_2O_2 diffuse into the lumen via water channels. However, when H_2O_2 leaks from the water channels, it might interact with manganese and formed HO^\bullet .

Hydroxyl Radical

Hydroxyl radical is formed by the one-electron reduction of H_2O_2 formed on the PSII electron donor side (Figure 4). It was demonstrated by the EPR spin trapping spectroscopy that the exposure of PSII membranes to high temperature results in HO^\bullet formation (Pospíšil et al., 2007). The authors showed that HO^\bullet production is completely suppressed by exogenous catalase and metal chelator desferal revealing that HO^\bullet is formed via the metal-catalyzed Fenton reaction. Furthermore, the observation that the addition of exogenous calcium and chloride prevented HO^\bullet formation reveals that HO^\bullet is produced by the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster. This proposal was confirmed by the observation that no HO^\bullet formation was observed in PSII membranes deprived by the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster (Yamashita et al., 2008). As the replacement of chloride by acetate at its binding site near to the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster and the blockage of water channel prevented HO^\bullet formation in a similar manner as H_2O_2 formation, it was assumed that chloride plays a crucial role in HO^\bullet formation (Yadav and Pospíšil, 2012b). The authors proposed that H_2O_2 formed by the incomplete H_2O oxidation is reduced to HO^\bullet via the Fenton reaction mediated by free manganese released from the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster. The release of manganese from its binding site at high temperature was reported using atomic absorption (Nash et al., 1985) and EPR (Coleman et al., 1988; Pospíšil et al., 2003) spectroscopy. Detailed study using X-ray absorption spectroscopy showed that decomposition of the $\text{Mn}_4\text{O}_5\text{Ca}$ cluster occurs in two steps (Pospíšil et al., 2003). In the first step, two manganese are released



from their binding sites into the lumen remaining two manganese connected by a di- μ -oxo bridge, whereas in the second phase the remaining two manganese are liberated from PSII.

PHYSIOLOGICAL RELEVANCE OF ROS FORMATION

Role of ROS in Retrograde Signaling

Both $^1\text{O}_2$ and H_2O_2 formed in the thylakoid membrane were proposed to be involved in retrograde signaling (Dietz et al., 2016). Role of $^1\text{O}_2$ in acclimation and programmed cell death was demonstrated in green algae (Erickson et al., 2015) and higher plants (Triantaphylides and Havaux, 2009; Laloi and Havaux, 2015). In higher plants, *fluorescent (flu)* and *chlorina 1 (chl)* Arabidopsis mutants were advantageously used due to their high capability to form $^1\text{O}_2$. It was proposed that the $^1\text{O}_2$ level determines whether acclimation response or programmed cell death is triggered (Laloi and Havaux, 2015).

At low $^1\text{O}_2$ level, acclimation response is mediated by β -cyclocitral formed by oxidation of β -carotene (Ramel et al., 2013a; Havaux, 2014). It was demonstrated that exposure of WT Arabidopsis plants to β -cyclocitral caused expression of $^1\text{O}_2$ related gene (Ramel et al., 2012). In agreement with this finding, it was shown that concentration of β -cyclocitral is enhanced in *chl* Arabidopsis plants under acclimation (Ramel et al., 2013b). Further, evidence was provided on the role of jasmonic acid in acclimation response. It was demonstrated that jasmonate-deficient Arabidopsis mutant (*delayed-dehiscence 2*) was more resistant to light and jasmonate biosynthesis was pronouncedly lowered under acclimation (Ramel et al., 2013b). Based on these observations, the authors proposed that downregulation of jasmonate biosynthesis plays a crucial role in the triggering of acclimation response (Ramel et al., 2013c).

At high $^1\text{O}_2$ level, programmed cell death is dependent on the plastid proteins EXECUTER1 (EX1) and EXECUTER2 (EX2; Lee et al., 2007) and OXIDATIVE SIGNAL INDUCIBLE1 (OXI1) encoding an AGC kinase (Shumbe et al., 2016). Several lines of evidence on the involvement of EX1 and EX2 in programmed cell death were provided using *flu* Arabidopsis mutant (Lee et al., 2007). In this mutant, $^1\text{O}_2$ is formed by triplet-triplet energy transfer from the triplet chlorophyll precursor protochlorophyllide to O_2 (op den Camp et al., 2003). Even if EX1 and EX2 are located in chloroplast, it was proposed that jasmonic acid formed by $^1\text{O}_2$ -initiated lipid peroxidation mediates genetically controlled programmed cell death response via these two plastid proteins (Przybyla et al., 2008). The initiation of $^1\text{O}_2$ signaling has been recently demonstrated close to EX1 in the grana margins nearby the site of chlorophyll synthesis and $^1\text{O}_2$ formation (Wang et al., 2016). As $^1\text{O}_2$ signaling depends on the FstH protease, the authors proposed that $^1\text{O}_2$ signaling is linked to D1 repair cycle. Apart to EX1 and EX2, it has been shown recently that OXI1 kinase is involved in $^1\text{O}_2$ signaling in *chl* Arabidopsis mutant (Shumbe et al., 2016). In this mutant, $^1\text{O}_2$ is formed by triplet-triplet energy transfer from the triplet chlorophyll formed in PSII to O_2 (Krieger-Liszkay, 2005). As OXI1 kinase is localized at the cytosol at the cell periphery or

in the nucleus, it seems to be likely that oxylipins mediate signal transduction from chloroplast to cytosol (Shumbe et al., 2016).

Hydrogen peroxide formed under high light was demonstrated to play a crucial role in signaling associated with acclimation and programmed cell death (Foyer and Noctor, 2009; Karpinski et al., 2013; Gollan et al., 2015). It is well established that H_2O_2 regulates expression of genes by the activation of protein kinase signaling pathways. It was proposed that precursor of jasmonic acid, 12-oxo phytodienoic acid (OPDA), mediates signal transduction from chloroplast to cytosol (Tikkanen et al., 2014). It has been recently demonstrated that H_2O_2 formed in PQ pool triggers signal transduction from the chloroplast to the nucleus via protein kinase signaling pathways leading to the regulation of the PSII antenna size during the acclimation response (Borisova-Mubarakshina et al., 2015).

Our knowledge on the involvement of ROS in retrograde signaling at high temperature is highly limited. While the physiological relevance of light-induced $^1\text{O}_2$ to acclimation and programmed cell death is described to some extent, no evidence was provided on the role of $^1\text{O}_2$ formed under high temperature to plant stress response. However, it seems to be likely that $^1\text{O}_2$ might oxidize lipid, protein or pigment forming specific oxidation products and thus initiates signal transduction from the chloroplast to the nucleus in the signaling cascade pathway. Contrary to $^1\text{O}_2$, H_2O_2 was shown to be an important component in heat stress-activated gene expression. Hydrogen peroxide was demonstrated to be involved in the synthesis of heat shock proteins (Volkov et al., 2006). More experimental data are required to pronouncedly progress our understanding of multiple signaling pathways involved in the response to heat stress.

Role of ROS in Oxidative Damage

At high light, proteins and lipids might be oxidized by ROS formed in PSII. PSII proteins were evidenced to be oxidatively modified in the following order D1 > D2 > Cyt b559 > CP43 > CP47 > $\text{Mn}_4\text{O}_5\text{Ca}$ cluster (Komenda et al., 2006). Amino acid oxidation at the lumen exposed AB-loop of D1 protein forms 24 kDa C-terminal and 9 kDa N-terminal fragments, whereas amino acid oxidation in the stromally exposed D-de loop of the D1 protein form 23-kDa N-terminal and 9-kDa C-terminal fragments (Edelman and Mattoo, 2008). Identification of naturally oxidized amino acid in D1 protein using mass spectrometry was shown nearby to the site of ROS production (Sharma et al., 1997; Frankel et al., 2012, 2013). Whereas D1 protein oxidation was pronouncedly studied *in vitro*, limited evidence was provided on D1 protein oxidation *in vivo* (Shipton and Barber, 1994; Lupinkova and Komenda, 2004). Regardless of a broad range of evidence on PSII protein oxidation obtained *in vitro*, the plausibility of these processes *in vivo* has to be clarified. An efficient repair cycle for D1 protein, which includes proteolytic degradation of damaged D1 protein and its replacement with a newly synthesized D1 copy is essential for maintaining the viability of PSII (Komenda et al., 2012; Mulo et al., 2012; Jarvi et al., 2015). Apart to involvement of ROS in PSII protein damage under high light, ROS were shown to suppress the synthesis *de novo* of proteins with the

elongation step of translation as primary target (Nishiyama et al., 2006). However, considering the limited ROS diffusion, it seems to be more likely that ROS produced in the stroma might oxidize the translational elongation factors involved in D1 repair cycle. Unbound chlorophylls released to the stroma from their binding sites during PSII protein damage or chlorophyll precursors during chlorophyll synthesis are likely candidates for $^1\text{O}_2$ formation due to the lack of effective quenching of triplet excitation energy by carotenoids. To avoid $^1\text{O}_2$ formation, unbound chlorophylls might be temporarily coordinated to early light-induced proteins (ELIPs). In agreement with this proposal, it was demonstrated that small CAB-like proteins prevent $^1\text{O}_2$ formation during PSII damage, most probably by the binding of unbound chlorophylls released from the damaged PSII complexes (Sinha et al., 2012). Lipids associated with membrane proteins were shown to be oxidized by ROS. The initiation of lipid peroxidation by $^1\text{O}_2$ comprises the insertion of $^1\text{O}_2$ to double bond of polyunsaturated fatty acid, whereas HO^\bullet initiates lipid peroxidation by hydrogen abstraction from polyunsaturated fatty acid. It has been demonstrated that primary (LOOH) and secondary (LOH, RCS, and electronically excited species) lipid peroxidation products are formed at high light. Formation of hydroxy fatty acid was demonstrated in Arabidopsis plants (Triantaphylides et al., 2008). The authors showed that oxidation of polyunsaturated fatty acid by $^1\text{O}_2$ leads to formation of LOOH which further forms LOH isomers (10-HOTE and 15-HOTE).

At high temperature, limited evidence was provided on the oxidation of proteins and lipids by ROS. It was demonstrated that exposure of thylakoid membranes to high temperature caused cleavage of D1 protein forming 9 kDa C-terminal and 23 kDa N-terminal fragments (Yoshioka et al., 2006). The authors demonstrated that FtsH protease is involved in the cleavage of the D1 protein at high temperature. Furthermore, it was reported that $^1\text{O}_2$ formed at Q_B site by the recombination of LOO^\bullet formed by the lipid peroxidation caused the D1 protein degradation by the interaction with D-de loop of the D1 protein in a similar manner as under high light (Yamashita et al., 2008). As experimental evidence for oxidative damage of PSII protein by endogenous ROS was obtained predominantly *in vitro*, it is unclear whether the PSII protein oxidation at high temperature occurs *in vivo*. Apart from involvement of ROS in PSII protein oxidation, the inhibition of *de novo* protein synthesis

by ROS was proposed at high temperature (Allakhverdiev et al., 2008). Lipid peroxidation is associated with formation of RCS. It was demonstrated that malondialdehyde is formed in Arabidopsis plants exposed to heat stress (Yamauchi et al., 2008).

CONCLUSION AND PERSPECTIVES

Under environmental conditions, abiotic stresses adversely affect plant growth and survival. The impact of high light on the photosynthetic apparatus is considered to be of particular significance as light reactions of photosynthesis are inhibited prior to other cell functions are impaired. However, under environmental conditions, plants are exposed to combination of multiple stresses. High light stress is often associated with high temperature causing global warming which is one of the most important characteristics of accelerated climatic changes. Extensive research over the last 10 years focused on the structural and functional changes of the photosynthetic complexes in response to high light, high temperature or their combination. The exploration of molecular mechanism of ROS production by PSII helps to understand the adaptive processes by which plants cope with high light and high temperature stresses.

AUTHOR CONTRIBUTIONS

The PP wrote and approved manuscript for publication.

FUNDING

This work was supported by the Ministry of Education, Youth and Sports of the Czech Republic through grant no. LO1204 (Sustainable development of research in the Centre of the Region Haná from the National Program of Sustainability I).

ACKNOWLEDGMENT

I thank to Ravindra Kale for his advice with molecular visualization system Pymol and Lenka Kuchařová for stimulating discussion.

REFERENCES

- Allakhverdiev, S. I., Kreslavski, V. D., Klimov, V. V., Los, D. A., Carpentier, R., and Mohanty, P. (2008). Heat stress: an overview of molecular responses in photosynthesis. *Photosynth. Res.* 98, 541–550. doi: 10.1007/s11120-008-9331-0
- Ananyev, G., Renger, G., Wacker, U., and Klimov, V. (1994). The photoproduction of superoxide radicals and the superoxide-dismutase activity of photosystem-II - the possible involvement of cytochrome B559. *Photosynth. Res.* 41, 327–338. doi: 10.1007/BF00019410
- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399. doi: 10.1146/annurev.arplant.55.031903.141701
- Arato, A., Bondarava, N., and Krieger-Liszka, A. (2004). Production of reactive oxygen species in chloride- and calcium-depleted photosystem II and their involvement in photoinhibition. *Biochim. Biophys. Acta* 1608, 171–180. doi: 10.1016/j.bbabi.2003.12.003
- Aro, E. M., Virgin, I., and Andersson, B. (1993). Photoinhibition of Photosystem II. Inactivation, protein damage and turnover. *Biochim. Biophys. Acta* 1143, 113–134. doi: 10.1016/0005-2728(93)90134-2
- Ballottari, M., Mozzo, M., Girardon, J., Hienerwadel, R., and Bassi, R. (2013). Chlorophyll triplet quenching and photoprotection in the higher plant monomeric antenna protein Lhcb5. *J. Phys. Chem. B* 117, 11337–11348. doi: 10.1021/jp402977y
- Barra, M., Haumann, M., and Dau, H. (2005). Specific loss of the extrinsic 18 Kda protein from Photosystem II upon heating to 47 degrees C causes inactivation

- of oxygen evolution likely due to Ca release from the Mn-complex. *Photosynth. Res.* 84, 231–237. doi: 10.1007/s11220-004-7158-x
- Borisova-Mubarakshina, M. M., Ivanov, B. N., Vetoshkina, D. V., Lubimov, V. Y., Fedorchuk, T. P., Naydov, I. A., et al. (2015). Long-term acclimatory response to excess excitation energy: evidence for a role of hydrogen peroxide in the regulation of photosystem II antenna size. *J. Exp. Bot.* 66, 7151–7164. doi: 10.1093/jxb/erv410
- Bradley, R. L., Long, K. M., and Frasch, W. D. (1991). The involvement of photosystem-ii-generated H₂O₂ in photoinhibition. *FEBS Lett.* 286, 209–213. doi: 10.1016/0014-5793(91)80975-9
- Cardona, T., Sedoud, A., Cox, N., and Rutherford, A. W. (2012). Charge separation in Photosystem II: a comparative and evolutionary overview. *Biochim. Biophys. Acta* 1817, 26–43. doi: 10.1016/j.bbabi.2011.07.012
- Chen, L. B., Jia, H. Y., Tian, Q., Du, L. B., Gao, Y. L., Miao, X. X., et al. (2012). Protecting effect of phosphorylation on oxidative damage of D1 protein by down-regulating the production of superoxide anion in photosystem II membranes under high light. *Photosynth. Res.* 112, 141–148. doi: 10.1007/s11220-012-9750-9
- Cleland, R. E., and Grace, S. C. (1999). Voltammetric detection of superoxide production by photosystem II. *FEBS Lett.* 457, 348–352. doi: 10.1016/S0014-5793(99)01067-4
- Coleman, W. J., Govindjee, and Gutowsky, H. S. (1988). The effect of chloride on the thermal inactivation of oxygen evolution. *Photosynth. Res.* 16, 261–276. doi: 10.1007/BF00028844
- Cupellini, L., Jurinovich, S., Prandi, I. G., Caprasecca, S., and Mennucci, B. (2016). Photoprotection and triplet energy transfer in higher plants: the role of electronic and nuclear fluctuations. *Phys. Chem. Chem. Phys.* 18, 11288–11296. doi: 10.1039/C6CP01437B
- Dall'Osto, L., Holt, N. E., Kaligotla, S., Fuciman, M., Cazzaniga, S., Carbonera, D., et al. (2012). Zeaxanthin protects plant photosynthesis by modulating chlorophyll triplet yield in specific light-harvesting antenna subunits. *J. Biol. Chem.* 287, 41820–41834. doi: 10.1074/jbc.M112.405498
- Dall'Osto, L., Lico, C., Alric, J., Giuliano, G., Havaux, M., and Bassi, R. (2006). Lutein is needed for efficient chlorophyll triplet quenching in the major LHCII antenna complex of higher plants and effective photoprotection in vivo under strong light. *BMC Plant Biol.* 6:32. doi: 10.1186/1471-2229-6-32
- Dau, H., Zaharieva, I., and Haumann, M. (2012). Recent developments in research on water oxidation by photosystem II. *Curr. Opin. Chem. Biol.* 16, 3–10. doi: 10.1016/j.cbpa.2012.02.011
- DeLano, W. L. (2002). *The PYMOL Molecular Graphics System. Software*. Available at: <http://www.pymol.org>
- Dexter, D. L. (1953). A theory of sensitized luminescence in solids. *J. Chem. Phys.* 21, 836–850. doi: 10.1063/1.1699044
- Dietz, K. J., Turkan, I., and Krieger-Liszskay, A. (2016). Redox- and reactive oxygen species-dependent signaling into and out of the photosynthesizing chloroplast. *Plant Physiol.* 171, 1541–1550. doi: 10.1104/pp.16.00375
- Domonkos, I., Kis, M., Gombos, Z., and Ughy, B. (2013). Carotenoids, versatile components of oxygenic photosynthesis. *Prog. Lipid Res.* 52, 539–561. doi: 10.1016/j.plipres.2013.07.001
- Edelman, M., and Mattoo, A. K. (2008). D1-protein dynamics in photosystem II: the lingering enigma. *Photosynth. Res.* 98, 609–620. doi: 10.1007/s11220-008-9342-x
- Enami, I., Kitamura, M., Tomo, T., Isokawa, Y., Ohta, H., and Katoh, S. (1994). Is the primary cause of thermal inactivation of oxygen evolution in spinach PS-II membranes release of the extrinsic 33 kDa protein or of MN. *Biochim. Biophys. Acta* 1186, 52–58. doi: 10.1016/0005-2728(94)90134-1
- Erickson, E., Wakao, S., and Niyogi, K. K. (2015). Light stress and photoprotection in *Chlamydomonas reinhardtii*. *Plant J.* 82, 449–465. doi: 10.1111/tj.12825
- Fine, P. L., and Frasch, W. D. (1992). The oxygen-evolving complex requires chloride to prevent hydrogen-peroxide formation. *Biochemistry* 31, 12204–12210. doi: 10.1021/bi00163a033
- Fischer, B. B., Hideg, E., and Krieger-Liszskay, A. (2013). Production, detection, and signaling of singlet oxygen in photosynthetic organisms. *Antioxid. Redox Signal.* 18, 2145–2162. doi: 10.1089/ars.2012.5124
- Fischer, B. B., Ledford, H. K., Wakao, S., Huang, S. G., Casero, D., Pellegrini, M., et al. (2012). Singlet oxygen resistant 1 links reactive electrophile signaling to singlet oxygen acclimation in *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. U.S.A.* 109, E1302–E1311. doi: 10.1073/pnas.1116843109
- Foyer, C. H., and Noctor, G. (2009). Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. *Antioxid. Redox Signal.* 11, 861–905. doi: 10.1089/ars.2008.2177
- Frankel, L. K., Sallans, L., Limbach, P. A., and Bricker, T. M. (2012). Identification of oxidized amino acid residues in the vicinity of the Mn₄CaO₅ cluster of Photosystem II: implications for the identification of oxygen channels within the photosystem. *Biochemistry* 51, 6371–6377. doi: 10.1021/bi300650n
- Frankel, L. K., Sallans, L., Limbach, P. A., and Bricker, T. M. (2013). Oxidized amino acid residues in the vicinity of Q(A) and Pheo(D1) of the photosystem II reaction center: putative generation sites of reducing-side reactive oxygen species. *PLoS ONE* 8:e58042. doi: 10.1371/journal.pone.0058042
- Gollan, P. J., Tikkanen, M., and Aro, E.-M. (2015). Photosynthetic light reactions: integral to chloroplast retrograde signalling. *Curr. Opin. Plant Biol.* 27, 180–191. doi: 10.1016/j.pbi.2015.07.006
- Havaux, M. (2014). Carotenoid oxidation products as stress signals in plants. *Plant J.* 79, 597–606. doi: 10.1111/tj.12386
- Havaux, M., and Niyogi, K. K. (1999). The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism. *Proc. Natl. Acad. Sci. U.S.A.* 96, 8762–8767. doi: 10.1073/pnas.96.15.8762
- Havaux, M., Triantaphylides, C., and Genty, B. (2006). Autoluminescence imaging: a non-invasive tool for mapping oxidative stress. *Trends Plant Sci.* 11, 480–484. doi: 10.1016/j.tplants.2006.08.001
- Jarvi, S., Suorsa, M., and Aro, E. M. (2015). Photosystem II repair in plant chloroplasts - regulation, assisting proteins and shared components with photosystem II biogenesis. *Biochim. Biophys. Acta* 1847, 900–909. doi: 10.1016/j.bbabi.2015.01.006
- Karpinski, S., Szechynska-Hebda, M., Wituszynska, W., and Burdiak, P. (2013). Light acclimation, retrograde signalling, cell death and immune defences in plants. *Plant Cell Environ.* 36, 736–744. doi: 10.1111/pce.12018
- Khorobrykh, S. A., Karonen, M., and Tyystjarvi, E. (2015). Experimental evidence suggesting that H₂O₂ is produced within the thylakoid membrane in a reaction between plastoquinol and singlet oxygen. *FEBS Lett.* 589, 779–786. doi: 10.1016/j.febslet.2015.02.011
- Komenda, J., Kuvikovi, S., Lupinkova, L., and Masojidek, J. (2006). “Biogenesis and structural dynamics of the photosystem II complex,” in *Biotechnological Applications of Photosynthetic Proteins: Biochips, Biosensors, and Biodevices*, eds M. T. Giardi and E. V. Piletska (New York, NY: Springer).
- Komenda, J., Sobotka, R., and Nixon, P. J. (2012). Assembling and maintaining the photosystem II complex in chloroplasts and cyanobacteria. *Curr. Opin. Plant Biol.* 15, 245–251. doi: 10.1016/j.pbi.2012.01.017
- Krieger-Liszskay, A. (2005). Singlet oxygen production in photosynthesis. *J. Exp. Bot.* 56, 337–346. doi: 10.1093/jxb/erh237
- Laloi, C., and Havaux, M. (2015). Key players of singlet oxygen-induced cell death in plants. *Front. Plant Sci.* 6:39. doi: 10.3389/fpls.2015.00039
- Laloi, C., Stachowiak, M., Pers-Kamczyc, E., Warzych, E., Murgia, I., and Apel, K. (2007). Cross-talk between singlet oxygen- and hydrogen peroxide-dependent signaling of stress responses in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 672–677. doi: 10.1073/pnas.0609063103
- Lambrev, M. D., Russo, D., Polticelli, F., Scognamiglio, V., Antonacci, A., Zobnina, V., et al. (2014). Structure/function/dynamics of photosystem II plastoquinone binding sites. *Curr. Protein Pept. Sci.* 15, 285–295. doi: 10.2174/1389203715666140327104802
- Lee, K. P., Kim, C., Landgraf, F., and Apel, K. (2007). EXECUTER1- and EXECUTER2-dependent transfer of stress-related signals from the plastid to the nucleus of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 10270–10275. doi: 10.1073/pnas.0702061104
- Liu, Z. F., Yan, H. C., Wang, K. B., Kuang, T. Y., Zhang, J. P., Gui, L. L., et al. (2004). Crystal structure of spinach major light-harvesting complex at 2.72 angstrom resolution. *Nature* 428, 287–292. doi: 10.1038/nature02373
- Lupinkova, L., and Komenda, J. (2004). Oxidative modifications of the Photosystem II D1 protein by reactive oxygen species: from isolated protein to cyanobacterial cells. *Photochem. Photobiol.* 79, 152–162. doi: 10.1562/0031-8655\yr{(2004)}079\$<\$0152:OMOTPI\$>\$2.0.CO;2
- Marutani, Y., Yamauchi, Y., Kimura, Y., Mizutani, M., and Sugimoto, Y. (2012). Damage to photosystem II due to heat stress without light-driven electron flow: involvement of enhanced introduction of reducing power into thylakoid membranes. *Planta* 236, 753–761. doi: 10.1007/s00425-012-1647-5

- Mathur, S., Agrawal, D., and Jadoo, A. (2014). Photosynthesis: response to high temperature stress. *J. Photochem. Photobiol. B Biol.* 137, 116–126. doi: 10.1016/j.jphotobiol.2014.01.010
- Mulo, P., Sakurai, I., and Aro, E.-M. (2012). Strategies for psbA gene expression in cyanobacteria, green algae and higher plants: from transcription to PSII repair. *Biochim. Biophys. Acta* 1817, 247–257. doi: 10.1016/j.bbabi.2011.04.011
- Najafpour, M. M., Renger, G., Holyńska, M., Moghaddam, A. N., Aro, E.-M., Carpentier, R., et al. (2016). Manganese compounds as water-oxidizing catalysts: from the natural water-oxidizing complex to nanosized manganese oxide structures. *Chem. Rev.* 116, 2886–2936. doi: 10.1021/acs.chemrev.5b00340
- Nash, D., Miyao, M., and Murata, N. (1985). Heat inactivation of oxygen evolution in photosystem-II particles and its acceleration by chloride depletion and exogenous manganese. *Biochim. Biophys. Acta* 807, 127–133. doi: 10.1016/0005-2728(85)90115-X
- Nelson, N., and Junge, W. (2015). Structure and energy transfer in photosystems of oxygenic photosynthesis. *Annu. Rev. Biochem.* 84, 659–683. doi: 10.1146/annurev-biochem-092914-041942
- Nishiyama, Y., Allakhverdiev, S. I., and Murata, N. (2006). A new paradigm for the action of reactive oxygen species in the photoinhibition of photosystem II. *Biochim. Biophys. Acta* 1757, 742–749. doi: 10.1016/j.bbabi.2006.05.013
- Noguchi, T., Tomo, T., and Kato, C. (2001). Triplet formation on a monomeric chlorophyll in the photosystem II reaction center as studied by time-resolved infrared spectroscopy. *Biochemistry* 40, 2176–2185. doi: 10.1021/bi0019848
- op den Camp, R. G. L., Przybyla, D., Ochsenbein, C., Laloi, C., Kim, C. H., Danon, A., et al. (2003). Rapid induction of distinct stress responses after the release of singlet oxygen in *Arabidopsis*. *Plant Cell* 15, 2320–2332. doi: 10.1105/tpc.014662
- Pinnola, A., Dall'osto, L., Gerotto, C., Morosinotto, T., Bassi, R., and Alboresi, A. (2013). Zeaxanthin binds to light-harvesting complex stress-related protein to enhance nonphotochemical quenching in *Physcomitrella patens*. *Plant Cell* 25, 3519–3534. doi: 10.1105/tpc.113.114538
- Pospíšil, P. (2009). Production of reactive oxygen species by photosystem II. *Biochim. Biophys. Acta* 1787, 1151–1160. doi: 10.1016/j.bbabi.2009.05.005
- Pospíšil, P. (2011). Enzymatic function of cytochrome b(559) in photosystem II. *J. Photochem. Photobiol. B Biol.* 104, 341–347. doi: 10.1016/j.jphotobiol.2011.02.013
- Pospíšil, P. (2012). Molecular mechanisms of production and scavenging of reactive oxygen species by photosystem II. *Biochim. Biophys. Acta* 1817, 218–231. doi: 10.1016/j.bbabi.2011.05.017
- Pospíšil, P., Arato, A., Krieger-Liszky, A., and Rutherford, A. W. (2004). Hydroxyl radical generation by Photosystem II. *Biochemistry* 43, 6783–6792. doi: 10.1021/bi036219i
- Pospíšil, P., Haumann, M., Dittmer, J., Sole, V. A., and Dau, H. (2003). Stepwise transition of the tetra-manganese complex of photosystem II to a binuclear Mn-2(μ -O)(2) complex in response to a temperature jump: a time-resolved structural investigation employing X-ray absorption spectroscopy. *Biophys. J.* 84, 1370–1386. doi: 10.1016/S0006-3495(03)74952-2
- Pospíšil, P., and Prasad, A. (2014). Formation of singlet oxygen and protection against its oxidative damage in Photosystem II under abiotic stress. *J. Photochem. Photobiol. B Biol.* 137, 39–48. doi: 10.1016/j.jphotobiol.2014.04.025
- Pospíšil, P., Snyrychova, I., Kruk, J., Strzalka, K., and Naus, J. (2006). Evidence that cytochrome b(559) is involved in superoxide production in photosystem II: effect of synthetic short-chain plastoquinones in a cytochrome b(559) tobacco mutant. *Biochem. J.* 397, 321–327. doi: 10.1042/BJ20060068
- Pospíšil, P., Šnyrychová, I., and Naus, J. (2007). Dark production of reactive oxygen species in photosystem II membrane particles at elevated temperature: EPR spin-trapping study. *Biochim. Biophys. Acta* 1767, 854–859. doi: 10.1016/j.bbabi.2007.02.011
- Pospíšil, P., and Tyystjärvi, E. (1999). Molecular mechanism of high-temperature-induced inhibition of acceptor side of Photosystem II. *Photosynth. Res.* 62, 55–66. doi: 10.1023/A:1006369009170
- Poudyal, R. S., Nath, K., Zulfugarov, I. S., and Lee, C. H. (2016). Production of superoxide from photosystem II-light harvesting complex II supercomplex in STN8 kinase knock-out rice mutants under photoinhibitory illumination. *J. Photochem. Photobiol. B Biol.* 162, 240–247. doi: 10.1016/j.jphotobiol.2016.06.050
- Prasad, A., Ferretti, U., Sedlářová, M., and Pospíšil, P. (2016). Singlet oxygen production in *Chlamydomonas reinhardtii* under heat stress. *Sci. Rep.* 6, 20094. doi: 10.1038/srep20094
- Prasad, A., Kumar, A., Suzuki, M., Kikuchi, H., Sugai, T., Kobayashi, M., et al. (2015). Detection of hydrogen peroxide in Photosystem II (PSII) using catalytic amperometric biosensor. *Front. Plant Sci.* 6:862. doi: 10.3389/fpls.2015.00862
- Przybyla, D., Gobel, C., Imboden, A., Hamberg, M., Feussner, I., and Apel, K. (2008). Enzymatic, but not non-enzymatic, O-1(2)-mediated peroxidation of polyunsaturated fatty acids forms part of the EXECUTER1-dependent stress response program in the flu mutant of *Arabidopsis thaliana*. *Plant J.* 54, 236–248. doi: 10.1111/j.1365-313X.2008.03409.x
- Ramel, F., Birtic, S., Ginies, C., Soubigou-Taconnat, L., Triantaphylides, C., and Havaux, M. (2012). Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *Proc. Natl. Acad. Sci. U.S.A.* 109, 5535–5540. doi: 10.1073/pnas.1115982109
- Ramel, F., Ksas, B., Akkari, E., Mialoundama, A. S., Monnet, F., Krieger-Liszky, A., et al. (2013b). Light-induced acclimation of the *Arabidopsis* chlorinal mutant to singlet oxygen. *Plant Cell* 25, 1445–1462. doi: 10.1105/tpc.113.109827
- Ramel, F., Ksas, B., and Havaux, M. (2013c). Jasmonate: a decision maker between cell death and acclimation in the response of plants to singlet oxygen. *Plant Signal. Behav.* 8, e26655. doi: 10.4161/psb.26655
- Ramel, F., Mialoundama, A. S., and Havaux, M. (2013a). Nonenzymic carotenoid oxidation and photooxidative stress signalling in plants. *J. Exp. Bot.* 64, 799–805. doi: 10.1093/jxb/ers223
- Ruban, A. V., Johnson, M. P., and Duffy, C. D. P. (2012). The photoprotective molecular switch in the photosystem II antenna. *Biochim. Biophys. Acta* 1817, 167–181. doi: 10.1016/j.bbabi.2011.04.007
- Sharma, J., Panico, M., Shipton, C. A., Nilsson, F., Morris, H. R., and Barber, J. (1997). Primary structure characterization of the photosystem II D1 and D2 subunits. *J. Biol. Chem.* 272, 33158–33166. doi: 10.1074/jbc.272.52.33158
- Shipton, C. A., and Barber, J. (1994). In-vivo and in-vitro photoinhibition reactions generate similar degradation fragments of D1 and D2 photosystem-ii reaction-center proteins. *Eur. J. Biochem.* 220, 801–808. doi: 10.1111/j.1432-1033.1994.tb18682.x
- Shumbe, L., Chevalier, A., Legeret, B., Taconnat, L., Monnet, F., and Havaux, M. (2016). Singlet oxygen-induced cell death in *Arabidopsis* under high-light stress is controlled by OXII kinase. *Plant Physiol.* 170, 1757–1771.
- Sinha, R. K., Komenda, J., Knoppova, J., Sedlářová, M., and Pospíšil, P. (2012). Small CAB-like proteins prevent formation of singlet oxygen in the damaged photosystem II complex of the cyanobacterium *Synechocystis* sp PCC 6803. *Plant Cell Environ.* 35, 806–818. doi: 10.1111/j.1365-3040.2011.02454.x
- Suga, M., Akita, F., Hirata, K., Ueno, G., Murakami, H., Nakajima, Y., et al. (2015). Native structure of photosystem II at 1.95 angstrom resolution viewed by femtosecond X-ray pulses. *Nature* 517, 99–103. doi: 10.1038/nature13991
- Sun, A. Z., and Guo, F. Q. (2016). Chloroplast retrograde regulation of heat stress responses in plants. *Front. Plant Sci.* 7:398. doi: 10.3389/fpls.2016.00398
- Telfer, A. (2014). Singlet oxygen production by PSII under light stress: mechanism, detection and the protective role of beta-carotene. *Plant Cell Physiol.* 55, 1216–1223. doi: 10.1093/pcp/pcu040
- Thompson, L. K., Blaylock, R., Sturtevant, J. M., and Brudvig, G. W. (1989). Molecular-basis of the heat denaturation of photosystem-II. *Biochemistry* 28, 6686–6695. doi: 10.1021/bi00442a023
- Tikkanen, M., Gollan, P. J., Mekala, N. R., Isojarvi, J., and Aro, E. M. (2014). Light-harvesting mutants show differential gene expression upon shift to high light as a consequence of photosynthetic redox and reactive oxygen species metabolism. *Philos. Trans. R. Soc. B Biol. Sci.* 369, 20130229. doi: 10.1098/rstb.2013.0229
- Tiwari, A., and Pospíšil, P. (2009). Superoxide oxidase and reductase activity of cytochrome b(559) in photosystem II. *Biochim. Biophys. Acta* 1787, 985–994. doi: 10.1016/j.bbabi.2009.03.017
- Triantaphylides, C., and Havaux, M. (2009). Singlet oxygen in plants: production, detoxification and signaling. *Trends Plant Sci.* 14, 219–228. doi: 10.1016/j.tplants.2009.01.008
- Triantaphylides, C., Krischke, M., Hoerberichts, F. A., Ksas, B., Gresser, G., Havaux, M., et al. (2008). Singlet oxygen is the major reactive oxygen species involved in photooxidative damage to plants. *Plant Physiol.* 148, 960–968. doi: 10.1104/pp.108.125690

- Umena, Y., Kawakami, K., Shen, J. R., and Kamiya, N. (2011). Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature* 473, 55–60. doi: 10.1038/nature09913
- van Amerongen, H., and Croce, R. (2013). Light harvesting in photosystem II. *Photosynth. Res.* 116, 251–263. doi: 10.1007/s11120-013-9824-3
- Vinyard, D. J., Ananyev, G. M., and Dismukes, G. C. (2013). Photosystem II: the reaction center of oxygenic photosynthesis. *Annu. Rev. Biochem.* 82, 577–606. doi: 10.1146/annurev-biochem-070511-100425
- Vogt, L., Vinyard, D. J., Khan, S., and Brudvig, G. W. (2015). Oxygen-evolving complex of Photosystem II: an analysis of second-shell residues and hydrogen-bonding networks. *Curr. Opin. Chem. Biol.* 25, 152–158. doi: 10.1016/j.cbpa.2014.12.040
- Volkov, R. A., Panchuk, I. I., Mullineaux, P. M., and Schoffl, F. (2006). Heat stress-induced H₂O₂ is required for effective expression of heat shock genes in *Arabidopsis*. *Plant Mol. Biol.* 61, 733–746. doi: 10.1007/s11103-006-0045-4
- von Sydow, L., Schwenkert, S., Meurer, J., Funk, C., Mamedov, F., and Schröder, W. P. (2016). The PsbY protein of *Arabidopsis* PhotosystemII is important for the redox control of cytochrome b559. *Biochim. Biophys. Acta* 1857, 1524–1533. doi: 10.1016/j.bbabi.2016.05.004
- Wang, L. S., Kim, C., Xu, X., Piskurewicz, U., Dogra, V., Singh, S., et al. (2016). Singlet oxygen- and EXECUTER1-mediated signaling is initiated in grana margins and depends on the protease FtsH2. *Proc. Natl. Acad. Sci. U.S.A.* 113, E3792–E3800. doi: 10.1073/pnas.1603562113
- Wei, X., Su, X., Cao, P., Liu, X.-Y., Chang, W., Li, M., et al. (2016). Structure of spinach photosystem II–LHCII supercomplex at 3.2 Å resolution. *Nature* 534, 69–74. doi: 10.1038/nature18020
- Yadav, D. K., and Pospíšil, P. (2012a). Evidence on the formation of singlet oxygen in the donor side photoinhibition of photosystem II: EPR spin-trapping study. *PLoS ONE* 7:e45883. doi: 10.1371/journal.pone.0045883
- Yadav, D. K., and Pospíšil, P. (2012b). Role of chloride ion in hydroxyl radical production in photosystem II under heat stress: electron paramagnetic resonance spin-trapping study. *J. Bioenerg. Biomembr.* 44, 365–372. doi: 10.1007/s10863-012-9433-4
- Yadav, D. K., Prasad, A., Kruk, J., and Pospíšil, P. (2014). Evidence for the involvement of loosely bound plastoquinones in superoxide anion radical production in photosystem II. *PLoS ONE* 9:e115466. doi: 10.1371/journal.pone.0115466
- Yamamoto, Y., Aminaka, R., Yoshioka, M., Khatoun, M., Komayama, K., Takenaka, D., et al. (2008). Quality control of photosystem II: impact of light and heat stresses. *Photosynth. Res.* 98, 589–608. doi: 10.1007/s11120-008-9372-4
- Yamashita, A., Nijo, N., Pospíšil, P., Morita, N., Takenaka, D., Aminaka, R., et al. (2008). Quality control of photosystem II - Reactive oxygen species are responsible for the damage to photosystem II under moderate heat stress. *J. Biol. Chem.* 283, 28380–28391. doi: 10.1074/jbc.M710465200
- Yamauchi, Y., Furutera, A., Seki, K., Toyoda, Y., Tanaka, K., and Sugimoto, Y. (2008). Malondialdehyde generated from peroxidized linolenic acid causes protein modification in heat-stressed plants. *Plant Physiol. Biochem.* 46, 786–793. doi: 10.1016/j.plaphy.2008.04.018
- Yoshioka, M., Uchida, S., Mori, H., Komayama, K., Ohira, S., Morita, N., et al. (2006). Quality control of photosystem II. Cleavage of reaction center D1 protein in spinach thylakoids by FtsH protease under moderate heat stress. *J. Biol. Chem.* 281, 21660–21669. doi: 10.1074/jbc.M602896200
- Zulfugarov, I. S., Tovuu, A., Eu, Y.-J., Dogsom, B., Poudyal, R. S., Nath, K., et al. (2014). Production of superoxide from Photosystem II in a rice (*Oryza sativa* L.) mutant lacking PsbS. *BMC Plant Biol.* 14:242. doi: 10.1186/s12870-014-0242-2

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Pospíšil. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.