



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in:

*Injury*

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa37128>

---

### **Paper:**

Eftaxiopoulou, T., Barnett-Vanes, A., Arora, H., Macdonald, W., Nguyen, T., Itadani, M., Sharrock, A., Britzman, D., Proud, W., et. al. (2016). Prolonged but not short-duration blast waves elicit acute inflammation in a rodent model of primary blast limb trauma. *Injury*, 47(3), 625-632.

<http://dx.doi.org/10.1016/j.injury.2016.01.017>

---

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

# 1 **Prolonged but not short duration blast waves elicit acute inflammation in a** 2 **rodent model of primary blast limb trauma**

3

4 Keywords: Rodent model; blast limb trauma; inflammatory response

## 5 **Abstract**

### 6 **Background**

7 Blast injuries from conventional and improvised explosive devices account for 75% of injuries  
8 from current conflicts; of these over 70% involve the limbs. Variable duration and magnitude  
9 of blast wave loading occurs in real-life explosions and is hypothesised to cause different  
10 injuries. While a number of *in-vivo* models report the inflammatory response to blast injuries,  
11 the extent of this response has not been investigated with respect to the duration of the primary  
12 blast wave. The relevance is that explosions in open air are of short duration compared to those  
13 in confined spaces.

### 14 **Methods**

15 Hind limbs of adult Sprague-Dawley rats were subjected to focal isolated primary blast waves  
16 of varying overpressure (1.8-3.65kPa) and duration (3.0-11.5ms), utilising a shock tube and  
17 purpose built experimental rig. Rats were monitored during and after blast. At 6 and 24hrs after  
18 exposure blood, lungs, liver and muscle tissue were collected and prepared for histology and  
19 flow cytometry.

### 20 **Results**

21 At 6hrs increases in circulating neutrophils and CD43Lo/His48Hi monocytes were observed in  
22 rats subjected to longer duration blast waves. This was accompanied by increases in circulating  
23 pro-inflammatory chemo/cytokines KC and IL-6. No changes were observed with shorter  
24 duration blast waves irrespective of overpressure. In all cases, no histological damage was  
25 observed in muscle, lung or liver. By 24hrs post-blast all inflammatory parameters had  
26 normalised.

### 27 **Conclusions**

28 We report the development of a rodent model of primary blast limb trauma that is the first to  
29 highlight an important role played by blast wave duration and magnitude in initiating acute  
30 inflammatory response following limb injury in the absence of limb fracture or penetrating

31 trauma. The combined biological and mechanical method developed can be used to further  
32 understand the complex effects of blast waves in a range of different tissues and organs *in-vivo*.

33

## 34 **Introduction**

35 Blast injuries from conventional and improvised explosive devices (IEDs) account for 75% of  
36 modern war injuries, over 70% of these involve the limbs [1]. Blast injuries remain a threat to  
37 civilians too; detonation of IEDs in the recent Boston marathon bombings in 2013, together  
38 with industrial accidents such as the 2013 west Texas fertilizer plant explosion caused many  
39 injuries with a similar pattern to that seen amongst military casualties [2].

40 Blast trauma may occur by four discrete mechanisms: primary injuries are due to the interaction  
41 between the blast wave and the human body, secondary injuries are caused by the impact of  
42 fragments thrown and energised during the explosion, tertiary injuries result from acceleration  
43 of the body against an obstruction, and quaternary injuries include other physical insults, such  
44 as burns and smoke inhalation [3-5]. The type and severity of the injury sustained depends on;  
45 the explosive system and the environment of the blast; the size of explosive device, the distance  
46 between the person and the explosion and the presence of obstacles or reflections. In an open  
47 space (free field blast) a blast wave spreads radially from its origin and quickly dissipates as a  
48 function of the cube of the distance [3]. In these cases the blast wave consists of a rapid rise to  
49 a positive overpressure followed by a negative under-pressure and return to ambient pressure.  
50 In an enclosed space, the explosive energy is contained leading to rises both in the peak  
51 overpressure and the duration of the positive-pressure phase of the blast wave [3]. Previous  
52 studies have shown that explosions within enclosed spaces are associated with a higher  
53 incidence of primary blast injuries and more severe injuries compared to open air explosions  
54 [6].

55 In both the civilian and military setting, blast injured patients are often poly-traumatised with  
56 the head, torso and soft tissues commonly affected [7-8]. The understanding within the trauma  
57 community of the relationship between injury, inflammation, sepsis and clinical outcome is  
58 growing [9-11]. It is important to note that blast injury survivors may suffer shock or  
59 hypoxemia in the absence of external signs of injury [12]. The onset of inflammation following  
60 injury is a common phenomenon, however, in severely compromised patients (particularly  
61 those with sepsis) systemic inflammation may contribute towards deleterious and life  
62 threatening changes, such as multi-organ failure, which are difficult to manage clinically [11].

63 *In - vivo* animal models are often used with simulated blast conditions in a controlled  
64 environment to investigate the mechanisms of blast injury. These studies enable greater

65 understanding of the nature of the injury, including mechanical failure thresholds of tissues,  
66 physiological or inflammatory responses, and the effect of therapeutic interventions. However,  
67 given the clinical burden of lower limb blast injuries [13], few experimental live models have  
68 been reported relating to blast limb trauma; those undertaken show that explosive limb injury  
69 may lead to systemic inflammatory changes affecting the limbs as well as distal organs [14-  
70 17]. However, the injury documented in these models is severe and encompasses several blast  
71 injury mechanisms. It is recognised that further research is needed to closely examine the  
72 interplay of different blast mechanisms in limb injury [18].

73 During IED explosion, blasts with peak pressures from 50 to 1000 kPa and 2-6 ms duration  
74 have been measured [19]. However, the majority of the experimental animal models involve  
75 blast waves with longer durations between 4 – 8 ms [20-22] and some with durations longer  
76 than 10 ms [23]. A large number of existing models also lack detail when reporting the pressure  
77 histories of the blasts produced, with many reporting only the peak overpressure or the distance  
78 from the outlet of the shock tube, often without details or schematics of animal positioning and  
79 orientation, thus limiting comparability between studies [24-25].

80 In this study we develop a model to investigate the inflammatory response to primary blast  
81 wave application to the limb, investigating the effect of changing the magnitude or duration of  
82 the blast wave, thus permitting the controlled delivery of primary blast to replicate durations  
83 associated with a range of open field and enclosed environments.

84

## 85 **Materials and Methods**

### 86 *Characterisation of the Injury Device*

87 A shock tube (Figure 1A) was employed in this study to generate pressure pulses of controlled  
88 intensity and duration. The shock tube used is a stainless steel tube 3.8 m long and 59 mm in  
89 internal diameter. It is comprised of two chambers; the first chamber, the driver section, is filled  
90 with compressed air [26] and the breech releases the pressure into the driven, low-pressure,  
91 section producing a shock wave [27].

92 In order to characterise the blast waves generated, three dynamic pressure gauges were installed  
93 along the shock tube's driven section. The first two pressure gauges were orientated radially at  
94 the middle (Sensor 1) and end of the driven section (Sensor 2) as shown in Figure 1A. The  
95 third (Sensor 3) was installed on a bull-nose probe of 2 cm in diameter sited axially facing  
96 upstream. Sensor 1 was used for triggering and monitoring the reproducibility of the blast.  
97 Sensor 2 for monitoring the output pressure, and Sensor 3 was used to characterise the pressure  
98 pulse at the sample position.

99 While the magnitude of the shock wave generated can be altered by using diaphragms of  
100 different burst pressure, the duration of the wave is controlled by changing the length in the  
101 driver tube. The 10% driver volume uses only the volume of the breech system (Figure 1A)  
102 while, 100% volume uses the full length of the driver tube. It should be noted that the 10%-  
103 volume driver generates the Friedlander waveform of open-air blasts, while full-volume driver  
104 produces blasts of longer pulse width (duration), similar to those seen in large-charge  
105 explosions and enclosed volumes such as inside a vehicle.

106 With the *in-vivo* model the sample replaced Sensor 3. Sensor 2 was then used to monitor the  
107 loading pressure. Three different experimental conditions were used:

108 200- $\mu\text{m}$  Mylar® diaphragm, bursting at  $\sim 15.7 \pm 0.5$  bar with 10% driver volume

109 75- $\mu\text{m}$  Mylar® diaphragm, bursting at  $\sim 6.2 \pm 0.2$  bar with 10% driver volume

110 50- $\mu\text{m}$  Mylar® diaphragm, bursting at  $\sim 6.0 \pm 0.1$  bar with full (100%) driver volume

111 *Induction of injury*

112 Sprague-Dawley (*Rattus norvegicus*) female rats weighing 250-320 grams were used. All  
113 animals were housed in clean cages and kept on a 12-hour light/- dark cycle with unlimited  
114 access to food and fresh water. All animals were anaesthetised with 60 mg/kg ketamine and  
115 0.20 mg/kg medetomidine [28] administered intraperitoneally and randomly allocated into four  
116 groups as summarised in Table 1. Sham animals were anaesthetised but not exposed to a blast  
117 wave. Animals from Groups I and II were anaesthetised and exposed to a single blast insult of  
118 6.0 and 16.0 bar overpressure respectively. Animals in the final Group III were exposed to a  
119 6.0 bar blast wave of a longer duration (Table 1) compared to animals in Groups I and II.

120 **Table 1: Experimental Groups used in the study**

121 A purpose-built subject support (Figure 1B) was developed to expose the rat's hind limbs to a  
122 focussed primary blast wave, whilst ensuring the remainder of the animal was not exposed.  
123 This was done to ensure local loading and to prevent the abdomen and chest sustaining injury.  
124 The animal's hind limbs were then strapped using adhesive tape to a steel bar in order to prevent  
125 limb movement and avoid tertiary blast effects.

126 **Figure 1: (A) Shock tube schematic. (B) Experimental rig mounted at the outlet of the shock**  
127 **tube to isolate the blast to the animal's left hind-limb. The distance from the shock tube's**  
128 **outlet to the left thigh is 5cm. The left leg is exposed to the shock wave from the pelvis to**  
129 **the ankle joint.**

130 All animals were closely monitored after recovery and received buprenorphine (0.03 mg/kg  
131 administered subcutaneously) for analgesia as a precaution. For the remainder of the study, the  
132 animals were monitored 3 times a day in their facility for several behavioural parameters,  
133 including appearance, weight changes, and response to handling. Six animals from the Sham  
134 group and Groups I, II and III were observed for 6 hrs to investigate the acute inflammatory  
135 response; in line with the 3R's principle; Shams, Group I and Group III were observed up to  
136 24 hrs to track the resolution of inflammation.

137 *Data collection and analysis*

138 Physiological parameters

139 Heart rate data were collected using a Veterinary Handheld Oximeter (Creative Medical,  
140 Shenzhen, China) at 1 min and 5 min after induction of anaesthesia, and then 1 min and 5 min

141 after blast or sham procedure. Finally, heart rate data were also collected once the subjects  
142 recovered (40 min after the initial induction of anaesthesia).

#### 143 Collection of blood and plasma

144 At specified time points of 6 or 24 hrs, rats were intraperitoneally (i.p) injected with an  
145 overdose of Pentobarbitone and blood was collected in citrated tubes on ice from the right  
146 femoral vein and centrifuged at 2500 rpm for 10 mins. Plasma was collected and stored at -80°C  
147 for further analysis.

#### 148 Blood processing and flow cytometry

149 Red blood cells were lysed in ammonium chloride buffer and centrifuged for 10 min at 2500  
150 RPM. Cell pellets were re-suspended and viable cells were counted using Trypan blue staining  
151 solution on a haemocytometer (Nikon, Tokyo Japan). **Flow cytometry was performed using a**  
152 **method adapted from Barnett-Vanes et al., (2015). Briefly, cells were stained** with live-dead  
153 stain (eBioscience), blocked with anti-cd32 blocking antibody and stained with antibodies  
154 CD43 PE (Biolegend) and His48 FITC (Biolegend) for neutrophils and monocytes, CD161  
155 APC (Biolegend) for NK Cells and CD3 VioGreen (Miltenyi Biotec) for T Cells, in buffer  
156 containing PBS, BSA and Azide [29]. Cells were fixed in BD Cell Fix and analysed using a  
157 multi-parameter flow cytometer (Fortessa LSR BD Biosciences New Jersey USA). Flow  
158 cytometric compensation was performed using fluorescent compensation beads (OneComp  
159 eBeads, eBioscience San Diego USA).

#### 160 Cytokine analysis

161 Blood plasma was thawed and analysed using a multiplex ELISA (MesoScaleDiscovery  
162 Maryland, USA) according to the manufacturer's instructions. A microtitre plate pre-coated  
163 with capture antibody was incubated with blocking buffer at room temperature (RT) with  
164 agitation for 1 hour. After washing, standards or samples were added and the plate was  
165 incubated at RT with shaking for 2 hours. The plate was then washed and incubated with  
166 detection antibody conjugated to electrochemiluminescent labels at RT for 2 hours. After  
167 washing, read buffer was added and the plate was read on an MSD plate reader.



168 Histology

169 Liver (Right Lateral Lobe), lung (left lobe) and left quadriceps muscle samples were harvested  
170 and fixed for 24hrs in 10% Buffered Formalin. Paraffin-embedded sections (4  $\mu$ m) were stained  
171 with hematoxylin and eosin (H&E). Images of the slides were captured using a Leica light  
172 microscope (Wetzlar, Germany).

173 Statistics

174 All data were collected from 2-4 individual experiments. Mechanical data are expressed as  
175 mean  $\pm$  standard deviations (SD). Physiological parameters were compared using a two way  
176 repeated measures Analysis of Variance (ANOVA) test using SPSS (version 22.0, Chicago,  
177 USA). All other data were analysed using a non-parametric Mann Whitney t-test expressed as  
178 mean  $\pm$  Standard Error of Mean (SEM) using GraphPad Prism v5 (San Diego, USA), \* $p < 0.05$ ,  
179 \*\* $p < 0.01$ .

180

181 **Results**

182 The characteristics of the blast waves produced in the shock tube are described by mean burst  
183 and corresponding shock wave peak pressures, plateau pressure, and impulse duration from  
184 Sensor 3 for all three test configurations (Table 2).

185 **Table 2: Average burst pressure, corresponding peak pressure, plateau pressure, and shock**  
186 **impulse (relative to ambient pressure) from Sensor 3 for three loading scenarios (mean and**  
187 **standard deviations for each condition are from 3 individual experiments).**

188 Sensor 2 traces show that all three waves exhibit a typical pattern consisting of a positive  
189 overpressure followed by a negative under-pressure and subsequent return to ambient pressure  
190 (Figure 2A). Sensor 3 gave slightly higher amplitudes, faster rise times and longer durations  
191 than the tracings of Sensor 2. The internal stress produced in the sample by a blast is influenced  
192 by the mechanical impedance of the sample, so measurements from the sensor inside the bull-  
193 nose that has a diameter one third the internal diameter of the shock tube<sup>27-26</sup> set the pressure  
194 trace's upper limit and enable readings more relevant to small samples than Sensor 2 regarding  
195 the conditions under which the specimens are loaded. In the absence of Sensor 3 the second  
196 lateral sensor was used to identify the loading pressure magnitude. A linear relationship  
197 between the peak pressures measured by Sensors 2 and 3 (Figure 2B) enabled estimation of the  
198 shock magnitude reaching the subject when only Sensor 2 was used.

199 **Figure 2: (A) Traces from Sensors 2 and 3 (relative to ambient pressure) of three loading**  
200 **configurations and (B) Relationship between the peak pressures (relative to ambient**  
201 **pressure) measured by Sensor 2 and Sensor 3.**

202 The repeated measures ANOVA test determined that there were no statistical differences in the  
203 heart rates measured between the four Groups ( $p=0.842$ ) across the different time points that  
204 the measurements were obtained.

205 At 6hrs significant increases in circulating blood neutrophils were seen in Group III (mean  
206  $2.6E+06$ , SEM  $0.360E+6$ ) compared to Shams (mean  $0.744E+6$ , SEM  $0.161E+6$ ), Groups I  
207 (mean  $0.674E+6$ , SEM  $0.164E+6$ ) and II (mean  $0.540E+3$ , SEM  $0.085E+6$ ). Circulating  
208 monocytes were significantly increased in Group III (mean  $162E+3$ , SEM  $21.8E+3$ ), in  
209 comparison to Shams (mean  $40E+3$ , SEM  $10.1E+3$ ) and Group II (mean  $38.8E+3$ , SEM  
210  $4.60E+3$ ). By 24hrs, no significant differences were observed between sham and blast groups  
211 (Figure 3A). No significant differences were observed in the proportion of CD43Hi/His48Int-  
212 Lo, NK Cell or CD3 T Cells between groups at either timepoint (Figure 3B).

213 At 6hrs significant increases in circulating IL-6 were seen in Group III (mean 223, SEM 74.5)  
214 compared to Shams (mean 34.4, SEM 8.94) and Group I (mean 40.5, SEM 7.19). Increases  
215 were also seen in circulating CXCL1 in Group III compared to Shams but this was not  
216 significant ( $p=0.125$ ). By 24hrs, no differences were observed in IL-6 or CXCL1 between sham  
217 and blast groups (Figure 3C).

218 **Figure 3: (A) Shows the number of circulating neutrophils and CD43Lo/His48Hi monocytes**  
219 **observed at 6hrs and 24hrs for all groups. Representative flow cytometry plots are shown**  
220 **for Sham and Group III at 6hrs. (B) Proportions of circulating CD43Hi/His48Int-Lo**  
221 **monocytes, NK Cells and T Cells were examined. (C) Representative bar graphs showing**  
222 **the concentration of circulating IL-6 and CXCL1 in the plasma. Data are n=4-6, performed**  
223 **in duplicate from 2-3 separate experiments, \* $p<0.05$ , \*\* $p<0.01$ .**

224 No macroscopic changes were seen by histology in limb muscle, liver or lung using H&E  
225 staining at 6hrs in Groups I or III (Figure 4).

226 **Figure 4: Histological sections of muscle, lung and liver tissue. Quadriceps muscle, left lung**  
227 **lobes and superior right lateral liver lobes were fixed in formalin, embedded in a wax block,**  
228 **cut and stained with Hematoxylin and eosin (H&E) stain. Slides were imaged under a light**  
229 **microscope at 10X magnification.**

## 230 **Discussion**

231 The development of a rodent model of primary blast limb trauma using a compression driven  
232 shock tube to deliver an isolated and controlled blast wave is reported. To our knowledge, it is  
233 the first study to examine the relationship between the effects of long versus short duration  
234 primary blast waves on the systemic inflammatory response. This has relevance in the field to  
235 open (free field) versus enclosed (in-vehicle or building) blast waves respectively.

### 236 *Blast Waves and Injury*

237 Blast waves generated by explosive devices may cause severe damage by coupling energy into  
238 human tissues, initiating onward stress wave propagation [30]. Due to the destructive capability  
239 of these waves, particularly at interfaces of differing tissue densities, organs such as the lungs  
240 and bowel are thought to be most susceptible to damage. Studies show the injuries sustained  
241 during explosions are significantly influenced by their environment. In open air blasts,  
242 fragments that are part of the explosive device (primary fragments) or from the explosion  
243 (secondary fragments) are responsible for the majority of injuries sustained [6]. However,

244 victims of bombings that occur in confined spaces suffer a higher incidence of primary blast  
245 injuries, which are more severe and result in clinically worse conditions [6]. For example, a  
246 significantly higher rate of primary blast lung injury (PBLI) is seen amongst mounted/in-  
247 vehicle casualties compared with those dismounted following an external explosion [31].

#### 248 *Animal Models*

249 In experimental studies of primary blast, animals are usually securely fixed to an animal holder,  
250 often using a metal mesh plane or plate [20]. This may be suspended within the main section  
251 [20-21, 32] or across the outlet of the shock tube at varying distances and body orientations,  
252 with or without a reflective plate [33-34] thus, leaving most of the subject's body exposed to  
253 the blast wave. In certain reports, particularly in studies of traumatic brain injury (TBI), a  
254 Kevlar vest [35] or 'body armour' [34] is utilised, wrapped around the animal's thorax to  
255 protect the animals from blast lung injury whilst leaving the head fully exposed. Because of  
256 concerns surrounding thoracic loading despite the presence of Kevlar protection, and the  
257 danger of blast waves pressuring an enclosed protective cylinder, our model utilises a purpose  
258 built experimental rig and harness to deliver isolated shock waves to the left hind limb of a  
259 rodent. However, whilst blast waves have a complex form which varies both in intensity and  
260 duration, most blast research on biological systems has focussed overwhelmingly on shock  
261 overpressure, neglecting the effect of pulse duration which remains poorly understood.

#### 262 *Physiological Response*

263 Studies of blast injury to the thorax show a transient bradycardia following blast as a result of  
264 a vagally driven depressor reflex [36-37]. Though previous studies have recorded shockwaves  
265 developed after missile extremity impact propagating to the abdomen, thorax and brain in  
266 anaesthetised animals [38-40], our observation of no significant differences in heart rate after  
267 blast limb injury - in addition to the lack of histological findings - supports our assertion that  
268 the thorax was not exposed to blast loading. This, therefore, is a true 'isolated blast' model.

#### 269 *Inflammation*

270 Flow cytometry was performed to dissect the inflammatory response and it was found that in  
271 in group III there was a significant increase in circulating neutrophils and CD43<sup>Lo</sup>/His48<sup>Hi</sup>  
272 (Classical) monocytes compared to Shams (Figure 3A) a response that was not seen in groups  
273 I and II. No significant changes were observed in the % of CD43<sup>Hi</sup>/His48<sup>Int-Lo</sup> (non-classical)

274 monocytes, NK cells or T Cells in the blood of rats following blast (Figure 3B). These results  
275 indicate that the systemic inflammatory response is correlated with the duration of the blast  
276 wave rather than the extent of the overpressure. Characterisation of the nature of the cellular  
277 inflammatory response showed a selective increase in neutrophils and CD43<sup>Lo</sup>/His48<sup>Hi</sup>  
278 monocytes consistent with previous reports of rats responding sub-acutely to damage or  
279 inflammatory signals [41-45].

280 Alongside cellular inflammation, at 6 hours we observed raised levels of the inflammatory  
281 cytokines CXCL1 and IL-6 in Group III compared to Shams that was not seen in Groups I and  
282 II suggesting again that the extent of inflammation is correlated with duration rather than the  
283 magnitude of the blast overpressure. Of note CXCL1 has previously been shown to mediate  
284 neutrophil mobilization from the bone marrow-directly in a model of hind limb perfusion [46]  
285 and in a model of blast lung injury [47-48], suggesting that the enhanced levels of CXCL1 seen  
286 in this model may account for the increase in circulating neutrophils. By 24hrs, both mediators  
287 had returned to basal levels. Of note and in contrast with other studies of blast limb trauma,  
288 [16] we observed no substantive macroscopic changes in limb muscle, liver or lung muscle at  
289 6hrs (Figure 4), supporting our method that blast waves were only applied to the limb and did  
290 not inadvertently damage other organs which would complicate the inflammatory response.  
291 Moreover, we saw no limb fractures in this model, likely due to the absence of flail against a  
292 hard surface.

293 Recent studies of more complex models [49-50] have outlined the importance of primary blast  
294 effects in exasperating other injuries and delaying recovery. Moreover, increasing use of novel  
295 explosives which generate blast waves of longer duration necessitates a better understanding  
296 of their impact on injury.

297 In this study, we have developed an experimental set up that allows us to accurately study the  
298 effects of short duration (free-field) and long duration (enclosed spaces) blast of specified  
299 magnitudes in an *in vivo* model. We have shown that focal application of prolonged but not  
300 short duration blast waves to a single hind limb promotes acute increases in circulating  
301 neutrophils and CD43<sup>Lo</sup>/His48<sup>Hi</sup> monocytes, together with rises in pro-inflammatory  
302 cytokines in the absence of any visible gross tissue damage. This research points to an  
303 important role played by blast waves in initiating inflammatory changes in the absence of limb  
304 fracture or penetrating injuries. It is well known that sub-lethal trauma can exacerbate the  
305 inflammatory response to secondary inflammatory stimuli including infectious agents or

306 mechanical stimuli, such as lung ventilation [11 51]. This study shows that the immune system  
307 is exquisitely sensitive to the greater damage induced by blast overpressures of prolonged  
308 duration.

### 309 *Conclusions*

310 This study demonstrates focal application of blast waves to the limb can elicit a systemic  
311 inflammatory response characterised by changes in circulating inflammatory cells and  
312 cytokines. The changes observed are; dependent on the characteristics (notably duration) of the  
313 wave, relatively short-lived (normalising by 24 hours), and occur in the absence of other blast  
314 wave injury mechanisms. Furthermore, the combined biological and mechanical method  
315 developed in this article is a significant step to further our understanding of the complex effect  
316 of blast waves in a range of different tissues and organs *in-vivo*.

### 317 **Acknowledgements**

318 This work was conducted under the auspices of the Royal British Legion Centre for Blast Injury  
319 Studies at Imperial College London. The authors would like to acknowledge the financial  
320 support of the Royal British Legion. The authors wish to additionally thank Mr Satpal Sangha  
321 for his technical support, Ms Lorraine Lawrence for her histological assistance and the Flow  
322 Cytometry facility at South Kensington Campus for their expert assistance and advice. The  
323 Institute of Shock Physics acknowledges the support of AWE, Aldermaston and Imperial  
324 College London. The sponsors had no involvement in the study design; collection, analysis and  
325 interpretation of data; the writing of the manuscript; or the decision to submit the manuscript  
326 for publication.

327

328 REFERENCES

- 329 1. Birch R, Misra P, Stewart MP, Eardley WG, Ramasamy A, Brown K, Shenoy R, Anand P,  
330 Clasper J, Dunn R et al. . Nerve injuries sustained during warfare: Part I--epidemiology. *J Bone*  
331 *Joint Surg Br.* 2012; 94:523-528.
- 332 2. Lesperance RN, Nunez TC. Blast injury: Impact on brain and internal organs. *Crit Care Nurs*  
333 *Clin North Am.* 2015; 27:277-287.
- 334 3. Wolf SJ, Bebarta VS, Bonnett CJ, Pons PT, Cantrill SV. Blast injuries. *Lancet.* 2009;  
335 374:405-415.
- 336 4. Kirkman E, Watts S. Characterization of the response to primary blast injury. *Philos Trans*  
337 *R Soc Lond B Biol Sci.* 2011; 366:286-290.
- 338 5. Mackenzie I, Keene D, Tunnicliffe B, Smith J, Mahoney P, Gibb I, Leason J, Kirkman E.  
339 Blast lung injuries: A clinical perspective from current operations. 2011.
- 340 6. Leibovici D, Gofrit ON, Stein M, Shapira SC, Noga Y, Heruti RJ, Shemer J. Blast injuries:  
341 Bus versus open-air bombings--a comparative study of injuries in survivors of open-air versus  
342 confined-space explosions. *J Trauma.* 1996; 41:1030-1035.
- 343 7. Frykberg ER, Tepas JJ,3rd. Terrorist bombings. lessons learned from belfast to beirut. *Ann*  
344 *Surg.* 1988; 208:569-576.
- 345 8. Singleton JAG, Gibb IE, Hunt NCA, Bull AMJ, Clasper JC. Identifying future 'unexpected'  
346 survivors: A retrospective cohort study of fatal injury patterns in victims of improvised  
347 explosive devices. *BMJ Open.* 2013;3.
- 348 9. Lenz A, Franklin GA, Cheadle WG. Systemic inflammation after trauma. *Injury.* 2007;  
349 38:1336-1345.
- 350 10. Manson J, Thiemermann C, Brohi K. Trauma alarmins as activators of damage-induced  
351 inflammation. *Br J Surg.* 2012; 99 Suppl 1:12-20.
- 352 11. Lord JM, Midwinter MJ, Chen Y, Belli A, Brohi K, Kovacs EJ, Koenderman L, Kubes P,  
353 Lilford RJ. The systemic immune response to trauma: An overview of pathophysiology and  
354 treatment. 2014; 384:1455-1465.

- 355 12. Irwin RJ, Lerner MR, Bealer JF, Mantor PC, Brackett DJ, Tuggle DW. Shock after blast  
356 wave injury is caused by a vagally mediated reflex. *J Trauma*. 1999; 47:105-110.
- 357 13. Ramasamy A, Hill AM, Masouros S, Gibb I, Bull AM, Clasper JC. Blast-related fracture  
358 patterns: A forensic biomechanical approach. *J R Soc Interface*. 2011; 8:689-698.
- 359 14. Ning JL, Mo LW, Lu KZ, Lai XN, Wang ZG, Ma D. Lung injury following lower extremity  
360 blast trauma in rats. *J Trauma Acute Care Surg*. 2012; 73:1537-1544.
- 361 15. Ning J, Mo L, Zhao H, Lu K, Lai X, Luo X, Zhao H, Ma D. Sodium hydrosulphide  
362 alleviates remote lung injury following limb traumatic injury in rats. *PLoS ONE*. 2013;  
363 8:e59100.
- 364 16. Ning J, Mo L, Zhao H, Lu K, Wang L, Lai X, Yang B, Zhao H, Sanders RD, Ma D.  
365 Transient regional hypothermia applied to a traumatic limb attenuates distant lung injury  
366 following blast limb trauma. *Crit Care Med*. 2014; 42:e68-78.
- 367 17. Zhao H, Ning J, Duan J, Gu J, Yi B, Lu K, Mo L, Lai X, Hennah L, Ma D. Regional  
368 traumatic limb hypothermia attenuates distant hepatic and renal injury following blast limb  
369 trauma in rats. *J Trauma Acute Care Surg*. 2014; 77:464-470.
- 370 18. Rizzo JA, Colombo CJ. Does blast limb trauma constitute a multisystem critical illness?\*.  
371 *Crit Care Med*. 2014; 42.
- 372 19. Panzer MB, Wood GW, Bass CR. Scaling in neurotrauma: How do we apply animal  
373 experiments to people? *Exp Neurol*. 2014; 261:120-126.
- 374 20. Cernak I, Merkle AC, Koliatsos VE, Bilik JM, Luong QT, Mahota TM, Xu L, Slack N,  
375 Windle D, Ahmed FA. The pathobiology of blast injuries and blast-induced neurotrauma as  
376 identified using a new experimental model of injury in mice. *Neurobiol Dis*. 2011; 41:538-551.
- 377 21. Goldstein LE, Fisher AM, Tagge CA, Zhang X, Velisek L, Sullivan JA, Upreti C, Kracht  
378 JM, Ericsson M, Wojnarowicz MW et al. . Chronic traumatic encephalopathy in blast-exposed  
379 military veterans and a blast neurotrauma mouse model. *Science Translational Medicine*. 2012;  
380 4:134ra60-134ra60.
- 381 22. Risling M, Plantman S, Angeria M, Rostami E, Bellander B-, Kirkegaard M, Arborelius U,  
382 Davidsson J. Mechanisms of blast induced brain injuries, experimental studies in rats.  
383 *Neuroimage*. 2011; 54, Supplement 1:S89-S97.



- 384 23. Rafaels KA, Bass CR, Panzer MB, Salzar RS. Pulmonary injury risk assessment for long-  
385 duration blasts: A meta-analysis. *J Trauma*. 2010; 69:368-374.
- 386 24. Cernak I, Merkle AC, Koliatsos VE, Bilik JM, Luong QT, Mahota TM, Xu L, Slack N,  
387 Windle D, Ahmed FA. The pathobiology of blast injuries and blast-induced neurotrauma as  
388 identified using a new experimental model of injury in mice. *Neurobiol Dis*. 2011; 41:538-551.
- 389 25. Cernak I, Noble-Haeusslein L. Traumatic brain injury: An overview of pathobiology with  
390 emphasis on military populations. *J Cereb Blood Flow Metab*. 2009; 30:255-266.
- 391 26. Nguyen T, Wilgeroth J, Macdonald W, Proud W. Methods of controlled shock wave  
392 generation in A shock tube for biological applications. 2013; 58.
- 393 27. Nguyen TN, Wilgeroth J, Proud W. Controlling blast wave generation in a shock tube for  
394 biological applications. 2014; 500:142025.
- 395 28. Jang HS, Choi HS, Lee SH, Jang KH, Lee MG. Evaluation of the anaesthetic effects of  
396 medetomidine and ketamine in rats and their reversal with atipamezole. *Vet Anaesth Analg*.  
397 2009; 36:319-327.
- 398 29. Barnett-Vanes A, Sharrock A, Birrel M, Rankin SM. A single 9-colour flow cytometric  
399 method to characterise major leukocyte populations in the rat: Validation in a model of LPS-  
400 induced pulmonary inflammation. *PloS One*, 2016 (In press)
- 401 30. Cooper GJ, Townend DJ, Cater SR, Pearce BP. The role of stress waves in thoracic visceral  
402 injury from blast loading: Modification of stress transmission by foams and high-density  
403 materials. *J Biomech*. 1991; 24:273-285.
- 404 31. Singleton JA, Gibb IE, Bull AM, Mahoney PF, Clasper JC. Primary blast lung injury  
405 prevalence and fatal injuries from explosions: Insights from postmortem computed  
406 tomographic analysis of 121 improvised explosive device fatalities. *J Trauma Acute Care Surg*.  
407 2013; 75:S269-74.
- 408 32. Skotak M, Wang F, Alai A, Holmberg A, Harris S, Switzer RC, Chandra N. Rat injury  
409 model under controlled field-relevant primary blast conditions: Acute response to a wide range  
410 of peak overpressures. *J Neurotrauma*. 2013; 30:1147-1160.
- 411 33. Long JB, Bentley TL, Wessner KA, Cerone C, Sweeney S, Bauman RA. Blast overpressure  
412 in rats: Recreating a battlefield injury in the laboratory. *J Neurotrauma*. 2009; 26:827-840.

- 413 34. Svetlov SI, Prima V, Kirk DR, Gutierrez H, Curley KC, Hayes RL, Wang KK. Morphologic  
414 and biochemical characterization of brain injury in a model of controlled blast overpressure  
415 exposure. *J Trauma*. 2010; 69:795-804.
- 416 35. Sundaramurthy A, Alai A, Ganpule S, Holmberg A, Plougonven E, Chandra N. Blast-  
417 induced biomechanical loading of the rat: An experimental and anatomically accurate  
418 computational blast injury model. *J Neurotrauma*. 2012; 29:2352-2364.
- 419 36. Ohnishi M, Kirkman E, Guy RJ, Watkins PE. Reflex nature of the cardiorespiratory  
420 response to primary thoracic blast injury in the anaesthetised rat. *Exp Physiol*. 2001; 86:357-  
421 364.
- 422 37. Skotak M, Wang F, Alai A, Holmberg A, Harris S, Switzer RC, Chandra N. Rat injury  
423 model under controlled field-relevant primary blast conditions: Acute response to a wide range  
424 of peak overpressures. *J Neurotrauma*. 2013; 30:1147-1160.
- 425 38. Suneson A, Hansson HA, Seeman T. Pressure wave injuries to the nervous system caused  
426 by high-energy missile extremity impact: Part II. distant effects on the central nervous system-  
427 a light and electron microscopic study on pigs. *J Trauma*. 1990; 30:295-306.
- 428 39. Berlin R. Energy transfer and regional blood flow changes following missile trauma. *J*  
429 *Trauma*. 1979; 19:170-176.
- 430 40. Tikka S, Cederberg A, Rokkanen P. Remote effects of pressure waves in missile trauma.  
431 the intra-abdominal pressure changes in anesthetized pigs wounded in one thigh. *Acta Chir*  
432 *Scand Suppl*. 1982; 508:167-173.
- 433 41. Scriba A, Schneider M, Grau V, van der Meide PH, Steiniger B. Rat monocytes up-regulate  
434 NKR-P1A and down-modulate CD4 and CD43 during activation in vivo: Monocyte  
435 subpopulations in normal and IFN-gamma-treated rats. *J Leukoc Biol*. 1997; 62:741-752.
- 436 42. Lu R, Zhou X, Ji W, Luo Y, Yang G, Sun H, Ge L, Jiang T, Li Y. Dynamic changes of  
437 peripheral blood monocyte subsets after myocardial ischemia/reperfusion in rats. *Xi Bao Yu*  
438 *Fen Zi Mian Yi Xue Za Zhi*. 2014; 30:131-134.
- 439 43. Ziegler-Heitbrock L. Monocyte subsets in man and other species. *Cell Immunol*. 2014;  
440 289:135-139.

441 44. Scriba A, Schneider M, Grau V, van der Meide PH, Steiniger B. Rat monocytes up-regulate  
442 NKR-P1A and down-modulate CD4 and CD43 during activation in vivo: Monocyte  
443 subpopulations in normal and IFN-gamma-treated rats. *Journal of Leukocyte Biology*. 1997;  
444 62:741-752.

445 45. Grau V, Scriba A, Stehling O, Steiniger B. Monocytes in the rat. *Immunobiology*. 2000;  
446 202:94-103.

447 46. Burdon PCE, Martin C, Rankin SM. Migration across the sinusoidal endothelium regulates  
448 neutrophil mobilization in response to ELR + CXC chemokines. *Br J Haematol*. 2008; 142:100-  
449 108.

450 47. Chavko M, Prusaczyk WK, McCarron RM. Lung injury and recovery after exposure to  
451 blast overpressure. *J Trauma*. 2006; 61:933-942.

452 48. Chavko M, Adeeb S, Ahlers ST, McCarron RM. Attenuation of pulmonary inflammation  
453 after exposure to blast overpressure by N-acetylcysteine amide. *Shock*. 2009; 32:325-331.

454 49. Claes L, Ignatius A, Lechner R, Gebhard F, Kraus M, Baumgartel S, Recknagel S, Krischak  
455 GD. The effect of both a thoracic trauma and a soft-tissue trauma on fracture healing in a rat  
456 model. *Acta Orthop*. 2011; 82:223-227.

457 50. Polfer EM, Hope DN, Elster EA, Qureshi AT, Davis TA, Golden D, Potter BK, Forsberg  
458 JA. The development of a rat model to investigate the formation of blast-related post-traumatic  
459 heterotopic ossification. *Bone Joint J*. 2015; 97-B:572-576.

460 51. van Wessem KJP, Hennis MP, van Wagenberg L, Koenderman L, Leenen LPH.  
461 Mechanical ventilation increases the inflammatory response induced by lung contusion. *J Surg*  
462 *Res*. 2013; 183:377-384.

463