

MAMORU NUNOKAWA, EMEL YAVUZ DUMAN  
and SHIGEYOSHI OWA**Properties of functions concerned with  
Carathéodory functions**

ABSTRACT. Let  $\mathcal{P}_n$  denote the class of analytic functions  $p(z)$  of the form  $p(z) = 1 + c_n z^n + c_{n+1} z^{n+1} + \dots$  in the open unit disc  $\mathbb{U}$ . Applying the result by S. S. Miller and P. T. Mocanu (J. Math. Anal. Appl. **65** (1978), 289–305), some interesting properties for  $p(z)$  concerned with Carathéodory functions are discussed. Further, some corollaries of the results concerned with the result due to M. Obradović and S. Owa (Math. Nachr. **140** (1989), 97–102) are shown.

**1. Introduction.** Let  $\mathcal{A}_n$  denote the class of functions  $f(z)$  of the form

$$(1.1) \quad f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k \quad (n = 1, 2, 3, \dots)$$

which are analytic in the open unit disc  $\mathbb{U} = \{z \in \mathbb{C} \mid |z| < 1\}$ . If a function  $f(z) \in \mathcal{A}_n$  satisfies

$$(1.2) \quad \operatorname{Re} \left( \frac{z f'(z)}{f(z)} \right) > 0 \quad (z \in \mathbb{U}),$$

then  $f(z)$  is said to be starlike with respect to the origin in  $\mathbb{U}$ . We denote by  $\mathcal{S}_n^*$  the subclass of  $\mathcal{A}_n$  consisting of functions  $f(z)$  which are starlike with respect to the origin in  $\mathbb{U}$ . From the definition of the class  $\mathcal{S}_n^*$ , we see that

---

2000 *Mathematics Subject Classification.* Primary 30C45.

*Key words and phrases.* Analytic, starlike, Carathéodory function.

if  $f(z) \in \mathcal{A}_n$  satisfies

$$(1.3) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| < 1 \quad (z \in \mathbb{U}),$$

then  $f(z) \in \mathcal{S}_n^*$ . We denote by  $\mathcal{T}_n^*$  the subclass of  $\mathcal{S}_n^*$  consisting of  $f(z)$  satisfying (1.3).

Obradović and Owa [5] have shown the following result:

**Theorem A.** *If  $f(z) \in \mathcal{A}_1$  satisfies  $f(z)f'(z) \neq 0$  for  $0 < |z| < 1$  and*

$$(1.4) \quad \left| 1 + \frac{zf''(z)}{f'(z)} \right| < \frac{5}{4} \left| \frac{zf'(z)}{f(z)} \right| \quad (z \in \mathbb{U}),$$

then  $f(z) \in \mathcal{T}_1^*$ .

In order to discuss our results, we have to recall here the following lemma due to Miller and Mocanu [3] (also due to Jack [2]):

**Lemma 1.1.** *Let*

$$w(z) = a_n z^n + a_{n+1} z^{n+1} + \dots \quad (a_n \neq 0)$$

be analytic in  $\mathbb{U}$ . If there exists a point  $z_0 \in \mathbb{U}$  on the circle  $|z| = r < 1$  such that

$$(1.5) \quad \max_{|z| \leq |z_0|} |w(z)| = |w(z_0)|,$$

then we can write

$$(1.6) \quad z_0 w'(z_0) = m w(z_0),$$

where  $m$  is real and  $m \geq n$ .

**Example 1.1.** We consider the function  $w(z)$  given by

$$(1.7) \quad w(z) = z^n + \frac{e^{i\theta}}{n+1} z^{n+1} \quad (n = 1, 2, 3, \dots).$$

Then, it follows that

$$(1.8) \quad \max_{|z| \leq |z_0|} |w(z)| = \max_{|z| \leq |z_0|} |z|^n \left| 1 + \frac{e^{i\theta} z}{n+1} \right| \leq r^n \left( 1 + \frac{r}{n+1} \right)$$

for  $z_0 = r e^{-i\theta} \in \mathbb{U}$ . This shows that  $|w(z)|$  attains its maximum value at a point  $z_0 \in \mathbb{U}$  on the circle  $|z| = r$ . For such a point  $z_0 = r e^{-i\theta}$ , we have that

$$(1.9) \quad \frac{z_0 w'(z_0)}{w(z_0)} = \frac{z_0^n (n + e^{i\theta} z_0)}{z_0^n \left( 1 + \frac{e^{i\theta} z_0}{n+1} \right)} = \frac{(n+1)(n+r)}{n+1+r} = m \geq n.$$

Let  $\mathcal{P}_n$  be the class of functions  $p(z)$  of the form

$$(1.10) \quad p(z) = 1 + \sum_{k=n}^{\infty} c_k z^k \quad (c_n \neq 0)$$

which are analytic in  $\mathbb{U}$ . We also denote by  $\mathcal{Q}_n$  the subclass of  $\mathcal{P}_n$  consisting of  $f(z)$  which satisfy

$$(1.11) \quad |p(z) - 1| < 1 \quad (z \in \mathbb{U}).$$

Since  $p(z) \in \mathcal{Q}_n$  shows that  $\operatorname{Re} p(z) > 0$  ( $z \in \mathbb{U}$ ),  $p(z) \in \mathcal{Q}_n$  is said to be a Carathéodory function in  $\mathbb{U}$  (see Carathéodory [1]).

**2. Conditions for the classes  $\mathcal{Q}_n$  and  $\mathcal{T}_n^*$ .** Applying Lemma 1.1, we discuss some conditions for  $p(z) \in \mathcal{P}_n$  to be in the class  $\mathcal{Q}_n$ .

**Theorem 2.1.** *If  $p(z) \in \mathcal{P}_n$  satisfies*

$$(2.1) \quad \operatorname{Re} \left( p(z) + \alpha \frac{zp'(z)}{p(z)} \right) < \sqrt{\alpha n} |p(z)| \quad (z \in \mathbb{U})$$

for some real  $\alpha > 0$ , then  $p(z) \in \mathcal{Q}_n$ .

**Proof.** Note that  $p(z) \neq 0$  ( $z \in \mathbb{U}$ ) with the condition (2.1). Let us define the function  $w(z)$  by

$$(2.2) \quad p(z) = 1 + w(z) \quad (z \in \mathbb{U})$$

for  $p(z) \in \mathcal{P}_n$ . Then  $w(z)$  is analytic in  $\mathbb{U}$  and

$$(2.3) \quad w(z) = c_n z^n + c_{n+1} z^{n+1} + \dots$$

It follows that

$$(2.4) \quad p(z) + \alpha \frac{zp'(z)}{p(z)} = 1 + w(z) + \frac{\alpha zw'(z)}{1 + w(z)}$$

and that

$$(2.5) \quad \begin{aligned} \frac{1}{|p(z)|} \operatorname{Re} \left( p(z) + \alpha \frac{zp'(z)}{p(z)} \right) \\ = \frac{1}{|1 + w(z)|} \operatorname{Re} \left( 1 + w(z) + \frac{\alpha zw'(z)}{1 + w(z)} \right) < \sqrt{\alpha n} \end{aligned}$$

for  $z \in \mathbb{U}$ .

We suppose that there exists a point  $z_0 \in \mathbb{U}$  such that

$$(2.6) \quad \max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1.$$

Then, Lemma 1.1 gives us that  $w(z_0) = e^{i\theta}$  and  $z_0 w'(z_0) = m e^{i\theta}$  ( $m \geq n$ ). For such a point  $z_0$ , we have that

$$\begin{aligned}
 \frac{1}{|p(z_0)|} \operatorname{Re} \left( p(z_0) + \alpha \frac{z_0 p'(z_0)}{p(z_0)} \right) &= \frac{1}{|1 + e^{i\theta}|} \operatorname{Re} \left( 1 + e^{i\theta} + \frac{\alpha m e^{i\theta}}{1 + e^{i\theta}} \right) \\
 (2.7) \qquad \qquad \qquad &= \frac{1}{\sqrt{2(1 + \cos \theta)}} \left( 1 + \cos \theta + \frac{\alpha m}{2} \right) \\
 &= \frac{1}{\sqrt{2}} \left( \sqrt{1 + \cos \theta} + \frac{\alpha m}{2\sqrt{1 + \cos \theta}} \right) \\
 &\geq \sqrt{\alpha m} \geq \sqrt{\alpha n}.
 \end{aligned}$$

This contradicts the condition (2.1). Therefore, there is no such point  $z_0 \in \mathbb{U}$ . This means that  $p(z) \in \mathcal{Q}_n$ .  $\square$

**Corollary 2.1.** *If  $f(z) \in \mathcal{A}_n$  satisfies  $f(z)f'(z) \neq 0$  for  $0 < |z| < 1$  and*

$$(2.8) \quad \operatorname{Re} \left\{ (1 - \alpha) \frac{z f'(z)}{f(z)} + \alpha \left( 1 + \frac{z f''(z)}{f'(z)} \right) \right\} < \sqrt{\alpha n} \left| \frac{z f'(z)}{f(z)} \right| \quad (z \in \mathbb{U})$$

for some real  $\alpha > 0$ , then  $f(z) \in \mathcal{T}_n^*$ .

**Proof.** Letting  $p(z) = \frac{z f'(z)}{f(z)}$  in Theorem 2.1, we have that

$$p(z) + \alpha \frac{z p'(z)}{p(z)} = (1 - \alpha) \frac{z f'(z)}{f(z)} + \alpha \left( 1 + \frac{z f''(z)}{f'(z)} \right).$$

The proof of the corollary follows from the above.  $\square$

Next we derive

**Theorem 2.2.** *If  $p(z) \in \mathcal{P}_n$  satisfies  $\operatorname{Re} p(z) \neq 0$  ( $z \in \mathbb{U}$ ) and*

$$(2.9) \quad \operatorname{Re} \left( p(z) + \alpha \frac{z p'(z)}{p(z)} \right) < \left( 1 + \frac{\alpha n}{4} \right) \operatorname{Re} p(z) \quad (z \in \mathbb{U})$$

for some real  $\alpha > 0$ , then  $p(z) \in \mathcal{Q}_n$ .

**Proof.** Define the function  $w(z)$  by (2.2) for  $p(z) \in \mathcal{P}_n$ . Then,  $w(z)$  is analytic in  $\mathbb{U}$ ,

$$w(z) = c_n z^n + c_{n+1} z^{n+1} + \dots,$$

and

$$(2.10) \quad \frac{\operatorname{Re} \left( p(z) + \alpha \frac{z p'(z)}{p(z)} \right)}{\operatorname{Re} p(z)} = \frac{\operatorname{Re} \left( 1 + w(z) + \frac{\alpha z w'(z)}{1 + w(z)} \right)}{\operatorname{Re}(1 + w(z))} < 1 + \frac{\alpha n}{4}$$

( $z \in \mathbb{U}$ ). If we suppose that there exists a point  $z_0 \in \mathbb{U}$  on the circle  $|z| = r < 1$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1,$$

we can write that  $w(z_0) = e^{i\theta}$  and  $z_0 w'(z_0) = m e^{i\theta}$ . This shows that

$$(2.11) \quad \frac{\operatorname{Re} \left( p(z_0) + \alpha \frac{z_0 p'(z_0)}{p(z_0)} \right)}{\operatorname{Re} p(z_0)} = \frac{1 + \cos \theta + \frac{\alpha m}{2}}{1 + \cos \theta} \geq 1 + \frac{\alpha m}{4} \geq 1 + \frac{\alpha n}{4}.$$

Since (2.11) contradicts our condition (2.9),  $|w(z)| < 1$  for all  $z \in \mathbb{U}$ . This means that  $p(z) \in \mathcal{Q}_n$ .  $\square$

If we take  $p(z) = \frac{z f'(z)}{f(z)}$  in Theorem 2.2, we have

**Corollary 2.2.** *If  $f(z) \in \mathcal{A}_n$  satisfies  $\operatorname{Re} \left( \frac{z f'(z)}{f(z)} \right) \neq 0$  ( $z \in \mathbb{U}$ ) and*

$$(2.12) \quad \operatorname{Re} \left\{ (1 - \alpha) \frac{z f'(z)}{f(z)} + \alpha \left( 1 + \frac{z f''(z)}{f'(z)} \right) \right\} < \left( 1 + \frac{\alpha n}{4} \right) \operatorname{Re} \left( \frac{z f'(z)}{f(z)} \right)$$

( $z \in \mathbb{U}$ ) for some real  $\alpha > 0$ , then  $f(z) \in \mathcal{T}_n^*$ .

**Corollary 2.3.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$(2.13) \quad \operatorname{Re} \left( \frac{z f''(z)}{f'(z)} \right) < \operatorname{Re} \left( \frac{z f'(z)}{f(z)} \right) + \frac{n-2}{n} \quad (z \in \mathbb{U}),$$

then  $f(z) \in \mathcal{T}_n^*$ .

**Proof.** If we write

$$\frac{z f'(z)}{f(z)} = 1 + w(z) \quad (f(z) \in \mathcal{A}_n),$$

we see that  $w(z)$  is analytic in  $\mathbb{U}$  and

$$w(z) = c_n z^n + c_{n+1} z^{n+1} + \dots$$

For such a function  $w(z)$ , we see that

$$(2.14) \quad \operatorname{Re} \left( \frac{z f''(z)}{f'(z)} - \frac{z f'(z)}{f(z)} \right) = \operatorname{Re} \left( \frac{z w'(z)}{1 + w(z)} - 1 \right) < \frac{n-2}{2} \quad (z \in \mathbb{U}).$$

Supposing that there exists a point  $z_0 \in \mathbb{U}$  on the circle  $|z| = r < 1$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1,$$

we can write that  $w(z_0) = e^{i\theta}$  and  $z_0 w'(z_0) = m e^{i\theta}$ . Therefore, we have

$$(2.15) \quad \operatorname{Re} \left( \frac{z_0 f''(z_0)}{f'(z_0)} - \frac{z_0 f'(z_0)}{f(z_0)} \right) = \operatorname{Re} \left( \frac{k e^{i\theta}}{1 + e^{i\theta}} - 1 \right) = \frac{k}{2} - 1 \geq \frac{n-2}{2},$$

which contradicts the condition (2.13). This implies that  $f(z) \in \mathcal{T}_n^*$ .  $\square$

**Example 2.1.** Let us consider the function  $p(z)$  given by

$$(2.16) \quad p(z) = 1 + a_n z^n \quad (z \in \mathbb{U})$$

for some  $n \in \mathbb{N} = \{1, 2, 3, \dots\}$ , where  $a_n$  satisfies

$$a_n^3 + 2a_n - 1 \leq 0 \quad (0 < a_n < 1).$$

Then  $p(z) \in \mathcal{P}_n$  and  $p(z) \neq 0$  ( $z \in \mathbb{U}$ ). It is clear that  $p(z)$  satisfies the condition (2.9) in Theorem 2.2 for  $z = 0$ .

Let us put  $z = e^{i\theta}$  for  $p(z)$ . Then we see that

$$(2.17) \quad \operatorname{Re} \left( p(z) + \alpha \frac{z p'(z)}{p(z)} \right) = 1 + a_n \cos n\theta + \frac{\alpha n a_n (a_n + \cos n\theta)}{a_n^2 + 1 + 2a_n \cos n\theta}$$

and

$$(2.18) \quad \left(1 + \frac{\alpha n}{4}\right) \operatorname{Re} p(z) = \left(1 + \frac{\alpha n}{4}\right) (1 + a_n \cos n\theta).$$

This gives us that

$$(2.19) \quad \begin{aligned} & \left(1 + \frac{\alpha n}{4}\right) \operatorname{Re} p(z) - \operatorname{Re} \left( p(z) + \alpha \frac{z p'(z)}{p(z)} \right) \\ &= \frac{\alpha n (1 + 2a_n \cos n\theta + a_n^3 \cos n\theta + 2a_n^2 \cos^2 n\theta)}{4(a_n^2 + 1 + 2a_n \cos n\theta)} \\ &\geq \frac{\alpha n (1 - 2a_n - a_n^3)}{4(a_n^2 + 1 + 2a_n \cos n\theta)} \geq 0. \end{aligned}$$

Therefore, the function  $p(z)$  satisfies the condition (2.9) for all  $z \in \mathbb{U}$ . Indeed, we see that

$$|p(z) - 1| = |a_n z^n| < a_n < 1 \quad (z \in \mathbb{U}).$$

Furthermore, if we define the function  $f(z) \in \mathcal{A}_n$  by

$$(2.20) \quad \frac{z f'(z)}{f(z)} = 1 + a_n z^n$$

with some real  $a_n$  ( $0 < a_n < 1$ ) satisfying

$$a_n^3 + 2a_n - 1 \leq 0,$$

then we have that

$$(2.21) \quad f(z) = z e^{\frac{\alpha n}{n} z^n}$$

which satisfies the condition (2.12) in Corollary 2.2.

If we consider the function

$$g(x) = x^3 + 2x - 1 \quad (0 < x < 1),$$

we see that  $g(0) = -1 < 0$  and  $g\left(\frac{1}{2}\right) = \frac{1}{8} > 0$ . Therefore, there exists some real  $x$  ( $0 < x < 1$ ) such that  $g(x) \leq 0$ . Indeed, we see that

$$0.4533 < x < 0.4534.$$

**3. Properties for the classes  $\mathcal{P}_n$  and  $\mathcal{A}_n$ .** We discuss some properties for functions in the classes  $\mathcal{P}_n$  and  $\mathcal{A}_n$ .

**Theorem 3.1.** *If  $p(z) \in \mathcal{P}_n$  satisfies*

$$(3.1) \quad \int_{|z|=r} \left| \operatorname{Re} \left( \frac{zp'(z)}{p(z)} \right) \right| d\theta < \pi$$

for  $z = re^{i\theta}$  ( $0 < r < 1$ ), then  $\operatorname{Re} p(z) > 0$  ( $z \in \mathbb{U}$ ).

**Proof.** It follows from (3.1) that

$$(3.2) \quad \int_{|z|=r} \left| \operatorname{Re} \left( \frac{zp'(z)}{p(z)} \right) \right| d\theta = \int_0^{2\pi} \left| \frac{d \arg p(z)}{d\theta} \right| d\theta = \int_{|z|=r} |d \arg p(z)| < \pi.$$

This implies that  $\operatorname{Re} p(z) > 0$  for  $|z| = r < 1$ . Applying the maximum principle for harmonic functions, we obtain that  $\operatorname{Re} p(z) > 0$  ( $z \in \mathbb{U}$ ).  $\square$

From Theorem 3.1, we have

**Corollary 3.1.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$(3.3) \quad \int_{|z|=r} \left| \operatorname{Re} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right| d\theta < \pi$$

for  $z = re^{i\theta}$  ( $0 < r < 1$ ), then  $f(z) \in \mathcal{S}_n^*$ .

Further, applying the same method as the proof by Umezawa [5] and Nunokawa [3], we derive the following result:

**Theorem 3.2.** *If  $f(z) \in \mathcal{A}_1$  satisfies*

$$(3.4) \quad -\frac{\beta}{4\beta - 1} < \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) < \beta \quad (z \in \mathbb{U})$$

for some real  $\beta \geq \frac{1}{4}$ , then  $\operatorname{Re} f'(z) > 0$  ( $z \in \mathbb{U}$ ).

**Proof.** We note that if  $f'(z_0) = 0$  for some  $z_0 \in \mathbb{U}$ , then  $f(z)$  does not satisfy the condition (3.4). This shows that  $f'(z) \neq 0$  for all  $z \in \mathbb{U}$ . Applying the same method by Umezawa [5] and Nunokawa [3], we have that

$$(3.5) \quad \int_{|z|=r} \frac{zf''(z)}{f'(z)} d\theta = \int_{|z|=r} \frac{zf''(z)}{f'(z)} \frac{dz}{iz} = -i \int_{|z|=r} \frac{zf''(z)}{f'(z)} dz = 0.$$

We denote by  $\mathcal{C}_1$  the part of the circle  $|z| = r$  on which

$$(3.6) \quad \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) \geq 0$$

and

$$(3.7) \quad \int_{\mathcal{C}_1} d \arg z = x.$$

On the other hand, let us denote by  $\mathcal{C}_2$  the part of the circle  $|z| = r$  on which

$$(3.8) \quad \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) < 0$$

and

$$(3.9) \quad \int_{\mathcal{C}_2} d \arg z = 2\pi - x.$$

Putting

$$(3.10) \quad y_1 = \int_{\mathcal{C}_1} \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) d\theta = \int_{\mathcal{C}_1} \left( \frac{d \arg f'(z)}{d\theta} \right) d\theta$$

and

$$(3.11) \quad -y_2 = \int_{\mathcal{C}_2} \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) d\theta = \int_{\mathcal{C}_2} \left( \frac{d \arg f'(z)}{d\theta} \right) d\theta,$$

we have that  $y_1 - y_2 = 0$ .

In view of the condition (3.4), we obtain that

$$y_1 < \beta x \quad \text{and} \quad y_2 < \frac{\beta}{4\beta - 1}(2\pi - x).$$

If  $y_1 \geq \frac{\pi}{2}$ , then  $y_2 = y_1 \geq \frac{\pi}{2}$  and  $\frac{\pi}{2} < \beta x$ . On the other hand, we have that

$$(3.12) \quad y_2 < \frac{\beta}{4\beta - 1}(2\pi - x) < \frac{2\pi\beta - \frac{\pi}{2}}{4\beta - 1} = \frac{\pi}{2}.$$

This contradicts the inequality  $y_2 \geq \frac{\pi}{2}$ . Therefore,  $y_1 = y_2 < \frac{\pi}{2}$ . Consequently, we obtain that

$$(3.13) \quad y_1 + y_2 = \int_{|z|=r} \left| \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) \right| d\theta = \int_{|z|=r} |d \arg f'(z)| < \pi,$$

which implies that  $\operatorname{Re} f'(z) > 0$  ( $z \in \mathbb{U}$ ).  $\square$

Finally, letting  $\beta \rightarrow \infty$ ,  $\beta = \frac{1}{4}$  and  $\beta = \frac{1}{2}$  in Theorem 3.2, we have the following corollary.

**Corollary 3.2.** *If  $f(z) \in \mathcal{A}_1$  satisfies one of the following conditions*

$$(3.14) \quad \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) > -\frac{1}{4} \quad (z \in \mathbb{U}),$$

$$(3.15) \quad \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) < \frac{1}{4} \quad (z \in \mathbb{U}),$$

$$(3.16) \quad \left| \operatorname{Re} \left( \frac{zf''(z)}{f'(z)} \right) \right| < 1 \quad (z \in \mathbb{U}),$$

then  $\operatorname{Re} f'(z) > 0$  ( $z \in \mathbb{U}$ ).



## REFERENCES

- [1] Carathéodory, C., *Über den Variabilitätsbereich der Koeffizienten von Potenzreihen, die gegebene Werte nicht annehmen*, Math. Ann. **64**(1907), 95–115.
- [2] Jack, I. S., *Functions starlike and convex of order  $\alpha$* , J. London Math. Soc. **3** (1971), 469–474.
- [3] Miller, S. S., Mocanu, P. T., *Second order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65** (1978), 289–305.
- [4] Nunokawa, M., *On the Bazilevič analytic functions*, Sci. Rep. Fac. Edu. Gunma Univ. **21** (1972), 9–13.
- [5] Obradović, M., Owa, S., *A criterion for starlikeness*, Math. Nachr. **140** (1989), 97–102.
- [6] Umezawa, T., *Analytic functions convex in one direction*, J. Math. Soc. Japan **4** (1952), 194–202.

Mamoru Nunokawa  
Emeritus Professor of University of Gunma  
Hoshikuki 798-8, Chuou-Ward  
Chiba, Chiba 260-0808  
Japan  
e-mail: [mamoru\\_nuno@doctor.nifty.jp](mailto:mamoru_nuno@doctor.nifty.jp)

Emel Yavuz Duman  
Department of Mathematics and Computer Science  
İstanbul Kültür University  
34156 Bakıröy, İstanbul  
Turkey  
e-mail: [e.yavuz@iku.edu.tr](mailto:e.yavuz@iku.edu.tr)

Shigeyoshi Owa  
Department of Mathematics  
Kinki University  
Higashi-Osaka, Osaka 577-8502  
Japan  
e-mail: [shige21@ican.zaq.ne.jp](mailto:shige21@ican.zaq.ne.jp)

Received April 11, 2012