

Properties of Some Classes of Structured Tensors

Yisheng Song & Liqun Qi

Journal of Optimization Theory and Applications

ISSN 0022-3239
Volume 165
Number 3

J Optim Theory Appl (2015) 165:854–873
DOI 10.1007/s10957-014-0616-5

Vol. 165, No. 3

June 2015

165(3) 693–1070 (2015)

ISSN 0022-3239

**JOURNAL OF OPTIMIZATION
THEORY AND APPLICATIONS**

 Springer

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media New York. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Properties of Some Classes of Structured Tensors

Yisheng Song · Liqun Qi

Received: 18 March 2014 / Accepted: 20 June 2014 / Published online: 11 July 2014
© Springer Science+Business Media New York 2014

Abstract In this paper, we extend some classes of structured matrices to higher-order tensors. We discuss their relationships with positive semi-definite tensors and some other structured tensors. We show that every principal sub-tensor of such a structured tensor is still a structured tensor in the same class, with a lower dimension. The potential links of such structured tensors with optimization, nonlinear equations, nonlinear complementarity problems, variational inequalities and the non-negative tensor theory are also discussed.

Keywords P tensor · P_0 tensor · B tensor · B_0 tensor · Principal sub-tensor · Eigenvalues

Mathematics Subject Classification (2010) 47H15 · 47H12 · 34B10 · 47A52 · 47J10 · 47H09 · 15A48 · 47H07

Yisheng Song's work was supported by the National Natural Science Foundation of P.R. China (Grant No. 11171094, 11271112). His work was partially done when he was visiting The Hong Kong Polytechnic University. Liqun Qi's work was supported by the Hong Kong Research Grant Council (Grant No. PolyU 502510, 502111, 501212 and 501913).

Y. Song (✉)
School of Mathematics and Information Science, Henan Normal University,
Xinxiang 453007, Henan, People's Republic of China
e-mail: songyisheng1@gmail.com

L. Qi
Department of Applied Mathematics, The Hong Kong Polytechnic University,
Hung Hom, Kowloon, Hong Kong
e-mail: maqilq@polyu.edu.hk

1 Introduction

P and P_0 matrices have a long history and wide applications in mathematical sciences. Fiedler and Pták first studied P matrices systematically in [1]. For the applications of P and P_0 matrices and functions in linear complementarity problems, variational inequalities and nonlinear complementarity problems, we refer readers to [2–4]. It is well known that a symmetric matrix is a P (P_0) matrix if and only if it is positive (semi-)definite [2, pp. 147, 153].

On the other hand, motivated by the discussion on positive definiteness of multivariate homogeneous polynomial forms [5–8], in 2005, Qi [9] introduced the concept of positive (semi-)definite symmetric tensors. In the same time, Qi also introduced eigenvalues, H-eigenvalues, E-eigenvalues and Z-eigenvalues for symmetric tensors. It was shown that an even order symmetric tensor is positive (semi-)definite if and only if all of its H-eigenvalues or Z-eigenvalues are positive (nonnegative) [9, Theorem 5]. Beside automatical control, positive semi-definite tensors also found applications in magnetic resonance imaging [10–13] and spectral hypergraph theory [14–16].

The following questions are natural. Can we extend the concept of P and P_0 matrices to P and P_0 tensors? If this can be done, is it true a symmetric tensor is a P (P_0) tensor if and only if it is positive (semi-)definite? Are there any odd order P (P_0) tensors?

In Sect. 3, we will extend the concept of P and P_0 matrices to P and P_0 tensors. We will show that a symmetric tensor is a P (P_0) tensor if and only if it is positive (semi-)definite. The close relationship between P (P_0) tensors and positive (semi-)definite tensors justifies our research on P and P_0 tensors. We will show that there does not exist an odd order symmetric P tensor. If an odd order non-symmetric P tensor exists, then it has no Z-eigenvalues. An odd order P_0 tensor has no nonzero Z-eigenvalues.

In Sect. 4, we will further study some properties of P and P_0 tensors. We will show that every principal sub-tensor of a P (P_0) tensor is still a P (P_0) tensor, and give some sufficient and necessary conditions for a tensor to be a P (P_0) tensor.

The class of B matrices is a subclass of P matrices [17, 18]. We will extend the concept of B matrices to B and B_0 tensors in Sect. 5. It is easily checkable if a given tensor is a B or B_0 tensor or not. We will show that a Z tensor is diagonally dominated if and only if it is a B_0 tensor. It was proved in [19] that a diagonally dominated Z tensor is an M tensor. Laplacian tensors of uniform hypergraphs, defined as a natural extension of Laplacian matrices of graphs, are M tensors [16, 20–23]. This justifies our research on B and B_0 tensors.

Some final remarks will be given in Sect. 6. The potential links of P , P_0 , B and B_0 tensors with optimization, nonlinear equations, nonlinear complementarity problems, variational inequalities and the non-negative tensor theory are discussed. These encourage further research on P , P_0 , B and B_0 tensors.

2 Preliminaries

In this section, we will define the notations and collect some basic definitions and facts, which will be used later on.

Denote $I_n := \{1, 2, \dots, n\}$ and $\mathbb{R}^n := \{(x_1, x_2, \dots, x_n)^T; x_i \in \mathbb{R}, i \in I_n\}$, where \mathbb{R} is the set of real numbers. The definitions of P and P_0 matrices are as follows.

Definition 2.1 Let $A = (a_{ij})$ be an $n \times n$ real matrix. We say that A is

- (i) a P_0 matrix iff for any nonzero vector \mathbf{x} in \mathbb{R}^n , there exists $i \in I_n$ such that $x_i \neq 0$ and

$$x_i(A\mathbf{x})_i \geq 0;$$

- (ii) a P matrix iff for any nonzero vector \mathbf{x} in \mathbb{R}^n ,

$$\max_{i \in I_n} x_i(A\mathbf{x})_i > 0.$$

A real m th order n -dimensional tensor (hypermatrix) $\mathcal{A} = (a_{i_1 \dots i_m})$ is a multi-array of real entries $a_{i_1 \dots i_m}$, where $i_j \in I_n$ for $j \in I_m$. Denote the set of all real m th order n -dimensional tensors by $T_{m,n}$. Then, $T_{m,n}$ is a linear space of dimension n^m . Let $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$. If the entries $a_{i_1 \dots i_m}$ are invariant under any permutation of their indices, then \mathcal{A} is called a *symmetric tensor*. Denote the set of all real m th order n -dimensional tensors by $S_{m,n}$. Then, $S_{m,n}$ is a linear subspace of $T_{m,n}$. Let $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$ and $\mathbf{x} \in \mathbb{R}^n$. Then $\mathcal{A}\mathbf{x}^m$ is a homogeneous polynomial of degree m , defined by

$$\mathcal{A}\mathbf{x}^m := \sum_{i_1, \dots, i_m=1}^n a_{i_1 \dots i_m} x_{i_1} \cdots x_{i_m}.$$

A tensor $\mathcal{A} \in T_{m,n}$ is called *positive semi-definite* if for any vector $\mathbf{x} \in \mathbb{R}^n$, $\mathcal{A}\mathbf{x}^m \geq 0$, and is called *positive definite* if for any nonzero vector $\mathbf{x} \in \mathbb{R}^n$, $\mathcal{A}\mathbf{x}^m > 0$. Clearly, if m is odd, there is no non-trivial positive semi-definite tensors.

In the following, we extend the definitions of eigenvalues, H-eigenvalues, E-eigenvalues and Z-eigenvalues of tensors in $S_{m,n}$ in [9] to tensors in $T_{m,n}$.

Denote $\mathbb{C}^n := \{(x_1, x_2, \dots, x_n)^T; x_i \in \mathbb{C}, i \in I_n\}$, where \mathbb{C} is the set of complex numbers. For any vector $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x}^{[m-1]}$ is a vector in \mathbb{C}^n with its i th component defined as x_i^{m-1} for $i \in I_n$. Let $\mathcal{A} \in T_{m,n}$. If and only if there is a nonzero vector $\mathbf{x} \in \mathbb{C}^n$ and a number $\lambda \in \mathbb{C}$ such that

$$\mathcal{A}\mathbf{x}^{m-1} = \lambda \mathbf{x}^{[m-1]}, \tag{1}$$

then λ is called an *eigenvalue* of \mathcal{A} and \mathbf{x} is called an *eigenvector* of \mathcal{A} , associated with λ . If the eigenvector \mathbf{x} is real, then the eigenvalue λ is also real. In this case, λ and \mathbf{x} are called an *H-eigenvalue* and an *H-eigenvector* of \mathcal{A} , respectively. For an even order symmetric tensor, H-eigenvalues always exist. An even order symmetric tensor is positive (semi-)definite if and only if all of its H-eigenvalues are positive (non-negative). Let $\mathcal{A} \in T_{m,n}$. If and only if there is a nonzero vector $\mathbf{x} \in \mathbb{C}^n$ and a number $\lambda \in \mathbb{C}$ such that

$$\mathcal{A}\mathbf{x}^{m-1} = \lambda \mathbf{x}, \quad \mathbf{x}^\top \mathbf{x} = 1, \tag{2}$$

then λ is called an *E-eigenvalue* of \mathcal{A} and \mathbf{x} is called an *E-eigenvector* of \mathcal{A} , associated with λ . If the E-eigenvector \mathbf{x} is real, then the E-eigenvalue λ is also real. In this case, λ and \mathbf{x} are called an *Z-eigenvalue* and an *Z-eigenvector* of \mathcal{A} , respectively. For a symmetric tensor, H-eigenvalues always exist. An even order symmetric tensor is positive (semi-)definite if and only if all of its H-eigenvalues or Z-eigenvalues are positive (non-negative) [9, Theorem 5].

Throughout this paper, we assume that $m \geq 2$ and $n \geq 1$. We use small letters x, u, v, α, \dots , for scalars, small bold letters $\mathbf{x}, \mathbf{y}, \mathbf{u}, \dots$, for vectors, capital letters A, B, \dots , for matrices, calligraphic letters $\mathcal{A}, \mathcal{B}, \dots$, for tensors. All the tensors discussed in this paper are real. We denote the zero tensor in $T_{m,n}$ by \mathcal{O} .

3 P and P₀ Tensors

Let $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$ and $\mathbf{x} \in \mathbb{R}^n$. Then, $\mathcal{A}\mathbf{x}^{m-1}$ is a vector in \mathbb{R}^n with its i th component as

$$\left(\mathcal{A}\mathbf{x}^{m-1}\right)_i := \sum_{i_2, \dots, i_m=1}^n a_{ii_2 \dots i_m} x_{i_2} \cdots x_{i_m}$$

for $i \in I_n$. We now give the definitions of P and P₀ tensors.

Definition 3.1 Let $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$. We say that \mathcal{A} is

- (i) a P₀ tensor iff for any nonzero vector \mathbf{x} in \mathbb{R}^n , there exists $i \in I_n$ such that $x_i \neq 0$ and

$$x_i \left(\mathcal{A}\mathbf{x}^{m-1}\right)_i \geq 0;$$

- (ii) a P tensor iff for any nonzero vector \mathbf{x} in \mathbb{R}^n ,

$$\max_{i \in I_n} x_i \left(\mathcal{A}\mathbf{x}^{m-1}\right)_i > 0.$$

Clearly, this definition is a natural extension of Definition 2.1.

We first prove a proposition.

Proposition 3.1 Let $\mathcal{A} \in S_{m,n}$. If $\mathcal{A}\mathbf{x}^m = 0$ for all $\mathbf{x} \in \mathbb{R}^n$, then $\mathcal{A} = \mathcal{O}$.

Proof Denote $f(\mathbf{x}) = \mathcal{A}\mathbf{x}^m$. Then, $f(\mathbf{x}) = 0$ for all $\mathbf{x} \in \mathbb{R}^n$. This implies all the partial derivatives of f are zero. Since the entries of \mathcal{A} are just some higher-order partial derivatives of f , we see that $\mathcal{A} = \mathcal{O}$. □

We now have the following theorem.

Theorem 3.2 Let $\mathcal{A} \in T_{m,n}$ be a P (P₀) tensor. Then, when m is even, all of its H-eigenvalues and Z-eigenvalues of \mathcal{A} are positive (non-negative). A symmetric tensor is a P (P₀) tensor if and only if it is positive (semi-)definite. There does not exist an odd order symmetric P tensor. If an odd order non-symmetric P tensor exists, then it has no Z-eigenvalues. An odd order P₀ tensor has no nonzero Z-eigenvalues.

Proof Let m be even and an H-eigenvalue λ of \mathcal{A} be given. If \mathcal{A} is a P tensor, then by the definition of H-eigenvalues, there is a nonzero $\mathbf{x} \in \mathbb{R}^n$ and a number $\lambda \in \mathfrak{R}$ such that (1) holds. Then by the definition of P tensors, there exists $i \in I_n$ such that

$$0 < x_i (\mathcal{A}\mathbf{x}^{m-1})_i = \lambda x_i^m.$$

Since m is an even number, we have $\lambda > 0$. Similarly, if \mathcal{A} is a P_0 tensor, we may prove that $\lambda \geq 0$. By [9, Theorem 5], if all H-eigenvalues of an even order symmetric tensor are positive (non-negative), then that tensor is positive (semi-)definite. We see now that an even order symmetric tensor is a P (P_0) tensor only if it is positive (semi-)definite. By the definitions of P (P_0) tensors and positive (semi-)definite tensors, it is easy to see that an even order symmetric tensor is a P (P_0) tensor if it is positive (semi-)definite. Thus, an even order symmetric tensor is a P (P_0) tensor if and only if it is positive (semi-)definite.

Now, let an Z-eigenvalue λ of \mathcal{A} be given. If \mathcal{A} is a P tensor, then by the definition of Z-eigenvalues, there is an $\mathbf{x} \in \mathbb{R}^n$ and a number $\lambda \in \mathfrak{R}$ such that (2) holds. Then by the definition of P tensors, there exists an $i \in I_n$ such that

$$0 < x_i (\mathcal{A}\mathbf{x}^{m-1})_i = \lambda x_i^2.$$

Thus, $\lambda > 0$. Note that for this, we do not assume that m is even. However, when m is odd, if λ is a Z-eigenvalue of a tensor in $T_{m,n}$ with a Z-eigenvector \mathbf{x} , by the definition of Z-eigenvalues, $-\lambda$ is also a Z-eigenvalue of that tensor with an Z-eigenvector $-\mathbf{x}$. Thus, if an odd order P tensor exists, then it has no Z-eigenvalues. However, by [9, Theorem 5], a symmetric tensor always has Z-eigenvalues. Thus, an odd order symmetric P tensor does not exist. Since an odd order symmetric positive definite tensor also does not exist and an even order symmetric tensor is a P tensor if and only if it is positive definite, we conclude that a symmetric tensor is a P tensor if and only if it is positive definite.

Similarly, if \mathcal{A} is a P_0 tensor, we may prove that all of its Z-eigenvalues are non-negative. When m is odd, this also means that all of its Z-eigenvalues are non-positive. Thus, an odd order P_0 tensor has no nonzero Z-eigenvalues. By [9, Theorem 5], a symmetric tensor always has Z-eigenvalues. Thus, both the largest Z-eigenvalue λ_{\max} and the smallest Z-eigenvalue λ_{\min} of an odd order symmetric P_0 tensor \mathcal{A} are zero. By [9, Theorem 5], we have

$$\lambda_{\max} = \max\{\mathcal{A}\mathbf{x}^m : \mathbf{x}^\top \mathbf{x} = 1\}$$

and

$$\lambda_{\min} = \min\{\mathcal{A}\mathbf{x}^m : \mathbf{x}^\top \mathbf{x} = 1\}.$$

Thus, if \mathcal{A} is an odd order symmetric P_0 tensor, $\mathcal{A}\mathbf{x}^m = 0$ for all $\mathbf{x} \in \mathbb{R}^n$. By Proposition 3.1, this implies that $\mathcal{A} = \mathcal{O}$. By the definition of positive semi-definite tensors, if \mathcal{A} is an odd order symmetric positive semi-definite tensor, then $\mathcal{A} = \mathcal{O}$. Since an even order symmetric tensor is a P_0 tensor if and only if it is positive semi-definite, we

conclude that a symmetric tensor is a P_0 tensor if and only if it is positive semi-definite. The theorem is proved. \square

4 Properties of P and P_0 Tensors

In this section, we will study some properties of P and P_0 tensors. Based on the definition of P matrices, Mathias and Pang [24] introduced a fundamental quantity $\alpha(A)$ corresponding to a P matrix A by

$$\alpha(A) := \min_{\|\mathbf{x}\|_\infty=1} \left\{ \max_{i \in I_n} x_i (A\mathbf{x})_i \right\} \tag{3}$$

and studied its properties and applications. Mathias [25] showed that $\alpha(A)$ has a lower bound that is larger than 0 whenever A is a P matrix. Xiu and Zhang [26] gave some extensions of such a quantity and obtained global error bounds for the vertical and horizontal linear complementarity problems. Also see García-Esnaola and Peña [27] for the error bounds for linear complementarity problems of B matrices.

In the following, we will show that every principal sub-tensor of a P (P_0) tensor is still a P (P_0) tensor and give some sufficient and necessary conditions for a tensor to be a P tensor. Let $\mathcal{A} \in T_{m,n}$. Define an operator $T_{\mathcal{A}} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by for any $\mathbf{x} \in \mathbb{R}^n$,

$$T_{\mathcal{A}}(\mathbf{x}) := \begin{cases} \|\mathbf{x}\|_2^{2-m} \mathcal{A}\mathbf{x}^{m-1}, & \mathbf{x} \neq 0 \\ 0, & \mathbf{x} = 0. \end{cases} \tag{4}$$

When m is even, define another operator $F_{\mathcal{A}} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by for any $\mathbf{x} \in \mathbb{R}^n$,

$$F_{\mathcal{A}}(\mathbf{x}) := (\mathcal{A}\mathbf{x}^{m-1})_{\left[\frac{1}{m-1}\right]}. \tag{5}$$

Here, for a vector $\mathbf{y} \in \mathbb{R}^n$, $\mathbf{y}_{\left[\frac{1}{m-1}\right]}$ is a vector in \mathbb{R}^n , with its i th component to be $y_i^{\frac{1}{m-1}}$. When m is even, this is well defined. Then, we define two quantities

$$\alpha(T_{\mathcal{A}}) := \min_{\|\mathbf{x}\|_\infty=1} \max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i \tag{6}$$

for any m , and

$$\alpha(F_{\mathcal{A}}) := \min_{\|\mathbf{x}\|_\infty=1} \max_{i \in I_n} x_i (F_{\mathcal{A}}(\mathbf{x}))_i \tag{7}$$

when m is even. When $m = 2$, $\alpha(T_{\mathcal{A}})$ and $\alpha(F_{\mathcal{A}})$ are simply $\alpha(A)$ defined by (3). We will establish monotonicity and boundedness of such two quantities when \mathcal{A} is a P (P_0) tensor. Furthermore, we will show that \mathcal{A} is a P (P_0) tensor if and only if $\alpha(T_{\mathcal{A}})$ is positive (non-negative), and when m is even, \mathcal{A} is a P tensor (P_0) if and only if $\alpha(F_{\mathcal{A}})$ is positive (non-negative).

4.1 Principal Sub-tensors of P (P₀) Tensors

Recall that a tensor $\mathcal{C} \in T_{m,r}$ is called a *principal sub-tensor* of a tensor $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$ ($1 \leq r \leq n$) iff there is a set J that composed of r elements in I_n such that

$$\mathcal{C} = (a_{i_1 \dots i_m}), \quad \text{for all } i_1, i_2, \dots, i_m \in J.$$

The concept was first introduced and used in [9] for symmetric tensor. We denote by \mathcal{A}_r^J the principal sub-tensor of a tensor $\mathcal{A} \in T_{m,n}$ such that the entries of \mathcal{A}_r^J are indexed by $J \subset I_n$ with $|J| = r$ ($1 \leq r \leq n$), and denoted by \mathbf{x}_J the r -dimensional sub-vector of a vector $\mathbf{x} \in \mathbb{R}^n$, with the components of \mathbf{x}_J indexed by J . Note that for $r = 1$, the principal sub-tensors are just the diagonal entries.

Theorem 4.1 *Let \mathcal{A} be a P (P₀) tensor. Then, every principal sub-tensor of \mathcal{A} is P (P₀) tensor. In particular, all the diagonal entries of a P (P₀) tensor are positive (non-negative).*

Proof Let a principal sub-tensor \mathcal{A}_r^J of \mathcal{A} be given. Then for each nonzero vector $\mathbf{x} = (x_{j_1}, \dots, x_{j_r})^\top \in \mathfrak{N}^r$, we may choose $\mathbf{x}^* = (x_1^*, x_2^*, \dots, x_n^*)^\top \in \mathbb{R}^n$ with $x_i^* = x_i$ for $i \in J$ and $x_i^* = 0$ for $i \notin J$. Suppose now that \mathcal{A} is a P tensor, then there exists $j \in I_n$ such that

$$0 < x_j^* (\mathcal{A}(\mathbf{x}^*)^{m-1})_j = x_j (\mathcal{A}_r^J \mathbf{x}_J^{m-1})_j.$$

By the definition of \mathbf{x}^* , we have $j \in J$, and hence, \mathcal{A}_r^J is a P tensor. The case for P₀ tensors can be proved similarly. □

4.2 A Necessary and Sufficient Condition for P Tensors

The following is a sufficient and necessary condition for a tensor to be a P tensor.

Theorem 4.2 *Let $\mathcal{A} \in T_{m,n}$. Then \mathcal{A} is a P tensor if and only if for each nonzero $\mathbf{x} \in \mathbb{R}^n$, there exists an n -dimensional positive diagonal matrix $D_{\mathbf{x}}$ such that $\mathbf{x}^\top D_{\mathbf{x}} (\mathcal{A}\mathbf{x}^{m-1})$ is positive.*

Proof First, we show the necessity. Take a nonzero $\mathbf{x} \in \mathbb{R}^n$. It follows from the definition of P tensors that there is $k \in I_n$ such that $x_k (\mathcal{A}\mathbf{x}^{m-1})_k > 0$. Then for an enough small $\varepsilon > 0$, we have

$$x_k (\mathcal{A}\mathbf{x}^{m-1})_k + \varepsilon \left(\sum_{j \in I_n, j \neq k} x_j (\mathcal{A}\mathbf{x}^{m-1})_j \right) > 0.$$

Take $D_{\mathbf{x}} = \text{diag}(d_1, d_2, \dots, d_n)$ with $d_k = 1$ and $d_j = \varepsilon$ for $j \neq k$. Then, we have

$$\mathbf{x}^\top D_{\mathbf{x}} (\mathcal{A}\mathbf{x}^{m-1}) > 0.$$

Now we show the sufficiency. Assume that for each nonzero $\mathbf{x} \in \mathbb{R}^n$, there exists an n -dimensional diagonal matrix $D_{\mathbf{x}} = \text{diag}(d_1, d_2, \dots, d_n)$ with $d_i > 0$ for all $i \in I_n$ such that

$$0 < \mathbf{x}^T D_{\mathbf{x}} (\mathcal{A}\mathbf{x}^{m-1}) = \sum_{i=1}^n d_i (x_i (\mathcal{A}\mathbf{x}^{m-1})_i).$$

Since $d_i > 0$ for all $i \in I_n$, there is an $i \in I_n$ such that $x_i (\mathcal{A}\mathbf{x}^{m-1})_i > 0$. Otherwise, $x_i (\mathcal{A}\mathbf{x}^{m-1})_i \leq 0$ for all i . Then, $\sum_{i=1}^n d_i (x_i (\mathcal{A}\mathbf{x}^{m-1})_i) \leq 0$, a contradiction. Hence, \mathcal{A} is a P tensor.

The desired conclusion follows. □

4.3 Monotonicity and Boundedness of $\alpha(F_{\mathcal{A}})$ and $\alpha(T_{\mathcal{A}})$

Recall that an operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called *positively homogeneous* iff $T(t\mathbf{x}) = tT(\mathbf{x})$ for each $t > 0$ and all $\mathbf{x} \in \mathbb{R}^n$. For $\mathbf{x} \in \mathbb{R}^n$, it is known well that

$$\|\mathbf{x}\|_{\infty} := \max\{|x_i|; i \in I_n\} \text{ and } \|\mathbf{x}\|_2 := \left(\sum_{i=1}^n |x_i|^2 \right)^{\frac{1}{2}}$$

are two main norms defined on \mathbb{R}^n . Then for a continuous, positively homogeneous operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$, it is obvious that

$$\|T\|_{\infty} := \max_{\|\mathbf{x}\|_{\infty}=1} \|T(\mathbf{x})\|_{\infty}$$

is an operator norm of T and $\|T(\mathbf{x})\|_{\infty} \leq \|T\|_{\infty} \|\mathbf{x}\|_{\infty}$ for any $\mathbf{x} \in \mathbb{R}^n$. For $\mathcal{A} \in T_{m,n}$, let $T_{\mathcal{A}}$ be defined by (4). When m is even, let $F_{\mathcal{A}}$ be defined by (5). Clearly, both $F_{\mathcal{A}}$ and $T_{\mathcal{A}}$ are continuous and positively homogeneous. The following upper bounds of the operator norm were established by Song and Qi [28].

Lemma 4.1 (Song and Qi [28, Theorem 4.3]) *Let $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$. Then*

- (i) $\|T_{\mathcal{A}}\|_{\infty} \leq \max_{i \in I_n} (\sum_{i_2, \dots, i_m=1}^n |a_{ii_2 \dots i_m}|)$;
- (ii) $\|F_{\mathcal{A}}\|_{\infty} \leq \max_{i \in I_n} (\sum_{i_2, \dots, i_m=1}^n |a_{ii_2 \dots i_m}|)^{\frac{1}{m-1}}$, when m is even.

Let $\alpha(F_{\mathcal{A}})$ and $\alpha(T_{\mathcal{A}})$ be defined by (7) and (6). We now establish their monotonicity and boundedness. The proof technique is similar to the proof technique of [26, Theorem 1.2]. For completeness, we give the proof here.

Theorem 4.3 *Let $\mathcal{D} = \text{diag}(d_1, d_2, \dots, d_n)$ be a non-negative diagonal tensor in $T_{m,n}$ and $\mathcal{A} = (a_{i_1 \dots i_m})$ be a P_0 tensor in $T_{m,n}$. Then,*

- (i) $\alpha(T_{\mathcal{A}}) \leq \alpha(T_{\mathcal{A}+\mathcal{D}})$ whenever m is even;
- (ii) $\alpha(T_{\mathcal{A}}) \leq \alpha(T_{\mathcal{A}_r^J})$ for all principal sub-tensors \mathcal{A}_r^J ;
- (iii) $\alpha(F_{\mathcal{A}}) \leq \alpha(F_{\mathcal{A}_r^J})$ for all principal sub-tensors \mathcal{A}_r^J , when m is even;
- (iv) $\alpha(T_{\mathcal{A}}) \leq \max_{i \in I_n} (\sum_{i_2, \dots, i_m=1}^n |a_{ii_2 \dots i_m}|)$;
- (v) $\alpha(F_{\mathcal{A}}) \leq \max_{i \in I_n} (\sum_{i_2, \dots, i_m=1}^n |a_{ii_2 \dots i_m}|)^{\frac{1}{m-1}}$, when m is even.

Proof (i) By the definition of P_0 tensors, clearly $\mathcal{A} + \mathcal{D}$ is a P_0 tensor. Then, $\alpha(T_{\mathcal{A} + \mathcal{D}})$ is well defined. Since m is even, then $x_i^m > 0$ for $x_i \neq 0$, and so

$$\begin{aligned} \alpha(T_{\mathcal{A}}) &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i \right\} \\ &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \|\mathbf{x}\|_2^{2-m} \max_{i \in I_n} x_i (\mathcal{A}\mathbf{x}^{m-1})_i \right\} \\ &\leq \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \|\mathbf{x}\|_2^{2-m} \max_{i \in I_n} \left\{ x_i (\mathcal{A}\mathbf{x}^{m-1})_i + d_i x_i^m \right\} \right\} \\ &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i \left(\|\mathbf{x}\|_2^{2-m} (\mathcal{A} + \mathcal{D})\mathbf{x}^{m-1} \right)_i \right\} \\ &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i (T_{\mathcal{A} + \mathcal{D}}(\mathbf{x}))_i \right\} \\ &= \alpha(T_{\mathcal{A} + \mathcal{D}}). \end{aligned}$$

(ii) Let a principal sub-tensor \mathcal{A}_r^J of \mathcal{A} be given. Then for each nonzero vector $\mathbf{z} = (z_1, \dots, z_r)^T \in \mathfrak{R}^r$, we may define $\mathbf{y}(\mathbf{z}) = (y_1(\mathbf{z}), y_2(\mathbf{z}), \dots, y_n(\mathbf{z}))^T \in \mathbb{R}^n$ with $y_i(\mathbf{z}) = z_i$ for $i \in J$ and $y_i(\mathbf{z}) = 0$ for $i \notin J$. Thus, $\|\mathbf{z}\|_{\infty} = \|\mathbf{y}(\mathbf{z})\|_{\infty}$ and $\|\mathbf{z}\|_2 = \|\mathbf{y}(\mathbf{z})\|_2$. Hence,

$$\begin{aligned} \alpha(T_{\mathcal{A}}) &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i \right\} \\ &= \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \|\mathbf{x}\|_2^{2-m} \max_{i \in I_n} x_i (\mathcal{A}\mathbf{x}^{m-1})_i \right\} \\ &\leq \min_{\|\mathbf{y}(\mathbf{z})\|_{\infty}=1} \left\{ \|\mathbf{y}(\mathbf{z})\|_2^{2-m} \max_{i \in I_n} \{ \mathbf{y}(\mathbf{z})_i (\mathcal{A}\mathbf{y}(\mathbf{z})^{m-1})_i \} \right\} \\ &= \min_{\|\mathbf{z}\|_{\infty}=1} \left\{ \max_{i \in I_n} z_i (\|\mathbf{z}\|_2^{2-m} \mathcal{A}_r^J \mathbf{z}^{m-1})_i \right\} \\ &= \min_{\|\mathbf{z}\|_{\infty}=1} \left\{ \max_{i \in I_n} z_i (T_{\mathcal{A}_r^J}(\mathbf{z}))_i \right\} \\ &= \alpha(T_{\mathcal{A}_r^J}). \end{aligned}$$

Similarly, we may show (iii).

(iv) Since for each nonzero vector $\mathbf{x} = (x_1, \dots, x_n)^T \in \mathbb{R}^n$ and each $i \in I_n$,

$$x_i (T_{\mathcal{A}}(\mathbf{x}))_i \leq \|\mathbf{x}\|_{\infty} \|T_{\mathcal{A}}(\mathbf{x})\|_{\infty} \leq \|T_{\mathcal{A}}\|_{\infty} \|\mathbf{x}\|_{\infty}^2,$$

Then,

$$\max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i \leq \|T_{\mathcal{A}}\|_{\infty} \|\mathbf{x}\|_{\infty}^2.$$

Therefore, we have

$$\alpha(T_{\mathcal{A}}) = \min_{\|\mathbf{x}\|_{\infty}=1} \{\max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i\} \leq \|T_{\mathcal{A}}\|_{\infty},$$

and hence, by Lemma 4.1, the desired conclusion follows.

Similarly, we may show (v). □

4.4 Necessary and Sufficient Conditions Based Upon $\alpha(F_{\mathcal{A}})$ and $\alpha(T_{\mathcal{A}})$

We now give necessary and sufficient conditions for a tensor $A \in T_{m,n}$ to be a P (P_0) tensor, based upon $\alpha(F_{\mathcal{A}})$ and $\alpha(T_{\mathcal{A}})$.

Theorem 4.4 *Let $\mathcal{A} \in T_{m,n}$. Then*

- (i) *\mathcal{A} is a P (P_0) tensor if and only if $\alpha(T_{\mathcal{A}})$ is positive (non-negative);*
- (ii) *when m is even, \mathcal{A} is a P (P_0) tensor if and only if $\alpha(F_{\mathcal{A}})$ is positive (non-negative).*

Proof We only prove the case for P tensors. The proof for the P_0 tensor case is similar.

- (i) Let \mathcal{A} be a P tensor. Then, it follows from the definition of P tensors that for each nonzero $\mathbf{x} \in \mathbb{R}^n$,

$$\max_{i \in I_n} x_i (\mathcal{A}\mathbf{x}^{m-1})_i > 0,$$

and so

$$\max_{i \in I_n} x_i (\|\mathbf{x}\|_2^{2-m} \mathcal{A}\mathbf{x}^{m-1})_i = \|\mathbf{x}\|_2^{2-m} \max_{i \in I_n} x_i (\mathcal{A}\mathbf{x}^{m-1})_i > 0.$$

Therefore, we have

$$\alpha(T_{\mathcal{A}}) = \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i (T_{\mathcal{A}}(\mathbf{x}))_i \right\} > 0.$$

If $\alpha(T_{\mathcal{A}}) > 0$, then it is obvious that for each nonzero $\mathbf{y} \in \mathbb{R}^n$,

$$\max_{i \in I_n} \left(\frac{\mathbf{y}}{\|\mathbf{y}\|_{\infty}} \right)_i \left(T_{\mathcal{A}} \left(\frac{\mathbf{y}}{\|\mathbf{y}\|_{\infty}} \right) \right)_i \geq \alpha(T_{\mathcal{A}}) > 0.$$

Hence,

$$\max_{i \in I_n} y_i (T_{\mathcal{A}}(\mathbf{y}))_i = \max_{i \in I_n} y_i \left(\|\mathbf{y}\|_2^{2-m} \mathcal{A}\mathbf{y}^{m-1} \right)_i > 0.$$

Thus, $y_j (\mathcal{A}\mathbf{y}^{m-1})_j > 0$ for some $j \in I_n$, i.e., \mathcal{A} is a P tensor.

- (ii) Assume that m is even.

Let \mathcal{A} be a P tensor. Then for each nonzero $\mathbf{x} \in \mathbb{R}^n$, there exists an $i \in I_n$ such that $x_i (\mathcal{A}\mathbf{x}^{m-1})_i > 0$, and so

$$0 < x_i^{\frac{1}{m-1}} (\mathcal{A}\mathbf{x}^{m-1})_i^{\frac{1}{m-1}} = x_i^{\frac{2-m}{m-1}} \left(x_i (\mathcal{A}\mathbf{x}^{m-1})_i^{\frac{1}{m-1}} \right).$$

Since m is even, we have $x_i^{\frac{2-m}{m-1}} > 0$ for $x_i \neq 0$, and so,

$$0 < x_i (\mathcal{A}\mathbf{x}^{m-1})_i^{\frac{1}{m-1}} = x_i (F_{\mathcal{A}}(\mathbf{x}))_i.$$

That is, for each nonzero $\mathbf{x} \in \mathbb{R}^n$, $\max_{i \in I_n} x_i (F_{\mathcal{A}}(\mathbf{x}))_i > 0$. Thus, we have

$$\alpha(F_{\mathcal{A}}) = \min_{\|\mathbf{x}\|_{\infty}=1} \left\{ \max_{i \in I_n} x_i (F_{\mathcal{A}}(\mathbf{x}))_i \right\} > 0.$$

If $\alpha(F_{\mathcal{A}}) > 0$, then it is obvious that for each nonzero $\mathbf{y} \in \mathbb{R}^n$,

$$\max_{i \in I_n} \left(\frac{\mathbf{y}}{\|\mathbf{y}\|_{\infty}} \right)_i \left(F_{\mathcal{A}} \left(\frac{\mathbf{y}}{\|\mathbf{y}\|_{\infty}} \right) \right)_i \geq \alpha(F_{\mathcal{A}}) > 0.$$

Hence, there exists a $j \in I_n$ such that

$$y_j (F_{\mathcal{A}}(\mathbf{y}))_j = y_j (\mathcal{A}\mathbf{y}^{m-1})_j^{\frac{1}{m-1}} > 0.$$

Thus,

$$y_j^{m-2} \left(y_j (\mathcal{A}\mathbf{y}^{m-1})_j \right) = y_j^{m-1} (\mathcal{A}\mathbf{y}^{m-1})_j > 0.$$

Since m is even, we have $y_j^{m-2} > 0$. Hence, $y_j (\mathcal{A}\mathbf{y}^{m-1})_j > 0$, i.e., \mathcal{A} is a P tensor.

□

5 B and B₀ Tensors

An n -dimensional B matrix $B = (b_{ij})$ is a square real matrix with its entries satisfying that for all $i \in I_n$

$$\sum_{j=1}^n b_{ij} > 0 \text{ and } \frac{1}{n} \sum_{j=1}^n b_{ij} > b_{ik}, \quad i \neq k.$$

Many nice properties and applications of such B matrices have been studied by Peña [17, 18]. It was proved that B matrices are a subclass of P matrices in [17].

As a natural extension of B matrices, we now give the definition of B and B₀ tensors.

Definition 5.1 Let $\mathcal{B} = (b_{i_1 \dots i_m}) \in T_{m,n}$. We say that \mathcal{B} is a B tensor iff for all $i \in I_n$

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} > 0$$

and

$$\frac{1}{n^{m-1}} \left(\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \right) > b_{ij_2j_3 \dots j_m} \quad \text{for all } (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i).$$

We say that \mathcal{B} is a B_0 tensor iff for all $i \in I_n$

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \geq 0$$

and

$$\frac{1}{n^{m-1}} \left(\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \right) \geq b_{ij_2j_3 \dots j_m} \quad \text{for all } (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i).$$

Unlike P and P_0 tensors, it is easily checkable if a given tensor in $T_{m,n}$ is a B or B_0 tensor or not.

5.1 Entries of B and B_0 Tensors

We first study some properties of entries of B and B_0 tensors.

Theorem 5.1. *Let $\mathcal{B} = (b_{i_1 \dots i_m}) \in T_{m,n}$. If \mathcal{B} is a B tensor, then for each $i \in I_n$,*

$$b_{ii \dots i} > \max\{0, b_{ij_2j_3 \dots j_m}; (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i), j_2, j_3, \dots, j_m \in I_n\}.$$

If \mathcal{B} is a B_0 tensor, then for each $i \in I_n$,

$$b_{ii \dots i} \geq \max\{0, b_{ij_2j_3 \dots j_m}; (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i), j_2, j_3, \dots, j_m \in I_n\}.$$

Proof Suppose that $\mathcal{B} \in T_{m,n}$ is a B tensor. By Definition 5.1 that for all $i \in I_n$

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} > 0 \tag{8}$$

and

$$\frac{1}{n^{m-1}} \left(\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \right) > b_{ij_2j_3 \dots j_m} \quad \text{for all } (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i). \tag{9}$$

Let $b_{ik_2k_3 \dots k_m} = \max\{b_{ij_2j_3 \dots j_m} : (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)\}$. Then, it follows from (9) that

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} > n^{m-1} b_{ik_2k_3 \dots k_m} \geq b_{ik_2k_3 \dots k_m} + \sum_{(j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)} b_{ij_2j_3 \dots j_m}.$$

Thus,

$$b_{iii \dots i} > b_{ik_2k_3 \dots k_m} = \max\{b_{ij_2j_3 \dots j_m} : (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)\}.$$

Therefore, $b_{iii \dots i} > 0$. In fact, suppose $b_{iii \dots i} \leq 0$. Then, $\max\{b_{ij_2j_3 \dots j_m} : (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)\} < b_{iii \dots i} \leq 0$, which implies that

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \leq 0.$$

This contradicts to (8). The case for B_0 tensors can be proved similarly. □

Let $\mathcal{B} = (b_{i_1 \dots i_m}) \in T_{m,n}$. For each $i \in I_n$, define

$$\beta_i(\mathcal{B}) = \max\{0, b_{ij_2j_3 \dots j_m}; (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i), j_2, j_3, \dots, j_m \in I_n\}. \tag{10}$$

With the help of the quantity $\beta_i(\mathcal{B})$, we will study further the properties of B tensors.

Theorem 5.2. *Let $\mathcal{B} = (b_{i_1 \dots i_m}) \in T_{m,n}$. Then, \mathcal{B} is B tensor if and only if for each $i \in I_n$,*

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} > n^{m-1} \beta_i(\mathcal{B});$$

and \mathcal{B} is B_0 tensor if and only if for each $i \in I_n$,

$$\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \geq n^{m-1} \beta_i(\mathcal{B}).$$

Proof Since $\beta_i(\mathcal{B}) \geq 0$, the desired conclusion directly follows from Definition 5.1. □

Theorem 5.3. *Let $\mathcal{B} = (b_{i_1 \dots i_m}) \in T_{m,n}$. If \mathcal{B} is a B tensor, then for each $i \in I_n$,*

- (i) $b_{ii\dots i} > \sum_{b_{ii_2\dots i_m} < 0} |b_{ii_2\dots i_m}|;$
 - (ii) $b_{ii\dots i} > |b_{ij_2j_3\dots j_m}|$ for all $(j_2, j_3, \dots, j_m) \neq (i, i, \dots, i), j_2, j_3, \dots, j_m \in I_n.$
- If \mathcal{B} is a B_0 tensor, then (i) and (ii) hold with “>” being replaced by “ \geq .”

Proof Suppose that \mathcal{B} is a B tensor. (i) It follows from Proposition 5.2. that for each $i \in I_n$

$$b_{ii\dots i} - \beta_i(\mathcal{B}) > \sum_{(j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)} (\beta_i(\mathcal{B}) - b_{ij_2j_3\dots j_m}). \tag{11}$$

It follows from Definition 5.1 that for all $i \in I_n,$

$$\beta_i(\mathcal{B}) \geq 0 \text{ and } \beta_i(\mathcal{B}) - b_{ij_2j_3\dots j_m} \geq 0 \text{ for all } (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i).$$

Then for all $b_{ii_2i_3\dots i_m} < 0,$

$$\beta_i(\mathcal{B}) - b_{ii_2i_3\dots i_m} \geq |b_{ii_2i_3\dots i_m}|$$

and

$$b_{ii\dots i} \geq b_{ii\dots i} - \beta_i(\mathcal{B}).$$

So by (11), we have

$$b_{ii\dots i} > \sum_{(j_2, j_3, \dots, j_m) \neq (i, i, \dots, i)} (\beta_i(\mathcal{B}) - b_{ij_2j_3\dots j_m}) \geq \sum_{b_{ii_2\dots i_m} < 0} |b_{ii_2i_3\dots i_m}|.$$

(ii) is an obvious conclusion by combining Theorem 5.1. with (i).

The case for B_0 tensors can be proved similarly. □

5.2 Principal Sub-tensors of B and B_0 Tensors

We now show that every principal sub-tensor of a B (B_0) tensor is a B (B_0) tensor.

Theorem 5.4. *Let $\mathcal{B} = (b_{i_1\dots i_m}) \in T_{m,n}.$ If \mathcal{B} is a B (B_0) tensor, then every principal sub-tensor of \mathcal{B} is also a B (B_0) tensor.*

Proof Suppose that \mathcal{B} is a B tensor. Let a principal sub-tensor \mathcal{B}_r^J of \mathcal{B} be given. Then, it follows from Theorem 5.3. (i) that for all $i \in J,$

$$\sum_{i_2, \dots, i_m \in J} b_{ii_2i_3\dots i_m} > 0.$$

Now it suffices to show that for all $i \in J,$

$$\sum_{i_2, \dots, i_m \in J} b_{ii_2i_3\dots i_m} > r^{m-1} b_{ij_2\dots j_m} \text{ for all } (j_2, j_3, \dots, j_m) \neq (i, i, \dots, i), j_2, \dots, j_m \in J.$$

Suppose not. Then, there is (i, j_2, \dots, j_m) such that $i, j_2, \dots, j_m \in J$ and

$$\sum_{i_2, \dots, i_m \in J} b_{ii_2i_3 \dots i_m} \leq r^{m-1} b_{ij_2 \dots j_m}.$$

Let $b_{ik_2k_3 \dots k_m} = \max\{b_{ii_2i_3 \dots i_m}; (i_2, i_3, \dots, i_m) \neq (i, i, \dots, i) \text{ and } i_2, i_3, \dots, i_m \in I_n\}$. Then, $b_{ik_2k_3 \dots k_m} \geq b_{ij_2 \dots j_m}$. Hence,

$$\begin{aligned} n^{m-1} b_{ik_2k_3 \dots k_m} &\geq r^{m-1} b_{ik_2k_3 \dots k_m} + \sum \{b_{ii_2i_3 \dots i_m} : \text{not all of } i_2, \dots, i_m \text{ are in } J\} \\ &\geq r^{m-1} b_{ij_2j_3 \dots j_m} + \sum \{b_{ii_2i_3 \dots i_m} : \text{not all of } i_2, \dots, i_m \text{ are in } J\} \\ &\geq \sum_{i_2, \dots, i_m \in J} b_{ii_2i_3 \dots i_m} + \sum \{b_{ii_2i_3 \dots i_m} : \text{not all of } i_2, \dots, i_m \text{ are in } J\} \\ &= \sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m}. \end{aligned}$$

Thus,

$$\frac{1}{n^{m-1}} \left(\sum_{i_2, \dots, i_m=1}^n b_{ii_2i_3 \dots i_m} \right) \leq b_{ik_2k_3 \dots k_m},$$

which obtains a contradiction since \mathcal{B} is a B tensor.

The case for B_0 tensors can be proved similarly. □

5.3 The Relationship with M Tensors

Recall that a tensor $\mathcal{A} = (a_{i_1 \dots i_m}) \in T_{m,n}$ is called a Z tensor iff all of its off-diagonal entries are non-positive, i.e., $a_{i_1 \dots i_m} \leq 0$ when never $(i_1, \dots, i_m) \neq (i, \dots, i)$ [19]; \mathcal{A} is called diagonally dominated iff for all $i \in I_n$,

$$a_{i \dots i} \geq \sum \{|a_{ii_2 \dots i_m}| : (i_2, \dots, i_m) \neq (i, \dots, i), i_j \in I_n, j = 2, \dots, m\};$$

\mathcal{A} is called strictly diagonally dominated iff for all $i \in I_n$,

$$a_{i \dots i} > \sum \{|a_{ii_2 \dots i_m}| : (i_2, \dots, i_m) \neq (i, \dots, i), i_j \in I_n, j = 2, \dots, m\}.$$

It was proved in [19] that a diagonally dominated Z tensor is an M tensor, and a strictly diagonally dominated Z tensor is a strong M tensor. The definition of M tensors may be found in [19, 29]. Strong M tensors are called non-singular tensors in [29]. Laplacian tensors of uniform hypergraphs, defined as a natural extension of Laplacian matrices of graphs, are M tensors [16, 21–23].

Now we give the properties of a B (B_0) tensor under the condition that it is a Z tensor.

Theorem 5.5. Let $\mathcal{B} = (b_{i_1 i_2 i_3 \dots i_m}) \in T_{m,n}$ be a Z tensor. Then, the following properties are equivalent:

- (i) \mathcal{B} is $B(B_0)$ tensor;
- (ii) for each $i \in n$, $\sum_{i_2, \dots, i_m=1}^n b_{i i_2 i_3 \dots i_m}$ is positive (non-negative);
- (iii) \mathcal{B} is strictly diagonally dominant (diagonally dominated).

Proof We now prove the case for B tensors. The proof for the B_0 tensor case is similar. It follows from Definition 5.1 that (i) implies (ii).

Since \mathcal{B} be a Z tensor, all of its off-diagonal entries are non-positive. Thus, for any of its off-diagonal entry $b_{i i_2 \dots i_m}$, we have $|b_{i i_2 i_3 \dots i_m}| = -b_{i i_2 i_3 \dots i_m}$. Thus, (ii) means that for $i \in I_n$,

$$\begin{aligned} b_{i i i \dots i} &> - \sum \{b_{i i_2 i_3 \dots i_m} : (i_2, \dots, i_m) \neq (i, \dots, i), i_j \in I_n, j = 2, \dots, m\} \\ &= \sum \{|b_{i i_2 i_3 \dots i_m}| : (i_2, \dots, i_m) \neq (i, \dots, i), i_j \in I_n, j = 2, \dots, m\} \\ &\geq 0. \end{aligned}$$

Thus, (ii) implies (iii).

From (iii), it is obvious that for each $i \in I_n$,

$$\sum_{i_2, \dots, i_m=1}^n b_{i i_2 i_3 \dots i_m} > 0.$$

Since all the off-diagonal entries of \mathcal{B} are non-positive, we have

$$\frac{1}{n^{m-1}} \sum_{i_2, \dots, i_m=1}^n b_{i i_2 i_3 \dots i_m} > 0 \geq b_{i i_2 i_3 \dots i_m} \quad \text{for all } (i_2, \dots, i_m) \neq (i, \dots, i).$$

This shows that (iii) implies (i). □

From this theorem, we see that if a Z tensor is also a B_0 (B) tensor, then it is a (strong) M tensor.

6 Questions and Perspectives

Question 6.1 Is there an odd order non-symmetric P tensor? Is there an odd order nonzero nonsymmetric P_0 tensor?

Question 6.2 For a P matrix P , Mathias [25] showed that $\alpha(A)$ has a strictly positive lower bound. Then for a P tensor $\mathcal{A} \in T_{m,n}$ ($m > 2$), does $\alpha(F_{\mathcal{A}})$ or $\alpha(T_{\mathcal{A}})$ have a strictly positive lower bound?

Question 6.3 It is well known that A is a P matrix if and only if the linear complementarity problem

$$\text{find } \mathbf{z} \in \mathbb{R}^n \text{ such that } \mathbf{z} \geq \mathbf{0}, \mathbf{q} + A\mathbf{z} \geq \mathbf{0}, \text{ and } \mathbf{z}^\top (\mathbf{q} + A\mathbf{z}) = 0$$

has a unique solution for all $\mathbf{q} \in \mathbb{R}^n$. Then for a P tensor $\mathcal{A} \in T_{m,n}$ ($m > 2$), does a similar property hold for the following nonlinear complementarity problem

$$\text{find } \mathbf{x} \in \mathbb{R}^n \text{ such that } \mathbf{x} \geq \mathbf{0}, \mathbf{q} + \mathcal{A}\mathbf{x}^{m-1} \geq \mathbf{0}, \text{ and } \mathbf{x}^\top(\mathbf{q} + \mathcal{A}\mathbf{x}^{m-1}) = 0?$$

Question 6.4 When $m = 2$, it is known that each B matrix is a P matrix. If m is odd, in general, a B (B_0) tensor is not a P (P_0) tensor. For example, let $a_{i\dots i} = 1$ and $a_{i_1\dots i_m} = 0$ otherwise. Then, $\mathcal{A} = (a_{i_1\dots i_m})$ is the identity tensor. When m is odd, the identity tensor is a B tensor, but not a P or P_0 tensor. But we still make ask, when $m \geq 4$ and is even, is a B (B_0) tensor a P (P_0) tensor?

Question 6.5 A symmetric P (P_0) tensor is positive (semi-)definite. When $m \geq 4$ and is even, is a symmetric B (B_0) tensor positive (semi-)definite? If the answer is “yes” to this question, then we will have another checkable sufficient condition for positive (semi-)definite tensors.

Question 6.6 What are the spectral properties of a B (B_0) tensor?

Question 6.7 When $m \geq 4$ and is even, is a symmetric non-negative B (B_0) tensor positive (semi-)definite? If the answer is “yes” to this question, then we will have more understanding on positive semi-definite, non-negative tensors.

We may also ask the following question:

Question 6.8 What is the relation between non-negative B (B_0) tensors and completely positive tensors introduced in [33]?

In this paper, we make an initial study on P, P_0 , B and B_0 tensors. We see that they are linked with positive (semi-)definite tensors and M tensors, which are useful in automatical control, magnetic resonance imaging and spectral hypergraph theory. The six questions at the ends of Sects. 3–5 pointed out some further research directions.

In the following, we point out the potential links between the above results and optimization, nonlinear equations, nonlinear complementarity problems, variational inequalities and the non-negative tensor theory.

(i) We now discuss the potential link between the above results and optimization, nonlinear equations, nonlinear complementarity problems and variational inequalities. Question 6.3 has also pointed out the potential link between P tensor and nonlinear complementarity problems. We may also consider the optimization problem

$$\min\{\mathcal{A}\mathbf{x}^m + \mathbf{q}^\top \mathbf{x}\},$$

the nonlinear equations [30]

$$\mathcal{A}\mathbf{x}^{m-1} = \mathbf{q}$$

and the variational inequality problem [3,4]

$$\text{find } \mathbf{x}_* \in X, \text{ such that } (\mathbf{x} - \mathbf{x}_*)^\top \mathcal{A}\mathbf{x}_*^{m-1} \geq 0, \quad \text{for all } \mathbf{x} \in X,$$

where X is a non-empty closed subset of \mathbb{R}^n . When \mathcal{A} is a P , P_0 , B or B_0 tensor, what properties we can obtain for the above problems?

- (ii) We now further discuss the potential link between the above results and the non-negative tensor theory. The non-negative tensor theory at least include two parts: the non-negative tensor decomposition [31] and the spectral theory of non-negative tensors [32]. The recent paper [33] linked these two parts. However, there are still many questions not answered in non-negative tensors. In the non-negative matrix theory [34], a doubly non-negative matrix is a positive semi-definite, non-negative matrix. The research on positive semi-definite, non-negative tensors is very little. Thus, we may ask a question weaker than Question 6.5.

In a word, this paper is only an initial study on P , P_0 , B and B_0 tensors. Many questions for these tensors are waiting for answers.

It should be pointed out that after the first version of this paper, two more papers [35,36] on P , P_0 , B and B_0 tensors appeared. In [35], it was proved that an even order symmetric B_0 tensor is positive semi-definite and an even order symmetric B tensor is positive definite. Some further properties of P , P_0 , B and B_0 tensors were obtained in [36]. These answered some questions raised in this paper and enriched the theory of P , P_0 , B and B_0 tensors.

7 Conclusions

In this paper, we extend some classes of structured matrices to higher-order tensors. We discuss their relationships with positive semi-definite tensors and some other structured tensors. We show that every principal sub-tensor of such a structured tensor is still a structured tensor in the same class, with a lower dimension. The potential links and applications of such structured tensors are also discussed.

There are more research topics on structured tensors. In particular, can one construct an efficient algorithm to compute the extreme eigenvalues of a special structured tensor, other than the largest eigenvalue of a non-negative tensor? It is well known [32] that there are efficient algorithms for computing the largest eigenvalue of a non-negative tensor. Until now, there are no polynomial-time algorithms for computing extreme eigenvalues of structured tensors in the other cases. The first challenging problem is to construct an efficient algorithm to compute the smallest real eigenvalue of a Hilbert tensor [37], with the condition that such a real eigenvalue has a real eigenvector. A further challenging problem is to address the above problem for a Cauchy tensor [38] instead of a Hilbert tensor. Note that the Hilbert tensor is a special case of the Cauchy tensor [38].

Acknowledgments The authors would like to thank the anonymous referees for their valuable suggestions which helped us to improve this manuscript.

References

1. Fiedler, M., Pták, V.: On matrices with non-positive off-diagonal elements and positive principal minors. *Czechoslov. Math. J.* **12**, 163–172 (1962)

2. Cottle, R.W., Pang, J.S., Stone, R.E.: *The Linear Complementarity Problem*. Academic Press, Boston (1992)
3. Facchinei, F., Pang, J.S.: *Finite Dimensional Variational Inequalities and Complementarity Problems*. Springer, New York (2003)
4. Qi, L., Sun, D., Zhou, G.: A new look at smoothing Newton methods for nonlinear complementarity problems and box constrained variational inequalities. *Math. Program.* **87**, 1–35 (2000)
5. Bose, N.K., Modares, A.R.: General procedure for multivariable polynomial positivity with control applications. *IEEE Trans. Autom. Control* **21**, 596–601 (1976)
6. Hasan, M.A., Hasan, A.A.: A procedure for the positive definiteness of forms of even-order. *IEEE Trans. Autom. Control* **41**, 615–617 (1996)
7. Jury, E.I., Mansour, M.: Positivity and nonnegativity conditions of a quartic equation and related problems. *IEEE Trans. Automat. Control* **26**, 444–451 (1981)
8. Wang, F., Qi, L.: Comments on ‘Explicit criterion for the positive definiteness of a general quartic form’. *IEEE Trans. Autom. Control* **50**, 416–418 (2005)
9. Qi, L.: Eigenvalues of a real supersymmetric tensor. *J. Symb. Comput.* **40**, 1302–1324 (2005)
10. Chen, Y., Dai, Y., Han, D., Sun, W.: Positive semidefinite generalized diffusion tensor imaging via quadratic semidefinite programming. *SIAM J. Imaging Sci.* **6**, 1531–1552 (2013)
11. Hu, S., Huang, Z., Ni, H., Qi, L.: Positive definiteness of diffusion kurtosis imaging. *Inverse Probl. Imaging* **6**, 57–75 (2012)
12. Qi, L., Yu, G., Wu, E.X.: Higher order positive semi-definite diffusion tensor imaging. *SIAM J. Imaging Sci.* **3**, 416–433 (2010)
13. Qi, L., Yu, G., Xu, Y.: Nonnegative diffusion orientation distribution function. *J. Math. Imaging Vis.* **45**, 103–113 (2013)
14. Hu, S., Qi, L.: Algebraic connectivity of an even uniform hypergraph. *J. Comb. Optim.* **24**, 564–579 (2012)
15. Li, G., Qi, L., Yu, G.: The Z-eigenvalues of a symmetric tensor and its application to spectral hypergraph theory. *Numer. Linear Algebra Appl.* **20**, 1001–1029 (2013)
16. Qi, L.: H^+ -eigenvalues of Laplacian and signless Laplacian tensors. *Commun. Math. Sci.* **12**, 1045–1064 (2014)
17. Peña, J.M.: A class of P-matrices with applications to the localization of the eigenvalues of a real matrix. *SIAM J. Matrix Anal. Appl.* **22**, 1027–1037 (2001)
18. Peña, J.M.: On an alternative to Gerschgorin circles and ovals of Cassini. *Numerische Mathematik* **95**, 337–345 (2003)
19. Zhang, L., Qi, L., Zhou, G.: M-tensors and some applications. *SIAM J. Matrix Anal. Appl.* **35**(2), 437–452 (2014)
20. Hu, S., Qi, L.: The eigenvectors associated with the zero eigenvalues of the Laplacian and signless Laplacian tensors of a uniform hypergraph. *Discrete Appl. Math.* **169**, 140–151 (2014)
21. Hu, S., Qi, L., Shao, J.: Cored hypergraphs, power hypergraphs and their Laplacian eigenvalues. *Linear Algebra Appl.* **439**, 2980–2998 (2013)
22. Hu, S., Qi, L., Xie, J.: The largest Laplacian and signless Laplacian eigenvalues of a uniform hypergraph. [arXiv:1304.1315](https://arxiv.org/abs/1304.1315) (2013)
23. Qi, L., Shao, J., Wang, Q.: Regular uniform hypergraphs, s -cycles, s -paths and their largest Laplacian H-eigenvalues. *Linear Algebra Appl.* **443**, 215–227 (2014)
24. Mathias, R., Pang, J.S.: Error bounds for the linear complementarity problem with a P-matrix. *Linear Algebra Appl.* **132**, 123–136 (1990)
25. Mathias, R.: An improved bound for a fundamental constant associated with a P-matrix. *Appl. Math. Lett.* **2**, 297–300 (1989)
26. Xiu, N., Zhang, J.: A characteristic quantity of P-matrices. *Appl. Math. Lett.* **15**, 41–46 (2002)
27. García-Esnaola, M., Peña, J.M.: Error bounds for linear complementarity problems for B-matrices. *Appl. Math. Lett.* **22**, 1071–1075 (2009)
28. Song, Y., Qi, L.: Spectral properties of positively homogeneous operators induced by higher order tensors. *SIAM J. Matrix Anal. Appl.* **34**, 1581–1595 (2013)
29. Ding, W., Qi, L., Wei, Y.: M-tensors and nonsingular M-tensors. *Linear Algebra Appl.* **439**, 3264–3278 (2013)
30. Ortega, J.M., Rheinboldt, W.C.: *Iterative Solutions of Nonlinear Equations in Several Variables*. Academic Press, New York (1970)

31. Cichocki, A., Zdunek, R., Phan, A.H., Amari, S.: *Nonnegative Matrix and Tensor Factorizations: Applications to Exploratory Multiway Data Analysis and Blind Source Separation*. Wiley, New York (2009)
32. Chang, K.C., Qi, L., Zhang, T.: A survey on the spectral theory of nonnegative tensors. *Linear Algebra Appl.* **20**, 891–912 (2013)
33. Qi, L., Xu, C., Xu, Y.: Nonnegative tensor factorization, completely positive tensors and an hierarchically elimination algorithm. to appear. *SIAM J. Matrix Anal. Appl.*
34. Berman, A., Plemmons, R.J.: *Nonnegative Matrices in the Mathematical Sciences*. SIAM, Philadelphia (1994)
35. Qi, L., Song, Y.: An even order symmetric B tensor is positive definite. *Linear Algebra Appl.* **457**, 303–312 (2014)
36. Yuan, P., You, L.: Some remarks on P , P_0 , B and B_0 tensors. [arXiv:1402.1288](https://arxiv.org/abs/1402.1288) (2014)
37. Song, Y., Qi, L.: Infinite and finite dimensional Hilbert tensors. *Linear Algebra Appl.* **451**, 1–14 (2014)
38. Chen, H., Qi, L.: Positive definiteness and semi-definiteness of even order symmetric Cauchy tensors. [arXiv:1405.6363](https://arxiv.org/abs/1405.6363) (2014)