

 Open access • Journal Article • DOI:10.2478/UMCSMATH-2014-0004

Properties of the determinant of a rectangular matrix — [Source link](#)

[Anna Makarewicz](#), [Piotr Pikuta](#), [Dominik Szalkowski](#)

Published on: 01 Jun 2014 - [Annales Umcs, Mathematica](#) (Wydawnictwo Uniwersytetu Marii Curie-Skłodowskiej)

Topics: [Minor \(linear algebra\)](#), [Square matrix](#), [Matrix \(mathematics\)](#) and [Laplace expansion](#)

Related papers:

- [The Determinant of a Triangular-Block Matrix](#)
- [A Method to Compute the Determinant of a \$5 \times 5\$ Matrix](#)
- [Matrices and determinoids](#)
- [Method of generating matrix factors for a finite-dimensional linear transform](#)
- [Method of generating a sequence of matrix factors for a transformation matrix](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/properties-of-the-determinant-of-a-rectangular-matrix-3uax4razzg>

ANNA MAKAREWICZ, PIOTR PIKUTA and DOMINIK SZAŁKOWSKI

Properties of the determinant of a rectangular matrix

ABSTRACT. In this paper we present new identities for the Radić's determinant of a rectangular matrix. The results include representations of the determinant of a rectangular matrix as a sum of determinants of square matrices and description how the determinant is affected by operations on columns such as interchanging columns, reversing columns or decomposing a single column.

1. Introduction. In [2] Radić introduced the following definition of the determinant of a rectangular matrix.

Definition 1.1. Let $A = [A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix with n columns A_1, \dots, A_n and $m \leq n$. The determinant of A is defined as

$$(1) \quad |A| = \sum_{1 \leq j_1 < \dots < j_m \leq n} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|,$$

where $r = 1 + 2 + \dots + m$.

The determinant of a square matrix and the determinant (1) of a $m \times n$ matrix, where $m \leq n$, have several common standard properties, including the following (see [2]):

- (1) If a row of A is identical to some other row or is a linear combination of other rows then $|A| = 0$.

- (2) If a row of A is multiplied by a number k , then the determinant of the resulting matrix is equal to $k|A|$.
- (3) Interchanging two rows of A results in changing the sign of the determinant.
- (4) The determinant $|A|$ can be calculated using the Laplace expansion.

The properties of the determinant (1) were investigated by Radić [3, 4, 5] and also by Radić and Sušanj [6]. In the papers cited, the results concerning $2 \times n$ matrices were applied in planar geometry.

Another approach was presented by Amiri, Fathy and Bayat in [1], where the authors proved determinant identities such as Dodgson Condensation Formula and Trahan Formula for rectangular matrices, as well as Cauchy–Binet Formula for non-square products of two matrices.

In this paper we present new identities for determinants of rectangular matrices. The results include representation of the determinant of a rectangular matrix as a sum of determinants of square matrices and description how the determinant is affected by operations on columns such as interchanging two columns, reversing columns or decomposing a single column.

2. Properties of the determinant.

2.1. Representation of the determinant of a rectangular matrix as a sum of determinants of square matrices. For $2 \times n$ matrices, where $n \geq 2$, Radić [3] proved the following theorem.

Theorem 2.1. *Let $A = [A_1, A_2, \dots, A_n]$ be a $2 \times n$ matrix with $n \geq 2$. Then*

$$\begin{aligned} |A| &= |A_1, A_2 - A_3 + A_4 - \dots + (-1)^n A_n| \\ &\quad + |A_2, A_3 - A_4 + \dots + (-1)^{n-1} A_n| \\ &\quad + \dots \\ &\quad + |A_{n-1}, A_n|. \end{aligned}$$

This theorem gives a representation of the determinant of a $2 \times n$ matrix, where $n \geq 2$, as a sum of determinants of square matrices other than the representation (1). We generalize this result to $m \times n$ matrices in the following way.

Theorem 2.2. *Let $A = [A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix, where m is a number of rows and n is a number of columns, $m \leq n$. Then we have*

$$\begin{aligned} |A| &= \sum_{1 \leq j_1 < \dots < j_{m-1} < n} (-1)^{r+j_1+j_2+\dots+j_{m-1}} \\ &\quad \times \left| A_{j_1}, A_{j_2}, \dots, A_{j_{m-1}}, \sum_{k=j_{m-1}+1}^n (-1)^k A_k \right|. \end{aligned}$$

Proof. Applying (1), we have

$$\begin{aligned}
 |A| &= \sum_{1 \leq j_1 < \dots < j_m \leq n} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}| \\
 &= \sum_{1 \leq j_1 < j_2 < \dots < j_{m-1} < n} \sum_{k=j_{m-1}+1}^n (-1)^{r+j_1+j_2+\dots+j_{m-1}+k} \\
 &\quad \times |A_{j_1}, A_{j_2}, \dots, A_{j_{m-1}}, A_k| \\
 &= \sum_{1 \leq j_1 < \dots < j_{m-1} < n} (-1)^{r+j_1+j_2+\dots+j_{m-1}} \\
 &\quad \times \left| A_{j_1}, A_{j_2}, \dots, A_{j_{m-1}}, \sum_{k=j_{m-1}+1}^n (-1)^k A_k \right|. \quad \square
 \end{aligned}$$

Using the same method, one can easily prove the following two theorems.

Theorem 2.3. Let $A = [A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix, $m \leq n$. Then we have

$$|A| = \sum_{1 < j_2 < \dots < j_m \leq n} (-1)^{r+j_2+j_3+\dots+j_m} \left| \sum_{k=1}^{j_2-1} (-1)^k A_k, A_{j_2}, \dots, A_{j_m} \right|,$$

where $r = 1 + 2 + \dots + m$.

Theorem 2.4. Let $A = [A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix, $m \leq n$. Then for each $p \in \{2, 3, \dots, m-1\}$ we have

$$\begin{aligned}
 |A| &= \sum_{\substack{1 \leq j_1 < \dots < j_{p-1} \\ j_{p+1} < \dots < j_m \leq n \\ j_{p+1} - j_{p-1} > 1}} (-1)^{r+j_1+j_2+\dots+j_{p-1}+j_{p+1}+\dots+j_m} \\
 &\quad \times \left| A_{j_1}, \dots, A_{j_{p-1}}, \sum_{k=j_{p-1}+1}^{j_{p+1}-1} (-1)^k A_k, A_{j_{p+1}}, \dots, A_{j_m} \right|,
 \end{aligned}$$

where $r = 1 + 2 + \dots + m$.

Example 1. Let $[A_1, A_2, A_3, A_4]$ be a 3×4 matrix. Then

$$\begin{aligned}
 |A_1, A_2, A_3, A_4| &= |A_1, A_2, A_3 - A_4| + |A_1, A_3, A_4| - |A_2, A_3, A_4| \\
 &= |A_1, A_2, A_3| - |A_1, A_2, A_4| + |A_1 - A_2, A_3, A_4| \\
 &= |A_1, A_2, A_3| - |A_1, A_2 - A_3, A_4| - |A_2, A_3, A_4|.
 \end{aligned}$$

2.2. Decomposing a column. If a column K in a square matrix A is a sum of two columns (eg. $K = K_1 + K_2$), then the determinant $|A|$ is a sum of two determinants of matrices obtained from A by replacing K by K_1 and K_2 respectively.

For rectangular matrices we have a similar property.

Theorem 2.5. *Let $A = [A_1, A_2, \dots, A_k, \dots, A_n]$ be a $m \times n$ matrix, $m \leq n$, and $A_k = B_k + C_k$ for some $k \in \{1, 2, \dots, n\}$. Then*

$$\begin{aligned} |A| &= |A_1, A_2, \dots, A_{k-1}, B_k, A_{k+1}, \dots, A_n| \\ &\quad + |A_1, A_2, \dots, A_{k-1}, C_k, A_{k+1}, \dots, A_n| \\ &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m+1} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|, \end{aligned}$$

where $r = 1 + 2 + \dots + m$.

Proof. After applying (1)

$$|A| = \sum_{1 \leq j_1 < \dots < j_m \leq n} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|$$

we separate the sum of determinants into two sums: the first one consisting of the determinants of matrices which contain the column $A_k = B_k + C_k$ and the second one consisting of other determinants.

$$\begin{aligned} |A| &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, A_k, \dots, A_{j_m}| \\ &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}| \\ &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, B_k, \dots, A_{j_m}| \\ &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, C_k, \dots, A_{j_m}| \\ &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|. \end{aligned}$$

Now the third sum is added and subtracted so that it can be included into both the first and the second sum:

$$\begin{aligned} |A| &= |A_1, A_2, \dots, A_{k-1}, B_k, A_{k+1}, \dots, A_n| \\ &\quad + |A_1, A_2, \dots, A_{k-1}, C_k, A_{k+1}, \dots, A_n| \\ &\quad - \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}| \end{aligned}$$

$$\begin{aligned}
 &= |A_1, A_2, \dots, A_{k-1}, B_k, A_{k+1}, \dots, A_n| \\
 &\quad + |A_1, A_2, \dots, A_{k-1}, C_k, A_{k+1}, \dots, A_n| \\
 &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ k \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m+1} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|. \quad \square
 \end{aligned}$$

Example 2. Let $[A_1, A_2, A_3]$ be a 2×3 matrix and $A_1 = B_1 + C_1$. Then according to Theorem 2.5 we have

$$\begin{aligned}
 |B_1 + C_1, A_2, A_3| &= |B_1, A_2, A_3| + |C_1, A_2, A_3| \\
 &\quad + \sum_{\substack{1 \leq j_1 < j_2 \leq 3 \\ 1 \notin \{j_1, j_2\}}} (-1)^{(1+2)+j_1+j_2+1} |A_{j_1}, A_{j_2}| \\
 &= |B_1, A_2, A_3| + |C_1, A_2, A_3| + (-1)^{3+2+3+1} |A_2, A_3| \\
 &= |B_1, A_2, A_3| + |C_1, A_2, A_3| - |A_2, A_3|.
 \end{aligned}$$

2.3. Interchanging columns. Interchanging columns in a square matrix results in changing the sign of the determinant. Rectangular matrices in which the number of columns is equal to the number of rows increased by one have the same property.

Theorem 2.6. Let $A = [A_1, A_2, \dots, A_m, A_{m+1}]$ be a $m \times (m+1)$ matrix. Then for each $i, j \in \{1, 2, \dots, m+1\}$ such that $i < j$, we have

$$|A| = -|A_1, A_2, \dots, A_{i-1}, A_j, A_{i+1}, \dots, A_{j-1}, A_i, A_{j+1}, \dots, A_m, A_{m+1}|.$$

Proof. Let $r = 1 + 2 + \dots + m$. Fix $i, j \in \{1, 2, \dots, m+1\}$ such that $i < j$. From all the determinants in the right-hand side of

$$|A| = \sum_{1 \leq j_1 < \dots < j_m \leq n} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, A_{j_2}, \dots, A_{j_m}|,$$

we distinguish determinants of two matrices which contain either A_i or A_j but not both of them. Thus we have

$$\begin{aligned}
 |A| &= (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - i\right]} \\
 &\quad \times |A_1, A_2, \dots, A_{i-1}, A_{i+1}, \dots, A_{j-1}, A_j, A_{j+1}, \dots, A_{m+1}| \\
 &\quad + (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - j\right]} \\
 &\quad \times |A_1, A_2, \dots, A_{i-1}, A_i, A_{i+1}, \dots, A_{j-1}, A_{j+1}, \dots, A_{m+1}| \\
 &\quad + \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i, j \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, A_i, \dots, A_j, \dots, A_{j_m}|.
 \end{aligned}$$

Notice that exactly $j - i - 1$ inversions are needed to move the column A_j to the position between A_{i-1} and A_{i+1} in the first summand. Similarly, in the second summand, also $j - i - 1$ inversions are needed to move the column A_i to the position between A_{j-1} and A_{j+1} .

In other summands we can simply interchange columns A_i and A_j with the sign change. Thus we have

$$\begin{aligned}
|A| &= (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - i + (j-i+1)\right]} \\
&\quad \times |A_1, A_2, \dots, A_{i-1}, A_j, A_{i+1}, \dots, A_{j-1}, A_{j+1}, \dots, A_{m+1}| \\
&+ (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - j + (j-i+1)\right]} \\
&\quad \times |A_1, A_2, \dots, A_{i-1}, A_{i+1}, \dots, A_{j-1}, A_i, A_{j+1}, \dots, A_{m+1}| \\
&- \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i, j \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, A_j, \dots, A_i, \dots, A_{j_m}| \\
&= - (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - j\right]} \\
&\quad \times |A_1, A_2, \dots, A_{i-1}, A_j, A_{i+1}, \dots, A_{j-1}, A_{j+1}, \dots, A_{m+1}| \\
&- (-1)^{\left[r + \frac{(m+1)(m+2)}{2} - i\right]} \\
&\quad \times |A_1, A_2, \dots, A_{i-1}, A_{i+1}, \dots, A_{j-1}, A_i, A_{j+1}, \dots, A_{m+1}| \\
&- \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i, j \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |A_{j_1}, \dots, A_j, \dots, A_i, \dots, A_{j_m}| \\
&= - |A_1, A_2, \dots, A_{i-1}, A_j, A_{i+1}, \dots, A_{j-1}, A_i, A_{j+1}, \dots, A_m, A_{m+1}|. \square
\end{aligned}$$

Consider a $m \times n$ matrix A with m rows and n columns, $m \leq n$. Let A' be a matrix obtained from A by interchanging two columns. Theorem 2.6 tells us that $|A| + |A'| = 0$ when $n - m = 1$. However, in general, if $n - m > 1$ the sum $|A| + |A'|$ is not zero.

For a $m \times n$ matrix $M = [M_1, M_2, \dots, M_n]$ and each $i, j \in \{1, 2, \dots, m\}$, such that $i < j$, denote

$$S_1(M, i, j) = \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i, j \notin \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |M_{j_1}, M_{j_2}, \dots, M_{j_m}|,$$

$$\begin{aligned}
 S_2(M, i, j) &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i, j \in \{j_1, \dots, j_m\}}} (-1)^{r+j_1+j_2+\dots+j_m} |M_{j_1}, M_{j_2}, \dots, M_{j_m}|, \\
 S_3(M, i, j) &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ (i \in J, j \notin J \text{ or } i \notin J, j \in J) \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 1 \pmod{2}}} (-1)^{r+j_1+j_2+\dots+j_m} |M_{j_1}, M_{j_2}, \dots, M_{j_m}|, \\
 S_4(M, i, j) &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ (i \in J, j \notin J \text{ or } i \notin J, j \in J) \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 0 \pmod{2}}} (-1)^{r+j_1+j_2+\dots+j_m} |M_{j_1}, M_{j_2}, \dots, M_{j_m}|,
 \end{aligned}$$

where $r = 1 + 2 + \dots + m$ and $\text{card}(X)$ stands for the cardinality of X .

Theorem 2.7. *Let $A = [A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix with m rows and n columns, $m \leq n$. For $i, j \in \{1, 2, \dots, n\}$ such that $i \neq j$ denote by $A_{A_i \leftrightarrow A_j}$ the matrix obtained from A by interchanging columns A_i and A_j . Then*

$$\begin{aligned}
 |A| + |A_{A_i \leftrightarrow A_j}| &= 2S_1(A, i, j) + 2S_4(A, i, j) \\
 &= 2S_1(A_{A_i \leftrightarrow A_j}, i, j) + 2S_4(A_{A_i \leftrightarrow A_j}, i, j).
 \end{aligned}$$

Proof. Fix $i, j \in \{1, 2, \dots, m\}$ such that $i < j$. (If $i > j$ we can proceed analogously). We have

$$|A| = S_1(A, i, j) + S_2(A, i, j) + S_3(A, i, j) + S_4(A, i, j).$$

It is easy to verify that

$$\begin{aligned}
 S_1(A_{A_i \leftrightarrow A_j}, i, j) &= S_1(A, i, j), \\
 S_2(A_{A_i \leftrightarrow A_j}, i, j) &= -S_2(A, i, j).
 \end{aligned}$$

Notice that each of the matrices in $S_3(A, i, j) + S_4(A, i, j)$ needs exactly $(j - i - \text{card}(J))$ column inversions to move the column A_i to the position where A_j would be, and also $(j - i - \text{card}(J))$ inversions are needed to move the column A_j to the position where A_i would be.

Therefore,

$$\begin{aligned}
S_3(A, i, j) &= \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i \in J, j \notin J \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 1 \pmod{2}}} (-1)^{r + \left(\sum_{k=1}^m j_k + j\right) - j + (j - i - \text{card}(J))} \\
&\quad \times |A_{j_1}, \dots, A_{j_p}, A_i, A_{j_q}, \dots, A_{j_m}| \\
&+ \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i \notin J, j \in J \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 1 \pmod{2}}} (-1)^{r + \left(\sum_{k=1}^m j_k + i\right) - i + (j - i - \text{card}(J))} \\
&\quad \times |A_{j_1}, \dots, A_{j_u}, A_j, A_{j_v}, \dots, A_{j_m}| \\
&= - \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i \in J, j \notin J \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 1 \pmod{2}}} (-1)^{r + \left(\sum_{k=1}^m j_k + j\right) - i} \\
&\quad \times |A_{j_1}, \dots, A_{j_p}, A_i, A_{j_q}, \dots, A_{j_m}| \\
&- \sum_{\substack{1 \leq j_1 < \dots < j_m \leq n \\ i \notin J, j \in J \\ J = \{i, \dots, j\} \setminus \{j_1, \dots, j_m\} \\ \text{card}(J) \equiv 1 \pmod{2}}} (-1)^{r + \left(\sum_{k=1}^m j_k + i\right) - j} \\
&\quad \times |A_{j_1}, \dots, A_{j_u}, A_j, A_{j_v}, \dots, A_{j_m}| \\
&= - S_3(A_{A_i \leftrightarrow A_j}, i, j),
\end{aligned}$$

where $r = 1 + 2 + \dots + m$ and $j_p < j < j_q$, $j_u < i < j_v$ for some p, q, u, v . Similarly, we have

$$S_4(A_{A_i \leftrightarrow A_j}, i, j) = S_4(A, i, j),$$

and finally,

$$\begin{aligned}
|A| + |A_{A_i \leftrightarrow A_j}| &= S_1(A, i, j) + S_2(A, i, j) + S_3(A, i, j) + S_4(A, i, j) \\
&\quad + S_1(A_{A_i \leftrightarrow A_j}, i, j) + S_2(A_{A_i \leftrightarrow A_j}, i, j) \\
&\quad + S_3(A_{A_i \leftrightarrow A_j}, i, j) + S_4(A_{A_i \leftrightarrow A_j}, i, j) \\
&= 2S_1(A, i, j) + 2S_4(A, i, j). \quad \square
\end{aligned}$$

Corollary 2.8. *Let A be a $m \times n$ matrix, $m \leq n$. If $i, j \in \{1, 2, \dots, n\}$ satisfy $|i - j| = 1$, then*

$$|A| + |A_{A_i \leftrightarrow A_j}| = 2S_1(A, i, j) = 2S_1(A_{A_i \leftrightarrow A_j}, i, j).$$

Example 3. Below we present a few identities obtained from Theorem 2.6, Theorem 2.7 and Corollary 2.8.

(a) Let $[A_1, A_2, A_3, A_4, A_5]$ be a 4×5 matrix. Then

$$|A_1, A_2, A_3, A_4, A_5| = -|A_5, A_2, A_3, A_4, A_1| = |A_5, A_4, A_3, A_2, A_1|.$$

(b) Let $[A_1, A_2, A_3, A_4]$ be a 2×4 matrix. Then

$$\begin{aligned} |A_1, A_2, A_3, A_4| + |A_2, A_1, A_3, A_4| &= 2|A_3, A_4|, \\ |A_1, A_2, A_3, A_4| + |A_1, A_4, A_3, A_2| &= 2(|A_1, A_2| - |A_1, A_3| + |A_1, A_4|), \\ |A_1, A_2, A_3, A_4| + |A_4, A_2, A_3, A_1| &= 2(|A_1, A_2| - |A_1, A_3| + |A_2, A_3| \\ &\quad - |A_2, A_4| + |A_3, A_4|). \end{aligned}$$

2.4. Reversing columns. Reversing columns in a $n \times n$ square matrix results in changing the sign of its determinant if and only if n is congruent to 2 or 3 (mod 4). Surprisingly, the determinant of a rectangular matrix also either changes or does not change the sign after column reversing, depending on the number of rows and the number of columns of the matrix.

Theorem 2.9. *Let $[A_1, A_2, \dots, A_n]$ be a $m \times n$ matrix, $m \leq n$. Then we have*

$$\begin{aligned} |A_n, A_{n-1}, \dots, A_2, A_1| &= |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1)^{\frac{m}{2}(2n+m+1)} \\ &= \begin{cases} |A_1, A_2, \dots, A_{n-1}, A_n| & \text{if } m \equiv 0 \pmod{4}, \\ |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1)^{n+1} & \text{if } m \equiv 1 \pmod{4}, \\ |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1) & \text{if } m \equiv 2 \pmod{4}, \\ |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1)^n & \text{if } m \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

Proof. Let $r = 1+2+\dots+m = \frac{m(m+1)}{2}$ and $B_k = A_{n+1-k}$, $k \in \{1, 2, \dots, n\}$. Since exactly $(m-1) + (m-2) + \dots + 1 = \frac{(m-1)m}{2}$ inversions of (adjacent) columns are needed to reverse the columns of a $m \times m$ matrix, we have

$$\begin{aligned} |B_1, B_2, \dots, B_n| &= \sum_{1 \leq i_1 < \dots < i_m \leq n} (-1)^{r+i_1+i_2+\dots+i_m} |B_{i_1}, B_{i_2}, \dots, B_{i_m}| \\ &= \sum_{1 \leq i_1 < \dots < i_m \leq n} (-1)^{r+i_1+i_2+\dots+i_m+\frac{(m-1)m}{2}} \\ &\quad \times |B_{i_m}, B_{i_{m-1}}, \dots, B_{i_1}| \\ &= \sum_{1 \leq i_1 < \dots < i_m \leq n} (-1)^{r+i_1+i_2+\dots+i_m+\frac{(m-1)m}{2}} \\ &\quad \times |A_{n+1-i_m}, A_{n+1-i_{m-1}}, \dots, A_{n+1-i_1}|. \end{aligned}$$

Applying the following change of variables: $j_k = n + 1 - i_{m-k+1}$ for each $k \in \{1, 2, \dots, m\}$, we get

$$\begin{aligned} |A_n, A_{n-1}, \dots, A_2, A_1| &= |B_1, B_2, \dots, B_n| \\ &= \sum_{1 \leq j_1 < \dots < j_m \leq n} (-1)^{r+m(n+1)-(j_1+j_2+\dots+j_m)+\frac{(m-1)m}{2}} |A_{j_1}, A_{j_2}, \dots, A_{j_m}| \\ &= |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1)^{m(n+1)+\frac{(m-1)m}{2}} \\ &= |A_1, A_2, \dots, A_{n-1}, A_n| \cdot (-1)^{\frac{m}{2}(2n+m+1)}. \end{aligned}$$

Finally, we state that

$$(-1)^{\frac{m}{2}(2n+m+1)} = \begin{cases} 1 & \text{if } m \equiv 0 \pmod{4}, \\ (-1)^{n+1} & \text{if } m \equiv 1 \pmod{4}, \\ (-1) & \text{if } m \equiv 2 \pmod{4}, \\ (-1)^n & \text{if } m \equiv 3 \pmod{4}, \end{cases}$$

which is easy to verify. □

Example 4. Let

$$[A_1, A_2, A_3, A_4, A_5, A_6, A_7, A_8, A_9]$$

be a 5×9 matrix. Then

$$|A_1, A_2, A_3, A_4, A_5, A_6, A_7, A_8, A_9| = |A_9, A_8, A_7, A_6, A_5, A_4, A_3, A_2, A_1|.$$

REFERENCES

- [1] Amiri, M., Fathy, M., Bayat, M., *Generalization of some determinantal identities for non-square matrices based on Radic's definition*, TWMS J. Pure Appl. Math. **1**, no. 2 (2010), 163–175.
- [2] Radić, M., *A definition of determinant of rectangular matrix*, Glas. Mat. Ser. III **1(21)** (1966), 17–22.
- [3] Radić, M., *About a determinant of rectangular $2 \times n$ matrix and its geometric interpretation*, Beiträge Algebra Geom. **46**, no. 2 (2005), 321–349.
- [4] Radić, M., *Areas of certain polygons in connection with determinants of rectangular matrices*, Beiträge Algebra Geom. **49**, no. 1 (2008), 71–96.
- [5] Radić, M., *Certain equalities and inequalities concerning polygons in \mathbb{R}^2* , Beiträge Algebra Geom. **50**, no. 1 (2009), 235–248.
- [6] Radić, M., Sušan, R., *Geometrical meaning of one generalization of the determinant of a square matrix*, Glas. Mat. Ser. III **29(49)**, no. 2 (1994), 217–233.

Anna Makarewicz
Lublin University of Technology
Department of Applied Mathematics
ul. Nadbystrzycka 38 D
20-618 Lublin
Poland
e-mail: anna_makarewicz@o2.pl

Piotr Pikuta
Institute of Mathematics
Maria Curie-Skłodowska University
pl. Marii Curie-Skłodowskiej 1
20-031 Lublin
Poland
e-mail: ppikuta@poczta.umcs.lublin.pl

Dominik Szalkowski
Institute of Mathematics
Maria Curie-Skłodowska University
pl. Marii Curie-Skłodowskiej 1
20-031 Lublin
Poland
e-mail: dominik.szalkowski@umcs.lublin.pl

Received February 21, 2013