

NCCN Guidelines® Insights

Prostate Cancer Early Detection, Version 2.2016

Featured Updates to the NCCN Guidelines

Peter R. Carroll, MD, MPH^{1,*}; J. Kellogg Parsons, MD, MHS^{2,*}; Gerald Andriole, MD³; Robert R. Bahnson, MD⁴; Erik P. Castle, MD^{5,*}; William J. Catalona, MD^{6,*}; Douglas M. Dahl, MD⁷; John W. Davis, MD^{8,*}; Jonathan I. Epstein, MD^{9,*}; Ruth B. Etzioni, PhD¹⁰; Thomas Farrington, BSEE^{11,*}; George P. Hemstreet III, MD, PhD¹²; Mark H. Kawachi, MD¹³; Simon Kim, MD, PhD¹⁴; Paul H. Lange, MD^{15,*}; Kevin R. Loughlin, MD¹⁶; William Lowrance, MD, MPH^{17,*}; Paul Maroni, MD^{18,*}; James Mohler, MD^{19,*}; Todd M. Morgan, MD²⁰; Kelvin A. Moses, MD, PhD²¹; Robert B. Nadler, MD^{6,*}; Michael Poch, MD²²; Chuck Scales, MD²³; Terrence M. Shaneyfelt, MD, MPH^{24,*}; Marc C. Saldone, MD, MSHP^{25,*}; Geoffrey Sonn, MD^{26,*}; Preston Sprenkle, MD²⁷; Andrew J. Vickers, PhD²⁸; Robert Wake, MD^{29,*}; Dorothy A. Shead, MS^{30,*}; and Deborah A. Freedman-Cass, PhD^{30,*}

Abstract

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for Prostate Cancer Early Detection provide recommendations for prostate cancer screening in healthy men who have elected to participate in an early detection program. The NCCN Guidelines focus on minimizing unnecessary procedures and limiting the detection of indolent disease. These NCCN Guidelines Insights summarize the NCCN Prostate Cancer Early Detection Panel's most significant discussions for the 2016 guideline update, which included issues surrounding screening in high-risk populations (ie, African Americans, *BRCA1/2* mutation carriers), approaches to refine patient selection for initial and repeat biopsies, and approaches to improve biopsy specificity.

J Natl Compr Canc Netw 2016;14(5):509–519

From ¹UCSF Helen Diller Family Comprehensive Cancer Center; ²UC San Diego Moores Cancer Center; ³Siteman Cancer Center at Barnes-Jewish Hospital and Washington University School of Medicine; ⁴The Ohio State University Comprehensive Cancer Center – James Cancer Hospital and Solove Research Institute; ⁵Mayo Clinic Cancer Center; ⁶Robert H. Lurie Comprehensive Cancer Center of Northwestern University; ⁷Massachusetts General Hospital Cancer Center; ⁸The University of Texas MD Anderson Cancer Center; ⁹The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins; ¹⁰Fred Hutchinson Cancer Research Center/Seattle Cancer Care Alliance; ¹¹Prostate Health Education Network; ¹²Fred & Pamela Buffett Cancer Center at The Nebraska Medical Center; ¹³City of Hope Comprehensive Cancer Center; ¹⁴Case Comprehensive Cancer Center/University Hospitals Seidman Cancer Center and Cleveland Clinic Taussig Cancer Institute; ¹⁵University of Washington Medical Center/Seattle Cancer Care Alliance; ¹⁶Dana-Farber/Brigham and Women's Cancer Center; ¹⁷Huntsman Cancer Institute at the University of Utah; ¹⁸University of Colorado Cancer Center; ¹⁹Roswell Park Cancer Institute; ²⁰University of Michigan Comprehensive Cancer Center; ²¹Vanderbilt-Ingram Cancer Center; ²²Moffitt Cancer Center; ²³Duke Cancer Institute; ²⁴University of Alabama at Birmingham Comprehensive Cancer Center; ²⁵Fox Chase Cancer Center; ²⁶Stanford Cancer Center; ²⁷Yale Cancer Center/Smilow Cancer Hospital; ²⁸Memorial Sloan Kettering Cancer Center; ²⁹St. Jude Children's Research Hospital/The University of Tennessee Health Science Center; and ³⁰National Comprehensive Cancer Network.

*Provided content development and/or authorship assistance.

Please Note

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) are a statement of consensus of the authors regarding their views of currently accepted approaches to treatment. **The NCCN Guidelines® Insights highlight important changes to the NCCN Guidelines® recommendations from previous versions. Colored markings in the algorithm show changes and the discussion aims to further the understanding of these changes by summarizing salient portions of the NCCN Guideline Panel discussion, including the literature reviewed.**

These NCCN Guidelines Insights do not represent the full NCCN Guidelines; further, the National Comprehensive Cancer Network® (NCCN®) makes no representation or warranties of any kind regarding the content, use, or application of the NCCN Guidelines and NCCN Guidelines Insights and disclaims any responsibility for their applications or use in any way.

The full and most current version of these NCCN Guidelines are available at NCCN.org.

© National Comprehensive Cancer Network, Inc. 2016, All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN.

Prostate Cancer Early Detection, Version 2.2016

NCCN: Continuing Education**Accreditation Statement**

This activity has been designed to meet the educational needs of physicians, nurses, and pharmacists involved in the management of patients with cancer. There is no fee for this article. The National Comprehensive Cancer Network (NCCN) is accredited by the ACCME to provide continuing medical education for physicians. NCCN designates this journal-based CE activity for a maximum of 1.0 *AMA PRA Category 1 Credit*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

NCCN is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation.

NCCN designates this educational activity for a maximum of 1.0 contact hour. Accreditation as a provider refers to recognition of educational activities only; accredited status does not imply endorsement by NCCN or ANCC of any commercial products discussed/displayed in conjunction with the educational activity. Kristina M. Gregory, RN, MSN, OCN, is our nurse planner for this educational activity.



National Comprehensive Cancer Network is accredited by the Accreditation Council for Pharmacy Education as a provider of continuing pharmacy education. NCCN designates this continuing education activity for 1.0 contact hour(s) (0.1 CEUs) of continuing education credit in states that recognize ACPE accredited providers. This is a knowledge-based activity. UAN: 0836-0000-16-005-H01-P

All clinicians completing this activity will be issued a certificate of participation. To participate in this journal CE activity: 1) review the learning objectives and author disclosures; 2) study the education content; 3) take the posttest with a 66% minimum passing score and complete the evaluation at <http://education.nccn.org/node/78544>; and 4) view/print certificate.

Release date: May 12, 2016; Expiration date: May 12, 2017

Learning Objectives:

Upon completion of this activity, participants will be able to:

- Integrate into professional practice the updates to NCCN Guidelines for Prostate Cancer Early Detection
- Describe the rationale behind the decision-making process for developing the NCCN Guidelines for Prostate Cancer Early Detection

Disclosure of Relevant Financial Relationships**Editor:**

Kerrin M. Green, MA, Assistant Managing Editor, *JNCCN—Journal of the National Comprehensive Cancer Network*, has disclosed that she has no relevant financial relationships.

CE Authors:

Deborah J. Moonan, RN, BSN, Director, Continuing Education, NCCN, has disclosed that she has no relevant financial relationships.

Kristina M. Gregory, RN, MSN, OCN, Vice President, Clinical Information Operations, NCCN, has disclosed that she has no relevant financial relationships.

Rashmi Kumar, PhD, Senior Manager, Clinical Content, NCCN, has disclosed that she has no relevant financial relationships.

Individuals Who Provided Content Development and/or Authorship Assistance:

Peter R. Carroll, MD, MPH, Panel Chair, has disclosed that he receives consulting fees and/or honoraria from Genomic Health, Inc. and Janssen Pharmaceutica Products, LP; and grant/research support from Genomic Health, Inc. and Myriad Genetic Laboratories, Inc.

J. Kellogg Parsons, MD, MHS, Panel Vice Chair, has disclosed that he receives consulting fees and/or honoraria from MDx Health.

Erik P. Castle, MD, Panel Member, has disclosed that he has no relevant financial relationships.

William J. Catalona, MD, Panel Member, has disclosed that he received grant/research support from and served as a scientific advisor for Beckman-Coulter Inc.

John W. Davis, MD, Panel Member, has disclosed that he receives consulting fees and/or honoraria from Intuitive Surgical, Inc. and Myriad Genetic Laboratories, Inc.

Jonathan I. Epstein, MD, Panel Member, has disclosed that he is a consultant for Metamark Genetics, Inc. and Dianon Pathology.

Thomas Farrington, BSEE, Panel Member, has disclosed that he has no relevant financial relationships.

Paul H. Lange, MD, Panel Member, has disclosed that he has no relevant financial relationships.

William Lowrance, MD, MPH, Panel Member, has disclosed that he receives grant/research support from Myriad Genetic Laboratories, Inc. and Argos Therapeutics, Inc.; and is on the advisory board for Myriad Genetic Laboratories, Inc. and MDx Health.

Paul Maroni, MD, Panel Member, has disclosed that he has no relevant financial relationships.

James Mohler, MD, Panel Member, has disclosed that he has no relevant financial relationships.

Robert B. Nadler, MD, Panel Member, has disclosed that he has no relevant financial relationships.

Terrence M. Shaneyfelt, MD, MPH, Panel Member, has disclosed that he has no relevant financial relationships.

Marc C. Smaldone, MD, MSHP, Panel Member, has disclosed that he has no relevant financial relationships.

Geoffrey Sonn, MD, Panel Member has disclosed that he has no relevant financial relationships.


Robert Wake, MD, Panel Member has disclosed that he has no relevant financial relationships.

Dorothy A. Shead, MS, Director, Patient & Clinical Information Operations, NCCN, has disclosed that she has no relevant financial relationships.

Deborah A. Freedman-Cass, PhD, Oncology Scientist/Senior Medical Writer, NCCN, has disclosed that she has no relevant financial relationships.

This activity is supported by educational grants from AstraZeneca, Bayer Healthcare Pharmaceuticals Inc., Bristol-Myers Squibb, Clovis Oncology, Foundation Medicine, Genentech, Novartis Oncology, Otsuka America Pharmaceutical, Inc., Seattle Genetics, Inc., and Takeda Oncology; support provided by Actelion Pharmaceuticals US, Inc.; and by an independent educational grant from Astellas and Medivation, Inc.

Prostate Cancer Early Detection, Version 2.2016



^aAfrican-American men have a higher incidence of prostate cancer, increased prostate cancer mortality, and earlier age of diagnosis compared to Caucasian-American men. However, the effects of earlier or more intensive screening on cancer outcomes and on screening-related harms in African-American men remain unclear. Therefore, although these men may require a higher level of vigilance and different considerations when analyzing the results of screening tests, the panel cannot provide separate screening recommendations for these men until more data become available.

^bThe best evidence supports the use of serum PSA for the early detection of prostate cancer. DRE should not be used as a stand-alone test, but should be performed in those with an elevated serum PSA. DRE may be considered as a baseline test in all patients as it may identify high-grade cancers associated with "normal" serum PSA values. Consider referral for biopsy, if DRE is very suspicious. Medications such as 5 α -reductase inhibitors (finasteride and dutasteride) are known to decrease PSA by approximately 50%, and PSA values in these men should be corrected accordingly.

^cTesting above the age of 75 years of age should be done with caution and only in very healthy men with little or no comorbidity as a large proportion may harbor cancer that would be unlikely to affect their life expectancy, and screening in this population would substantially increase rates of over-detection. However, a clinically significant number of men in this age group may present with high-risk cancers that pose a significant risk if left undetected until signs or symptoms develop. One could consider increasing the PSA threshold for biopsy in this group (i.e., >4 ng/mL). Very few men above the age of 75 years benefit from PSA testing.

^dThe reported median PSA values for men aged 40-49 y range from 0.5-0.7 ng/mL, and the 75th percentile values range from 0.7-0.9 ng/mL. Therefore, the PSA value of 1.0 ng/mL selects for the upper range of PSA values. Men who have a PSA above the median for their age group are at a higher risk for prostate cancer and for the aggressive form of the disease. The higher above the median, the greater the risk.

^eMen age \geq 60 years with serum PSA <1.0 ng/mL have a very low risk of metastases or death due to prostate cancer and may not benefit from further testing. A PSA cut point of 3.0 ng/mL at age 75 years also low risk of poor outcome.

Version 2.2016 © National Comprehensive Cancer Network, Inc. 2016. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

PROSD-2

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.


Clinical trials: NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

Overview

In men, prostate cancer is the most commonly diagnosed cancer and the second leading cause of cancer death. Prostate cancer represents a spectrum of disease that ranges from nonaggressive and slow-growing disease that does not require treatment to aggressive and fast-growing disease that does necessitate treatment. Early detection strategies should maximize the detection of aggressive prostate cancer that is potentially curable, while minimizing unnecessary procedures and the detection of indolent disease. Population-based prostate cancer screening studies have shown survival benefits using prostate-specific antigen (PSA) levels, sometimes in combination with a digital rectal examination or other ancillary tests.¹⁻⁵ Therefore, the NCCN Prostate Cancer Early Detection Panel supports the use of PSA testing for the early detection of prostate cancer in informed, healthy men in specific age groups and encourages active surveillance or observation for appropriate candidates in whom low-risk disease is

Prostate Cancer Early Detection, Version 2.2016

INDICATIONS FOR BIOPSY



TRUS-GUIDED BIOPSY
Initial and Repeat
Extended-pattern biopsy (12 cores)

- Number of cores:
 - ▶ Sextant (6),
 - ▶ Lateral peripheral zone (6), and
 - ▶ Lesion-directed at palpable nodule or suspicious image
- Anteriorly directed biopsy is not supported in routine biopsy. However, the addition of a transition zone biopsy to an extended biopsy protocol may be considered in a repeat biopsy if PSA is persistently elevated.
- Multiparametric MRI followed by lesion targeting may maximize the detection of higher risk disease and limit the detection of lower risk disease.ⁱ
- Local anesthesia can decrease pain/discomfort associated with prostate biopsy and should be offered to all patients.

^fThe level of PSA correlates with the risk of prostate cancer. The Prostate Cancer Prevention Trial (PCPT) demonstrated that 15% of men with a PSA level of ≤4.0 ng/mL and a normal DRE had prostate cancer diagnosed on end-of-study biopsies. Approximately 30% to 35% of men with serum PSA between 4 to 10 ng/mL will be found to have cancer. Total PSA levels >10 ng/mL confer a greater than 67% likelihood of prostate cancer.

^gFor patients with abnormal DRE, biopsy or additional testing should be considered based on concern for cancer.

^hBiomarkers that improve the specificity of detection are not recommended as firstline screening tests. However, there may be some patients who meet PSA standards for consideration of prostate biopsy, but for whom the patient and/or the physician wish to further define the probability of high-grade cancer. A percent free PSA <10%, PHI >35 or 4Kscore (which provides an estimate of the probability of high-grade prostate cancer) are potentially informative in patients who have never undergone biopsy or after a negative biopsy; a PCA3 score >35 is potentially informative after a negative biopsy.

ⁱMRI is not recommended routinely prior to initial prostate biopsy, but emerging data suggest that, in men undergoing initial biopsy, targeting using MRI/ultrasound fusion may increase the detection of clinically significant, higher-risk (Gleason grade ≥ 4+3=7) disease while lowering the detection of lower-risk (Gleason sum 6 or lower-volume Gleason grade 3+4=7) disease. Siddiqui M, Rais-Bahrami S, Turkbey B, et al. Comparison of MRI/Ultrasound Fusion-Guided Biopsy With Ultrasound-Guided Biopsy for the Diagnosis of Prostate Cancer. *JAMA* 2015;313:390-7.

Version 2.2016 © National Comprehensive Cancer Network, Inc. 2016. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

PROSD-3

identified (see the full NCCN Guideline recommendations online at NCCN.org).

The issues discussed by the panel this year are described in these NCCN Guidelines Insights. Changes to the NCCN Guidelines algorithms are indicated with blue font herein.

Patient Selection for Biopsies

PSA testing results in the diagnosis of high-grade tumors at a potentially earlier, more curable stage than clinical detection. However, PSA testing has poor specificity and often produces false-positive results, which contribute to patient anxiety and increased costs and potential complications associated with unnecessary biopsies (eg, drug-resistant *Escherichia coli* infections). PSA testing also leads to the detection of much indolent disease (overdetection), which contributes to overtreatment, patient stress, possible complications, and inconvenience of active surveillance. Additional techniques used with PSA

testing have the potential to refine patient selection for biopsies, decrease unnecessary biopsies, reduce overdetection, and increase biopsy specificity, without missing a substantial number of aggressive cancers. Such improvements to PSA screening could make it a more valuable tool for men who choose to participate in an early detection program by increasing the benefits while decreasing the risks.


Therefore, the panel discussed the roles of biomarker testing and advanced imaging to refine patient selection for initial and repeat biopsies, as described herein.

Biomarker Tests

When a patient meets the standards for biopsy, patients and physicians may wish to further define the probability of cancer before proceeding to biopsy with its associated risks. Several biomarker tests have been developed with the goals of refining patient selection for biopsies, decreasing unnecessary biopsies, and increasing the specificity of cancer

Prostate Cancer Early Detection, Version 2.2016

MANAGEMENT OF BIOPSY RESULTS



^jIt is well known that a negative prostate biopsy does not preclude a diagnosis of prostate cancer on subsequent biopsy. Those patients with negative prostate biopsies should be followed with DRE and PSA. Tests that improve specificity in the post-biopsy state—including 4Kscore, PHI, percent free PSA, PCA3, and ConfirmMDx—should be considered in patients thought to be higher risk despite a negative prostate biopsy (See PROSD-3).

^kPSA testing may be discontinued at certain ages and PSA cutpoints, as noted in the discussion section.

^lEmerging evidence suggests that use of multiparametric MRI and/or use of refined prostate biopsy techniques (image guidance using MRI/ultrasound fusion, transperineal, or saturation prostate biopsies) may be of value. These techniques may help identify regions of cancer missed on prior prostate biopsies and should be considered in selected cases after at least 1 negative prostate biopsy. Multiparametric MRI followed by lesion targeting may maximize the detection of higher risk disease and limit the detection of lower risk disease.

Version 2.2016 © National Comprehensive Cancer Network, Inc. 2016. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

PROSD-4

detection, without missing a substantial number of higher-grade (Gleason score ≥ 7) cancers. These tests may be especially useful in men with PSA levels between 3 and 10 ng/mL. Most often, these tests have been used in patients with one negative biopsy to determine whether repeat biopsy is an appropriate consideration.

In the 2015 version of these guidelines, the panel recommended consideration of percent free PSA, 4Kscore, or Prostate Health Index (PHI) in patients with PSA levels greater than 3 ng/mL who have not had a biopsy. For men who have had at least one prior negative biopsy and are thought to be at higher risk, the panel recommended the same tests or PCA3.⁶⁻¹⁶

Multiple institutional review comments requested that the panel consider adding ConfirmMDx in the repeat biopsy setting based on data showing that it has potential value for patients with an elevated PSA and prior negative biopsy. ConfirmMDx is a tissue-based, multiplex epigenetic assay that aims to improve the stratification of men being considered

for repeat biopsy. Hypermethylation of the promoter regions of *GSTP1*, *APC*, and *RASSF1* are assessed in core biopsy tissue samples.^{17,18} The panel found the available data convincing and unanimously agreed (with several abstentions) to add consideration of ConfirmMDx for men contemplating repeat biopsy (see PROSD-4; above).

The panel then discussed how to compare the various recommended tests and whether they could recommend one or more tests over the others. As the list of potential assays grows, the NCCN Guidelines Panel believes that clinicians and patients will need more guidance. These tests have different objectives (eg, improving sensitivity, reducing false-positives), different impacts on downstream outcomes (eg, mortality reduction, reduced overdiagnosis), and are used in different settings (ie, initial or repeat biopsy). Ideally, the negative predictive value and positive predictive value (PPV) for any cancer and for high-grade cancer; the number of biopsies avoided; and the number of aggressive

Prostate Cancer Early Detection, Version 2.2016

cancers missed for each test could be compared. However, studies for the different assays have been performed with different cohorts, settings, time points, and reported outcomes. Head-to-head comparisons have been performed in Europe for some of these tests, used independently or in combinations in the initial or repeat biopsy settings, but sample sizes were small and results varied.^{19–26}

Therefore, the panel emphasized the following points: no biomarker test can be recommended over any other; a biomarker assay can be performed in addition to multiparametric MRI (mp-MRI)/refined biopsy techniques in the repeat biopsy setting (discussed in the following section); the optimal order of biomarker tests and imaging is unknown; and it remains unclear how to interpret results of multiple tests in individual patients, especially when results are contradictory. The panel further emphasized that results of any of these tests, when performed, should be used as part of discussions between clinician and patient to assist in decisions regarding whether to proceed with biopsy.

Multiparametric MRI

Considerable interest exists for the use of novel MRI, most notably mp-MRI, to select patients who need a prostate biopsy or to guide needle placement during the procedure, although data supporting its use in these settings are currently limited.^{27–29} The goals of using MRI to inform the decision of whether to perform a biopsy include reducing the number of men undergoing biopsy, reducing the detection of indolent disease (and thus the risks of overdiagnosis and overtreatment), and improving the detection of clinically significant disease through targeted biopsies (see “Advanced Biopsy Techniques,” this page).

The panel discussed the use of MRI before initial biopsy based on the available data analyzing the clinical utility of mp-MRI for patient selection in the initial biopsy setting. In a prospective study of 223 biopsy-naïve men with an elevated PSA level, all men underwent standard transrectal ultrasound (TRUS) biopsies in addition to mp-MRI.²⁸ Not performing a biopsy in men with PI-RADS 1/2 lesions on MRI would have reduced the number of men requiring biopsy by 36%, reduced the identification of low-risk prostate cancer by 87%, and increased the yield of intermediate/high-risk tumors by 18%, but would have missed 15 intermediate/high-risk tumors (6.7% of study population). A trial at a single

hospital-based practice randomized 130 biopsy-naïve men with elevated PSA levels to either TRUS-guided random biopsy alone or to prebiopsy mp-MRI, TRUS-guided random biopsy, and cognitive MRI/TRUS fusion targeted biopsy.²⁹ In this study, 13 tumors with Gleason score 3 + 4 would have been missed in the 53 evaluable men in the mp-MRI group based on MRI results (24.5%). Additional clinical trials are underway to assess the value of MRI for diagnosis in the prebiopsy setting (ClinicalTrials.gov identifiers: NCT02131207, NCT02485379, and NCT02450266).

Based on these studies, the panel agreed that the use of MRI to exclude men from biopsies could lead to many clinically significant cancers being missed. Overall, the panel believes that MRI alone should not be used to determine whether to perform an initial biopsy, and emphasizes that a negative MRI is not a reason to forego biopsy in men with strong indications for a first-time biopsy. All men with indications for biopsy should receive the standard 12-core TRUS-guided biopsy regardless of MRI results, with possible additional targeted biopsies. The panel therefore added a footnote stating that MRI is not currently routinely recommended before initial biopsy to accentuate these points (see PROSD-3; page 512). Additionally, the panel acknowledged emerging data showing that mp-MRI followed by lesion targeting may increase the detection of higher-risk disease while lowering the detection of lower-risk disease (see “Advanced Biopsy Techniques,” below).³⁰

Advanced Biopsy Techniques

Targeted, template, and saturation biopsy techniques are being studied in hope of improving detection of clinically significant prostate cancer while minimizing the detection of indolent disease.^{31–35} Targeted biopsy techniques include cognitive or visual targeting (ultrasound-guided, based on an MRI image), MRI/TRUS fusion platforms (merging a stored MRI image with a real-time ultrasound image), and direct in-bore magnetic resonance-guided biopsy (performed by an interventional radiologist while the patient is in the scanner).^{31–33} In saturation biopsies, cores are systematically collected every few millimeters across the entire prostate to improve detection over that of a standard 12-core biopsy. Saturation

Prostate Cancer Early Detection, Version 2.2016

biopsies can be performed via transrectal or transperineal approaches. Another advanced technique is transperineal template biopsy, which does not saturate every single area but instead systematically samples anterior, mid, posterior, and basal zones for approximately 24 to 32 cores.³⁶

For the 2016 guideline update, the NCCN Guidelines Panel discussed the role of targeted biopsies in the initial and repeat settings, as described in these Insights.

Targeted Biopsies in the Initial Biopsy Setting

The panel discussed the data on targeted biopsy techniques for biopsy-naïve patients. In a prospective study of 223 biopsy-naïve men with elevated PSA levels, all men had standard TRUS biopsies and mp-MRI.²⁸ Participants with suspicious or equivocal lesions (PI-RADS 3–5) then underwent MRI-guided biopsy. TRUS biopsies detected 126 of 142 cases (88.7%), including 47 cases classified as low risk. The MRI-guided biopsies identified an additional 16 cases of intermediate/high-risk prostate cancer and led to the reclassification of 13 cases from low risk to intermediate/high risk. Therefore, the addition of mp-MRI with targeted biopsies for suspicious or equivocal lesions to standard biopsy allowed the identification of clinically significant disease in an additional 13% of the study population. A single-center trial randomized 130 biopsy-naïve men to receive either TRUS-guided random biopsy alone or prebiopsy mp-MRI, TRUS-guided random biopsy, and cognitive MRI/TRUS fusion targeted biopsy.²⁹ Similar rates of detection of prostate cancer (64% vs 57%; $P=.5$) and clinically significant cancer (55% vs 45%; $P=.8$) were seen in the 2 arms.

In a single-institution prospective cohort study, 1,003 men with elevated PSA levels or an abnormal digital rectal examination and lesions visible on mp-MRI underwent both MRI/ultrasound fusion targeted and standard biopsy.³⁰ In this study, 196 men had no prior biopsy, and results appear to be similar in the biopsy-naïve subgroup compared with the entire cohort. In the full cohort, 170 men had pathology results available after radical prostatectomy: 8 men (4.7%) had intermediate- or high-risk cancers that would have been missed based on targeted biopsy results of no or low-risk cancer, and 44 men (26%) had intermediate- or high-risk cancers that would have been missed based on standard biopsy results of no or low-risk cancer. The sensitivities of targeted

and standard biopsies for detecting intermediate- or high-grade cancer were 77% and 53%, respectively, whereas the specificities of the approaches were similar, at 68% and 66%, respectively. Combining both biopsy techniques increased sensitivity to 85% but decreased specificity to 49%. The effect of targeted biopsies on clinical outcomes is still unknown.

The panel did not believe that the current data for the use of targeted biopsies in the initial biopsy setting were sufficient to recommend them over standard biopsies in this setting. However, the panel agreed that it was important to acknowledge the emerging data showing that mp-MRI followed by lesion targeting may increase the detection of clinically significant, higher-risk disease (Gleason $\geq 4 + 3$) while lowering the detection of lower-risk disease (Gleason 6 or lower-volume 3 + 4) (see PROSD-3; page 512).

Targeted Biopsies in the Repeat Biopsy Setting

A negative biopsy does not preclude a diagnosis of prostate cancer on subsequent biopsy, and repeat biopsies are often considered in men for whom clinical suspicion of high-risk disease persists. Reviewers requested that the NCCN Guidelines Panel consider the addition of mp-MRI-guided biopsy techniques for patients with prior negative biopsies and for patients in whom a high suspicion exists for clinically significant prostate cancer. These individuals pointed out that increasing data are demonstrating the superiority of MRI-targeted biopsy techniques over standard biopsy for the detection of clinically significant/high-grade prostate cancer after a negative biopsy, and therefore requested that the panel consider putting more emphasis on these approaches by adding them to the algorithm. The panel was also asked to consider where targeted biopsy belonged in the diagnostic algorithm relative to the recommended biomarkers assays in this repeat biopsy setting.

In a prospective study that included 347 patients with findings suspicious for prostate cancer, many of whom had 1 or more previous negative biopsies, all patients received mp-MRI and those with abnormal findings proceeded to MRI/TRUS fusion targeted biopsies.²⁷ The outcome was defined as improved detection in targeted cores, with significantly more cancer detected in targeted cores than in systematic biopsies (30% vs 8.2%). Approximately 12% of men without MRI-suspicious lesions were diagnosed with intermediate-risk tumors. In this study, the cancer

Prostate Cancer Early Detection, Version 2.2016

detection rate was 51% in men with previous negative biopsies. In a prospective cohort study of 1,003 men undergoing both MRI-targeted and standard biopsy (see “Targeted Biopsies in the Initial Biopsy Setting,” page 515), Siddiqui et al³⁰ noted that targeted biopsies were associated with the detection of high-risk cancer and decreased detection of low-risk cancer.³⁰ A recent meta-analysis of 16 studies (1,926 men) also showed that MRI-targeted biopsy improved detection of clinically significant prostate cancer in biopsy-naïve men compared with standard TRUS biopsy.³⁷

Although the NCCN Guidelines Panel found the increasing evidence convincing, they also pointed out that the publications come from only a few centers with a small number of clinicians and radiologists. No data to date show that these techniques will work in other oncology settings (eg, community hospitals). Overall, the panel believes that the level of evidence has risen to the level needed for targeted biopsies to be considered in the repeat biopsy setting. The panel agreed to add mp-MRI and/or refined biopsy techniques for consideration in men with benign findings or focal high-grade prostatic intraepithelial neoplasia (see PROSD-4; page 513). These techniques include saturation biopsy strategies and the use of mp-MRI followed by a targeted biopsy technique based on the results, and these techniques can be considered before or after biomarker tests (discussed previously) to aid in patient/clinician discussions.

Screening in High-Risk Populations

Certain populations (eg, African Americans, men with a first-degree relative with prostate cancer) have a higher risk of developing prostate cancer.^{38–42} At the meeting for the 2016 guideline update, the panel discussed the role that race and genetic susceptibility play in prostate cancer and whether or not screening recommendations should be altered for high-risk men.

Race

African American men have a 64% higher incidence of prostate cancer and 2.3-fold increase in prostate cancer mortality compared with white men, and are approximately 4 times more likely to be diagnosed with metastatic disease.^{38,43} Furthermore, autopsy data indicate that prostate cancer may un-

dergo transformation to aggressive disease earlier in African American men than in white men.⁴³ Factors that contribute to this racial disparity may include differences in genetic risk factors, environmental exposures, and patient and physician behaviors; decreased access to high-quality health care, including cancer early detection and follow-up care; delays in diagnosis; and suboptimal treatment.^{44–47}

Two institutional reviewers introduced the topic of prostate cancer early detection in African Americans. In particular, they questioned whether discussions about screening should be initiated earlier with this population, if screening should start earlier in this group, and whether these men should be screened more intensively. The panel discussion that ensued found a lack of data to inform the best strategy for prostate cancer screening in these men. Prostate cancer screening has been best studied in white men; data on screening in diverse and high-risk populations are lacking. In the PLCO trial, approximately 4.4% of the participants were African American and 6.9% had a positive family history, but no subset analyses were performed for black men.⁴⁸ In the ERSPC trial, no information on race or family history was reported.³

The panel discussed data suggesting that African American men have an earlier onset of prostate cancer, which might support earlier screening initiation. An analysis of SEER data from 2010 found that non-Hispanic black men were diagnosed with prostate cancer an adjusted average of 1.2 years earlier than non-Hispanic white men,⁴⁹ whereas an older SEER analysis found that African American men were diagnosed at an average of 3 years younger than white men.⁵⁰ A retrospective population-based cohort study in the United Kingdom found that black men were diagnosed an adjusted average of 5.1 years earlier than white men.⁵¹ Another study estimated that African American men have an almost 2-fold higher risk of being diagnosed with prostate cancer before the age of 45 years compared with white men.⁵⁰

Although the panel members agreed that this topic cannot and should not be ignored, they believe that data do not support earlier or more intensive screening at this time. Race is, however, included as a baseline evaluation factor for risk assessment that can be used to help decide when a man should begin the early detection process within the NCCN recommended ages of 45 to 75 years. Clearly, African

Prostate Cancer Early Detection, Version 2.2016

American men are at a higher risk for prostate cancer and prostate cancer mortality, but the effects of earlier or more intensive screening on cancer outcomes and screening-related harms in this population are not clear. The panel also noted that a baseline PSA value is a stronger predictive factor than a positive family history or race.⁵² Therefore, although these individuals may require a higher level of vigilance and potentially different considerations when analyzing the results of screening tests, the panel did not provide separate screening recommendations for this patient population at this time. Instead, they added a footnote to emphasize these points (see PROSD-2; page 511).

BRCA1 and BRCA2 Mutations

Germline *BRCA1* and *BRCA2* mutations (associated with hereditary breast and/or ovarian cancer syndrome) have been associated with an increased risk for prostate.^{53–61} In particular, *BRCA2* mutations have been associated with a 2- to 6-fold increase in the risk for prostate cancer, whereas the association between *BRCA1* mutations and an increased risk for prostate cancer is less consistent.^{54,56,57,61–63} Furthermore, prostate cancer in men with germline *BRCA* mutations appears to have a more aggressive phenotype and is associated with significantly reduced survival times than prostate cancer in those who do not have the gene.^{64–68}

The panel received an external submission requesting that genetic risk assessment be recommended in the NCCN Guidelines for Prostate Cancer Early Detection based on recommendations in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian. These latter guidelines include personalized risk assessment, genetic counseling, and possible genetic testing for individuals with a personal history of prostate cancer and 1 or more close blood relatives with breast cancer (age ≤ 50 years) or ovarian cancer, or 2 relatives with breast, pancreatic, or prostate cancer (Gleason score ≥ 7). Individuals meeting these criteria are considered at risk of harboring a *BRCA1/2* mutation.⁶⁹ These guidelines note that individuals with a family history only can also be considered for genetic testing if no living affected family member is available, but that the limitations of testing in unaffected relatives should be discussed.⁶⁹

Although the panel did not believe that the NCCN Guidelines for Prostate Cancer Early Detec-

tion should list recommendations for genetic risk assessment, they did agree that men should be asked about the presence of known *BRCA1/2* mutations in their families. Therefore, the panel added “Family history of *BRCA1/2* mutations” to the baseline evaluation that precedes discussions on the risks and benefits of prostate cancer screening (see PROSD-2; page 511). If there is a known or suspected cancer susceptibility gene, referral to a cancer genetics professional is recommended. Information regarding *BRCA1/2* gene status should be used as part of the discussion about prostate cancer screening; patients may not be aware of the increased risk of prostate cancer associated with these mutations. Some panel members believe that men with known germline *BRCA1/2* mutations should be encouraged to undergo PSA testing.

The panel then discussed whether a known *BRCA1* or *BRCA2* mutation would impact prostate cancer screening. The NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian (available at NCCN.org) recommend that men with *BRCA2* mutations begin prostate cancer screening at age 40 years and that those with *BRCA1* mutations consider the same.⁶⁹ Results were recently reported from the first round of screening in the IMPACT study, which enrolled men aged 40 to 69 years with germline *BRCA1/2* mutations and a control group of men with wild-type *BRCA1/2* related to mutation carriers.⁷⁰ Although no difference was shown between carriers and controls in the rate of prostate cancer detection or the PPV of biopsy for detecting cancer in men with PSA levels greater than 3.0 ng/mL, a significant difference was seen in the PPV of biopsy for detecting intermediate/high-grade cancer in *BRCA2* carriers with PSA levels greater than 3.0 ng/mL (2.4% vs 0.7%; $P=.04$). Future rounds of screening in this trial may help inform the best strategy for screening in this high-risk population. Therefore, the panel noted that, although these mutations are clearly risk factors, data supporting a change in the PSA screening and biopsy recommendations for these men relative to those without mutations are insufficient at this time. The panel also noted that, as with race, baseline PSA value is a stronger predictive factor than a positive family history.⁵²

Prostate Cancer Early Detection, Version 2.2016

Conclusions

The NCCN Guideline revisions for the 2016 update highlight several techniques designed to improve the identification of significant cancer while avoiding the detection of indolent disease. Advanced biopsy techniques, including targeted biopsies, are one promising technique to help advance these goals. In addition, patient selection for biopsies can be refined through the appropriate use of biomarker assays and advanced imaging, thereby minimizing unnecessary procedures and limiting the detection of indolent disease. In the future, data may inform more individualized, risk-based approaches to prostate cancer early detection that take into account race, genetic predisposition, and family history.

References

- Andriole GL. Update of the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Recent Results Cancer Res* 2014;202:53–57.
- Andriole GL, Crawford ED, Grubb RL III, et al. Prostate cancer screening in the randomized Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial: mortality results after 13 years of follow-up. *J Natl Cancer Inst* 2012;104:125–132.
- Schroder FH, Hugosson J, Roobol MJ, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med* 2009;360:1320–1328.
- Schroder FH, Hugosson J, Roobol MJ, et al. Prostate-cancer mortality at 11 years of follow-up. *N Engl J Med* 2012;366:981–990.
- Schroder FH, Hugosson J, Roobol MJ, et al. Screening and prostate cancer mortality: results of the European Randomised Study of Screening for Prostate Cancer (ERSPC) at 13 years of follow-up. *Lancet* 2014;384:2027–2035.
- Catalona WJ, Partin AW, Sanda MG, et al. A multicenter study of [-2]prostate specific antigen combined with prostate specific antigen and free prostate specific antigen for prostate cancer detection in the 2.0 to 10.0 ng/ml prostate specific antigen range. *J Urol* 2011;185:1650–1655.
- Filella X, Gimenez N. Evaluation of [-2] proPSA and Prostate Health Index (phi) for the detection of prostate cancer: a systematic review and meta-analysis. *Clin Chem Lab Med* 2013;51:729–739.
- Lazzeri M, Haese A, Abrate A, et al. Clinical performance of serum prostate-specific antigen isoform [-2]proPSA (p2PSA) and its derivatives, %p2PSA and the prostate health index (PHI), in men with a family history of prostate cancer: results from a multicentre European study, the PROMEtheuS project. *BJU Int* 2013;112:313–321.
- Loeb S. Prostate cancer: Prostate Health Index—improving screening in men with family history. *Nat Rev Urol* 2013;10:497–498.
- Partin AW, Brawer MK, Subong EN, et al. Prospective evaluation of percent free-PSA and complexed-PSA for early detection of prostate cancer. *Prostate Cancer Prostatic Dis* 1998;1:197–203.
- Bryant RJ, Sjoberg DD, Vickers AJ, et al. Predicting high-grade cancer at ten-core prostate biopsy using four kallikrein markers measured in blood in the ProtecT Study. *J Natl Cancer Inst* 2015;107:pii: djv095.
- Parekh DJ, Punnen S, Sjoberg DD, et al. A multi-institutional prospective trial in the USA confirms that the 4Kscore accurately identifies men with high-grade prostate cancer. *Eur Urol* 2015;68:464–470.
- Vickers A, Cronin A, Roobol M, et al. Reducing unnecessary biopsy during prostate cancer screening using a four-kallikrein panel: an independent replication. *J Clin Oncol* 2010;28:2493–2498.
- Vickers AJ, Gupta A, Savage CJ, et al. A panel of kallikrein marker predicts prostate cancer in a large, population-based cohort followed for 15 years without screening. *Cancer Epidemiol Biomarkers Prev* 2011;20:255–261.
- Vickers AJ. Markers for the early detection of prostate cancer: some principles for statistical reporting and interpretation. *J Clin Oncol* 2014;32:4033–4034.
- Wei JT, Feng Z, Partin AW, et al. Can urinary PCA3 supplement PSA in the early detection of prostate cancer? *J Clin Oncol* 2014;32:4066–4072.
- Stewart GD, Van Neste L, Delvenne P, et al. Clinical utility of an epigenetic assay to detect occult prostate cancer in histopathologically negative biopsies: results of the MATLOC study. *J Urol* 2013;189:1110–1116.
- Partin AW, Van Neste L, Klein EA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol* 2014;192:1081–1087.
- Auprich M, Augustin H, Budaus L, et al. A comparative performance analysis of total prostate-specific antigen, percentage free prostate-specific antigen, prostate-specific antigen velocity and urinary prostate cancer gene 3 in the first, second and third repeat prostate biopsy. *BJU Int* 2012;109:1627–1635.
- Boegemann M, Stephan C, Cammann H, et al. The percentage of prostate-specific antigen (PSA) isoform [-2]proPSA and the Prostate Health Index improve the diagnostic accuracy for clinically relevant prostate cancer at initial and repeat biopsy compared with total PSA and percentage free PSA in men aged ≤ 65 years. *BJU Int* 2016;117:72–79.
- De Luca S, Passera R, Bollito E, et al. Comparison of prostate cancer gene 3 score, prostate health index and percentage free prostate-specific antigen for differentiating histological inflammation from prostate cancer and other non-neoplastic alterations of the prostate at initial biopsy. *Anticancer Res* 2014;34:7159–7165.
- De Luca S, Passera R, Fiori C, et al. Prostate health index and prostate cancer gene 3 score but not percent-free prostate specific antigen have a predictive role in differentiating histological prostatitis from PCa and other nonneoplastic lesions (BPH and HG-PIN) at repeat biopsy. *Urol Oncol* 2015;33:424.e417–423.
- Ferro M, Bruzzese D, Perdoni S, et al. Prostate Health Index (Phi) and prostate cancer antigen 3 (PCA3) significantly improve prostate cancer detection at initial biopsy in a total PSA range of 2-10 ng/ml. *PLoS One* 2013;8:e67687.
- Perdoni S, Bruzzese D, Ferro M, et al. Prostate health index (phi) and prostate cancer antigen 3 (PCA3) significantly improve diagnostic accuracy in patients undergoing prostate biopsy. *Prostate* 2013;73:227–235.
- Porpiglia F, Russo F, Manfredi M, et al. The roles of multiparametric magnetic resonance imaging, PCA3 and prostate health index-which is the best predictor of prostate cancer after a negative biopsy? *J Urol* 2014;192:60–66.
- Scattoni V, Lazzeri M, Lughezzi G, et al. Head-to-head comparison of prostate health index and urinary PCA3 for predicting cancer at initial or repeat biopsy. *J Urol* 2013;190:496–501.
- Kuru TH, Roethke MC, Seidenader J, et al. Critical evaluation of magnetic resonance imaging targeted, transrectal ultrasound guided transperineal fusion biopsy for detection of prostate cancer. *J Urol* 2013;190:1380–1386.
- Pokorny MR, de Rooij M, Duncan E, et al. Prospective study of diagnostic accuracy comparing prostate cancer detection by transrectal ultrasound-guided biopsy versus magnetic resonance (MR) imaging with subsequent MR-guided biopsy in men without previous prostate biopsies. *Eur Urol* 2014;66:22–29.
- Tonttila PP, Lantto J, Paakko E, et al. Prebiopsy multiparametric magnetic resonance imaging for prostate cancer diagnosis in biopsy-naive men with suspected prostate cancer based on elevated prostate-specific antigen values: results from a randomized prospective blinded controlled trial. *Eur Urol* 2016;69:419–425.
- Siddiqui MM, Rais-Bahrami S, Turkbey B, et al. Comparison of MR/ultrasound fusion-guided biopsy with ultrasound-guided biopsy for the diagnosis of prostate cancer. *JAMA* 2015;313:390–397.
- Rastinehad AR, Turkbey B, Salami SS, et al. Improving detection of clinically significant prostate cancer: MRI/TRUS fusion-guided prostate biopsy. *J Urol* 2014;191:1749–1754.
- Robertson NL, Emberton M, Moore CM. MRI-targeted prostate biopsy: a review of technique and results. *Nat Rev Urol* 2013;10:589–597.
- Puech P, Rouviere O, Renard-Penna R, et al. Prostate cancer diagnosis: multiparametric MR-targeted biopsy with cognitive and transrectal US-MR fusion guidance versus systematic biopsy—prospective multicenter study. *Radiology* 2013;268:461–469.
- Abdollah F, Novara G, Briganti A, et al. Trans-rectal versus trans-perineal saturation rebiopsy of the prostate: is there a difference in cancer detection rate? *Urology* 2011;77:921–925.
- Nelson AW, Harvey RC, Parker RA, et al. Repeat prostate biopsy strategies after initial negative biopsy: meta-regression comparing cancer detection of transperineal, transrectal saturation and MRI guided biopsy. *PLoS One* 2013;8:e57480.
- Sivaraman A, Sanchez-Salas R, Barret E, et al. Transperineal template-guided mapping biopsy of the prostate. *Int J Urol* 2015;22:146–151.
- Schoots IG, Roobol MJ, Nieboer D, et al. Magnetic resonance imaging-targeted biopsy may enhance the diagnostic accuracy of significant prostate cancer detection compared to standard transrectal ultrasound-guided biopsy: a systematic review and meta-analysis. *Eur Urol* 2015;68:438–450.

Prostate Cancer Early Detection, Version 2.2016

38. SEER Stat Fact Sheets: Prostate Cancer. 2015. Available at: <http://seer.cancer.gov/statfacts/html/prost.html>. Accessed April 28, 2015.
39. Bratt O. Hereditary prostate cancer: clinical aspects. *J Urol* 2002;168:906–913.
40. Carter BS, Beaty TH, Steinberg GD, et al. Mendelian inheritance of familial prostate cancer. *Proc Natl Acad Sci U S A* 1992;89:3367–3371.
41. Chen YC, Page JH, Chen R, Giovannucci E. Family history of prostate and breast cancer and the risk of prostate cancer in the PSA era. *Prostate* 2008;68:1582–1591.
42. Grill S, Fallah M, Leach RJ, et al. Incorporation of detailed family history from the Swedish Family Cancer Database into the PCPT risk calculator. *J Urol* 2015;193:460–465.
43. Powell IJ, Bock CH, Ruterbusch JJ, Sakr W. Evidence supports a faster growth rate and/or earlier transformation to clinically significant prostate cancer in black than in white American men, and influences racial progression and mortality disparity. *J Urol* 2010;183:1792–1796.
44. Barocas DA, Grubb R III, Black A, et al. Association between race and follow-up diagnostic care after a positive prostate cancer screening test in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening trial. *Cancer* 2013;119:2223–2229.
45. Mahal BA, Aizer AA, Ziehr DR, et al. Trends in disparate treatment of African American men with localized prostate cancer across National Comprehensive Cancer Network risk groups. *Urology* 2014;84:386–392.
46. Yamoah K, Johnson MH, Choerung V, et al. Novel biomarker signature that may predict aggressive disease in african american men with prostate cancer. *J Clin Oncol* 2015;33:2789–2796.
47. Zhang H, Messing EM, Travis LB, et al. Age and racial differences among PSA-detected (AJCC stage T1cN0M0) prostate cancer in the U.S.: a population-based study of 70,345 men. *Front Oncol* 2013;3:312.
48. Andriole GL, Crawford ED, Grubb RL III, et al. Mortality results from a randomized prostate-cancer screening trial. *N Engl J Med* 2009;360:1310–1319.
49. Robbins HA, Engels EA, Pfeiffer RM, Shiels MS. Age at cancer diagnosis for blacks compared with whites in the United States. *J Natl Cancer Inst* 2015;107: pii: dju489.
50. Karami S, Young HA, Henson DE. Earlier age at diagnosis: another dimension in cancer disparity? *Cancer Detect Prev* 2007;31:29–34.
51. Metcalfe C, Evans S, Ibrahim F, et al. Pathways to diagnosis for black men and white men found to have prostate cancer: the PROCESS cohort study. *Br J Cancer* 2008;99:1040–1045.
52. Vertosick EA, Poon BY, Vickers AJ. Relative value of race, family history and prostate specific antigen as indications for early initiation of prostate cancer screening. *J Urol* 2014;192:724–728.
53. Cancer risks in BRCA2 mutation carriers. The Breast Cancer Linkage Consortium. *J Natl Cancer Inst* 1999;91:1310–1316.
54. Agalliu I, Gern R, Leanza S, Burk RD. Associations of high-grade prostate cancer with BRCA1 and BRCA2 founder mutations. *Clin Cancer Res* 2009;15:1112–1120.
55. Ford D, Easton DF, Bishop DT, et al. Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Lancet* 1994;343:692–695.
56. Gallagher DJ, Gaudet MM, Pal P, et al. Germline BRCA mutations denote a clinicopathologic subset of prostate cancer. *Clin Cancer Res* 2010;16:2115–2121.
57. Kirchoff T, Kauff ND, Mitra N, et al. BRCA mutations and risk of prostate cancer in Ashkenazi Jews. *Clin Cancer Res* 2004;10:2918–2921.
58. Leongamornlert D, Mahmud N, Tymrakiewicz M, et al. Germline BRCA1 mutations increase prostate cancer risk. *Br J Cancer* 2012;106:1697–1701.
59. Liede A, Karlan BY, Narod SA. Cancer risks for male carriers of germline mutations in BRCA1 or BRCA2: a review of the literature. *J Clin Oncol* 2004;22:735–742.
60. Tulinius H, Olafsdottir GH, Sigvaldason H, et al. The effect of a single BRCA2 mutation on cancer in Iceland. *J Med Genet* 2002;39:457–462.
61. van Asperen CJ, Brohet RM, Meijers-Heijboer EJ, et al. Cancer risks in BRCA2 families: estimates for sites other than breast and ovary. *J Med Genet* 2005;42:711–719.
62. Mersch J, Jackson MA, Park M, et al. Cancers associated with BRCA1 and BRCA2 mutations other than breast and ovarian. *Cancer* 2015;121:269–275.
63. Moran A, O'Hara C, Khan S, et al. Risk of cancer other than breast or ovarian in individuals with BRCA1 and BRCA2 mutations. *Fam Cancer* 2012;11:235–242.
64. Castro E, Goh C, Olmos D, et al. Germline BRCA mutations are associated with higher risk of nodal involvement, distant metastasis, and poor survival outcomes in prostate cancer. *J Clin Oncol* 2013;31:1748–1757.
65. Mitra A, Fisher C, Foster CS, et al. Prostate cancer in male BRCA1 and BRCA2 mutation carriers has a more aggressive phenotype. *Br J Cancer* 2008;98:502–507.
66. Narod SA, Neuhausen S, Vichodez G, et al. Rapid progression of prostate cancer in men with a BRCA2 mutation. *Br J Cancer* 2008;99:371–374.
67. Thorne H, Willems AJ, Niedermayr E, et al. Decreased prostate cancer-specific survival of men with BRCA2 mutations from multiple breast cancer families. *Cancer Prev Res (Phila)* 2011;4:1002–1010.
68. Tryggvadottir L, Vidarsdottir L, Thorgeirsson T, et al. Prostate cancer progression and survival in BRCA2 mutation carriers. *J Natl Cancer Inst* 2007;99:929–935.
69. Daly MB, Pilarski R, Axilbund JE, et al. NCCN Clinical Practice Guidelines in Oncology for Genetic/Familial High-Risk Assessment: Breast and Ovarian. Version 2.2016. Accessed February 12, 2016. To view the most recent version of these guidelines, visit NCCN.org.
70. Bancroft EK, Page EC, Castro E, et al. Targeted prostate cancer screening in BRCA1 and BRCA2 mutation carriers: results from the initial screening round of the IMPACT study. *Eur Urol* 2014;66:489–499.

Instructions for Completion

To participate in this journal CE activity: 1) review the learning objectives and author disclosures; 2) study the education content; 3) take the posttest with a 66% minimum passing score and complete the evaluation at <http://education.nccn.org/node/78544>; and 4) view/print certificate. After reading the article, you should be able to answer the following multiple-

choice questions. Credit cannot be obtained for tests completed on paper. You must be a registered user on NCCN.org. If you are not registered on NCCN.org, click on “New Member? Sign up here” link on the left hand side of the Web site to register. Only one answer is correct for each question. Once you successfully answer all posttest questions you will be able to view and/or print your certificate. Software requirements: Internet

Posttest Questions

1. According to the 2016 NCCN Guidelines for Prostate Cancer Early Detection, which of the following techniques should NOT be used when deciding whether to proceed with first-time biopsy for men with PSA levels >3 ng/mL?
 - a. Multiparametric MRI
 - b. Percent free PSA
 - c. 4Kscore
 - d. Prostate Health Index (PHI)

2. Targeted biopsies may
 - a. Improve detection of clinically significant prostate cancer
 - b. Minimize the detection of indolent disease
 - c. Both a and b
3. True or False: African American men, men with a first degree relative with prostate cancer, and *BRCA1/2* mutation carriers have an increased risk of prostate cancer compared with the general population.

