



OPEN

Protective effect of snail secretion filtrate against ethanol-induced gastric ulcer in mice

Enrico Gugliandolo^{3,4}, Marika Cordaro^{1,4}, Roberta Fusco^{1,4}, Alessio Filippo Peritore¹, Rosalba Siracusa¹, Tiziana Genovese¹, Ramona D'Amico¹, Daniela Impellizzeri¹, Rosanna Di Paola¹✉, Salvatore Cuzzocrea^{1,2}✉ & Rosalia Crupi³

Gastric ulcer or peptic ulcer is a common disease worldwide. Basically, it develops when there is an imbalance between the protective and aggressive factors, especially at the luminal surface of epithelial cells. Thus, there is a constant interest in research new drugs for treatment of gastric ulcer. The snail secretion is a dense mucous, that covers the external surface of the snails, with important functions for the survival of snails. The biological proprieties of snail *Helix Aspersa Muller* mucus it has been known for centuries to treat human disorders in particular for skin disease. Recently the use of snail mucus has seen a worldwide increase, as a component in cosmetic product and it has been used in particular for the management of wound and skin disorders. In this study we use a murine model of ethanol intragastric administration which has been widely used to test the drugs efficacies and to explore the underlying mechanism for gastric ulcer development. The intragastric ethanol administration causes several mucosal damages and an induction of a severe inflammatory response. Our results show a significant protective effect of snail secretion filtrate in reducing macroscopic and histological lesions, as well the protective effect on mucus content, oxidative stress and inflammatory response. In conclusion this study demonstrate the protective effect of intragastrical snail secretion filtrate, in a model of ethanol-induced gastric ulcer in mice, suggesting its possible useful use in the treatment or prevention of gastric ulcer.

The gastric ulcer or peptic ulcer is a disease that affect a significative number of people worldwide. Basically, it develops when there is an imbalance between the protective and aggressive factors at the luminal surface of epithelial cells. The most common factors contributes to the development of gastric ulcer are the *Helicobacter pylori* infection, long term use of aspirin or other NSAID, and other factors such as alcohol drinking, smoking and dietary habits¹⁻³. The causes mentioned above cause an imbalance in normal mucosal barrier. In fact, normally there is a layer of adherent mucus on gastric luminal surface. Furthermore, the gastric mucus act as protective layer for the underlying epithelium, from gastric juice and pepsin. The gastric mucus content is normally balanced by the secretion of new mucus to maintain a continuous barrier, against the continuous action of gastric juice and pepsin⁴. For the treatment of gastric ulcer several drugs are available, the common approach is based on the use of H receptor antagonism and proton pump inhibitors, or the use of drugs with mechanical protective action on gastric mucosa⁵. However a prolonged use of these drugs may cause serious adverse effect⁶. Thus, there is a constant interest in research in new drugs for treatment of gastric ulcer, with a particular interest in substances of natural origin. The snail secretion or mucus is a dense mucous that covers the external surface of the snail. This mucus is produced by a particularly salivary epidermal glands in snail. The mucus has various function for the life of snails with his adhesive, emollient, protective and reparative proprieties⁷. The *Helix Aspersa Muller* mucus, is still not well characterized composition⁷. In fact, the bioactive substances present in this peculiar natural product, make it a unique product not replicable in the laboratory and with synthetic compounds. The biological proprieties of *Helix Aspersa Muller* snail mucus it has been known for centuries to treat human disorders in particular for skin disease. Recently the use of snail mucus has seen a worldwide increase, as a component in cosmetic product and it has been used in particular for the management of wound and treatment of chronic

¹Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Via F. Stagno D'alcontres 31, 98166 Messina, Italy. ²School of Medicine, 1402 South Grand Blvd, St Louis, MO 63104, USA. ³Department of Veterinary Science, University of Messina, Viale Annunziata, 98168 Messina, Italy. ⁴These authors contributed equally: Enrico Gugliandolo, Marika Cordaro and Roberta Fusco. ✉email: dipaolar@unime.it; salvator@unime.it

bronchitis^{8,9}. It has been seen that the snail mucus components are able to stimulate the formation of dermal components, in particular the formation of collagen and elastin, and to minimize the damage generated by oxidative stress and free radicals¹⁰. It has shown to possess also other important biological properties such as antimicrobial activity and protective effect in wound repair¹¹. In fact, the effectiveness of *Helix Aspersa* extract has been demonstrated as a safe and effective alternative treatment in open wound management of partial thickness burns in adults, favoring its re-epithelialization⁹. These important properties of snail slime and in particular pro-epithelizing and wound repair are due to the peculiar chemical composition of this compound¹². Moreover, many of the components present in snail slime play a fundamental role in homeostasis and protection of the gastric mucosa. In particular a protective effect in the gastric mucosa has already been demonstrated for some of the components most present in the Snail Secretion Filtrate (SSF) such as, collagen, elastin, glycolic acid and allantoin^{13,14}. However, there are numerous other compounds present in snail slime that contribute to its peculiar protective action, such as the mineral component¹⁵, due for example to the presence of copper in the SSF. In fact, copper has been shown to have an anti-ulcer action¹⁶. Finally, also the mucopolysaccharide component naturally present in snail slime plays a key role in the protective activity on mucous tissues^{17,18}. Therefore, on the basis of the known properties of snail slime and the pathophysiology of gastric ulcer, the hypothesis of this study is that oral treatment with SSF may be a useful tool in maintaining homeostasis of the gastric mucosa, and in the prevention or treatment of inflammation of the gastric mucosa and therefore of the gastric ulcer. The murine model of intragastric ethanol administration has been widely used to test the drugs efficacies and to explore the underlying mechanism for gastric ulcer development, the intragastric ethanol administration causes several mucosal damage and an induction of a severe inflammatory response^{19,20}. Alcohol also induce an important increase in oxidative stress by the generation of reactive oxygen species and lipid peroxidation, the oxidative stress play a key role in gastric ulcer disease²¹. In this study for the first time we study the potentially protective effect of Snail Secretion Filtrate in an experimental model of ethanol induced gastric ulcer in mice. Evaluating the effect on gastric mucosal, oxidative stress and inflammatory response.

Results

Snail secretion filtrate (SSF) chemical characterization. Figure 1 panel a, reports the qualitative and quantitative properties and composition of SSF. The SSF used in this study showed a high content in glycolic acid and collagen followed by allantoin and elastin as showed by the chemical characterization showed in Fig. 1 and supplementary figure S1-S8. The dose of SSF used in this experimental model of ethanol-induced acute ulcer in mice was calculated based on maximum volume in ml/kg that can be given by o.s. in mice, so as to gradually obtain a greater surface of gastric mucosa in contact with SSF. In particular were chosen as a high dose 15 ml/kg, medium 7.5 ml/kg and low 3 ml/kg. The total protein content in SSF was 25,6 mg/ml. The Sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) profiles of proteins pattern of SSF from *Helix Aspersa Muller* are shown in Supplementary Figure S1.

Effect of SSF on macroscopic examination. One hour after EtOH gastric ulcer induction, as shown in Fig. 2 the stomachs from EtOH group (the negative control group) showed a significant presence in mucosa hyperemia and mucosal damage with large ulcer formation. Instead the treatment with omeprazole (used as positive control) at dose of 20 mg/kg significantly prevented the ulceration and mucosal damage induced by EtOH. The treatment with SSF showed a dose dependent protective effect as shown in Fig. 2 by the graphs of gastric ulcer index and the preventive index. In particular the dose of 15 ml/kg and 7.5 ml/kg showed an important reduction in mucosal damage and ulceration compared to EtOH group.

Effect of SSF on histological damage. As shown in Fig. 3a, compared to stomach tissue from control group the EtOH group showed a significant increase in tissue damage characterized by a significant epithelial cell loss, edema, hemorrhagic damage and abundant presence of inflammatory cells. The treatment with SSF showed a dose dependent protective effect in particular with a significant protective effect on epithelial cell loss. Lower magnification of histological images on both upper and lower mucosa are shown in supplementary figure S8-S13. The presence of inflammatory cells has been confirmed through MPO assay that showed a significant increase in MPO in EtOH group compared to control group. As shown in Fig. 3b, the treatment with omeprazole significantly prevented the increase in MPO, for the SSF treatment only the dose of 7.5 ml/kg and 15 ml/kg showed a significant protective effect on increase in MPO induced by EtOH.

Effect of SSF on mucosa glycoproteins and collagen contents. Mucosa glycoprotein evaluation was performed using PAS staining. As shown in Fig. 4, one hour after the EtOH administration, in the stomach from mice of EtOH group there was a significant decrease in PAS staining compared to sham group. The loss in mucosa glycoprotein was significantly reduced by pre-treatment with omeprazole. Compared to EtOH group the treatment with SSF at dose of 3 ml/kg did not show any notable protective effect, instead the dose of 7.5 and 15 ml/kg of SSF showed a significant protective effect in a dose dependent manner compared to the EtOH group. Through Masson staining collagen can be dyed in blue, by which we can roughly evaluate the healing effect at the macro level and stains also showed the formation of granulation tissue and collagen disorganization in the gastric wall. As shown in Fig. 5, in the EtOH group stain indicated fragmented and disorganized collagen fibers, while omeprazole group showed a significant protective effect. For the SSF treatment only the dose of 15 ml/kg showed a significant protective effect.

Effect of SSF on mucus contents and oxidative stress. The mucus contents assay was performed by alcian blue binding assay. As shown by the graph in Fig. 6a, the mucus contents was significantly reduced in the

Specification	Values	Measure unit
Aspect	Clear fluid	
Color	slightly amber	
pH	4.80	
Refractive index	1,3405	
Density	1,02	g\ml
Dry residual	3	g\100g
Protein	1,54	g\100g
Gram+	0	CFU
Gram-	0	CFU
Fungi	0	CFU
Vitamin E	<0,10	mg\Kg
Vitamin C	0,13	mg\Kg
Allantoin	0,081	g\100g
Glicolic acid	0,99	g\100g
Hyaluronic acid	<0,1	g\100g
Elastin	0,092	g\100g
Collagen	0,32	g\100g
All-trans-retinol	<10	μg\100g
13-cis-retinol	<10	μg\100g
Vitamin B12	<10	μg\100g
Vitamin B3	<10	μg\100g
Minerals		
Cr	0,007	mg\Kg
Cu	5,04	mg\Kg
Hg	0,22	μg\Kg
Cd	0,012	μg\Kg
Co	<0,001	μg\Kg
Ni	0,820	μg\Kg

Figure 1. Qualitative and quantitative analysis of crude SSF. Full chemical characterization are shown in supplementary figure S1-S7.

EtOH group compared to the control group. When compared to EtOH group, the omeprazole group showed a significant decrease in mucus contents loss, the treatment with SSF showed a dose dependent protective effect on mucus contents loss due to EtOH administration. To assess the effect of on SSF on oxidative stress we evaluated the levels of MDA as index of lipid peroxidation, as show in Fig. 6b intragastrical EtOH administration induced a significant increase MDA levels, while the treatment with omeprazole prevent this increase in MDA in a significant manner, the treatment with SSF showed a dose dependent protective effect on mucus. Next we observed a significant reduction in CAT and SOD induced by EtOH administration that has been significantly antagonized by omeprazole pre-treatment. As show in Fig. 6c,d treatment with SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose dependent manner.

Effect of SSF on PGE₂ and inflammatory response. As show in Fig. 7a, through ELISA assay we evaluated the levels PGE₂ in gastric tissue homogenates, compared to control group EtOH administration induced a significant decrease in PGE₂, significantly antagonized by omeprazole pre-treatment. The treatment with SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose dependent manner. Figure 7b–d, showed the results for the ELISA assays for mayor pro inflammatory cytokines IL-6, IL-1β, TNF-α, respectively. EtOH administration induced a significant increase in IL-6, IL-1β, TNF-α levels significantly antagonized by omeprazole pre-treatment. The treatment SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose dependent manner.




Figure 2. The macroscopic picture of stomach from different experimental groups. stomachs from EtOH group (the negative control group) showed a significant presence in mucosa hyperemia and mucosal damage with large ulcer formation, instead the treatment with omeprazole at dose of 20 mg/kg significant prevent the ulceration and mucosal damage induced by EtOH. The treatment with SSF showed a dose dependent protective effect. Data are presented as means \pm SEM, or median with interquartile range for non-parametric data of 10 mice for each group. *** $p < 0.001$ versus control; ° $p < 0.05$ versus EtOH; °°° $p < 0.001$ versus EtOH; ### $p < 0.001$ versus omeprazole.

Discussion

Several factors contributes to gastric ulcer development, such as drug abuse, alcoholism, endocrine dyscrasia and bacterial helicobacter pylori infection, which can cause mucosal barrier damage through alteration in gastric acid secretion and characterized by hemorrhage and severe inflammatory response^{22,23}. A key event in gastric ulcer formation is the imbalance in the gastric mucosa between the defensive factor such as mucin, prostaglandin, bicarbonate nitric oxide and growth factor, and the offensive factor such as increased secretion of pepsin and gastric acid. Thus inflammatory response, oxidative stress, and neutrophilic infiltration has been show a key role in pathophysiology of gastric ulcer, on the other hand endogenous antioxidant, mucus layer acts as protective agents^{24,25}. The common approach in treating the gastric ulcer is based on H receptor antagonism and proton pump inhibitors⁵. These class of drugs may cause serious adverse effect when used for prolonged time⁶. Recently, several studies has highlighted the ability of snail secretion (or snail mucus), to improve skin conditions thanks




Figure 3. Stomach tissue section from: Control group healthy mice; EtOH group negative control; omeprazole 20 mg/kg positive control group; Histological section was assessed for: a) epithelial cell loss; b) edema in the upper mucosa; c) hemorrhagic damage; d) presence of inflammatory cells. The treatment with SSF showed a dose dependent protective effect SSF 3 ml/kg by os; SSF 7.5 ml/kg by os; SSF 15 ml/kg by os. Panel b showed the results of MPO assay. Data are presented as means \pm SEM, or median with interquartile range for non-parametric data of 10 mice for each group. *** $p < 0.001$ versus control; ° $p < 0.05$ versus EtOH; °° $p < 0.01$ versus EtOH; °°° $p < 0.001$ versus EtOH.

to its peculiar proprieties such as emollient, moisturizing, lubricating and protective, especially to prevent skin damage, and thus there is a growing interest in the use of this compound in cosmetology²⁶. Furthermore the *Helix Aspersa Muller* secretion has shown to possess other important biological properties such as, antimicrobial activity¹¹ and wound repair²⁷. Several studies have highlighted the absence of cytotoxic effects in different cell lines such as in human keratinocytes, human dermal fibroblasts (MRC-5) and murine embryo fibroblasts (NIH-3T3)^{12,27}. These effect of Snail mucus is due also to his peculiar mechanical proprieties due mucopolysaccharide contents, on the other hand these effects are due to the peculiar contents of active molecules¹². As previously demonstrated, the two molecules the glycolic acid and allantoin, most present in snail slime were believed to be an essential component for the biological activities of snail slime⁹, however recently has been demonstrated that the mucus *in toto* has a greater effect than that of the individual molecules²⁷. Thus, a synergism in activity of several molecules present in snail secretion cannot be excluded, factor that make a snail secretion a peculiar compound with important biological proprieties. In fact, beyond all molecules contained in snail secretion also the specific ratio of these components in the natural snail secretion is a key factor for the biological activity. So, in this study we consider the crude Snail Secretion Filtrate (SSF) form *Helix Aspersa Muller* as compound “*in toto*” and it is not possible to understand which one the molecule is responsible for the biological activity. In this




Figure 4. PAS staining from control group showed the normal contents of mucosa glycoprotein (magenta color); EtOH group showed a significant reduction in PAS staining, significantly inhibited by treatment with omeprazole 20 mg/kg; SSF treatment showed a dose-dependent protective effect as shown by the increase of PAS staining. Image J software (1.49v, <https://imagej.nih.gov/ij/>, 1997–2018. Schneider, C.A., Rasband, W.S., Eliceiri, K.W.) was used for the quantification of glycoprotein as the positively stained area (pixel/field). Data are presented as means \pm SEM of 10 mice for each group. *** $p < 0.001$ versus control; °°° $p < 0.001$ versus EtOH.

study due to the route of administration and the maximum volume in ml/kg that can be given by o.s. in mice²⁸, we consider three doses of SSF as high dose of 15 ml/kg, medium of 7.5 ml/kg and low 3 ml/kg, to gradually obtain a greater surface of gastric mucosa in contact with SSF²¹. Considering the protein content in the SSF the respective dosage as mg/kg were respectively 384 mg/kg, 192 mg/kg and 76.8 mg/kg. We choose an experimental model of ethanol intragastric administration in mice, that has been widely used to explore the underlying mechanism of gastric ulcer development and thus to test the efficacy of new drugs. The intragastric administration of ethanol causes several mucosa damages and an induction of a severe inflammatory response and oxidative stress^{19–21}. In our study, the results of macroscopic gastric evaluation after EtOH administration showed a significant protective effect of SSF on gastric mucosa. In particular, the treatment with SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect, instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose-dependent manner. Also, on histological evaluation of gastric mucosa we confirmed the protective effect of SSF, in agreement with macroscopic scores. A fundamental step involved in the complex event of gastric ulcer formation is the infiltration of neutrophils, in particular activated neutrophils are responsible for an increase in oxidative stress and inflammatory response^{29,30}. Our results on MPO evaluation as an index of inflammatory cell infiltration, showed that doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose-dependent manner on neutrophil infiltration. A standard goal in treating gastric ulcers is preventing the decrease or degradation of gastric mucus, that acts as a natural protective agent on gastric mucosa³¹. Gastric mucus consists of mucin-type glycoproteins and the depletion in gastric wall mucus has been seen to be closely related to the gastric ulcer pathology^{32,33}. EtOH-induced gastric ulcers are also responsible for the reduction of collagen within gastric tissue³⁴. Therefore, we evaluated both the presence of glycoproteins (PAS staining) and collagen (Masson




Figure 5. Representative image of collagen within mucosa layer using Masson's trichrome staining. Control group showed the staining in healthy gastric tissue. EtOH group stain indicated fragmented and disorganized collagen fibers, while omeprazole group showed a significant protective effect. For the SSF treatment only the dose of 15 ml/Kg showed a significant protective effect.

staining) following the administration of EtOH, and our results show that this has produced a significant reduction in both cases, while the treatment with SSF at doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect. These results are also confirmed by the results of mucus content performed by Alcian blue staining. According to the results mentioned so far we have found that the treatment with SSF at doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect on oxidative stress induced by EtOH, and in particular SSF treatment significantly prevent the increase in MDA levels, and the depletion in CAT and SOD levels. Oxidative stress in related to both initial stage of gastric ulcer development and with the worsening of the pathology³⁵. Has previously demonstrated the gastric mucosa homeostasis is regulated by both the control of oxidative stress status and by secretion of several mediators such as cytokines and prostaglandin. In particular, prostaglandin play a key role on the state of health of the mucosa by regulating several factors such as the stimulation of mucus and bicarbonate secretion, maintaining the integrity of epithelial cells and improving mucosal mucosal blood flow, in particular PGE₂ showed an important role in regulation of these function, and thus PGE₂ act as a gastro protective agent^{36,37}. Gastric ulcer and an gastric ulcer healing has been seen previously are regulated by the levels of prostaglandin an PGE₂ in particular³⁸. Our results indicate that gastric ulcer induced by EtOH is related with a significant reduction on PGE₂ levels, the treatment SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose dependent manner. Inflammatory response is responsible for the progressive trigger and worsening of the gastric ulcer³⁹. Then we evaluated the levels of the mayor inflammatory cytokines responsible for driving the inflammatory response in gastric ulcer such as IL-1 β , IL-6 and TNF- α ⁴⁰. Ours results showed a significant increased levels of these cytokines in EtOH group while the treatment SSF at the dose of 3 ml/kg did not demonstrate a significant protective effect instead the doses of 7.5 ml/kg and 15 ml/kg showed a significant protective effect in a dose dependent manner in reducing the IL-1 β , IL-6 and TNF- α levels. In conclusion this study demonstrate for the first time the protective effect of intragastrical snail secretion filtrate, in a model of ethanol-induced gastric ulcer in mice. The gastric cytoprotective action of snail secretion filtrate might be attributed to an increase or preservation of gastric mucus with preservation of mucosal mucopolysaccharides and collagen that which are also reflected in a reduction of oxidative stress and induced inflammatory response. surely future studies are necessary to clarify themechanism of action of this peculiar compound, however the proven beneficial effects suggest its possible useful use in the treatment or prevention of gastric ulcer.

Methods

Snail secretion filtrate (SSF) collection and sterilization. *Helix aspersa muller* mucus was kindly provided by Snail S.R.L.S (Messina, Italy). Briefly the breeding is cruelty free. In particular the mucus was obtained mechanically manually by stimulating snails by sterile cotton swab tip. The mucus is filtered in a first step with a coarse filter stabilizing the pH, after this phase the mucus is passed in a filtration train consisting of 3




Figure 6. (a) The graph showed the mucus contents evaluation performed by alcian blue binding assay. (b) MDA assay levels. (c, d) CAT and SOD determination. Data are presented as means \pm SEM of 10 mice for each group; *** $p < 0.001$ versus control; ^{oo} $p < 0.01$ versus EtOH; ^{ooo} $p < 0.001$ versus EtOH;

different filters (10 micron-1micron -0.22-micron, Pall) and then stored at 4 °C. The use of the 0.22 micron filter is necessary to eliminate impurities and endotoxins in order to make the product injectable.

Snail secretion filtrate (SSF) chemical characterization. The crude Snail Secretion Filtrate extract from different batches, was qualitative and quantitative analyzed by professional service for Snail S.R.L.S (Messina, Italy), (Science4Life srl, Messina Italy. and Sialab srl. Avola, Italy) using standard analytical techniques such as IR/UV vis spectrometry and HPLC analysis to evaluate the protein quality and the allantoin and glycolic acid content respectively, and to obtain a qualitative determination of the total protein content and to a Bradford assay (Bio-Rad) to evaluate the protein amount. Elastin and collagen were analyzed as seen previously^{41,42}. All-trans-retinol, 13-cis-retinol, Vitamin B12, Vitamin B3 were analyzed according EN 12,823-1:2014, ISO20634:2015 and UNI EN 15,652:2009 respectively. All slime doses in experiments were subsequently calculated on microgram of proteins per milliliter (mg/mL) content.

Animals. CD1 mice (male, 20–30 g; Envigo) were accommodated in a standard location (room 22 °C and 12-h light/dark cycles) with standard rodent chow and water ad libitum. The animals were adapted to these conditions for 1 week. Messina University Review Board for the care of animals approved the research. All animal




Figure 7. ELISA assays for (a) PGE₂; (b) IL-6; (c) IL-1β; (d) TNF-α. Data are presented as means ± SEM of 10 mice for each group; ***p < 0.001 versus control; °°p < 0.01 versus EtOH; °°°p < 0.001 versus EtOH;

experiments agree with the new regulations in Italy (D.Lgs 2014/26), EU regulations (EU Directive 2010/63). All experimental were conducted in according to ARRIVE guidelines.

Ethanol-induced acute ulcer. Ethanol-induced acute ulcer in mice were performed as seen previously^{21,43}. Briefly the mice were fasted for 24 h and divided into the follow group:

- Control (n = 10): mice were treated by oral gavage with distilled water (15 ml/Kg)
- EtOH (n = 10): mice were treated by oral gavage with distilled water (15 ml/Kg) 1 h before ethanol-induced acute ulcer
- Omeprazole 20 mg/Kg: mice were treated by oral gavage with omeprazole at 20 mg/kg 1 h before ethanol-induced acute ulcer
- SSF 3 ml/Kg: mice were treated by oral gavage with Snail Secretion Filtrate (SSF) at 3 ml/kg 1 h before ethanol-induced acute ulcer
- SSF 7.5 ml/kg: mice were treated by oral gavage with Snail Secretion Filtrate (SSF) at 7.5 before ethanol-induced acute ulcer
- SSF 15 ml/Kg: mice were treated by oral gavage with Snail Secretion Filtrate (SSF) at 15 ml/kg 1 h before ethanol-induced acute ulcer

One hour after the oral treatment, acute ulcer was induced by orally EtOH (98% ethanol containing 150 mM HCl) administration at 5 ml/kg. Control mice were treated with an equal volume of distilled water instead of EtOH solution. Animals were euthanized 1 h after treatment with the EtOH mixture, under isoflurane inhalation anesthesia (5% in air. Baxter) by cervical dislocation. In this study due to the route of administration and maximum volume in ml/kg that can be given by o.s. in mice^{21,28}, we consider three dose of Snail Secretion Filtration (SSF) as high dose of 15 ml/kg, medium of 7.5 ml/kg and low 3 ml/kg, consider the proteins content in the SSF the respective dosage as mg/kg were respectively 384 mg/kg, 192 mg/kg and 76.8 mg/kg.

The gastric ulcer index and preventive index. After sacrificing the animals, the stomachs were quickly removed and washed with 0.9% saline. This was followed by macroscopic examination of the stomach for the detection of any hemorrhagic lesions on the glandular mucosa. The length in mm of each lesion was measured to determine the mean ulcer index (UI)⁴⁴. The length (mm) and the width (mm) of each band was measured by Vernier caliper. The degree of gastric mucosal lesions was evaluated from digital pictures, and the severity of mucosal lesions was scored as follows: no ulcer (0 mm), 1–5 petechiae (<1 mm) (1), 6–10 petechiae (<1 mm) (2), >10 petechiae (<1 mm) (3), small linear ulcer (<2 mm) (2), medium linear ulcer (2–4 mm) (3), and large linear ulcer (>4 mm) (4). If the width was >1 mm, then the points were multiplied by two. The GU index (UI) was determined by adding the sum of the total of the scores and divided by the number of animals. The preventive index (PI) of pretreatments against ulceration was calculated according to the following equation:

$$PI = (UI_{\text{ethanol}} - UI_{\text{pretreated}}) / UI_{\text{ethanol}} \times 100.$$

Histological analysis. For histological assessment, the stomach was fixed in 10% (V/V) neutral buffered formalin solution, and then embedded in paraffin, sliced into 5 µm thicknesses, stained with hematoxylin-eosin (H&E)⁴⁵. The specimens were examined under an optical microscope (DM5500, Leica), and were assessed according to what has been previously⁴⁶. Histological section was assessed for epithelial cell loss (score: 0–3), edema in the upper mucosa (score: 0–4), hemorrhagic damage (score: 0–4), and the presence of inflammatory cells (score: 0–3), yielding a maximum total score of 14. The sections were assessed by an experienced pathologist who was blinded to the study.

Measurement of mucosa glycoproteins and collagen. To further evaluate the mucosal lesions in the gastric tissues, we performed mucosa glycoprotein measurements with periodic acid-Schiff (PAS) staining according to the manufacturer's instructions (Bio-optica, Italy). The positive glycoprotein site will appear as magenta color. Image J software was used for the determination of the positively stained area (pixel/field)^{47,48}. Masson trichrome staining were performed according to the manufacturer's protocol (Bio-Optica, Milan, Italy).

Measurement of gastric mucus contents. Alcian blue (Sigma-Aldrich) binding assay was performed according to what seen previously⁴⁹. After removing the stomach, some parts of the stomach were weighted and immersed in 10 mL of 0.02% Alcian blue and 0.16 M sucrose/0.05 M sodium acetate solution (pH 5.8) and incubated at 25 °C for 24 h. The Alcian blue binding extract was centrifuged at 3000 × g for 10 min at 4 °C. The absorbance of the supernatant was measured at 620 nm on a spectrophotometer. Free mucus in the gastric content was calculated based on the amount of Alcian blue binding to the gastric mucus (mg/g of tissue).

Myeloperoxidase activity. Gastric tissues, were homogenized and MPO activity was detected^{50,51}.

Determination of malondialdehyde (MDA) level. Thiobarbituric acid-reactant substances measurement was determined as a marker of lipid peroxidation. Thiobarbituric acid-reactant substances were calculated by comparison the O.D. to a standard mixture of 1,1,3,3-tetramethoxypropan/99% malondialdehyde bis (dimethylacetal)/99% (MDA) (Sigma, Milan, Italy). The absorbance of the supernatant was measured spectrophotometrically at 532 nm as seen previously⁵².

Determination of oxidative stress markers and biochemical determinations. SOD activity in the gastric tissue was determined using the Superoxide Dismutase Assay Kit (Cayman Chemicals) according to manufacturer's protocols. CAT activity in the gastric tissue was determined using the Catalase Assay Kit (Cayman Chemicals) according to manufacturer's protocols. The supernatant of stomach tissue homogenate was subjected to measurement of PGE2 level using an ELISA kit (Cayman chemicals) according to manufacturer protocols. IL-1β, IL-6, TNF-α were measured using ELISA kit (Invitrogen, Thermo Fisher) according to manufacturer protocols.

Analysis of protein pattern. Analysis of protein pattern were performed as previously seen⁵³. Total protein concentrations in extracts were determined by the Bradford method with bovine serum albumin (BSA) as the standard. Then, SDS-PAGE gel electrophoresis was performed on samples with equalized concentrations of total protein. In Particular 20 µl of samples was mixed with 20 µl of Laemmli Sample Buffer with β-mercaptoethanol (Bio-Rad) and heating (95 °C, 5 min). then 40 µl of mixed samples and 5 µl of protein marker (Precision Plus Protein Dual Color Standards, Bio-Rad) were loaded onto the gel and resolved, using the Mini-PROTEAN electrophoresis system (Bio-Rad). The protein bands, separated on gel, were fixed, stained in QC Colloidal Coomassie Stain and destained solution according to the producer procedure (Bio-Rad).

Statistical analysis. All values are shown as the mean \pm standard error of the mean (SEM) of N observations (N = 10) or median with interquartile range for non-parametric data. Data were analyzed by Non-parametric Kruskal–Wallis one-way ANOVA followed by Dunn's multiple comparisons test, or one-way ANOVA followed by a Bonferroni post hoc test for multiple comparisons. A p-value of less than 0.05 was considered significant: *p < 0.05 versus control; \circ p < 0.05 versus EtOH; **p < 0.01 versus control; $\circ\circ$ p < 0.01 versus EtOH; ***p < 0.001 versus control; $\circ\circ\circ$ p < 0.001 versus EtOH; #p < 0.05 versus omeprazole; ##p < 0.01 versus omeprazole; ###p < 0.001 versus omeprazole.

Data availability

All data generated or analyzed during this study are included in this published article.

Received: 14 May 2020; Accepted: 29 January 2021

Published online: 11 February 2021

References

- Kusters, J. G., van Vliet, A. H. & Kuipers, E. J. Pathogenesis of *Helicobacter pylori* infection. *Clin. Microbiol. Rev.* **19**, 449–490. <https://doi.org/10.1128/CMR.00054-05> (2006).
- Li, L. F. *et al.* Cigarette smoking and gastrointestinal diseases: the causal relationship and underlying molecular mechanisms (review). *Int. J. Mol. Med.* **34**, 372–380. <https://doi.org/10.3892/ijmm.2014.1786> (2014).
- Ma, S. H. *et al.* Impact of alcohol drinking on gastric cancer development according to *Helicobacter pylori* infection status. *Br. J. Cancer* **113**, 1381–1388. <https://doi.org/10.1038/bjc.2015.333> (2015).
- Allen, A. *et al.* Pepsins and the mucus barrier in peptic ulcer disease. *Scand. J. Gastroenterol. Suppl.* **146**, 50–57. <https://doi.org/10.3109/00365528809099130> (1988).
- LiverTox: Clinical and Research Information on Drug-Induced Liver Injury* (2012).
- Thomson, A. B., Sauve, M. D., Kassam, N. & Kamitakahara, H. Safety of the long-term use of proton pump inhibitors. *World J. Gastroenterol.* **16**, 2323–2330. <https://doi.org/10.3748/wjg.v16.i19.2323> (2010).
- Greistorfer, S. *et al.* Snail mucus - glandular origin and composition in *Helix pomatia*. *Zoology (Jena)* **122**, 126–138. <https://doi.org/10.1016/j.zool.2017.05.001> (2017).
- Pons, F., Koenig, M., Michelot, R., Mayer, M. & Frossard, N. The bronchorelaxant effect of helicidine, a *Helix pomatia* extract, interferes with prostaglandin E2. *Pathol. Biol. (Paris)* **47**, 73–80 (1999).
- Tsoutsos, D., Kakagia, D. & Tamparopoulos, K. The efficacy of *Helix aspersa* Muller extract in the healing of partial thickness burns: a novel treatment for open burn management protocols. *J. Dermatol. Treat.* **20**, 219–222. <https://doi.org/10.1080/09546630802582037> (2009).
- Brieva, A. *et al.* Molecular basis for the regenerative properties of a secretion of the mollusk *Cryptomphalus aspersa*. *Skin Pharmacol. Physiol.* **21**, 15–22. <https://doi.org/10.1159/000109084> (2008).
- Bortolotti, D., Trapella, C., Bernardi, T. & Rizzo, R. Letter to the Editor: Antimicrobial properties of mucus from the brown garden snail *Helix aspersa*. *Br. J. Biomed. Sci.* **73**, 49–50. <https://doi.org/10.1080/09674845.2016.1155377> (2016).
- Genitli, V. *et al.* HelixComplex snail mucus as a potential technology against O3 induced skin damage. *PLoS ONE* **15**, e0229613. <https://doi.org/10.1371/journal.pone.0229613> (2020).
- Rusanova, A. V. *et al.* Thrombin receptor agonist Peptide immobilized in microspheres stimulates reparative processes in rats with gastric ulcer. *Bull. Exp. Biol. Med.* **142**, 35–38. <https://doi.org/10.1007/s10517-006-0285-y> (2006).
- da Silva, D. M. *et al.* Effect of allantoin on experimentally induced gastric ulcers: Pathways of gastroprotection. *Eur. J. Pharmacol.* **821**, 68–78. <https://doi.org/10.1016/j.ejphar.2017.12.052> (2018).
- Interdonato, M. *et al.* Levels of heavy metals in adolescents living in the industrialised area of Milazzo-Valle del Mela (northern Sicily). *J. Environ. Public Health* **2014**, 326845. <https://doi.org/10.1155/2014/326845> (2014).
- Alberghina, M. *et al.* Cytoprotective effect of copper(II) complexes against ethanol-induced damage to rat gastric mucosa. *J. Inorg. Biochem.* **45**, 245–259. [https://doi.org/10.1016/0162-0134\(92\)84013-d](https://doi.org/10.1016/0162-0134(92)84013-d) (1992).
- Bonnemain, B. Helix and drugs: snails for western health care from antiquity to the present. *Evid. Based Complem. Alternat. Med.* **2**, 25–28. <https://doi.org/10.1093/ecam/neh057> (2005).
- Nj, N., Okonta, M., Co, E. & Aa, A. Anti-ulcer potentials of phylum mollusca (tropical snail) slime. *Asian Pac. J. Trop. Med.* **2**, 23–28 (2009).
- Salama, S. M. *et al.* A zinc morpholine complex prevents HCl/ethanol-induced gastric ulcers in a rat model. *Sci. Rep.* **6**, 29646. <https://doi.org/10.1038/srep29646> (2016).
- Zhao, W. *et al.* Protective effects of DIDS against ethanol-induced gastric mucosal injury in rats. *Acta Biochim. Biophys. Sin. (Shanghai)* **41**, 301–308. <https://doi.org/10.1093/abbs/gmp014> (2009).
- Yoo, J. H., Lee, J. S., Lee, Y. S., Ku, S. & Lee, H. J. Protective effect of bovine milk against HCl and ethanol-induced gastric ulcer in mice. *J. Dairy Sci.* **101**, 3758–3770. <https://doi.org/10.3168/jds.2017-13872> (2018).
- Erin, N., Turker, S., Elpek, O. & Yildirir, B. ADAM proteases involved in inflammation are differentially altered in patients with gastritis or ulcer. *Exp. Ther. Med.* **15**, 1999–2005. <https://doi.org/10.3892/etm.2017.5619> (2018).
- Drini, M. Peptic ulcer disease and non-steroidal anti-inflammatory drugs. *Aust. Prescr.* **40**, 91–93. <https://doi.org/10.18773/austprescr.2017.037> (2017).
- Arab, H. H., Salama, S. A., Omar, H. A., el Arafa, S. & A. & Maghrabi, I. A. . Diosmin protects against ethanol-induced gastric injury in rats: novel anti-ulcer actions. *PLoS ONE* **10**, e0122417. <https://doi.org/10.1371/journal.pone.0122417> (2015).
- Thomas, D. *et al.* Cyperus rotundus L. prevents non-steroidal anti-inflammatory drug-induced gastric mucosal damage by inhibiting oxidative stress. *J. Basic Clin. Physiol. Pharmacol.* **26**, 485–490. <https://doi.org/10.1515/jbcpp-2014-0093> (2015).
- Lopez Angulo, D. E. & do Amaral Sobral, P. J. Characterization of gelatin/chitosan scaffold blended with aloe vera and snail mucus for biomedical purpose. *Int. J. Biol. Macromol.* **92**, 645–653. <https://doi.org/10.1016/j.ijbiomac.2016.07.029> (2016).
- Trapella, C. *et al.* HelixComplex snail mucus exhibits pro-survival, proliferative and pro-migration effects on mammalian fibroblasts. *Sci. Rep.* **8**, 17665. <https://doi.org/10.1038/s41598-018-35816-3> (2018).
- Hull, R. M. Guideline limit volumes for dosing animals in the preclinical stage of safety evaluation: Toxicology Subcommittee of the Association of the British Pharmaceutical Industry. *Hum. Exp. Toxicol.* **14**, 305–307. <https://doi.org/10.1177/096032719501400312> (1995).
- Rawlingson, A. *et al.* Functional significance of inducible nitric oxide synthase induction and protein nitration in the thermally injured cutaneous microvasculature. *Am. J. Pathol.* **162**, 1373–1380. [https://doi.org/10.1016/S0002-9440\(10\)63933-8](https://doi.org/10.1016/S0002-9440(10)63933-8) (2003).
- Lowe, P. P. *et al.* Correction: alcohol-related changes in the intestinal microbiome influence neutrophil infiltration, inflammation and steatosis in early alcoholic hepatitis in mice. *PLoS ONE* **12**, e0179070. <https://doi.org/10.1371/journal.pone.0179070> (2017).
- Lima, Z. P. *et al.* Effects of the ethyl acetate fraction of alchornea triplinervia on healing gastric ulcer in rats. *Pharmaceuticals (Basel)* **4**, 1423–1433. <https://doi.org/10.3390/ph4111423> (2011).

32. Koo, M. W., Ogle, C. W. & Cho, C. H. Effects of verapamil, carbenoxolone and N-acetylcysteine on gastric wall mucus and ulceration in stressed rats. *Pharmacology* **32**, 326–334. <https://doi.org/10.1159/000138188> (1986).
33. Indran, M., Mahmood, A. A. & Kuppusamy, U. R. Protective effect of *Carica papaya* L leaf extract against alcohol induced acute gastric damage and blood oxidative stress in rats. *West Indian Med. J.* **57**, 323–326 (2008).
34. Liu, W. *et al.* Gastroprotective effects of chebulagic acid against ethanol-induced gastric injury in rats. *Chem. Biol. Interact.* **278**, 1–8. <https://doi.org/10.1016/j.cbi.2017.09.019> (2017).
35. Tandon, R., Khanna, H. D., Dorababu, M. & Goel, R. K. Oxidative stress and antioxidants status in peptic ulcer and gastric carcinoma. *Indian J. Physiol. Pharmacol.* **48**, 115–118 (2004).
36. Wallace, J. L. Prostaglandins, NSAIDs, and gastric mucosal protection: why doesn't the stomach digest itself?. *Physiol. Rev.* **88**, 1547–1565. <https://doi.org/10.1152/physrev.00004.2008> (2008).
37. Yu, C., Mei, X. T., Zheng, Y. P. & Xu, D. H. Gastroprotective effect of taurine zinc solid dispersions against absolute ethanol-induced gastric lesions is mediated by enhancement of antioxidant activity and endogenous PGE2 production and attenuation of NO production. *Eur. J. Pharmacol.* **740**, 329–336. <https://doi.org/10.1016/j.ejphar.2014.07.014> (2014).
38. Konturek, S. J., Konturek, P. C. & Brzozowski, T. Prostaglandins and ulcer healing. *J. Physiol. Pharmacol.* **56**(Suppl 5), 5–31 (2005).
39. Sipponen, P. & Maaroo, H. I. Chronic gastritis. *Scand. J. Gastroenterol.* **50**, 657–667. <https://doi.org/10.3109/00365521.2015.1019918> (2015).
40. Li, W. *et al.* The gastroprotective effect of nobiletin against ethanol-induced acute gastric lesions in mice: impact on oxidative stress and inflammation. *Immunopharmacol. Immunotoxicol.* **39**, 354–363. <https://doi.org/10.1080/08923973.2017.1379088> (2017).
41. Akin, C. Mast cell activation syndromes. *J. Allergy Clin. Immunol.* **140**, 349–355. <https://doi.org/10.1016/j.jaci.2017.06.007> (2017).
42. Neuman, R. E. & Logan, M. A. The determination of collagen and elastin in tissues. *J. Biol. Chem.* **186**, 549–556 (1950).
43. Oyagi, A., Ogawa, K., Kakino, M. & Hara, H. Protective effects of a gastrointestinal agent containing Korean red ginseng on gastric ulcer models in mice. *BMC Complement. Altern. Med.* **10**, 45. <https://doi.org/10.1186/1472-6882-10-45> (2010).
44. Li, Q. *et al.* Kaempferol protects ethanol-induced gastric ulcers in mice via pro-inflammatory cytokines and NO. *Acta Biochim. Biophys. Sin. (Shanghai)* **50**, 246–253. <https://doi.org/10.1093/abbs/gmy002> (2018).
45. Pizzino, G. *et al.* Blockade of the JNK signalling as a rational therapeutic approach to modulate the early and late steps of the inflammatory cascade in polymicrobial sepsis. *Mediat. Inflamm.* **2015**, 591572. <https://doi.org/10.1155/2015/591572> (2015).
46. Li, W. *et al.* Anti-ulcerogenic effect of cavidine against ethanol-induced acute gastric ulcer in mice and possible underlying mechanism. *Int. Immunopharmacol.* **38**, 450–459. <https://doi.org/10.1016/j.intimp.2016.06.016> (2016).
47. Collins, T. J. ImageJ for microscopy. *Biotechniques* **43**, 25–30. <https://doi.org/10.2144/000112517> (2007).
48. Coelho, I. P. *et al.* Chemical profile and gastroprotective effect of *Jatropha elliptica* (Pohl) Oken roots. *Fitoterapia* **146**, 104707. <https://doi.org/10.1016/j.fitote.2020.104707> (2020).
49. Ribeiro, A. R. *et al.* The effects of baicalin on gastric mucosal ulcerations in mice: protective pathways and anti-secretory mechanisms. *Chem. Biol. Interact.* **260**, 33–41. <https://doi.org/10.1016/j.cbi.2016.10.016> (2016).
50. Fusco, R. *et al.* Absence of formyl peptide receptor 1 causes endometriotic lesion regression in a mouse model of surgically-induced endometriosis. *Oncotarget* **9**, 31355–31366. <https://doi.org/10.18632/oncotarget.25823> (2018).
51. Irrera N., *et al.* Activation of A2A receptor by PDRN reduces neuronal damage and stimulates WNT/ β -CATENIN driven neurogenesis in spinal cord. *Front Pharmacol.* **9**, 506. <https://doi.org/10.3389/fphar.2018.00506> (2018).
52. Cordaro, M. *et al.* Adelmidrol, a palmitoylethanolamide analogue, as a new pharmacological treatment for the management of inflammatory bowel disease. *Mol. Pharmacol.* **90**, 549–561. <https://doi.org/10.1124/mol.116.105668> (2016).
53. Matusiewicz, M. *et al.* In vitro influence of extracts from snail *Helix aspersa* Muller on the colon cancer cell line Caco-2. *Int J. Mol. Sci.* <https://doi.org/10.3390/ijms19041064> (2018).

Author contributions

E.G. conceived of the study and drafted the manuscript. M.C. drafted the manuscript. R.F., A.F.P. and D.I. carried out the experiments, revised the manuscript and performed the statistical analysis. R.D.P. R.C. and S.C. designed the study and critically revised the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-83170-8>.

Correspondence and requests for materials should be addressed to R.P. or S.C.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021