

Protein cargo encapsulation by virus-like particles: Strategies and applications

Author

McNeale, Donna, Dashti, Noor, Cheah, Li Chen, Sainsbury, Frank

Published

2022

Journal Title

WIREs Nanomedicine and Nanobiotechnology

Version

Version of Record (VoR)

DOI

<https://doi.org/10.1002/wnan.1869>

Copyright Statement

© 2022 The Authors. WIREs Nanomedicine and Nanobiotechnology published by Wiley Periodicals LLC. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Downloaded from

<http://hdl.handle.net/10072/422337>

Griffith Research Online

<https://research-repository.griffith.edu.au>

Protein cargo encapsulation by virus-like particles: Strategies and applications

Donna McNeale¹  | Noor Dashti²  | Li Chen Cheah²  | Frank Sainsbury^{1,2} 

¹Centre for Cell Factories and Biopolymers, Griffith Institute for Drug Discovery, Griffith University, Nathan, Queensland, Australia

²Australian Institute for Bioengineering and Nanotechnology, The University of Queensland, St Lucia, Queensland, Australia

Correspondence

Frank Sainsbury, Centre for Cell Factories and Biopolymers, Griffith Institute for Drug Discovery, Griffith University, Nathan, QLD 4111, Australia.
 Email: f.sainsbury@griffith.edu.au

Edited by: Emily Frieben, Managing Editor and Nils Walter, Co-Editor-in-Chief

Abstract

Viruses and the recombinant protein cages assembled from their structural proteins, known as virus-like particles (VLPs), have gained wide interest as tools in biotechnology and nanotechnology. Detailed structural information and their amenability to genetic and chemical modification make them attractive systems for further engineering. This review describes the range of non-enveloped viruses that have been co-opted for heterologous protein cargo encapsulation and the strategies that have been developed to drive encapsulation. Spherical capsids of a range of sizes have been used as platforms for protein cargo encapsulation. Various approaches, based on native and non-native interactions between the cargo proteins and inner surface of VLP capsids, have been devised to drive encapsulation. Here, we outline the evolution of these approaches, discussing their benefits and limitations. Like the viruses from which they are derived, VLPs are of interest in both biomedical and materials applications. The encapsulation of protein cargo inside VLPs leads to numerous uses in both fundamental and applied biocatalysis and biomedicine, some of which are discussed herein. The applied science of protein-encapsulating VLPs is emerging as a research field with great potential. Developments in loading control, higher order assembly, and capsid optimization are poised to realize this potential in the near future.

This article is categorized under:

Nanotechnology Approaches to Biology > Nanoscale Systems in Biology
 Biology-Inspired Nanomaterials > Protein and Virus-Based Structures

KEYWORDS

biocatalysis, encapsulation, protein therapeutics, virus capsids, virus-like particles

1 | INTRODUCTION

Virus-like particles (VLPs) mimic the structure of viruses, albeit devoid of native genetic material. They retain many of the inherent properties of virus capsids, including precise structure and uniformity of size, biocompatibility, stability, immunogenicity, and cell entry behavior (Lua et al., 2014). These features make them attractive starting materials for a

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *WIREs Nanomedicine and Nanobiotechnology* published by Wiley Periodicals LLC.

range of diverse applications in health and manufacturing. As biomimetic materials, VLPs and virus-derived nanoparticles have been engineered to both display and encapsulate functional biomolecules since the 1990s (Plummer & Manchester, 2011; Wen & Steinmetz, 2016). At first this was achieved by genetically modifying replicating viruses in their natural or experimental hosts. However, recombinant VLPs are produced using scalable and biocompatible processes using various production hosts, which may or may not be the same as the source host for the cognate virus (J. Glasgow & Tullman-Ereck, 2014; Le & Müller, 2021; Sainsbury, 2020). Moreover, VLPs are amenable to both genetic and chemical modification (Z. Chen et al., 2016; Schoonen & van Hest, 2014; Szyszka et al., 2022), which can be made with molecular precision to impart new functionalities, further diversifying the potential applications of VLPs.

This review focuses on the use of VLPs as protein cages and the capacity for VLPs to encapsulate heterologous protein cargos. The interior cavity of VLPs have been used for the sequestration of many different protein cargos, most prominently fluorescent proteins (FPs) and various enzymes (Wilkerson et al., 2018). The former has been instrumental in the design and testing of protein-encapsulating VLPs (Dashti et al., 2018; Minten et al., 2009; O'Neil et al., 2011), as well as their use as intracellular delivery vehicles (Brillault et al., 2017; Dashti et al., 2018) and in vivo imaging agents (Das, Zhao, Crooke, et al., 2020; Herbert et al., 2020). The latter results in the creation of biocatalytic nanocompartments for application in vitro and in vivo, and their development has revealed information on the fundamental behavior of enzymes in confined environments (Jordan et al., 2016; Minten et al., 2011; Wang et al., 2020). Here, we discuss the diversity of VLP systems developed for protein encapsulation. Numerous other types of protein cages have been developed in nanotechnology, and some of these are indeed virus-like. We point the reader to several useful reviews for further information on these nonviral protein cages (Bhaskar & Lim, 2017; Edwardson et al., 2022; Edwardson & Hilvert, 2019; Stupka & Heddle, 2020).

Viruses have evolved a variety of mechanisms for the encapsulation of their genome, minor coat proteins (CPs), and nonstructural proteins, and researchers have explored these strategies for the encapsulation of foreign cargos. From adapting native biomolecular interactions to capsid protein translational fusions, we discuss the various strategies for protein cargo encapsulation within non-enveloped VLPs. The source of these VLPs are viruses that infect prokaryotes, plants, and animals, and the VLPs are diverse in terms of size and capsid protein organization (Figure 1). Properties of the particles and their assembly mechanisms dictate the available encapsulation approaches that may be used for each VLP. The advantages and limitations of these approaches are considered, and we highlight several studies that demonstrate advancements in heterologous cargo encapsulation within VLPs. These advancements are accompanied by applications in biomedicine and biocatalysis, and we discuss a range of applications and progress made within these areas.

2 | NONSPECIFIC ENCAPSULATION

VLP assembly in the presence of high bulk concentrations of cargo proteins can result in encapsulation by nonspecific interactions. To achieve specificity with this type of encapsulation approach, assembly must be carried out in vitro from purified components. While this does allow a degree of control, in general, loading density of cargo proteins is low.

2.1 | Statistical encapsulation

Statistical encapsulation is achieved when capsid protein subunits are assembled in vitro in the presence of a high concentration of free protein cargo (Figure 2), leading to encapsulation of cargo at a density that is dependent on the cargo concentration. This approach led to the first demonstration of protein encapsulation by a capsid-based VLP, which was assembled from the CP of cowpea chlorotic mottle virus (CCMV) (Comellas-Aragonès et al., 2007). CCMV VLPs exhibit reversible assembly and disassembly, which is pH dependent. Although this can limit encapsulation of pH-sensitive cargos, this study reported a novel means of investigating single enzyme kinetics via the encapsulation of horseradish peroxidase (Comellas-Aragonès et al., 2007). This approach has also been taken for green FP (GFP) encapsulation by core particles of hepatitis B virus (HBV) (K. W. Lee & Tan, 2008). Hepatitis B core antigen (HBcAg) VLPs dissociate in the presence of denaturants and reassemble upon their removal (Wingfield et al., 1995).

Amenability to in vitro assembly characterizes the VLPs used for statistical encapsulation. This provides specific advantages related to the chemical control over assembly, for example, avoiding the encapsulation of host proteins and potentially some control over the average number of proteins encapsulated per particle. However, statistical encapsulation requires a large excess of cargo that needs to be separately expressed and purified before loading and subsequently




FIGURE 1 Examples of non-enveloped viruses used for the encapsulation of heterologous proteins. PDB IDs: MS2, 2MS2; CCMV, 1ZA7; Qβ, 1QBE; PP7, 1DWN; CMV, 1F15; HBcAg, 1QGT; PyV, 1SIE; P22, 2XYX; BTV core, 2BTV minus PCD chains; and RV core, 3KZ4. Murine polyomavirus is shown as a representative of the Polyomaviridae family. Average external diameters were obtained from VIPERdb (<https://viperdbscripps.edu>). Figure lists maximum determined cargo numbers, see the text for details. CCMV, cowpea chlorotic mottle virus; CMV, cucumber mosaic virus; HBcAg, hepatitis B core antigen; PyV, polyomavirus; RV, rotavirus

removed from reassembled VLPs before quantitative assays on encapsulation can be performed. Furthermore, loading density is limited by the concentration of cargo proteins required to achieve loading by statistical encapsulation alone. The following strategies use protein engineering to shift the equilibrium towards partitioning of cargo proteins within VLPs assembled both *in vitro* and *in vivo*.

2.2 | Electrostatic interactions

Encapsulation by electrostatic interactions relies on charge complementarity between the interior surface of the VLP and the cargo protein (Figure 2). Many capsid-based viruses carry positive charges on their interior surface, which facilitates encapsulation of their viral genome, and this feature can be exploited to encapsulate other negatively charged




FIGURE 2 Cargo protein–capsid protein interactions used to direct the encapsulation of heterologous protein cargos in VLPs. VLPs, virus-like particles

cargo. For example, a far-red FP, small-ultrared FP (smURFP), was encapsulated within two structurally similar VLPs from the Leviviridae family, bacteriophage Q β , and pseudomonas phage seven (PP7) (Herbert et al., 2020). Both viral capsids are composed of 180 copies of a CP and assembly is normally directed by interactions with RNA. Allophycocyanin α -subunit, from which smURFP is derived, has an isoelectric point of 4.64, allowing smURFP to mimic the role of RNA during assembly and thus be encapsulated at neutral pH. Herbert et al. (2020) achieved packaging of approximately three smURFP per particle, producing brightly fluorescent VLPs which, when injected subcutaneously and intravenously, were shown to be effective noninvasive imaging agents (Herbert et al., 2020).

Charge-mediated encapsulation can also be achieved by imparting negative charges on the cargo protein through the conjugation of nucleic acids (Brasch et al., 2017; J. E. Glasgow et al., 2012; Lu et al., 2012). In the case of CCMV, the N-terminal region of the CP is enriched in arginine residues, and it was found that chemically linking nucleic acid adapters to cargo proteins can trigger assembly around the cargo protein (Brasch et al., 2017). This represented a significant development in the use of CCMV for the encapsulation of proteins sensitive to the changes in pH previously used to drive assembly. Moreover, it provides a mechanism that is potentially tunable by varying the adapter length or salt concentration during assembly. Using this strategy, Brasch et al. (2017) demonstrated the confinement of two separate enzyme pathways, achieving an average encapsulation of one glucose oxidase (GOx), or one GOx and one or two gluconokinase (GCK) per particle. Cucumber mosaic virus (CMV) belongs to the same viral family as CCMV and is structurally similar; however, unlike CCMV, CMV requires CP–RNA interactions for assembly. This interaction was mimicked by a DNA oligomer conjugated to streptavidin and a fluorophore to induce assembly of CMV, leading to incorporation of the protein cargo into VLPs (Lu et al., 2012). The Bacteriophage MS2 VLP was one of the first non-enveloped VLPs used for protein encapsulation and this was mediated by a sequence-specific RNA–CP interaction (Wu et al., 1995). However, in the presence of the osmolyte, trimethylamine N-oxide, assembly can proceed efficiently around cargo modified with nonspecific polyanions (J. E. Glasgow et al., 2012), obviating the need to work with relatively unstable RNA aptamers. The conjugation of a DNA tag to GFP, for example, resulted in the encapsulation of approximately 6.5 GFPs per capsid. The subsequent use of a genetically fused polyanionic tag (16 acidic amino acid residues) also resulted in capsid assembly in the presence of osmolyte, though the number of encapsulated GFPs or alkaline phosphatases was not reported (J. E. Glasgow et al., 2012).

The use of complementary charges to direct assembly provides a relatively simple method for cargo encapsulation, albeit with the requirement of separate cargo purification for *in vitro* assembly and, in some cases, the chemical modification of cargo molecules with polyanions such as nucleic acids. This strategy has been of particular use in VLPs that require a charged cargo for assembly. Although cargo loading efficiencies are lower compared to encapsulation methods that use specific interactions, the approach is more flexible with regard to polyanion conjugate and loading densities can be somewhat tunable through the variation of adapter length or ionic strength.

3 | SPECIFIC NON-COVALENT INTERACTIONS

Encapsulation via specific interactions can be mediated by either protein–protein or protein–nucleic acid interactions. Furthermore, these interactions can be native, borrowing from the assembly of virions around internal components, or engineered non-native interactions. These approaches may be used for encapsulation during *in vitro* assembly, but particularly in the case of native interactions, the specificity of the interactions affords the possibility of *in vivo* self-sorting of protein cargos into VLPs that assemble in their host cells.

3.1 | Native

3.1.1 | Protein

Early demonstration of heterologous protein encapsulation within virus capsids used viable Bacteriophage T4 and the interaction between the scaffold protein (SP) and the interior surface of the CP (Hong & Black, 1993). This work showed that active proteins, including GFP and several enzymes, could be encapsulated high density, despite the presence of the genome (Mullaney & Black, 1996, 1998), and was a key demonstration that native protein–protein interactions could be engineered for specific *in vivo* cargo packaging. This approach has since been adapted to recombinant VLPs of numerous viruses as discussed in the following paragraphs.

Bacteriophage P22 VLPs have been widely investigated for their encapsulation abilities (Esquirol et al., 2022; O'Neil et al., 2011, 2012; D. Patterson et al., 2012; D. P. Patterson et al., 2014; D. P. Patterson, LaFrance, et al., 2013; Qazi et al., 2016; Wang et al., 2020). The particle is composed of 420 copies of CP forming an icosahedral capsid and is assembled with the support of 100 to 330 copies of P22 SP. A truncated SP, which retains its ability to template the assembly of P22 VLPs, can be used to encapsulate cargo proteins via genetic fusion to the cargo N or C termini (Figure 3a). O'Neil et al. (2011) reported a method for the *in vivo* encapsulation of FPs, GFP and mCherry, genetically fused to an N-terminally truncated SP and co-expressed with P22 CP (O'Neil et al., 2011). P22 capsids showed high loading capacity with 281 GFP molecules or 233 mCherry molecules within each capsid. This high loading capacity enables the co-encapsulation of cargo proteins via sequential fusion to SP in a 1:1 ratio (O'Neil et al., 2012). Although the *in vivo* assembly and encapsulation in P22 does not require the separate purification of cargo and CP components, it provides little control over cargo stoichiometry and loading density. To overcome these limitations, an *in vitro* assembly approach was developed (Sharma et al., 2017) where packaging stoichiometry and density of cargos can be modified by varying the ratio of cargo-fused SP and wildtype SP (Sharma & Douglas, 2020).

Polyomavirus (PyV) VLPs have also been widely explored for both *in vivo* (Bouřa et al., 2005; Catrice & Sainsbury, 2015; Cheah et al., 2021) and *in vitro* (Abbing et al., 2004; Dashti et al., 2018) encapsulation. The capsid of murine polyomavirus (MPyV) is comprised of 360 copies of the major CP VP1 arranged as 72 pentamers. Each of these capsomeres can accommodate the binding of one minor CP VP2/3, which share the same C-terminus. Abbing et al. (2004) reported a method for *in vitro* encapsulation by fusion of GFP to a 49 amino acid C-terminal fragment of VP2 (VP2C) (Abbing et al., 2004), which is sufficient for VP1 capsomere binding (X. S. Chen et al., 1998). GFP-loaded VLPs were assembled by mixing GFP-VP2C with VP1 pentamers resulting in an estimated 64 GFPs per VLP. However, due to the hydrophobicity of VP2, a major drawback to this method is the poor solubility of the anchor protein. To address this limitation, an approach combining *in vivo* cargo capture with *in vitro* assembly was established (Dashti et al., 2018). In contrast to self-assembly in eukaryotic cells, MPyV VP1 capsomeres do not self-assemble into VLPs when expressed in prokaryotic cells. Co-expression of VP1 and VP2C-EGFP resulted in the formation of soluble VP1/VP2C-EGFP complexes which were subsequently assembled *in vitro* (Figure 3b). The co-encapsulation of two FPs, EGFP and mRuby3, was used to demonstrate Förster resonance energy transfer between the separately tagged yet co-confined cargo molecules (Dashti et al., 2018). This work reduced the length of the VP2 C-terminal fragment used as a capsomere-binding anchor to 31 aa and the *in vivo* capture approach enabled fusing the anchor to cargo protein N-termini. This orientation destabilizes the fusion protein *in vitro* (Abbing et al., 2004) and *in vivo* (Cheah et al., 2021); however, interaction with VP1 capsomeres masks the hydrophobic side chains of the conserved alpha-helix that mediates binding (X. S. Chen et al., 1998). Work on the similar polyomavirus, Simian virus 40 (SV40), showed that placing VP2C fragments at the N-terminus of cargo proteins avoids the external presentation of cargo proteins that sometimes results from C-terminal fusion of VP2C (Inoue et al., 2008). Another polyomavirus, JC polyomavirus, is able to self-assemble in prokaryotic cells, and has also been explored in protein encapsulation studies




FIGURE 3 Examples of specific interactions mediating cargo protein interactions in VLPs. (a) SP-mediated in vivo encapsulation in P22 adapted from D. Patterson et al. (2012). (b) VP2 peptide-mediated in vivo cargo capture and in vitro encapsulation in MPyV VLPs adapted from Dashti et al. (2018). (c) RNA adapter-mediated in vivo encapsulation in Qβ VLPs adapted from Das, Zhao, Elofson, et al. (2020). (d) Coiled-coil interactions mediating in vitro CP purification followed by in vitro encapsulation in CCMV VLPs adapted from Minten et al. (2009). CCMV, Cowpea chlorotic mottle virus; CP, coat protein; MPyV, murine polyomavirus; SP, scaffold protein; VLPs, virus-like particles

(Ohtake et al., 2010). GFP fused to the N-terminus of VP2 and co-expressed with VP1 in *Escherichia coli* was encapsulated in vivo. When GFP was replaced with a 6x histidine tag, small molecules containing nitrilotriacetic acid (NTA) could be encapsulated via diffusion through the capsid pores and binding to the engineered cargo protein. Release of the cargo was shown at low pH and protonation of the histidines, showing potential as a controlled release drug carrier (Ohtake et al., 2010). Human papillomavirus (HPV) is structurally similar to PyV VLPs and has also been used for in vivo encapsulation in insect cells by an orthologous approach, with fusion of GFP to the minor capsid protein (Windram et al., 2008).

3.1.2 | Nucleic acid

Sequence or structure-specific protein–RNA interactions are necessary for the capsid assembly of many ssRNA viruses (Twarock et al., 2018). These secondary structure elements, known as packaging signals (PSs), have been used to trigger

the assembly and encapsulation of foreign cargo within VLPs (Ashley et al., 2011; Das, Zhao, Crooke, et al., 2020; Fiedler et al., 2010; Rhee et al., 2011; Wu et al., 1995). Genome encapsulation and assembly of the Bacteriophage MS2 capsid is mediated by a 19-nucleotide RNA stem-loop (*pac* site) (Pickett & Peabody, 1993). The encapsulation of foreign protein cargo has been achieved through the conjugation of the *pac* site RNA to the cargo (Ashley et al., 2011; Wu et al., 1995). Wu et al. (1995) first demonstrated the in vitro encapsulation of a glycoprotein toxin, ricin toxin A-chain into MS2 VLPs using this approach, which were successfully delivered into mammalian cells.

Similar to MS2, bacteriophage Q β assembly uses an RNA hairpin structure that interacts with the interior of the CP to form viral capsids (Witherell & Uhlenbeck, 1989). To test the in vivo encapsulation of a protein cargo within Q β VLPs, a bifunctional RNA molecule containing an α -Rev RNA aptamer and the genome packaging hairpin flanking the CP RNA sequence was designed, while an N-terminal Rev peptide tag was fused to the cargo enzyme (Fiedler et al., 2010). Co-expression of this dual-plasmid system led to the binding of the Rev-tagged enzyme to the α -Rev aptamer, which was subsequently tethered to the interior of the Q β capsid via the RNA hairpin (Figure 3c). Three enzymes were tested with this system, and it was shown that the average number of encapsulated cargos could be somewhat controlled by modifying expression conditions, achieving a variable loading of PepE, a 24 kDa enzyme, between 2 and 18 copies per particle. This encapsulation strategy was also used to package FPs separately into Q β VLPs, with up to 15 proteins per particle (Rhee et al., 2011), and near-infrared FPs with either five or nine copies per particle (Das, Zhao, Crooke, et al., 2020).

3.2 | Non-native (engineered)

3.2.1 | Protein

Non-covalent encapsulation strategies also include the use of non-native binding domains to generate specific interactions between the capsid inner surface and cargo, directing encapsulation of cargo proteins upon assembly. WW domains are small protein modules that mediate protein–protein interactions through the binding of proline-rich ligands (Ingham et al., 2005). Schmidt et al. (2001) used the first WW domain of the mouse formin-binding protein 11 (FBP11) and the proline-rich motif PPLP to direct the in vitro encapsulation of PPLP-tagged proteins within polyomavirus (MPyV) VLPs (Schmidt et al., 2001). The FBP11 WW domain was fused to the inward-facing N-terminus of VP1, while GFP was modified with the PPLP tag. Increase in cargo-to-capsomere ratio during assembly led to an increase in cargo loading, reaching a maximum of 260 GFP molecules per particle. Although only demonstrated for in vitro loading, the capacity is greater than encapsulation via the minor CP VP2, which has a theoretical loading maximum of 72 monomeric cargo molecules per particle.

Coiled-coils are protein structural motifs composed of two to six alpha helices arranged either parallel or antiparallel to each other (Truebestein & Leonard, 2016). They can be homo- or hetero-oligomers and contain repeating hydrophobic residues which drives the association of the helices resulting in the coiled-coil formation. Heterodimeric E- and K-coil peptides were chosen for the directed non-covalent protein encapsulation inside CCMV particles (Minten et al., 2009, 2011; Rurup et al., 2014). Through genetic modification, the K-coil was fused to the interior-facing N-terminus of the CCMV CP and the E-coil to the C-terminus of EGFP. Purification of EGFP–capsid protein complexes was achieved through immobilization of the GFP and capture of the CP directly from lysates using K-/E-coil dimerization (Figure 3d; Minten et al., 2009). In vitro assembly of the EGFP-capsid protein complexes required mixing with unmodified capsid protein in various ratios, enabling the encapsulation of up to 15 EGFP molecules per capsid. This approach allowed for higher loading with more control over loading density than by nonspecific encapsulation in CCMV and was further explored for the encapsulation of *Pseudozyma antartica* lipase B (PalB) at various loading densities to enable investigation of reaction rate as a function of PalB loading density (Minten et al., 2011). The use of non-native binding domains provides an alternative to natural targeting peptides, such as scaffolding proteins, although it has only been applied to in vitro assembly and encapsulation.

4 | COVALENT INTERACTIONS

Encapsulation via covalent interactions with capsid proteins results from both translational and posttranslational, enzymatic, fusion of cargo proteins. Due to the considerable changes to capsid protein properties that result from cargo

TABLE 1 Recombinant non-enveloped virus capsids that have been used to encapsulate heterologous proteins

Host kingdom	Family	Virus	Abbr.	Minimal protein composition (no. CP)	Internal diameter	Assembly	Encapsulation strategy
Bacteria	Leviviridae	Enterobacteria phage MS2	MS2	180 CP	21 nm	In vivo or in vitro	DNA aptamer RNA aptamer Posttranslational CP fusion
		Enterobacteria phage Qbeta	Qβ	180 CP	21.4 nm	In vivo or in vitro	Electrostatic RNA aptamer
		Pseudomonas phage seven	PP7	180 CP	21.6 nm	In vitro	Electrostatic
	Podoviridae	Enterobacteria phage P22	P22	420 CP, 100–330 SP	46.4 nm to > 53.4 nm ^a	In vivo or in vitro	Native protein tag
Animalia	Hepadnaviridae	Hepatitis B virus (core antigen)	HBcAg	180 CP, or 240 CP	25.4 nm	In vivo or in vitro	Statistical
	Polyomaviridae	JC Polyomavirus	JCPyV	360 CP	35 nm	In vivo	Native protein tag
		Murine Polyomavirus	MPyV	360 CP	35 nm	In vivo or in vitro	Native and non-native protein tags
	Reoviridae	Simian virus 40	SV40	360 CP	35.8 nm	In vivo	Native protein tag
		Bluetongue virus	BTV	600 VP7 ^b , 120 VP3	46.2 nm	In vivo	Translational CP fusion
		Rotavirus	RV	780 VP6, 120 VP2	45.4 nm	In vivo	Translational CP fusion
	Birnaviridae	Infectious bursal disease virus	IBDV	780 CP ^c	50.4 nm	In vivo	Translational CP fusion
Parvoviridae	Canine parvovirus	CPV	60 CP	16 nm	In vivo	Translational CP fusion	
Plantae	Bromoviridae	Cowpea chlorotic mottle virus	CCMV	180 CP	18.6 nm	In vivo or in vitro	Statistical Electrostatic DNA aptamer Non-native protein tag Posttranslational CP fusion
		Cucumber mosaic virus	CMV	180 CP	17 nm	In vitro	DNA aptamer
	Secoviridae	Grapevine fanleaf virus	GFLV	60 CP	22 nm	In vivo	Translational CP fusion

^aP22 VLPs can vary in size with thermal or chemically inducible maturation from the procapsid to expanded and wiffle ball forms (Selivanovitch et al., 2019).

^bBTV core particles derived from the virion possess a full complement of 780 VP7 proteins making up the T = 13 second layer. However, it has been found that recombinant cores lack VP7 trimers at the fivefold axis of symmetry (Brillault et al., 2017). See Figure 1.

^cWild-type VP2 assembles into a T = 1 capsid whereas the truncated form that enables protein encapsulation assembles into a T = 13 capsid (Pascual et al., 2015).

fusion, this approach is generally used for in vivo encapsulation. Nearly all non-enveloped VLPs that have been used for heterologous protein encapsulation can be assembled in vivo (Table 1); however, success of this strategy depends on the organization of capsid proteins and size of the capsid. In larger VLPs that can accommodate the full complement of fusion proteins while maintaining high-fidelity assembly, cargo proteins can be loaded with predictable stoichiometries. In smaller VLPs, encapsulation is achieved in mosaic particles containing both fused and unfused capsid proteins.

4.1 | Translational CP fusion

Genetic fusion directly to CPs is a conceptually simple method for the encapsulation of foreign protein cargo (Brillault et al., 2017; Charpilienne et al., 2001). In cases where encapsulation is efficient, this approach can allow high loading of cargo proteins at known stoichiometries. However, it is limited by steric constraints on cargo protein size and assembly of capsid proteins (Belval et al., 2016; Beterams et al., 2000; Cortes-Perez et al., 2010; Gilbert et al., 2004; Kar et al., 2005; Pascual et al., 2015). Infectious bursal disease virus (IBDV) capsids are formed by 260 trimers of VP2 protein (Luque et al., 2009). In the absence of the scaffolding protein VP3, precursor VP2 (pVP2) with an N-terminal histidine tag can assemble into $T = 13$ VLPs (Pascual et al., 2015). Both the N and C termini of pVP2 reside within the interior of the capsid and while direct fusion of EGFP to the N-terminus of pVP2 via the histidine-tag did not result in VLPs, this was overcome by co-expression of cargo-CP and unmodified CP in insect cells. Stable particles containing approximately 240 copies of EGFP (~ 1 per pVP2 trimer) were formed when the EGFP fusion was co-expressed with unfused pVP2, indicating that steric hindrance limited capsid formation (Pascual et al., 2015).

The reoviruses, rotavirus (RV) and bluetongue virus (BTV) have also been used for encapsulation by translational fusion (Brillault et al., 2017; Charpilienne et al., 2001; Kar et al., 2005; Thuenemann et al., 2021). The VLP is a triple layer particle composed of four different proteins, although recombinant double layer core particles composed of two proteins and single layer subcore particles composed of only one protein, are also possible. The different particles exhibit a range of physical properties, with the core-like particles the most stable under physiological conditions (Jiménez-Zaragoza et al., 2018; Thuenemann et al., 2013), although the VLP possesses the most effective cell uptake ability (Thuenemann et al., 2021). Encapsulation of protein cargo is achieved by fusion to each of the 120 copies of the scaffolding protein that make up the subcore particle, and fusion to the N-terminus has enabled the encapsulation of 120 GFP molecules in both RV (Charpilienne et al., 2001) and BTV (Brillault et al., 2017).

Although this approach best suits larger VLPs with higher cargo capacity, there are examples of smaller capsids that have been used for cargo encapsulation by direct fusion. Canine parvovirus (CPV) is one such example (Gilbert et al., 2004). Sixty copies of VP2 are sufficient for VLP assembly and GFP fusion to the internal facing N-terminus yielded fluorescent VLPs, although a protein product similar in apparent molecular weight to unfused VP2 was observed. This suggested the formation of mosaic particles resulting from the proteolytic cleavage of the fusion product, which may be necessary for the assembly of VLPs. Another smaller VLP used for encapsulation by translational fusion is Grapevine fanleaf virus (GFLV). Fluorescent protein fusion showed that the N-terminus of GFLV CP is also internal, with recombinant expression in plants leading to red FP (RFP) encapsulation in VLPs (Belval et al., 2016). Cargo-loaded VLPs appeared to be comprised of a majority intact CP-cargo fusion. Encapsulation by CP fusion within another smaller VLP, HBV VLPs, has also been demonstrated (Beterams et al., 2000). Two-hundred and forty copies of a 17 kDa *Staphylococcus aureus* nuclease were encapsulated via fusion to the C-terminus of the CP. CP-cargo fusions remained intact; however, the frequency of aberrant particles was higher than that of VLPs formed by the co-expression of cargo-fused and wildtype CP.

4.2 | Posttranslational fusions

4.2.1 | Sortase-mediated

Surface proteins are anchored to the bacterial cell wall of *S. aureus* via the action of the membrane protein Sortase A (SrtA) (Mazmanian et al., 2001). Proteins harboring the C-terminal sorting signal, an LPXTG motif, are recognized by SrtA, which cleaves between the threonine and glycine within the LPXTG motif, liberating the carboxyl of the threonine, which is subsequently coupled to an N-terminal glycine residue of peptidoglycan cross-bridges. Schoonen et al. (2015) developed a method for the site-specific covalent modification of CCMV capsids catalyzed by SrtA. The N-terminus of the CP, located within the interior of the capsid, was engineered to display a glycine residue while GFP was modified with a LPETG-tag at its C-terminus. In vitro assembly was conducted following the Sortase-mediated coupling, resulting in VLPs-encapsulating 16–18 GFP molecules (Schoonen et al., 2015). The peptide tags required for this SrtA-based strategy are relatively short and therefore modification of capsid and cargo proteins are more likely to be tolerated, making this a potentially broadly applicable strategy for protein cargo encapsulation, and could theoretically be used to mediate in vivo encapsulation.

4.2.2 | SpyTag/SpyCatcher-mediated

The SpyTag/SpyCatcher system is another example of a bacterial-derived protein linkage system from *Streptococcus pyogenes* (Zakeri et al., 2012). The SpyTag (13 amino acids), harboring a reactive aspartate, and SpyCatcher (12.3 kDa) which contains a reactive lysine, rapidly form a covalent isopeptide bond upon interaction. This system has been used for the in vivo encapsulation of two enzymes within bacteriophage MS2 capsids (Giessen & Silver, 2016). The SpyTag was inserted within an internal loop of the MS2 CP, while the SpyCatcher was genetically fused to two cargo proteins. SpyCatcher-tagged enzymes and SpyTag-CP were co-expressed, resulting in the covalent attachment of two to four molecules of each enzyme within the MS2 VLPs. In contrast to Sortase-mediated cargo capture, SpyCatcher fusions are considerably larger, possibly leading to steric constraints on cargo loading density; however, this method does not require the addition or co-expression of a ligating enzyme, making it relatively simpler than Sortase-mediated encapsulation, and easier to implement for in vivo encapsulation.

5 | BIOMEDICAL APPLICATIONS

Engineered viruses and VLPs have had clinical success as vaccines and gene delivery vectors. As nanoparticles, both modified viruses and recombinant VLPs are showing preclinical promise as useful scaffolds and delivery vehicles for various inorganic, organic and biomolecular therapeutics (Chung et al., 2020). Size, biocompatibility, and engineerability are key features of VLPs that make them attractive as delivery vehicles in biomedical applications. The ability to chemically or genetically modify the exterior of VLPs carrying heterologous cargo proteins has been used to enhance or modify cell uptake, tune cell tropism, or present antigens (Steinmetz et al., 2020). Specific uses for protein loaded VLPs include the delivery of prodrug-converting enzymes, diagnostic imaging, and the encapsulation of T-cell epitopes.

Enzyme prodrug therapy is a strategy to increase the specificity and efficacy of a therapeutic by delivering a prodrug converting enzyme to a specific site. Both CCMV and P22 have been used for the encapsulation of cytochrome P450 BM3 (CYP_{BM3}) for the activation of anticancer drugs (Sanchez-Sanchez et al., 2014; Sánchez-Sánchez et al., 2015; Tapia-Moreno et al., 2017). P22 VLPs loaded with an average of 109 enzymes were transfected into human cervical carcinoma cells using lipofectamine, where they conferred 10-fold higher CYP activity compared to endogenous CYP activity alone (Sanchez-Sanchez et al., 2015). Chemical modification of the exterior of P22 VLPs carrying CYP_{BM3} to present folic acid increased uptake into a breast cancer cell line and considerably increased their sensitivity to tamoxifen (Tapia-Moreno et al., 2017). In another study, multifunctional P22 VLPs containing CYP were decorated with photosensitizer and targeting moiety on the surface to allow the combination of enzyme prodrug therapy and photodynamic therapy (Chauhan et al., 2018). BTV VLPs and CLPs have also been developed for this application to take advantage of their natural ability for cell entry (Thuenemann et al., 2021). Here, both the triple layer VLP and double layer core-like particles were loaded with Herpes simplex virus 1 thymidine kinase, which is widely used in the conversion of prodrugs to cytotoxic agents. Although both particles are capable of efficient cell entry, the VLP is able to mediate endosomal escape and localization in the cytosol, resulting in the death of human glioblastoma cells in the presence of ganciclovir (Thuenemann et al., 2021).

Near-infrared FPs (NIR-FPs) are receiving growing attention for their use in noninvasive deep-tissue and whole-body imaging. Encapsulation within VLPs has the added benefits of increasing the stability of the cargo and potential for tunable cell and tissue tropism by exterior modification. To demonstrate the capacity for encapsulated NIR-FPs in in vivo imaging, monomeric mIFP and a red-shifted dimeric iRFP720 variant were separately packaged within Q β VLPs (Das, Zhao, Crooke, et al., 2020). Particles produced similar photochemical properties to unencapsulated NIR-FP but exhibited increased stabilization toward denaturation and proteolytic digestion. Systemic administration resulted in localization of the particles, and NIR fluorescence, in the liver. Similarly, smURFP was encapsulated within Q β and PP7 VLPs (Herbert et al., 2020). Encapsulated smURFP showed different tissue and organ localization and remained visible for longer than free smURFP. Although VLPs based on bacteriophages have not yet been shown to have a natural tropism for human cell receptors, they are readily engineered to confer cell binding. Cargo encapsulation within Q β VLPs via bifunctional RNA was combined with the external display of targeting ligands through CP-fusion, showing that external modification did not adversely affect cargo packaging efficiency (Fiedler et al., 2018). GFP-loaded Q β VLPs have also been decorated with a CD22 receptor ligand via Cu-catalyzed azide-alkyne cycloaddition and showed strong

and selective affinity to CD22⁺ cells with internalization (Rhee et al., 2011). Together this shows the potential of NIR-FP-loaded VLPs for diagnostic imaging.

VLPs are well established as effective vaccines against both the cognate virus and as presentation platforms for heterologous antigens (K. L. Lee et al., 2016; Lua et al., 2014). As a scaffold for the design of vaccines, the repetitive organization of the CP is particularly advantageous for external presentation of immunogenic moieties to elicit antibody responses. However, the encapsulation of immunogens can also be effective. Immunization of mice with IBDV VLPs loaded with one or both of the influenza virus antigens HA2 and M2 raised HA-specific or M2-specific antibodies and subsequent mice challenge studies resulted in protection against mortality (Pascual et al., 2015). This approach could serve to protect antigens as well as mediate their delivery to appropriate immune cells. The delivery to professional antigen presenting cells to stimulate humoral immunity is a promising use case for protein-encapsulating VLPs. P22 VLPs loaded with the conserved influenza nucleoprotein elicited a robust T-cell response that provided broad protection in a multistrain challenge assay in mice (D. P. Patterson, Rynda-Apple, et al., 2013). Similarly, respiratory syncytial virus (RSV) M and M2 proteins induced a humoral response that reduced RSV titers following challenge (Schwarz et al., 2016). Combination with external modifications has the potential to enhance or modify the immune response. For example, external presentation of CD40L trimers led to association of FP-loaded P22 VLPs with B lymphocytes (Schwarz et al., 2015).

In addition to infectious disease applications, the encapsulation of immunogens can also be used to stimulate therapeutic immune responses against chronic diseases. Several studies have demonstrated the use of antigen loaded MPyV VLPs for cancer immune therapy in mouse model systems (Eriksson et al., 2011; Tegerstedt et al., 2005, 2007). Protection against the development of Her2 expressing tumor in mice was shown following immunization with MPyV VLPs containing the extracellular and transmembrane domains of human Her2 (Tegerstedt et al., 2007). The antitumor effect was more efficient when VLPs were loaded into murine dendritic cells (DCs) *ex vivo*, whereas the unencapsulated antigen, directly administered or *ex vivo*-loaded on DCs, was unable to protect mice against a lethal dose of D2F2/E2 cells. Similarly, MPyV VLPs loaded with human prostate-specific antigen (PSA) were loaded onto DCs and PSA-specific CD4⁺ and CD8⁺ cells, which were sufficient to protect mice from PSA-expressing tumor outgrowth (Eriksson et al., 2011). Although these examples clearly demonstrate the utility of VLPs in stimulating the immune system, this inherent property of VLPs remains a barrier to their use as *in vivo* diagnostics or therapeutic delivery vehicles.

6 | BIOCATALYTIC APPLICATIONS

Over the past decade, VLPs have attracted increasing interest as containers for *in vitro* and *in vivo* biocatalysis. Pioneering work on the development of CCMV-based and then P22-based nanoreactors discussed in this review, have demonstrated the potential for fundamental studies of the effect of confinement on enzyme activity kinetics. Despite the fact that the impact of encapsulation on the activity of a given enzyme is not predictable (Esquirol et al., 2022), VLPs have been particularly useful tools for studying the effect of macromolecular crowding on individual enzyme activity due to their structural homogeneity and potential for precise control of the cargo packing density. Control over the number of encapsulated lipase molecules per CCMV VLP showed specific activity decreased at higher packing densities despite higher activity of the encapsulated enzyme compared to free enzyme (Minten et al., 2011). The effect of macromolecular crowding was studied in P22 VLPs by co-encapsulation of alcohol dehydrogenase with varying ratios of wild type SP, which can be selectively removed by treatment with a mild chaotrope (Sharma & Douglas, 2020). The specific activity of alcohol dehydrogenase was inversely related to the concentration of SP, suggesting that high macromolecular crowding could restrict enzyme conformational dynamics. Similarly, expansion of a P22 nanoreactor by heat treatment was found to increase the activity of encapsulated alcohol dehydrogenase, an effect attributed to reduction in macromolecular crowding (D. Patterson et al., 2012).

Protein cages are often seen to stabilize encapsulated cargo against proteases, high temperatures, and/or chemical denaturants *in vitro*. This has been demonstrated with VLPs of P22 (Jordan et al., 2016; Wang et al., 2020), MS2 (Giessen & Silver, 2016), and Q β (Das, Zhao, Elofson, et al., 2020; Fiedler et al., 2018). Moreover, *in vivo* loading into P22 VLPs has been shown to be a promising strategy to mitigate enzyme misfolding and aggregation during high-level prokaryote expression (D. P. Patterson, LaFrance, et al., 2013). Applied to cellular biocatalysis in baker's yeast, *Saccharomyces cerevisiae*, the use of MPyV VLPs to stabilize myoinositol oxygenase led to increased flux through a recombinant metabolic pathway (Cheah et al., 2021).

VLPs can function as a scaffold to spatially organize successive enzymes in a biocatalytic cascade, generating artificial metabolons. The goal is to improve pathway efficiency and minimize intermediate loss to competing reactions, mimicking the function of natural catalytic compartments such as bacterial microcompartments and eukaryotic organelles. Up to three enzymes in a glycolytic cascade have been co-encapsulated in P22 VLPs by fusing them in series with SP (D. P. Patterson et al., 2014). In CCMV VLPs, complementary ssDNA tags served to control GOx and GCK stoichiometry or to act at a secondary biocatalyst with GOx when the ssDNA sequence encoded a DNzyme (Brasch et al., 2017). Posttranslational fusion was used to co-encapsulate two enzymes in the indigo biosynthetic pathway within MS2 VLPs in vivo. Even with minimal control over loading stoichiometry, indigo bioproduction in *E. coli* increased by ~60% (Giessen & Silver, 2016). Going further to mimic natural compartmentalization, a nested protein cage system has been constructed with P22 VLPs (Waghvani et al., 2020). Ferritin (Fn) protein cages and the cellobiose-hydrolyzing enzyme were co-encapsulated within P22 via SP fusion. This work demonstrated the co-encapsulation of two distinct macromolecules, with Fn-cages themselves serving as subcompartments within P22. These in vitro studies and others demonstrate that controlled co-encapsulation of multiple enzymes is central to organizing biocatalytic cascades. The ability to achieve this in vivo, to organize metabolic pathways with defined stoichiometry in cellular biocatalysis, presents an ongoing challenge in the development of VLP-based biomimetic biocatalysts.

A promising direction for VLP-based nanoreactors is the potential to generate higher-order assemblies such as sheets, gels, and ordered 3D superlattices. Viruses, and VLPs in particular, have long been explored to develop novel nanomaterials due to their natural propensity to form ordered arrays (Sun et al., 2007; Yoshimura et al., 2016). VLPs can be assembled into 3D composites using combinations of covalent and non-covalent interactions mediated by additional proteins, ions, or organic compounds. Examples of hybrid composites include CCMV with photosensitive dendrons (Kostiainen et al., 2010) or avidin (Liljestrom et al., 2014), P22 VLPs with Dec proteins (Uchida et al., 2015) or PAMAM dendrimers (Uchida et al., 2018), and DNA-modified Q β VLPs (Cigler et al., 2010). For in vitro applications, such assemblies could facilitate efficient recycling and recovery of encapsulated enzymes. However, this use case is dependent on reaching a catalytic advantage that justifies the purification and processing of in vitro biocatalysts. The emergence of more interesting properties in the literature point to this possibility. Higher-order assemblies of P22 VLPs have been tuned to massively enhance partitioning of substrates, increasing the catalytic rates of encapsulated alcohol dehydrogenase (Selivanovitch et al., 2021).

7 | OUTLOOK

Recent results from heterologous protein encapsulation in VLPs cover an impressive array of applications, particularly in biomedicine and biocatalysis. On one hand, protein cargo loading has enabled fundamental studies on enzyme activity and confinement. On the other hand, these are synthetic systems developed as biocompatible and biomimicking vehicles with aspirations in biotechnology and nanotechnology. In this regard, significant barriers to clinical or commercial success remain to be addressed. While there is great potential for enhanced in vivo biocatalysis using VLPs as scaffolds for metabolic pathways, challenges remain for the translation of in vitro VLP engineering progress to in vivo settings. These include the implementation of stoichiometrically controlled cargo loading and directed assembly of higher-order structures. As intracellular delivery vehicles, VLPs are capable of long circulation and highly efficient uptake, though the specific pathway of uptake and inherent immunogenicity are important considerations that require more attention. Chemical modification is one approach to addressing some of these concerns. A more elegant approach might be to take advantage of the inherent mutability of virus capsids to improve their physical or chemical properties for a given application. Directed evolution has been shown to generate MS2 capsids with different morphology (Asensio et al., 2016), pH sensitivity (Hartman et al., 2018) as well as permissibility to site-specific modification (Brauer et al., 2019). Viral vectors for use in gene delivery have been evolved into immune-orthologous variants to evade immune system recognition during repeat delivery (Moreno et al., 2019). In a nonviral protein cage, both the capsid protein and the cargo have been co-evolved to improve capsid stability and encapsulation of a nucleic acid cargo (Tetter et al., 2021). These examples provide inspiration for refining protein-encapsulating VLPs tailored to specific conditions or applications and point to a general approach applicable to the various platforms discussed in this review.

AUTHOR CONTRIBUTIONS

Donna McNeale: Conceptualization (equal); writing – original draft (lead); writing – review and editing (equal).
Noor Dashti: Writing – original draft (supporting); writing – review and editing (supporting). **Li Chen Cheah:**

Writing – original draft (supporting); writing – review and editing (supporting). **Frank Sainsbury**: Conceptualization (lead); writing – original draft (supporting); writing – review and editing (equal).

ACKNOWLEDGMENTS

This work was carried out on the traditional lands of the Yugarabul, Yuggera, Jagera, and Turrbal peoples. Frank Sainsbury acknowledges support from the Commonwealth Scientific and Industrial Research Organisation (CSIRO) in the form of a Synthetic Biology Future Science Platform Fellowship. Donna McNeale and Li Chen Cheah were supported by CSIRO Synbio PhD Top-up Scholarships. Open access publishing facilitated by Griffith University, as part of the Wiley - Griffith University agreement via the Council of Australian University Librarians.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Donna McNeale  <https://orcid.org/0000-0002-3342-3216>

Noor Dashti  <https://orcid.org/0000-0002-9195-8286>

Li Chen Cheah  <https://orcid.org/0000-0002-0072-2885>

Frank Sainsbury  <https://orcid.org/0000-0001-8152-3820>

RELATED WIREs ARTICLES

[Viral chemistry: the chemical functionalization of viral architectures to create new technology](#)

[Virus-based nanoparticles as platform technologies for modern vaccines](#)

[Plant viral and bacteriophage delivery of nucleic acid therapeutics](#)

[Physical, chemical, and synthetic virology: Reprogramming viruses as controllable nanodevices](#)

REFERENCES

- Abbing, A., Blaschke, U. K., Grein, S., Kretschmar, M., Stark, C. M. B., Thies, M. J. W., Walter, J., Weigand, M., Woith, D. C., Hess, J., & Reiser, C. O. A. (2004). Efficient intracellular delivery of a protein and a low molecular weight substance via recombinant polyomavirus-like particles. *The Journal of Biological Chemistry*, 279, 27410–27421.
- Asensio, M. A., Morella, N. M., Jakobson, C. M., Hartman, E. C., Glasgow, J. E., Sankaran, B., Zwart, P. H., & Tullman-Ercek, D. (2016). A selection for assembly reveals that a single amino acid mutant of the bacteriophage MS2 coat protein forms a smaller virus-like particle. *Nano Letters*, 16, 5944–5950.
- Ashley, C. E., Carnes, E. C., Phillips, G. K., Durfee, P. N., Buley, M. D., Lino, C. A., Padilla, D. P., Phillips, B., Carter, M. B., Willman, C. L., Brinker, C. J., do Carmo Caldeira, J., Chackerian, B., Wharton, W., & Peabody, D. S. (2011). Cell-specific delivery of diverse cargos by bacteriophage MS2 virus-like particles. *American Chemical Society Nano*, 5(7), 5729–5745.
- Belval, L., Hemmer, C., Sauter, C., Reinbold, C., Fauny, J. D., Berthold, F., Ackerer, L., Schmitt-Keichinger, C., Lemaire, O., Demangeat, G., & Ritzenthaler, C. (2016). Display of whole proteins on inner and outer surfaces of grapevine fanleaf virus-like particles. *Plant Biotechnology Journal*, 14, 2288–2299.
- Beterams, G., Böttcher, B., & Nassal, M. (2000). Packaging of up to 240 subunits of a 17 kDa nuclease into the interior of recombinant hepatitis B virus capsids. *FEBS Letters*, 481, 169–176.
- Bhaskar, S., & Lim, S. (2017). Engineering protein nanocages as carriers for biomedical applications. *Nature Publishing Group Asia Materials*, 9, e371.
- Bouřa, E., Liebl, D., Špiřek, R., Frič, J., Marek, M., Štokrová, J., Holáň, V., & Forstová, J. (2005). Polyomavirus EGFP-pseudocapsids: Analysis of model particles for introduction of proteins and peptides into mammalian cells. *Federation of European Biochemical Societies Letters*, 579, 6549–6558.
- Brasch, M., Putri, R. M., de Ruiter, M. V., Luque, D., Koay, M. S., Caston, J. R., & Cornelissen, J. J. (2017). Assembling enzymatic cascade pathways inside virus-based nanocages using dual-tasking nucleic acid tags. *Journal of the American Chemical Society*, 139, 1512–1519.
- Brauer, D. D., Hartman, E. C., Bader, D. L. V., Merz, Z. N., Tullman-Ercek, D., & Francis, M. B. (2019). Systematic engineering of a protein nanocage for high-yield, site-specific modification. *Journal of the American Chemical Society*, 141, 3875–3884.
- Brillault, L., Jutras, P. V., Dashti, N., Thuenemann, E. C., Morgan, G., Lomonosoff, G. P., Landsberg, M. J., & Sainsbury, F. (2017). Engineering recombinant virus-like nanoparticles from plants for cellular delivery. *American Chemical Society Nano*, 11, 3476–3484.
- Catrice, E. V. B., & Sainsbury, F. (2015). Assembly and purification of polyomavirus-like particles from plants (Report). *Molecular Biotechnology*, 57, 904–913.

- Charpillienne, A., Nejmeddine, M., Berois, M., Cohen, J., Trugnan, G., Parez, N., Neumann, E., & Hewat, E. (2001). Individual rotavirus-like particles containing 120 molecules of fluorescent protein are visible in living cells. *Journal of Biological Chemistry*, 276, 29361–29367.
- Chauhan, K., Hernandez-Meza, J. M., Rodriguez-Hernandez, A. G., Juarez-Moreno, K., Sengar, P., & Vazquez-Duhalt, R. (2018). Multi-functionalized biocatalytic P22 nanoreactor for combinatory treatment of ER plus breast cancer. *Journal of Nanobiotechnology*, 16, 17. <https://doi.org/10.1186/s12951-018-0345-2>
- Cheah, L. C., Stark, T., Adamson, L. S. R., Abidin, R. S., Lau, Y. H., Sainsbury, F., & Vickers, C. E. (2021). Artificial self-assembling nanocompartment for organizing metabolic pathways in yeast. *American Chemical Society Synthetic Biology*, 10, 3251–3263.
- Chen, X. S., Stehle, T., & Harrison, S. C. (1998). Interaction of polyomavirus internal protein VP2 with the major capsid protein VP1 and implications for participation of VP2 in viral entry. *The European Molecular Biology Organization Journal*, 17, 3233–3240.
- Chen, Z., Li, N., Li, S., Dharmarwardana, M., Schlimme, A., & Gassensmith, J. J. (2016). Viral chemistry: The chemical functionalization of viral architectures to create new technology. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 8, 512–534.
- Chung, Y. H., Cai, H., & Steinmetz, N. F. (2020). Viral nanoparticles for drug delivery, imaging, immunotherapy, and theranostic applications. *Advanced Drug Delivery Reviews*, 156, 214–235.
- Cigler, P., Lytton-Jean, A. K. R., Anderson, D. G., Finn, M. G., & Park, S. Y. (2010). DNA-controlled assembly of a NaTl lattice structure from gold nanoparticles and protein nanoparticles. *Nature Materials*, 9, 918–922.
- Comellas-Aragonès, M., Engelkamp, H., Claessen, V. I., Sommerdijk, N. A. J. M., Rowan, A. E., Christianen, P. C. M., Maan, J. C., Verduin, B. J. M., Cornelissen, J. J. L. M., & Nolte, R. J. M. (2007). A virus-based single-enzyme nanoreactor. *Nature Nanotechnology*, 2, 635–639.
- Cortes-Perez, N. G., Sapin, C., Jaffrelo, L., Daou, S., Grill, J. P., Langella, P., Seksik, P., Beaugerie, L., Chwetzoff, S., & Trugnan, G. (2010). Rotavirus-like particles: A novel nanocarrier for the gut. *Journal of Biomedicine and Biotechnology*, 2010, 317545.
- Das, S., Zhao, L., Crooke, S. N., Tran, L., Bhattacharya, S., Gaucher, E. A., & Finn, M. G. (2020). Stabilization of near-infrared fluorescent proteins by packaging in virus-like particles. *Biomacromolecules*, 21, 2432–2439.
- Das, S., Zhao, L., Eloffson, K., & Finn, M. G. (2020). Enzyme stabilization by virus-like particles. *Biochemistry*, 59, 2870–2881.
- Dashti, N. H., Abidin, R. S., & Sainsbury, F. (2018). Programmable in vitro coencapsidation of guest proteins for intracellular delivery by virus-like particles. *American Chemical Society Nano*, 12, 4615–4623.
- Edwardson, T. G. W., & Hilvert, D. (2019). Virus-inspired function in engineered protein cages. *Journal of the American Chemical Society*, 141, 9432–9443.
- Edwardson, T. G. W., Lévassieur, M. D., Tetter, S., Steinauer, A., Hori, M., & Hilvert, D. (2022). Protein cages: From fundamentals to advanced applications. *Chemical Reviews*, 122, 9145–9197.
- Eriksson, M., Andreasson, K., Weidmann, J., Lundberg, K., Tegerstedt, K., Dalianis, T., & Ramqvist, T. (2011). Murine polyomavirus virus-like particles carrying full-length human PSA protect BALB/c mice from outgrowth of a PSA expressing tumor. *PLoS One*, 6, e23828.
- Esquirol, L., McNeale, D., Douglas, T., Vickers, C. E., & Sainsbury, F. (2022). Rapid assembly and prototyping of biocatalytic virus-like particle nanoreactors. *American Chemical Society Synthetic Biology*, 11, 2709–2718.
- Fiedler, J. D., Brown, S. D., Lau, J. L., & Finn, M. G. (2010). RNA-directed packaging of enzymes within virus-like particles. *Angewandte Chemie International Edition*, 49, 9648–9651.
- Fiedler, J. D., Fishman, M. R., Brown, S. D., Lau, J., & Finn, M. G. (2018). Multifunctional enzyme packaging and catalysis in the Qbeta protein nanoparticle. *Biomacromolecules*, 19, 3945–3957.
- Giessen, T. W., & Silver, P. A. (2016). A catalytic nanoreactor based on in vivo encapsulation of multiple enzymes in an engineered protein nanocompartment. *ChemBiochem*, 17, 1931–1935.
- Gilbert, L., Toivola, J., Lehtomäki, E., Donaldson, L., Käpylä, P., Vuento, M., & Oker-Blom, C. (2004). Assembly of fluorescent chimeric virus-like particles of canine parvovirus in insect cells. *Biochemical and Biophysical Research Communications*, 313, 878–887.
- Glasgow, J., & Tullman-Ercek, D. (2014). Production and applications of engineered viral capsids. *Applied Microbiology and Biotechnology*, 98, 5847–5858.
- Glasgow, J. E., Capehart, S. L., Francis, M. B., & Tullman-Ercek, D. (2012). Osmolyte-mediated encapsulation of proteins inside MS2 viral capsids. *ACS Nano*, 6, 8658–8664.
- Hartman, E. C., Jakobson, C. M., Favor, A. H., Lobba, M. J., Álvarez-Benedicto, E., Francis, M. B., & Tullman-Ercek, D. (2018). Quantitative characterization of all single amino acid variants of a viral capsid-based drug delivery vehicle. *Nature Communications*, 9, 1385.
- Herbert, F. C., Brohlin, O. R., Galbraith, T., Benjamin, C., Reyes, C. A., Luzuriaga, M. A., Shahrivarkevishahi, A., & Gassensmith, J. J. (2020). Supramolecular encapsulation of small-ultrared fluorescent proteins in virus-like nanoparticles for noninvasive in vivo imaging agents. *Bioconjugate Chemistry*, 31, 1529–1536.
- Hong, Y.-R., & Black, L. W. (1993). Protein folding studies in vivo with a bacteriophage T4 expression-packaging-processing vector that delivers encapsidated fusion proteins into bacteria. *Virology*, 194, 481–490.
- Ingham, R. J., Colwill, K., Howard, C., Dettwiler, S., Lim, C. S. H., Yu, J., Hersi, K., Raaijmakers, J., Gish, G., Mbamalu, G., Taylor, L., Yeung, B., Vassilovski, G., Amin, M., Chen, F., Matskova, L., Winberg, G., Ernberg, I., Linding, R., ... Pawson, T. (2005). WW domains provide a platform for the assembly of multiprotein networks. *Molecular and Cellular Biology*, 25, 7092–7106.
- Inoue, T., Kawano, M.-A., Takahashi, R.-U., Tsukamoto, H., Enomoto, T., Imai, T., Kataoka, K., & Handa, H. (2008). Engineering of SV40-based nano-capsules for delivery of heterologous proteins as fusions with the minor capsid proteins VP2/3. *Journal of Biotechnology*, 134, 181–192.

- Jiménez-Zaragoza, M., Yubero, M. P., Martín-Forero, E., Castón, J. R., Reguera, D., Luque, D., de Pablo, P. J., & Rodríguez, J. M. (2018). Biophysical properties of single rotavirus particles account for the functions of protein shells in a multilayered virus. *eLife*, 7, e37295.
- Jordan, P. C., Patterson, D. P., Saboda, K. N., Edwards, E. J., Miettinen, H. M., Basu, G., Thielges, M. C., & Douglas, T. (2016). Self-assembling biomolecular catalysts for hydrogen production. *Nature Chemistry*, 8, 179–185.
- Kar, A. K., Iwatani, N., & Roy, P. (2005). Assembly and intracellular localization of the bluetongue virus core protein VP3. *The Journal of Virology*, 79, 11487–11495.
- Kostiainen, M. A., Kasyutich, O., Cornelissen, J. J. L. M., & Nolte, R. J. M. (2010). Self-assembly and optically triggered disassembly of hierarchical dendron-virus complexes. *Nature Chemistry*, 2, 394–399.
- Le, D. T., & Müller, K. M. (2021). In vitro assembly of virus-like particles and their applications. *Life*, 11, 334.
- Lee, K. L., Twyman, R. M., Fiering, S., & Steinmetz, N. F. (2016). Virus-based nanoparticles as platform technologies for modern vaccines. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 8, 554–578.
- Lee, K. W., & Tan, W. S. (2008). Recombinant hepatitis B virus core particles: Association, dissociation and encapsidation of green fluorescent protein. *Journal of Virological Methods*, 151, 172–180.
- Liljestrom, V., Mikkila, J., & Kostiainen, M. A. (2014). Self-assembly and modular functionalization of three-dimensional crystals from oppositely charged proteins. *Nature Communications*, 5, 4445.
- Lu, X., Thompson, J. R., & Perry, K. L. (2012). Encapsidation of DNA, a protein and a fluorophore into virus-like particles by the capsid protein of cucumber mosaic virus. *Journal of General Virology*, 93, 1120–1126.
- Lua, L. H. L., Connors, N. K., Sainsbury, F., Chuan, Y. P., Wibowo, N., & Middelberg, A. P. J. (2014). Bioengineering virus-like particles as vaccines. *Biotechnology and Bioengineering*, 111, 425–440.
- Luque, D., Rivas, G., Alfonso, C., Carrascosa, J. L., Rodriguez, J. F., & Caston, J. R. (2009). Infectious bursal disease virus is an icosahedral polyloid dsRNA virus. *Proceedings of the National Academy of Sciences*, 106, 2148–2152.
- Mazmanian, S. K., Hung, I. T., & Schneewind, O. (2001). Sortase-catalysed anchoring of surface proteins to the cell wall of *Staphylococcus aureus*. *Molecular Microbiology*, 40, 1049–1057.
- Minten, I. J., Claessen, V. I., Blank, K., Rowan, A. E., Nolte, R. J. M., & Cornelissen, J. J. L. M. (2011). Catalytic capsids: The art of confinement. *Chemical Science*, 2, 358–362.
- Minten, I. J., Hendriks, L. J. A., Nolte, R. J. M., & Cornelissen, J. J. L. M. (2009). Controlled encapsulation of multiple proteins in virus capsids. *Journal of the American Chemical Society*, 131, 17771–17773.
- Moreno, A. M., Palmer, N., Alemán, F., Chen, G., Pla, A., Jiang, N., Leong Chew, W., Law, M., & Mali, P. (2019). Immune-orthogonal orthologues of AAV capsids and of Cas9 circumvent the immune response to the administration of gene therapy. *Nature Biomedical Engineering*, 3, 806–816.
- Mullaney, J. M., & Black, L. W. (1996). Capsid targeting sequence targets foreign proteins into bacteriophage T4 and permits proteolytic processing. *Journal of Molecular Biology*, 261, 372–385.
- Mullaney, J. M., & Black, L. W. (1998). Activity of foreign proteins targeted within the bacteriophage T4 head and prohead: Implications for packaged DNA structure. *Journal of Molecular Biology*, 283, 913–929.
- Ohtake, N., Niikura, K., Suzuki, T., Nagakawa, K., Mikuni, S., Matsuo, Y., Kinjo, M., Sawa, H., & Ijiro, K. (2010). Low pH-triggered model drug molecule release from virus-like particles. *Chembiochem*, 11, 959–962.
- O’Neil, A., Prevelige, P. E., Basu, G., & Douglas, T. (2012). Coconfinement of fluorescent proteins: Spatially enforced communication of GFP and mCherry encapsulated within the P22 capsid. *Biomacromolecules*, 13, 3902–3907.
- O’Neil, A., Reichhardt, C., Johnson, B., Prevelige, P. E., & Douglas, T. (2011). Genetically programmed in vivo packaging of protein cargo and its controlled release from bacteriophage P22. *Angewandte Chemie International Edition*, 50, 7425–7428.
- Pascual, E., Mata, C. P., Gómez-Blanco, J., Moreno, N., Bárcena, J., Blanco, E., Rodríguez-Frandsen, A., Nieto, A., Carrascosa, J. L., & Castón, J. R. (2015). Structural basis for the development of avian virus capsids that display influenza virus proteins and induce protective immunity. *Journal of Virology*, 89, 2563–2574.
- Patterson, D., Prevelige, P., & Douglas, T. (2012). Nanoreactors by programmed enzyme encapsulation inside the capsid of the bacteriophage P22. *American Chemical Society Nano*, 6, 5000–5009.
- Patterson, D. P., LaFrance, B., & Douglas, T. (2013). Rescuing recombinant proteins by sequestration into the P22 VLP. *Chemical Communications*, 49, 10412–10414.
- Patterson, D. P., Rynda-Apple, A., Harmsen, A. L., Harmsen, A. G., & Douglas, T. (2013). Biomimetic antigenic nanoparticles elicit controlled protective immune response to influenza. *American Chemical Society Nano*, 7, 3036–3044.
- Patterson, D. P., Schwarz, B., Waters, R. S., Gedeon, T., & Douglas, T. (2014). Encapsulation of an enzyme cascade within the bacteriophage P22 virus-like particle. *ACS Chemical Biology*, 9, 359–365.
- Pickett, G. G., & Peabody, D. S. (1993). Encapsidation of heterologous RNAs by bacteriophage-Ms2 coat protein. *Nucleic Acids Research*, 21, 4621–4626.
- Plummer, E. M., & Manchester, M. (2011). Viral nanoparticles and virus-like particles: Platforms for contemporary vaccine design. *Wiley Interdisciplinary Reviews. Nanomedicine and Nanobiotechnology*, 3, 174–196.
- Qazi, S., Miettinen, H. M., Wilkinson, R. A., McCoy, K., Douglas, T., & Wiedenheft, B. (2016). Programmed self-assembly of an active P22-Cas9 nanocarrier system. *Molecular Pharmaceutics*, 13, 1191–1196.
- Rhee, J.-K., Hovlid, M., Fiedler, J. D., Brown, S. D., Manzenrieder, F., Kitagishi, H., Nycholat, C., Paulson, J. C., & Finn, M. G. (2011). Colorful virus-like particles: Fluorescent protein packaging by the Q β capsid. *Biomacromolecules*, 12, 3977–3981.

- Rurup, W. F., Verbij, F., Koay, M. S. T., Blum, C., Subramaniam, V., & Cornelissen, J. J. L. M. (2014). Predicting the loading of virus-like particles with fluorescent proteins. *Biomacromolecules*, *15*, 558–563.
- Sainsbury, F. (2020). Innovation in plant-based transient protein expression for infectious disease prevention and preparedness. *Current Opinion in Biotechnology*, *61*, 110–115.
- Sanchez-Sanchez, L., Cadena-Nava, R. D., Palomares, L. A., Ruiz-Garcia, J., Koay, M. S., Cornelissen, J. J., & Vazquez-Duhalt, R. (2014). Chemotherapy pro-drug activation by biocatalytic virus-like nanoparticles containing cytochrome P450. *Enzyme and Microbial Technology*, *60*, 24–31.
- Sanchez-Sanchez, L., Tapia-Moreno, A., Juarez-Moreno, K., Patterson, D. P., Cadena-Nava, R. D., Douglas, T., & Vazquez-Duhalt, R. (2015). Design of a VLP-nanovehicle for CYP450 enzymatic activity delivery. *Journal of Nanobiotechnology*, *13*, 66.
- Schmidt, U., Guenther, C., Rudolph, R., & Boehm, G. (2001). Protein and peptide delivery via engineered polyomavirus-like particles. *FASEB Journal*, *15*, 1646–1648.
- Schoonen, L., Pille, J., Borrmann, A., Nolte, R. J., & van Hest, J. C. (2015). Sortase A-mediated N-terminal modification of cowpea chlorotic mottle virus for highly efficient cargo loading. *Bioconjugate Chemistry*, *26*, 2429–2434.
- Schoonen, L., & van Hest, J. C. (2014). Functionalization of protein-based nanocages for drug delivery applications. *Nanoscale*, *6*, 7124–7141.
- Schwarz, B., Madden, P., Avera, J., Gordon, B., Larson, K., Miettinen, H. M., Uchida, M., LaFrance, B., Basu, G., Rynda-Apple, A., & Douglas, T. (2015). Symmetry controlled, genetic presentation of bioactive proteins on the P22 virus-like particle using an external decoration protein. *American Chemical Society Nano*, *9*, 9134–9147.
- Schwarz, B., Morabito, K. M., Ruckwardt, T. J., Patterson, D. P., Avera, J., Miettinen, H. M., Graham, B. S., & Douglas, T. (2016). Viruslike particles encapsidating respiratory syncytial virus M and M2 proteins induce robust T cell responses. *American Chemical Society Biomaterials Science & Engineering*, *2*, 2324–2332.
- Selivanovitch, E., Koliyatt, R., & Douglas, T. (2019). Chemically induced morphogenesis of P22 virus-like particles by the surfactant sodium dodecyl sulfate. *Biomacromolecules*, *20*, 389–400.
- Selivanovitch, E., Uchida, M., Lee, B., & Douglas, T. (2021). Substrate partitioning into protein macromolecular frameworks for enhanced catalytic turnover. *American Chemical Society Nano*, *15*, 15687–15699.
- Sharma, J., & Douglas, T. (2020). Tuning the catalytic properties of P22 nanoreactors through compositional control. *Nanoscale*, *12*, 336–346.
- Sharma, J., Uchida, M., Miettinen, H. M., & Douglas, T. (2017). Modular interior loading and exterior decoration of a virus-like particle. *Nanoscale*, *9*, 10420–10430.
- Steinmetz, N. F., Lim, S., & Sainsbury, F. (2020). Protein cages and virus-like particles: From fundamental insight to biomimetic therapeutics. *Biomaterials Science*, *8*, 2771–2777.
- Stupka, I., & Hedde, J. G. (2020). Artificial protein cages—Inspiration, construction, and observation. *Current Opinion in Structural Biology*, *64*, 66–73.
- Sun, J., DuFort, C., Daniel, M.-C., Murali, A., Chen, C., Gopinath, K., Stein, B., De, M., Rotello, V. M., Holzenburg, A., Cheng Kao, C., & Dragnea, B. (2007). Core-controlled polymorphism in virus-like particles. *Proceedings of the National Academy of Sciences*, *104*, 1354–1359.
- Szyska, T. N., Jenner, E. N., Tasneem, N., & Lau, Y. H. (2022). Molecular display on protein nanocompartments: Design strategies and systems applications. *ChemSystemsChem*, *4*, e202100025.
- Tapia-Moreno, A., Juarez-Moreno, K., Gonzalez-Davis, O., Cadena-Nava, R. D., & Vazquez-Duhalt, R. (2017). Biocatalytic virus capsid as nanovehicle for enzymatic activation of Tamoxifen in tumor cells. *Biotechnology Journal*, *12*, 1600706.
- Tegerstedt, K., Franzén, A., Ramqvist, T., & Dalianis, T. (2007). Dendritic cells loaded with polyomavirus VP1/VP2Her2 virus-like particles efficiently prevent outgrowth of a Her2/neu expressing tumor. *Cancer Immunology, Immunotherapy*, *56*, 1335–1344.
- Tegerstedt, K., Lindencrona, J. A., Curcio, C., Andreasson, K., Tullus, C., Forni, G., Dalianis, T., Kiessling, R., & Ramqvist, T. (2005). A single vaccination with polyomavirus VP1/VP2Her2 virus-like particles prevents outgrowth of HER-2/neu-expressing tumors. *Cancer Research*, *65*, 5953–5957.
- Tetter, S., Terasaka, N., Steinauer, A., Bingham, R. J., Clark, S., Scott, A. J. P., Patel, N., Leibundgut, M., Wroblewski, E., Ban, N., Stockley, P. G., Twarock, R., & Hilvert, D. (2021). Evolution of a virus-like architecture and packaging mechanism in a repurposed bacterial protein. *Science*, *372*, 1220–1224.
- Thuenemann, E. C., Le, D. H. T., Lomonosoff, G. P., & Steinmetz, N. F. (2021). Bluetongue virus particles as nanoreactors for enzyme delivery and cancer therapy. *Molecular Pharmaceutics*, *18*, 1150–1156.
- Thuenemann, E. C., Meyers, A. E., Verwey, J., Rybicki, E. P., & Lomonosoff, G. P. (2013). A method for rapid production of heteromultimeric protein complexes in plants: Assembly of protective bluetongue virus-like particles. *Plant Biotechnology Journal*, *11*, 839–846.
- Truebestein, L., & Leonard, T. A. (2016). Coiled-coils: The long and short of it. *BioEssays*, *38*, 903–916.
- Twarock, R., Bingham, R. J., Dykeman, E. C., & Stockley, P. G. (2018). A modelling paradigm for RNA virus assembly. *Current Opinion in Virology*, *31*, 74–81.
- Uchida, M., LaFrance, B., Broomell, C. C., Prevelige, P. E., & Douglas, T. (2015). Higher order assembly of virus-like particles (VLPs) mediated by multi-valent protein linkers. *Small*, *11*, 1562–1570.
- Uchida, M., McCoy, K., Fukuto, M., Yang, L., Yoshimura, H., Miettinen, H. M., LaFrance, B., Patterson, D. P., Schwarz, B., Karty, J. A., Prevelige, P. E., Jr., Lee, B., & Douglas, T. (2018). Modular self-assembly of protein cage lattices for multistep catalysis. *ACS Nano*, *12*, 942–953.

- Waghvani, H. K., Uchida, M., Fu, C. Y., LaFrance, B., Sharma, J., McCoy, K., & Douglas, T. (2020). Virus-like particles (VLPs) as a platform for hierarchical compartmentalization. *Biomacromolecules*, *21*, 2060–2072.
- Wang, Y., Uchida, M., Waghvani, H. K., & Douglas, T. (2020). Synthetic virus-like particles for glutathione biosynthesis. *ACS Synthetic Biology*, *9*, 3298–3310.
- Wen, A. M., & Steinmetz, N. F. (2016). Design of virus-based nanomaterials for medicine, biotechnology, and energy. *Chemical Society Reviews*, *45*, 4074–4126.
- Wilkerson, J. W., Yang, S. O., Funk, P. J., Stanley, S. K., & Bundy, B. C. (2018). Nanoreactors: Strategies to encapsulate enzyme biocatalysts in virus-like particles. *New Biotechnology*, *44*, 59–63.
- Windram, O. P., Weber, B., Jaffer, M. A., Rybicki, E. P., Shepherd, D. N., & Varsani, A. (2008). An investigation into the use of human papillomavirus type 16 virus-like particles as a delivery vector system for foreign proteins: N- and C-terminal fusion of GFP to the L1 and L2 capsid proteins. *Archives of Virology*, *153*, 585–589.
- Wingfield, P. T., Stahl, S. J., Williams, R. W., & Steven, A. C. (1995). Hepatitis core antigen produced in *Escherichia coli*: Subunit composition, conformational analysis, and in vitro capsid assembly. *Biochemistry*, *34*, 4919–4932.
- Witherell, G. W., & Uhlenbeck, O. C. (1989). Specific RNA binding by Q beta coat protein. *Biochemistry*, *28*, 71–76.
- Wu, M., Brown, W. L., & Stockley, P. G. (1995). Cell-specific delivery of bacteriophage-encapsidated ricin A chain. *Bioconjugate Chemistry*, *6*, 587–595.
- Yoshimura, H., Edwards, E., Uchida, M., McCoy, K., Roychoudhury, R., Schwarz, B., Patterson, D., & Douglas, T. (2016). Two-dimensional crystallization of P22 virus-like particles. *The Journal of Physical Chemistry B*, *120*, 5938–5944.
- Zakeri, B., Fierer, J. O., Celik, E., Chittock, E. C., Schwarz-Linek, U., Moy, V. T., & Howarth, M. (2012). Peptide tag forming a rapid covalent bond to a protein, through engineering a bacterial adhesin. *Proceedings of the National Academy of Sciences United States of America*, *109*, E690–E697.

How to cite this article: McNeale, D., Dashti, N., Cheah, L. C., & Sainsbury, F. (2022). Protein cargo encapsulation by virus-like particles: Strategies and applications. *WIREs Nanomedicine and Nanobiotechnology*, e1869. <https://doi.org/10.1002/wnan.1869>