

Showcasing discussion on the protein networks maturing human Fe–S proteins by Simone Ciofi-Baffoni, Lucia Banci and colleagues from the University of Florence, Italy.

Protein networks in the maturation of human iron–sulfur proteins

Despite the large amount of data reporting on the function of the human proteins involved in Fe–S protein biogenesis, how these proteins work at the molecular level remains poorly defined. Our review surveys the literature on Fe–S protein maturation processes in humans with the specific goal of providing a molecular picture of the currently known protein–protein interaction networks. From this picture, it emerges that future studies defining interactions among the “Fe–S players” at atomic or near-atomic resolution will be needed to understand how Fe–S protein machineries work. Image designed by Simone Ciofi-Baffoni and realized by Veronica Nasta.

As featured in:



See Simone Ciofi-Baffoni, Veronica Nasta and Lucia Banci, *Metallomics*, 2018, 10, 49.



Cite this: *Metallomics*, 2018, 10, 49

Protein networks in the maturation of human iron–sulfur proteins

Simone Ciofi-Baffoni, *^{ab} Veronica Nasta ^{ab} and Lucia Banci *^{ab}

The biogenesis of iron–sulfur (Fe–S) proteins in humans is a multistage process occurring in different cellular compartments. The mitochondrial iron–sulfur cluster (ISC) assembly machinery composed of at least 17 proteins assembles mitochondrial Fe–S proteins. A cytosolic iron–sulfur assembly (CIA) machinery composed of at least 13 proteins has been more recently identified and shown to be responsible for the Fe–S cluster incorporation into cytosolic and nuclear Fe–S proteins. Cytosolic and nuclear Fe–S protein maturation requires not only the CIA machinery, but also the components of the mitochondrial ISC assembly machinery. An ISC export machinery, composed of a protein transporter located in the mitochondrial inner membrane, has been proposed to act in mediating the export process of a still unknown component that is required for the CIA machinery. Several functional and molecular aspects of the protein networks operative in the three machineries are still largely obscure. This Review focuses on the Fe–S protein maturation processes in humans with the specific aim of providing a molecular picture of the currently known protein–protein interaction networks. The human ISC and CIA machineries are presented, and the ISC export machinery is discussed with respect to possible molecules being the substrates of the mitochondrial protein transporter.

Received 19th September 2017,
Accepted 24th November 2017

DOI: 10.1039/c7mt00269f

rsc.li/metallomics

^a *Magnetic Resonance Center-CERM, University of Florence, Via Luigi Sacconi 6, 50019, Sesto Fiorentino, Florence, Italy. E-mail: ciofi@cerm.unifi.it, banci@cerm.unifi.it; Fax: +39 055 4574923; Tel: +39 055 4574192, +39 055 4574273*

^b *Department of Chemistry, University of Florence, Via della Lastruccia 3, 50019 Sesto Fiorentino, Florence, Italy*

Introduction

Metal ions are essential for life, with more than one third of proteins encoded by the human genome depending on metals for their function.^{1–3} Nature has built several pathways responsible for metal handling and metalloprotein maturation in human cells. A large portion of metalloproteins themselves in the



Simone Ciofi-Baffoni

Simone Ciofi Baffoni, born in 1971, graduated in Chemistry in 1996 and received his PhD in Chemistry in 2000 at the University of Florence. He is now an Associate Professor at the University of Florence and has been since 2012. His research interest is focused on the molecular characterization of cellular pathways responsible for metal trafficking and metalloprotein maturation, through a multidisciplinary approach that

integrates structural, interaction and dynamical information. His current research activities are largely devoted to the investigation of the molecular mechanisms at the basis of iron–sulfur protein biogenesis in humans.



Veronica Nasta

Veronica Nasta graduated in Pharmaceutical Chemistry and Technology at the University of Naples “Federico II” in 2013, and received her PhD in Structural Biology at CERM (Magnetic Resonance Center), the University of Florence, in 2016. She is now a postdoctoral fellow in Prof. Banci’s group at the University of Florence. Her research interest is focused on the structural and functional characterization of metalloproteins through a variety of biophysical

and spectroscopic techniques, with an emphasis on protein structure and dynamics through NMR spectroscopy. Her current research focuses on the elucidation of the molecular mechanisms of the Fe–S protein maturation pathway.

human proteome are involved in pathways dedicated to maintaining the delicate balance between metal uptake and usage, as essential nutrients, and the excretion of excess metals thus preventing any potential damage.^{4–6} Altered or lacking regulation of these cellular processes can lead to cell toxicity and death.^{7–9} Despite the extensive progress in the area of “metals in biology”, we still have only a partial understanding of: (i) how metalloproteins acquire the correct metals/metal cofactors, (ii) how cells establish and maintain the required metal levels, (iii) how the maturation of a metalloprotein occurs, and (iv) how these three aspects intersect each other and with other cellular processes. A molecular understanding of the cellular pathways related to all these processes needs to be thoroughly investigated through the identification and description of the individual components, of their modes of interaction to form networks, and of their interlinks among other cellular pathways. This knowledge provides an inescapable step to obtain a complete and realistic description of the cellular processes related to metals, and is particularly relevant because several human diseases, including a number of neurodegenerative diseases, cardiovascular disorders and certain types of cancer, have been associated with the malfunction of metal-based cellular mechanisms.^{10–14} A molecular description of the pathways responsible for obtaining the mature, active forms of metalloproteins thus represents an essential step in understanding the origin of the related human diseases and in developing effective treatments.

Extensive studies in the past few decades have addressed, at a molecular level, the cellular pathways related to copper homeostasis and copper enzyme maturation in human cells.^{5,15–20} Those studies focused on the investigation of the structural and dynamic properties of proteins involved in copper transport, of their modes of interaction and of the factors driving copper transfer between protein partners along cellular routes.^{21,22}



Lucia Banci

Lucia Banci is a Professor of Chemistry at the University of Florence. She has extensive expertise and has provided original contributions and breakthroughs in Structural Biology and in biological NMR. She has addressed and unraveled many aspects of the biology of metal ions in biological systems. The innovative in cell NMR approach developed by Lucia Banci and her group allows for the detection of human individual proteins in living human cells with

atomic level resolution. She also exploited the extensive knowledge of structural biology approaches through NMR expertise to develop an absolutely innovative approach to vaccine design, based on the knowledge of the structure of the pathogen antigens and of the interaction patterns with antibodies, to design structure-based vaccines.

All these studies contributed to a reliable picture of the copper molecular systems biology in humans. Iron handling in human cells is by far much more complex than that of copper. The human iron proteome is, indeed, larger than that of copper, reflecting its extensive usage in human physiological processes,^{23–25} and, as a further degree of complexity with respect to copper, iron is physiologically present in various forms, *i.e.* either as individual metal ions or as bound to cofactors, mainly hemes and iron–sulfur (Fe–S) clusters. The latter are inorganic cofactors that bind to protein ligands, which, in the majority of the cases, are cysteines. Common Fe–S clusters include a rhomboid cluster composed of two iron ions and two inorganic sulfide ions ([2Fe–2S]),²⁶ and a cubane composed of four iron and four sulfide ions ([4Fe–4S]).²⁷ The [2Fe–2S] cluster can function as the building block to form [4Fe–4S] clusters. The capability of two rhombic [2Fe–2S] clusters to fuse to form a cubane [4Fe–4S] cluster has been, indeed, documented *in vitro*²⁸ and *in vivo*.²⁹ Much larger and more complex Fe–S clusters are found in enzymes such as hydrogenase and nitrogenase, and often also contain other metals, such as molybdenum.³⁰ The biosynthesis of such complex Fe–S clusters requires specific maturation machinery for their proper synthesis and insertion into the enzyme.³¹

Fe–S clusters are present in all three kingdoms of life. The most common function of Fe–S clusters is based on their ability to accept or donate single electrons to carry out complex enzymatic reactions. This feature differentiates them from non-metal-containing prosthetic groups, such as NAD⁺, FMN or FAD, which are less versatile and are limited to accepting electron pairs once involved in a redox enzymatic reaction cycle. Moreover, the fact that electron transfer can occur without undergoing energetically costly protein reorganization³² favours Fe–S proteins to be responsible for fast electron transfer processes between different metal centres in large protein complexes. For example, in complex I of the mitochondrial respiratory chain, eight Fe–S clusters are positioned close to each other in a quasi-linear arrangement throughout the protein section that extends into the mitochondrial matrix.³³ The increasing reduction potentials along the eight Fe–S clusters drive the electron transfer direction from NADH, a strong electron donor in the mitochondrial matrix, to ubiquinone, an electron acceptor and a carrier in the inner mitochondrial membrane.

The function of Fe–S clusters is not limited to electron transfer. Another well-characterized function of Fe–S clusters is enzymatic catalysis; a classic example being aconitase, in which a non-protein-coordinated Fe at one edge of a [4Fe–4S] cluster serves as a Lewis acid to assist H₂O abstraction from citrate (the substrate), which is converted to isocitrate. Other well-known Fe–S enzymes are biotin synthase and lipoate synthase, which bind two Fe–S clusters each. One of these clusters is disassembled during the formation of the products biotin and lipoic acid, respectively, thus serving as a sulfur donor.³⁴ Numerous other catalytic functions are known for eukaryotic Fe–S enzymes involved in metabolism.³⁵ However, in many cases, the precise role of the Fe–S cluster is still unclear, and it is therefore possible that in some proteins the Fe–S cluster simply plays a structural role.

A third general role of Fe–S clusters is in sensing environmental or intracellular conditions to regulate gene expression.³⁶ The best characterized example in mammals is provided by the cytosolic iron regulatory protein 1 (IRP1, also known as ACO1). Under iron-replete conditions, IRP1 holds a [4Fe–4S] cluster and functions as an aconitase. When the protein loses its labile cluster under iron deprivation, it can bind to stem-loop structures (termed iron-responsive elements (IREs)) in certain messenger RNAs of proteins involved in iron uptake, storage and distribution in the cell.³⁷ Binding of apo-IRP1 to 5'-located IREs blocks translation by inhibiting ribosome scanning to the start AUG codon, whereas the association with 3'-located IREs protects the mRNAs from nucleolytic degradation, leading to increased translation.

A sequence-based bioinformatic approach³⁸ has been applied to the human genome to identify the Fe–S proteome. This search mapped 70 unique genes (*i.e.* 0.35% of the human genes), with 61% of Fe–S proteins binding [4Fe–4S] clusters, 39% of them binding [2Fe–2S] clusters and only one (*i.e.* succinate dehydrogenase) binding a [3Fe–4S] cluster.³⁹ Experimental verification of their ability to bind an iron–sulfur cofactor is available for most of these proteins either directly or by similarity to some close homologue. From this analysis it also emerges that the number of mitochondrial Fe–S proteins that bind a [2Fe–2S] cluster is very close to that of those binding the [4Fe–4S] cofactor.³⁹ Instead, there is a higher number of [4Fe–4S] binding proteins in the cytoplasm, with the ratio of the [4Fe–4S]/[2Fe–2S] cluster-binding proteins being close to 2:1. Nuclear Fe–S proteins are almost exclusively of the [4Fe–4S] type, with the above ratio nearly 5:1.³⁹ Finally, the distribution of the identified human Fe–S proteins within cell compartments shows that the mitochondrial proteins are inherited from prokaryotic proteins of aerobes, whereas cytosolic and nuclear Fe–S proteins are inherited from anaerobic organisms.⁴⁰

Initially, researchers thought that the Fe–S clusters assemble spontaneously *in vivo*, given the fact that chemists are able to synthesize and to incorporate into proteins various Fe–S clusters *in vitro* using elemental iron and sulfur components as building blocks.²⁸ However, the discovery of a bacterial operon (nitrogen fixation (*nif*)) that contains genes associated with the synthesis of the Fe–S clusters of nitrogenase in the bacterial plant symbiont *Azotobacter vinelandii*⁴¹ revolutionized the studies on cellular Fe–S cluster biogenesis. In bacterial DNA, the genes that function in a pathway are often grouped together in an operon that facilitates the co-expression of functionally related proteins. This is the case of the *nif* operon of *A. vinelandii* and of two related operons of *Escherichia coli*, the iron–sulfur cluster (*isc*) and the sulfur formation (*suf*) operons.^{26,42} In bacteria, the Fe–S protein biogenesis machineries were thus straightforwardly identified and showed how Fe–S clusters are synthesized and then provided to client proteins in a single cellular compartment, *i.e.* the cytoplasm. In mammalian cells, the synthesis and distribution of Fe–S clusters are more complex. Mitochondria are the major sites containing Fe–S clusters; the twelve Fe–S clusters of the respiratory chain complexes I–III and the Fe–S cluster of aconitase, a component of the citric acid

cycle, are synthesized and incorporated into their respective mitochondrial proteins by the iron–sulfur cluster (ISC) assembly machinery.^{43,44} More recently, several cytosolic and nuclear proteins have been recognized as Fe–S proteins, including DNA primases, polymerases, and glycosylases, involved in base excision repair, and helicases, involved in the maintenance of genome stability.^{45,46} A cytosolic iron–sulfur assembly (CIA) machinery has been identified and shown to be responsible for the Fe–S cluster incorporation into cytosolic and nuclear Fe–S proteins.⁴⁷ From the studies of yeast and human cells,^{48–51} it has been shown that their maturation, however, requires not only the CIA machinery, but also components of the mitochondrial ISC assembly machinery. From this finding, it was proposed that a sulfur-containing compound of still unknown identity (X–S) is exported from the mitochondria to the cytoplasm for use in the cytosolic and nuclear compartments of eukaryotic cells.^{52–54} An ISC export machinery has been proposed to act in mediating the export process of the X–S compound.⁵¹ The export mechanism is based on a protein transporter located in the mitochondrial inner membrane, which has been shown to have a key role for the maturation of the cytosolic and nuclear proteins.⁴⁸ However, whether the mitochondrial and cytosolic Fe–S cluster assembly pathways are truly connected in mammals is still debated, and the paradigm that Fe–S clusters are assembled solely in the mitochondrial matrix does not necessarily extrapolate to mammalian cells.^{55,56} Indeed, a small amount of extra-mitochondrial isoforms of several components of the ISC assembly machinery responsible for assembling Fe–S clusters have been identified,^{57–61} and there is functional evidence that one of these components in its cytosolic isoform functions as a scaffold and a source for extra-mitochondrial Fe–S clusters in mammalian cells.⁵⁹

This Review focuses on the Fe–S protein maturation processes in humans with the specific aim of providing a molecular picture of the currently known protein–protein interaction networks. The human ISC and CIA machineries are presented, and the ISC export machinery is discussed with respect to possible molecules being substrates of the mitochondrial protein transporter.

The mitochondrial iron–sulfur cluster (ISC) assembly machinery

The mitochondrial ISC assembly machinery was inherited from endosymbiotic bacteria during evolution and is highly conserved in all eukaryotes.^{62,63} It is the crucial cellular component required for the maturation of mitochondrial and cytosolic/nuclear Fe–S proteins.⁵¹ In humans, this process encompasses 17 known proteins (Table 1), but the way in which they cooperate is still not well understood. The overall process can be dissected into three major functional steps: (i) the assembly of an initial [2Fe–2S] cluster on a scaffold protein, ISCU2; (ii) a chaperone-assisted [2Fe–2S] cluster transfer from ISCU2 to glutaredoxin-5 (GLRX5), which works as a [2Fe–2S] chaperone transferring the [2Fe–2S] clusters to several possible acceptors; and (iii) the assembly of a [4Fe–4S] cluster followed by its

Table 1 Components of the mitochondrial ISC assembly machinery in humans

Short name	Full name	Functional cofactors	Proposed main function
ISC assembly factors involved in <i>de novo</i> Fe-S cluster synthesis			
ISCU2	Iron-sulfur cluster assembly enzyme 2	[2Fe-2S]	Scaffold protein
NFS1	Cysteine desulfurase	PLP	Sulfur donor
ISD11	Iron-sulfur protein biogenesis desulfurase-interacting protein 11	—	Stabilizer of NFS1
ACP	Acyl carrier protein	<i>S</i> -dodecanoyl-4'-PPT	Metabolic sensor
FXN	Frataxin	Fe ²⁺	Iron donor, regulator of NFS1
FDX	Ferredoxin	[2Fe-2S]	Electron transfer
FDXR	Ferredoxin reductase	FAD	Electron transfer
ISC factors involved in cluster transfer			
GRP75	75 kDa glucose-regulated protein	ATP	Fe-S cluster transfer
HSCB	Iron-sulfur cluster co-chaperone protein	—	Fe-S cluster transfer
GRPEL1/2	GrpE protein homolog 1/2	—	Nucleotide exchange
GLRX5	Glutaredoxin-5	[2Fe-2S], GSH	Fe-S cluster transfer
Late-acting ISC factors involved in assembly and the insertion of Fe-S clusters into the target proteins			
ISCA1	Iron-sulfur cluster assembly 1 homolog	[2Fe-2S], [4Fe-4S]	[4Fe-4S] cluster assembly
ISCA2	Iron-sulfur cluster assembly 2 homolog	[2Fe-2S], [4Fe-4S]	[4Fe-4S] cluster assembly
IBA57	Iron-sulfur cluster assembly factor	—	[4Fe-4S] cluster assembly
NFU1	Iron-sulfur cluster scaffold	[4Fe-4S]	Dedicated ISC targeting factor
BOLA3	BoLA-like protein 3	—	Dedicated ISC targeting factor
NUBPL	Nucleotide-binding protein-like	[4Fe-4S]	Dedicated ISC targeting factor

target-selective insertion into mitochondrial [4Fe-4S] target proteins.^{51,64,65}

Assembly of a [2Fe-2S] cluster on the scaffold protein ISCU2

In humans, seven proteins act in the first step of the mitochondrial ISC assembly machinery (Table 1), five of them (ISCU2, NFS1, ISD11, ACP and FXN) forming a stable complex devoted to forming a [2Fe-2S]²⁺ cluster by assembling two Fe²⁺ ions and two sulfide ions (Fig. 1). Ferrous ions are imported into the mitochondrion by the intermembrane transporters mitoferrin-1 and mitoferrin-2 (MFRN1/2) to form a bioavailable iron pool in the mitochondrial matrix (Fig. 1), in which ferrous ions are believed to be bound to the still unidentified ligands.⁶⁶ A recent approach based on liquid chromatography coupled with inductively coupled plasma mass spectrometry was used to analyse non-proteinaceous low-molecular-mass metal complexes in mitochondria.⁶⁷ It was found that three iron complexes with masses of 580, 1100 and ~1500 Da constitute the mitochondrial labile iron pool in mammals.⁶⁸ This iron pool is used by the scaffold protein ISCU2 to assemble a [2Fe-2S] cluster (Fig. 1).⁶⁹ It has been proposed that frataxin is responsible of supplying the iron to ISCU2, but concrete *in vivo* evidence of frataxin supplying iron ions is still lacking.⁷⁰ It was also shown that the role of frataxin can go well beyond that of an iron transporter by working as a regulator of Fe-S cluster assembly.⁷¹ The origin of the two sulfide ions required to build a [2Fe-2S] cluster is instead clearer. They are indeed provided by the conversion of two cysteines into alanines. This process is performed by a sub-complex formed by the cysteine desulfurase NFS1, ISD11 and ACP proteins (Fig. 1) and requires four electrons. Assuming that two electrons are supplied by the oxidation of two ferrous ions,⁷² two additional electrons from an external source are required to make a [2Fe-2S]²⁺ cluster. They are provided by an

electron donor chain formed by two other components acting in this first step, FDX and FDXR (Fig. 1).

The sub-complex formed by NFS1, ISD11 and ACP proteins is a symmetric dimer in solution (Fig. 2),⁷³ which is responsible for the conversion of a cysteine into alanine per each NFS1 subunit. The sulfur atom released from the cysteine is transiently bound as a persulfide group on a conserved cysteine residue of NFS1.^{74,75} NFS1 is formed by two domains, the larger one having a PLP binding site.⁷³ NFS1 is inactive on its own and needs further proteins for its functioning, *i.e.* ISD11 and ACP.^{57,76,77} ISD11, also known as LYRM4, belongs to the LYR protein family characterized by a Leu-Tyr-Arg (LYR) motif near the N-terminus. It has been shown that the LYR motif and the N-terminus are essential for the role of ISD11 in the formation of a stable and active complex with NFS1.⁷⁸ ISD11 is a small α -helical protein with three helix-forming regions (Fig. 2);⁷³ and it has been demonstrated that it stabilizes NFS1 by preventing its self-aggregation through hydrophobic interactions.⁷⁶ Recently, the mitochondrial acyl carrier protein (ACP) has been identified as part of this sub-complex, *i.e.* in the absence of ACP, the sub-complex is destabilized resulting in a profound depletion of Fe-S clusters throughout the cell.^{77,79,80} This role of ACP depends upon its covalently bound *S*-dodecanoyl-4'-phosphopantetheine (4'-PPT) to support maximal cysteine desulfurase activity.⁷⁷ Crystallographic and EM structures of the co-expressed human NFS1-ISD11 and the *E. coli* ACP (ACP_{ec} 44% identical to the human mitochondrial ACP) dimeric complex, also called the SDA_{ec} complex, have been solved⁷³ (Fig. 2). They are different from any previously determined prokaryotic cysteine desulfurase structures, identifying the bases for the differences in prokaryotic and eukaryotic cysteine desulfurase mechanisms. The SDA_{ec} complex shows an unprecedented architecture in which a pair of ISD11 subunits forms the dimeric core of the SDA_{ec} complex, explaining the critical role of ISD11 in eukaryotic assemblies.




Fig. 1 A model for the maturation of human mitochondrial and cytosolic/nuclear Fe-S proteins. The mitochondrial ISC assembly machinery can be dissected into three major steps. In the first step, a $[2\text{Fe}-2\text{S}]^{2+}$ cluster is assembled on the scaffold protein ISCU2. This reaction requires the complex formed by NFS1, ISD11, FXN and ACP, and the electron transfer chain composed by FDX and FDXR, which receives two electrons from NADPH. The conversion of two ($\times 2$) cysteines to alanines provides the two sulfur atoms required to assemble the $[2\text{Fe}-2\text{S}]^{2+}$ cluster. Two ferrous ions imported into the mitochondrion by the intermembrane transporters MFRN1/2 are used to assemble the $[2\text{Fe}-2\text{S}]^{2+}$ cluster. In the second step, the $[2\text{Fe}-2\text{S}]^{2+}$ cluster is released from ISCU2 to GLRX5, an event that is mediated by a dedicated chaperone system (GRP75, HSCB, and GRPEL1-2). The $[2\text{Fe}-2\text{S}]^{2+}$ cluster can then be inserted into mitochondrial $[2\text{Fe}-2\text{S}]$ target proteins. An alternative model suggests that ISCU2 can directly insert the $[2\text{Fe}-2\text{S}]^{2+}$ cluster on the mitochondrial $[2\text{Fe}-2\text{S}]$ target proteins without involving GLRX5, and that it can assemble and insert a $[4\text{Fe}-4\text{S}]^{2+}$ cluster into the mitochondrial $[4\text{Fe}-4\text{S}]$ target proteins (bold dashed lines). In the third step, ISCA1, ISCA2 and IBA57 proteins contribute to convert a $[2\text{Fe}-2\text{S}]^{2+}$ cluster into a $[4\text{Fe}-4\text{S}]^{2+}$ cluster, which then is inserted into the mitochondrial $[4\text{Fe}-4\text{S}]$ target proteins with the possible help of various ISC targeting factors, i.e. NFU1, NUBPL and BOLA3. $[[2\text{Fe}-2\text{S}] \text{BOLA1} \text{GLRX5}]$ and $[[2\text{Fe}-2\text{S}] \text{BOLA3} \text{GLRX5}]$ complexes can potentially play a role in the maturation of mitochondrial $[4\text{Fe}-4\text{S}]$ target proteins. The CIA machinery requires the mitochondrial ISC assembly machinery, which produces a still unidentified compound X-S that is exported from mitochondria by the ISC export machinery (composed of ABCB7 and GSH). This compound might be used to assemble $[2\text{Fe}-2\text{S}]^{2+}$ clusters on the GLRX3 homodimer (or the $[[2\text{Fe}-2\text{S}]^{2+} \text{GLRX3} \text{BOLA2}]_2$ heterotrimer). The $[2\text{Fe}-2\text{S}]^{2+}$ GLRX3₂ homodimers (and/or the $[[2\text{Fe}-2\text{S}]^{2+} \text{GLRX3} \text{BOLA2}]_2$ heterotrimers) insert their $[2\text{Fe}-2\text{S}]^{2+}$ clusters into cytosolic $[2\text{Fe}-2\text{S}]$ target proteins, such as Anamorsin, or, possibly, into the $[[\text{NUBP1} \text{NUBP2}]]$ heterotetrameric or heterodimeric complex to assemble a $[4\text{Fe}-4\text{S}]^{2+}$ cluster. In such a molecular model, the electron transfer chain composed of NADPH, NDOR1 and Anamorsin provides the two electrons required for the reductive coupling of two $[2\text{Fe}-2\text{S}]^{2+}$ clusters on the $[[\text{NUBP1} \text{NUBP2}]]$ complex. This complex then transfers the assembled $[4\text{Fe}-4\text{S}]^{2+}$ cluster to the carrier NARFL, which transfers it into cytosolic and nuclear $[4\text{Fe}-4\text{S}]$ target proteins. The latter process is assisted by the CIA targeting complex $[[\text{CIAO1} \text{CIA2B} \text{MMS19}]]$ or by the $[[\text{CIA2A} \text{CIAO1}]]$ complex, and by the $[[\text{YAE1D1} \text{ORAOV1}]]$ complex. Protein colour code: grey, Fe-S target proteins; yellow, scaffold proteins; blue, membrane carriers; green, proteins involved in electron transfer and sulfide production; pink, proteins involved in Fe-S cluster trafficking to assist Fe-S insertion/assembly processes; brown, unknown function.

The structure also reveals that the 4'-PPT-conjugated acyl-group of ACP occupies a hydrophobic pocket of ISD11, explaining the basis of ACP stabilization (Fig. 2).⁷³

The $[[\text{NFS1-ISD11-ACP}]]$ sub-complex can bind ISCU2 in the presence or absence of FXN.^{81,82} Binding of FXN to the sub-complex accelerates desulfurase activity⁸¹ and persulfide formation on NFS1, and is also thought to induce a conformational change in ISCU2 that enhances the transfer of sulfur from NFS1 to Cys 138 of ISCU2, the primary persulfide acceptor of ISCU2.⁸³⁻⁸⁵ The mechanism of sulfur transfer from the persulfide group on

the conserved cysteine residue of NFS1 to the Fe-S cluster assembly site of ISCU2 involves the flexible loop of NFS1 containing the invariant cysteine residue.^{81,86,87} Ferrous ions further stimulate the cysteine desulfurase activity.⁸¹

The human ISCU protein has two different isoforms, ISCU1 and ISCU2, with different subcellular locations. It has been found by immunofluorescence microscopy that ISCU1 is localized to the cytosol and nucleus, whereas ISCU2 is localized to the mitochondria, being part of the ISC assembly machinery.⁸⁸ Also human frataxin has two different isoforms, FXN⁴²⁻²¹⁰




Fig. 2 Structural overview of the early steps of the human mitochondrial iron-sulfur cluster (ISC) assembly machinery. The $[2\text{Fe}-2\text{S}]^{2+}$ cluster biosynthesis is performed by a protein complex composed of NFS1, ISD11, ACP, FXN and ISCU2. The structure of this complex has been solved in pieces: the crystal structure of the [NFS1 ISD11 ACP] complex (PDB ID 5USR) and the cryo-EM structure of the [NFS1 FXN ISCU2] complex (PDB ID 5KZ5). For the electron transfer chain composed of FDX and FDXR, the X-ray structure of FDX is available (PDB ID 2Y5C). For the subsequent step consisting of the chaperone-mediated transfer of the assembled $[2\text{Fe}-2\text{S}]^{2+}$ cluster from ISCU2 to GLRX5, structural information are available on some single players only: the crystal structures of the Zn-bound HSCB protein (PDB ID 3BVO), of the SBD (PDB ID 3N8E) and ATPase domains (PDB ID 4KBO) of GRP75, and of the homodimeric $[2\text{Fe}-2\text{S}]^{2+}$ GLRX5 (PDB ID 2WUL).

and FXN⁸¹⁻²¹⁰, with the 81-210 isoform being the most abundant species in normal individuals.⁸⁹ Both are generated and localized within the mitochondria,⁸⁹ and participate in the mitochondrial Fe-S cluster assembly using different modes of interaction with NFS1, ISD11 and ISCU2 as well as different iron binding capacities.⁹⁰ These isoforms are generated through the sequential cleavage of the FXN protein precursor (FXN 1-210) by the mitochondrial processing peptidase upon import of FXN 1-210 to the mitochondrial matrix.⁹¹⁻⁹³ As determined by size-exclusion chromatography of normal human cell extracts, the native state of FXN⁴²⁻²¹⁰ is a monomer-oligomer equilibrium, whereas the native state of FXN⁸¹⁻²¹⁰ is monomeric.⁹⁴ Complexes containing oligomeric FXN⁴²⁻²¹⁰, NFS1, ISD11 and ISCU2 could be isolated upon fractionation of normal human cell extracts by size-exclusion chromatography followed by co-immunoprecipitation,⁹⁴⁻⁹⁶ underscoring the stability of these complexes. In contrast, native FXN⁸¹⁻²¹⁰ was only recovered in fractions that contained no detectable NFS1 and only traces of ISCU2. However, a [FXN⁸¹⁻²¹⁰·NFS1-ISD11-ISCU2] complex has been isolated from bacterial cells upon the co-expression of all four components of the complex, and it could also be reconstituted *in vitro* from the purified [NFS1-ISD11-ISCU2] complex and the FXN⁸¹⁻²¹⁰ monomer. However, in both cases, a large excess of monomeric FXN⁸¹⁻²¹⁰ was required for complex formation,^{82,97} which may explain why this complex could not be detected under native conditions in human cells.^{94,95} On the basis of differentiated iron binding abilities of the two FXN isoforms under aerobic conditions,^{81,95,97} it has been proposed that, under physiological conditions, FXN⁸¹⁻²¹⁰ may bind low levels of iron and support basal levels of Fe-S cluster assembly *via* transient interactions with the [NFS1-ISD11-ISCU2] complex. While the stable [FXN⁴²⁻²¹⁰·NFS1-ISD11-ISCU2] complex may provide a mechanism to increase the rate of Fe-S cluster assembly when the demand exceeds the low iron-binding capacity of FXN⁸¹⁻²¹⁰.⁹⁰

Recently, isolation, biochemical and structural characterization of the human Fe-S cluster assembly complex containing most of the proteins of the first step of the mitochondrial ISC assembly machinery (ISCU2, NFS1, ISD11 and FXN⁴²⁻²¹⁰) has been reported (Fig. 2).⁹⁸ The structural data provided a first model of a coordinated mechanism for the transfer of iron and sulfur to ISCU2 for the synthesis of a $[2\text{Fe}-2\text{S}]^{2+}$ cluster. It was shown that these four human proteins, once co-expressed in *E. coli* cell extracts, form a stable, active complex with a 1:1:1:1 stoichiometry. This complex may also contribute to explain how human frataxin may serve as both the activator of NFS1 and the iron donor, an argument still largely debated in the literature.^{54,99} Indeed, in the context of the human [FXN⁸¹⁻²¹⁰·NFS1-ISD11-ISCU2] complex having the shorter FXN isoform,⁹⁷ one monomer of FXN⁸¹⁻²¹⁰ is thought to bind in a pocket between NFS1 and ISCU2 through the conserved frataxin iron-binding surface,^{97,100} in such a way to block the ability of FXN⁸¹⁻²¹⁰ to bind iron and thus contrasting with its role in iron delivery. In contrast, the location of ISCU2 relative to FXN⁴²⁻²¹⁰ and NFS1 in the human [FXN⁴²⁻²¹⁰·NFS1-ISD11-ISCU2] structure allows FXN⁴²⁻²¹⁰ and NFS1 to simultaneously bind to ISCU2 and to stimulate Fe-S cluster synthesis, consistent with the previous observation that ISCU2 and [NFS1-ISD11] can independently or simultaneously bind to oligomeric FXN⁴²⁻²¹⁰ in solution.¹⁰¹

The electron transfer chain formed by ferredoxin (FDX) and ferredoxin reductase (FDXR), which receives an electron from NADH, accomplishes the task of providing the two electrons required to form a $[2\text{Fe}-2\text{S}]^{2+}$ cluster on ISCU2 (Fig. 1).⁸⁰ In human mitochondria, there are two different isoforms of ferredoxins, FDX1 and FDX2. Their X-ray structures are very similar even if the two ferredoxins share only 33% protein sequence identity (Fig. 2).⁸⁰ By NMR spectroscopy, it has been demonstrated that both FDX1 and FDX2 interact directly with the [NFS1 ISD11 ACP] sub-complex and that their reduced forms, when added to

a solution of the complex and L-cysteine, become oxidized, indicating that both ferredoxins are able to donate electrons leading to the reduction of the sulfur atom. Despite the fact that both FDX isoforms interact with the [NFS1 ISD11 ACP] sub-complex, FDX2 exhibits both a higher binding affinity for the complex and a much higher efficiency in assisting the Fe-S cluster assembly. Functional data indicated that only FDX2 is part of the electron transport chain required for *de novo* mitochondrial Fe-S protein biogenesis, while FDX1 is essentially devoted to the steroid metabolism in the adrenal gland.¹⁰² In contrast to this view, the RNA interference-mediated depletion of either FDX1 or FDX2 was reported to impair Fe-S protein maturation.¹⁰³ An *in vitro* reconstitution system containing the yeast ISC members of the first step of the ISC assembly machinery showed that the yeast orthologue of FDX2, Yah1, in a reduced but not oxidized state tightly interacts with apo-Isu1 indicating a dynamic interaction between Yah1-apo-Isu1.¹⁰⁴ Nuclear magnetic resonance structural studies identify the Yah1-apo-Isu1 interaction surface and suggest a pathway for the electron flow from reduced Yah1 to Isu1.¹⁰⁴

A chaperone-assisted [2Fe-2S] cluster transfer from ISCU2 to glutaredoxin-5

In the second step, the *de novo* assembled [2Fe-2S]²⁺ cluster is released from the scaffold protein ISCU2 to the monothiol glutaredoxin GLRX5. This event is mediated by a dedicated chaperone system comprising the GRP75 chaperone (also named mortalin, HSPA9 or PBP74), which is a member of the ATP-dependent 70-kDa heat shock protein family Hsp70, its co-chaperone HSCB (also named HSC20), and the nucleotide exchange factors GRPEL1/2 (Table 1 and Fig. 1).¹⁰⁵ The actual molecular model is based on the initial binding of the HSCB protein to [2Fe-2S] ISCU2, which then forms a complex with the chaperone GRP75.¹⁰⁵⁻¹⁰⁷ NMR data indicated that a conformational change of ISCU2 occurs upon HSCB binding, which specifically weakens the [2Fe-2S] ISCU2 interaction in the [FXN-ACP-NFS1-ISD11-ISCU2] complex, and strengthens its interaction with the co-chaperone HSCB, which targets the [2Fe-2S] [ISCU2 HSCB] complex to the chaperone GRP75.¹⁰⁸ The C-terminal α -helical domain of the HSCB co-chaperone (Fig. 2) is directly responsible for binding ISCU2, with three highly conserved non-contiguous hydrophobic residues being of crucial importance for the HSCB-ISCU2 interaction.^{44,109-112} The N-terminal domain of HSCB (Fig. 2), which contains an invariant histidine, proline, and aspartate motif, is responsible for stimulating the ATPase activity of the GRP75 chaperone. Overall, the HSCB co-chaperone might have a dual function in stimulating the ATPase activity of the chaperone GRP75 and in promoting its recognition between ISCU2 and the chaperone GRP75. The N-terminus of the human HSCB co-chaperone shows distinctive features compared to the homologous co-chaperone proteins of bacteria and fungi (DnaJ type III), as it contains, downstream of the mitochondrial targeting sequence (residues 1-26), an additional domain, which harbors two CxxC stretches (C41/C44 and C58/C61) that were found to coordinate a zinc ion *in vitro* (Fig. 2).¹¹² The physiological relevance of the

unique N-terminal domain remains to be elucidated although it may facilitate protein dimerization.

As for all Hsp70 family chaperones, GRP75 is composed of two domains, the crystal structures of them being determined: a ~42 kDa N-terminal ATP-binding domain and a ~25 kDa substrate-binding domain (SBD) involved in the ISCU2 recognition (Fig. 2).¹¹³ The two domains of GRP75 do not interact with each other but are tethered by a short interdomain linker. The sequence of the ATP-binding domain is highly conserved in all Hsp70 family members, while the SBD has several differences among family members. The ADP dissociation step on GRP75 is facilitated by the nucleotide exchange factors GRPEL1/2. Such dissociation enables GRP75 to undergo the next round of Fe-S cluster transfer. The ATP-binding domain of GRP75 is found to interact with the nucleotide exchange factors.¹¹⁴

On the basis of studies on the yeast Grx5 homologue,¹¹⁵⁻¹¹⁷ and considering that the Grx5 function is conserved throughout the eukaryotic kingdom,¹¹⁸⁻¹²⁰ it has been proposed that GLRX5 mediates [2Fe-2S] cluster transfer from the [HSCB GRP75 [2Fe-2S] ISCU2] complex to [2Fe-2S] target proteins and to intermediate proteins responsible for the assembly of [4Fe-4S] clusters (Fig. 1). GLRX5 in solution binds a [2Fe-2S] cluster forming a [2Fe-2S]²⁺ cluster-bridged homodimer with two glutathione (GSH) molecules involved in cluster binding (Fig. 2).¹²¹ Gel filtration and ¹⁵N NMR relaxation data on the holo and apo states of GLRX5 showed that cluster binding induces a change in the quaternary structure from an apo-monomer to a holo dimer.¹²¹ The dimeric state of [2Fe-2S] GLRX5 prevents the cluster from being released in the presence of physiological concentrations of GSH, suggesting that GLRX5 can specifically transfer the cluster.

Recent studies have shown that the HSCB co-chaperone in the [HSCB GRP75 [2Fe-2S] ISCU2] complex can guide the selection of specific Fe-S recipient proteins for cluster delivery by binding to a conserved LYR motif present in specific proteins.¹²² In this regard, the best-characterized example is the succinate dehydrogenase subunit b (SDHB), which is the Fe-S cluster-containing subunit of the respiratory complex II.¹⁰⁶ This protein contains three independent LYR binding sites, all close to the cysteines that coordinate the three Fe-S clusters of SDHB and in a position where the binding of the chaperone/co-chaperone [2Fe-2S] transfer apparatus can guide the release of the cluster from [2Fe-2S]-ISCU2 into the Fe-S binding sites of SDHB. The LYR motif family is present only in eukaryotes (conserved domains accession: cl05087),¹²³ which lists other proteins related to the mitochondrial Fe-S protein biogenesis in humans, thus potentially being the HSCB-dependent recognition mechanism conserved in other Fe-S target insertion processes. One of the members of the LYR family is SDHAF1 (LYRM8), a succinate dehydrogenase complex assembly factor.¹²⁴ SDHAF1 was shown to recruit the [HSCB GRP75 ISCU2] complex to the C-terminus of SDHB through direct binding of its N-terminal LYR motif to the HSCB co-chaperone.¹²⁵ Another LYR protein is LYRM7, a complex III assembly factor,¹²⁶ which interacts with the [HSCB GRP75 ISCU2] complex,¹⁰⁶ suggesting that the LYRM7-HSCB interaction might guide the insertion of the [2Fe-2S] cluster into the Rieske Fe-S protein of complex III. LYR-containing subunits are also

present in the mitochondrial complex I, and one of these has been suggested to be required for the proper incorporation of the N2 [4Fe–4S] center and for the function of complex I.¹²³ A second consensus sequence, in addition to the LYR motif, is the KKX_{6–10}KK motif, which has been found to interact with the HSCB co-chaperone.¹⁰⁶ GLRX5 has a similar pattern of lysines at its C-terminus (K₁₃₉K₁₄₀X₁₀K₁₅₁K₁₅₂) and has been found to interact with HSCB *in vivo*.¹⁰⁶ All these findings suggest that the model in which GLRX5 is the central component for distributing [2Fe–2S] clusters to final [2Fe–2S] acceptors and to the proteins involved in the late steps of the mitochondrial ISC assembly machinery for maturing [4Fe–4S] target proteins might have to be revisited. Indeed, GLRX5 could be just one among the LYR and KKX_{6–10}KK targets of the [HSCB GRP75 [2Fe–2S] ISCU2] complex, which, in contrast, could be the real distributor of [2Fe–2S] clusters (Fig. 1). However, there are several pieces of evidence that this new possible model is not able to answer. Among them, how [4Fe–4S] clusters are formed and inserted into the mitochondrial target proteins is fully undefined by this model. It has been shown, indeed, that the mammalian Fe–S assembly complex (*i.e.* the [FXN·ACP·NFS1·ISD11·ISCU2] complex) is able to synthesize exclusively [2Fe–2S] clusters.¹²⁷ Therefore, the way that the [4Fe–4S] and [3Fe–4S] clusters of LYR-containing targets of the [HSC20 GRP75 [2Fe–2S] ISCU2] complex (*i.e.* mitochondrial respiratory complexes I–III) are formed, is still not addressed in this possible alternative model.

Reductive coupling of two [2Fe–2S]²⁺ clusters to form a [4Fe–4S]²⁺ cluster on ISCA proteins

Among the GLRX5 protein partners,^{121,128,129} two A-type Fe–S proteins named ISCA1 (iron–sulfur cluster assembly 1) and ISCA2 (iron–sulfur cluster assembly 2) are required for mitochondrial [4Fe–4S] protein maturation.^{130,131} ISCA1 and ISCA2

have been first proposed to function in concert with IBA57 to mature [4Fe–4S] proteins.¹³⁰ However, the molecular relation between IBA57 and ISCA proteins has been recently debated. Indeed, while ISCA1 and ISCA2 are protein partners *in vivo*, IBA57 has been found to interact *in vivo* with ISCA2 but not with ISCA1, in contrast to what occurs for the yeast homologues, which both interact with Iba57.¹³² Surprisingly, it has been recently observed that only ISCA1 is required for the maturation of mitochondrial [4Fe–4S] proteins.¹²⁸ A possible explanation for the differently observed ISCA2 phenotypes is that different mammalian cells and conditions were used in the two phenotyping experiments,^{128,130} and thus further *in vivo* data are required to resolve this matter. *In vitro* data^{133,134} support a model where a [ISCA1 ISCA2] complex converts [2Fe–2S]²⁺ clusters, received by GLRX5, into a [4Fe–4S]²⁺ cluster, with IBA57 not being required in this process (Fig. 1). In agreement with this model, it has been shown that GLRX5 can transfer a [2Fe–2S] cluster to a heterodimeric complex formed by human ISCA1 and ISCA2 through a cluster-mediated protein–protein interaction event,^{121,133} and that this complex acts as an assembler of a [4Fe–4S]²⁺ cluster through a reductive coupling process of two [2Fe–2S]²⁺–GLRX5 donated clusters (Fig. 3).¹³³ In the latter reaction, dithiothreitol provides the two electrons required for the reductive coupling process. How these electrons are provided *in vivo* is still an open question. Finally, although ISCA2 alone can also bind a [4Fe–4S] cluster in a dimeric state, it has been observed that the formation of a heterodimeric [ISCA1 ISCA2] complex is thermodynamically favoured compared to the corresponding homodimeric forms, indicating that, at the cellular level, the heterodimeric complex should be the preferential species responsible of assembling [4Fe–4S]²⁺ clusters. This is in full agreement with the strong, specific interaction between ISCA1 and ISCA2 observed *in vivo*.¹²⁸




Fig. 3 Structural overview of the late steps of the human mitochondrial iron–sulfur cluster (ISC) assembly machinery. In the late steps of the mitochondrial ISC assembly machinery, the [ISCA2 ISCA1] heterodimeric complex assembles a [4Fe–4S]²⁺ cluster by the reductive coupling of two [2Fe–2S]²⁺ clusters received by GLRX5.¹³³ No evidence on the functional role of IBA57, which is part of this process, is available, nor is the physiological donor of the two electrons required in the [2Fe–2S] coupling process. NUF1 and NUBPL assist in the maturation of specific [4Fe–4S] target proteins, but how they work mechanistically in such a process is still undefined. [2Fe–2S] GLRX5 by interacting with BOLA1 and BOLA3 can be converted into [[2Fe–2S] BOLA1 GLRX5] and [[2Fe–2S] BOLA3 GLRX5] complexes, respectively, but their potential role in the maturation of specific [4Fe–4S] target proteins is not defined (dotted lines). The solution NMR structures of the C-domain (PDB ID 2M5O) and the N-domain (PDB ID 2LTM) of apo-NUF1, BOLA1 (PDB ID 5LCI) and BOLA3 (PDB ID 2NCL) have been solved. Experimentally-driven docking models of the two heterodimeric [BOLAs GLRX5] complexes are available in ref. 145.

Recently, a molecular picture of how the [4Fe–4S] cluster has been assembled on the heterodimeric [ISCA1 ISCA2] complex has been provided.¹³⁴ In this model, the two fully conserved C-terminal cysteines, located in the unstructured and flexible C-terminal tail of the ISCA proteins,¹³³ are required to extract the [2Fe–2S]²⁺ cluster from GLRX5, while the other fully conserved cysteine of ISCA, Cys 79, is not essential for the cluster transfer step. This cysteine is required to promote the acquisition of the second [2Fe–2S]²⁺ cluster from another GLRX5 molecule *via* C-terminal cysteines of ISCA, in such a way that the reductive coupling between the two [2Fe–2S]²⁺ clusters can occur on the [ISCA1 ISCA2] complex. This molecular model agrees with the *in vivo* data on yeast, which showed that the three conserved cysteines of yeast Isa1 and Isa2 are essential for the maturation of [4Fe–4S] proteins.^{135,136}

Target-selective insertion of the assembled [4Fe–4S] cluster into mitochondrial [4Fe–4S] target proteins

The [4Fe–4S] cluster assembled on the [ISCA1 ISCA2] complex is inserted into mitochondrial [4Fe–4S] target proteins with the possible assistance of dedicated ISC targeting factors, *i.e.* NFU1, BOLA3 and NUBPL (also named IND1) (Table 1). Specifically, some mitochondrial [4Fe–4S] target proteins could acquire their [4Fe–4S] cluster directly from the [ISCA1 ISCA2] complex, whereas others may derive their clusters after a prior transfer of the [4Fe–4S] cluster from the [ISCA1 ISCA2] complex to NFU1 and NUBPL. The latter can bind a [4Fe–4S] cluster, which facilitates the insertion of the cluster into specific mitochondrial [4Fe–4S] target proteins (Fig. 1).^{137–140} In particular, NFU1 acts as a late-acting factor that transfers the [4Fe–4S] clusters to the lipoic acid synthase and to the subunits of respiratory complexes I and II, facilitating their maturation,^{137,138} while NUBPL works as an assembly factor for the human respiratory complex I only.¹⁴⁰ NFU1 was detected as an ISCA1 specific interacting partner only.¹²⁸ However, in yeast, Nfu1 physically interacts with both Isa1 and Isa2 in the [Isa1 Isa2 Iba57] complex and with mitochondrial target proteins that need [4Fe–4S] clusters to function.¹⁴¹ The yeast data suggest that human NFU1 might specifically bind to the mitochondrial [4Fe–4S] target proteins independent of the [ISCA1 ISCA2] complex. Then, the binding of this complex with the [ISCA1 ISCA2] complex may serve to recruit the NFU1-associated [4Fe–4S] target proteins to the [ISCA1 ISCA2] complex, specifically assisting cluster transfer and insertion into these target proteins. As other possible model, NFU1 was proposed to be an alternative scaffold to ISCU2, thanks to its ability to assemble [4Fe–4S] clusters and transfer them to apo-proteins, thus operating upstream of the [ISCA1 ISCA2] complex.⁵⁸ Human NFU1 comprises two domains: an N-terminal domain and a C-terminal domain, which contains the CXXC motif involved in [4Fe–4S] cluster binding (Fig. 3).^{58,142} While apo-NFU1 is largely monomeric in solution, [4Fe–4S] NFU1 consists of a trimer of dimers, with three [4Fe–4S] clusters bound by the two conserved cysteine residues in the CTDs of two NFU1 subunits forming the dimer.¹⁴² The functional relevance of the binding of three [4Fe–4S] clusters in a single aggregate is still unknown.

The phenotypic similarity between the pathogenic mutations of NFU1 and of another mitochondrial protein named BOLA3 suggests that the two proteins function together in the late steps of the mitochondrial Fe–S protein maturation pathway (Fig. 1).^{137,138} Using yeast as a model system, it has been proposed that the yeast ortholog of human BOLA3 functions in the late step of transfer of [4Fe–4S] clusters to specific target proteins, together with yeast Nfu1.¹⁴¹ Despite the fact that yeast BOLA3 and Nfu1 both act late in the mitochondrial Fe–S protein biogenesis, they appear to fulfil different tasks that cannot be taken over by the other protein.¹⁴³ Proteomic studies did not detect any stable *in vivo* interaction between yeast Nfu1 and BOLA3, suggesting a transient interaction in nature; *in vitro* this interaction has been found to have a μM affinity.¹⁴³ Overall, these results in yeast might suggest that BOLA3 might function with NFU1 in the cluster insertion into specific mitochondrial [4Fe–4S] target proteins (Fig. 1), but this mechanism is still largely hypothetical. BOLA3 belongs to the highly conserved BOLA-like protein family, which also includes the human mitochondrial BOLA1.¹⁴⁴ Recently, it has been shown in yeast that the yeast homolog BOLA1 also facilitates the insertion of [4Fe–4S] clusters into a subset of mitochondrial target proteins, being unable to replace Nfu1, similarly to what occurs with BOLA3 (Fig. 1).¹⁴³ Even though BOLA1 and BOLA3 play some overlapping roles, their functions are not identical.¹⁴³ The solution structures of BOLA3 and BOLA1 showed that they have a similar fold, but with local structural differences in the regions containing conserved sequence patterns comprising cluster ligands.¹⁴³ It has been demonstrated that both BOLA1 and BOLA3 interact with apo-GLRX5 with K_d values of $3 \mu\text{M}$ and are able to form [2Fe–2S] cluster-bridged dimeric hetero-complexes with GLRX5.^{141,143} The [[2Fe–2S] BOLA1 GLRX5] complex is preferentially formed over the [[2Fe–2S] BOLA3 GLRX5] complex, because of a higher cluster-binding affinity.¹⁴⁵ Experimentally driven structural models and spectroscopic data of these two heterodimeric holo complexes¹⁴⁵ (Fig. 3) showed that: (i) cluster solvent accessibility is higher in the BOLA3 hetero-complex with respect to that in the BOLA1 hetero-complex; (ii) the [BOLA1 GLRX5] complex stabilizes a reduced, Rieske-type [2Fe–2S]¹⁺ cluster, while an oxidized, ferredoxin-like [2Fe–2S]²⁺ cluster is present in the [BOLA3 GLRX5] complex; (iii) the [2Fe–2S] center in the [BOLA1 GLRX5] complex behaves like a stable redox active center able to perform electron transfer reactions, at variance with what occurs for the [[2Fe–2S] BOLA3 GLRX5] complex. This experimental evidence suggests that the [[2Fe–2S] BOLA1 GLRX5] complex might work in electron transfer reactions, while the [[2Fe–2S] BOLA3 GLRX5] complex might be involved in iron–sulfur cluster transfer processes *versus* target proteins along the Fe–S protein assembly pathway.¹⁴⁵ In conclusion, the molecular scenarios of NFU1 and BOLA3 (alone or complexed with GLRX5) proteins acting late in the ISC pathway are still largely undefined, as well as the role of BOLA1 (alone or complexed with GLRX5) as the possible further ISC late-acting factor is still indefinite.

The other known targeting factor, NUBPL, is another [4Fe–4S] cluster binding protein, specifically required for the respiratory complex I assembly (Fig. 3).¹⁴⁰ It is closely related in

sequence to two cytosolic P-loop NTPases, NUBP1 and NUBP2, which act as a heteromeric scaffold complex in the synthesis of cytosolic [4Fe–4S] clusters (see later). It contains a highly conserved nucleotide binding domain and a Fe–S binding CPXC motif, which is required for the function of NUBPL and is proposed to provide the ligands for a transiently bound [4Fe–4S] cluster, probably inducing protein dimerization.¹⁴⁰

The cytosolic iron–sulfur assembly (CIA) machinery

Up to now, thirteen CIA components participating in the maturation of human cytosolic and nuclear Fe–S proteins have been identified (Table 2).^{47,146,147} Most of these components are conserved in almost all eukaryotes.^{39,148} Six of these proteins act in the early stage of the CIA machinery, while all the others are late-acting CIA factors (Table 2).

The assembly of a [4Fe–4S] cluster in the early stage of the CIA machinery

The early stage of the CIA machinery involves two P-loop NTPase proteins, named NUBP1 and NUBP2 in humans (also known as NBP35 and CFD1, respectively). NUBP1 and NUBP2 are nuclear and cytosolic localized proteins containing several cysteine residues at their C- and N-termini.^{149,150} Sequence alignment of NUBP1 and NUBP2 with their respective eukaryotic homologues identified, in NUBP1, an invariant N-terminal Cx₁₃Cx₂Cx₅C motif, which is absent in the NUBP2 family, and, in NUBP2, an invariant C-terminal Cx₁₈Cx₂Cx₂C motif, which is partially conserved or totally absent in the NUBP1 family depending on the organism. In the yeast homologues, each of these two motifs binds a [4Fe–4S] cluster, such that the N-terminal domain is tightly bound, while that in the C-terminus is labile.^{151–153} No structural models of the cluster bound forms of NUBP1 and NUBP2 or of any eukaryotic homologues are available. This piece of

information would be fundamental to address definitively how the two proteins bind the [4Fe–4S] cluster(s). Co-immunoprecipitation shows that the two proteins interact in the cellular context, suggesting that NUBP1 and NUBP2 proteins cooperate in the same molecular process of the CIA machinery.¹⁴⁹ *In vitro* and *in vivo* studies on the yeast homologous heterocomplex suggest the presence of bridging [4Fe–4S] clusters in a heterodimeric or heterotetrameric structural arrangement and that this complex performs a scaffold function by assembling a [4Fe–4S] cluster on the C-terminal motif in the early stage of the CIA machinery (Fig. 4).^{152,154} This function should be conserved in humans. Indeed, depletion of NUBP1 by RNA interference in HeLa cells resulted in a severe defect in the assembly of cytosolic and nuclear Fe–S proteins without an apparent effect on mitochondrial Fe–S protein maturation.¹⁴⁹ However, how this heterocomplex works in the CIA machinery as a scaffold protein is unknown. The donors of Fe²⁺ and sulfide ions required to assemble a [4Fe–4S] cluster on the heterocomplex are not defined yet, nor are the cluster binding properties of the human heterocomplex well addressed. Since some early ISC components and the membrane transporter of the ISC export machinery are required for the functioning of the CIA machinery in yeast,^{48,50,155,156} it might be speculated that the sulfide ions for assembling the [4Fe–4S] cluster on the heterocomplex require the mitochondria-exported X–S compound.

The formation of the [4Fe–4S] cluster on the heterocomplex also requires electrons, and NADPH might provide these *via* the NDOR1 and Anamorsin CIA components (Fig. 1). Similar to the [NUBP1 NUBP2] complex, they operate indeed in the early stage of the CIA pathway by forming a stable complex.^{157,158} According to this proposal, a relatively weak, but specific interaction between NUBP1 and Anamorsin, compared to the much stronger interaction between NDOR1 and Anamorsin, has been detected from yeast-two-hybrid and surface plasmon resonance experiments in plants.¹⁵⁹ Although plants lack the NUBP2 gene, thus not fully representing a fully reliable model for the molecular

Table 2 Components of the CIA machinery in humans

Short name	Full name	Functional cofactors	Proposed main function
Early-acting CIA factors involved in Fe–S cluster assembly and transfer			
NUBP1	Nucleotide-binding protein 1	[4Fe–4S]	Scaffold protein
NUBP2	Nucleotide-binding protein 2	[4Fe–4S]	Scaffold protein
NDOR1	NADPH-dependent diflavin oxidoreductase 1	FAD, FMN	Electron transfer
Anamorsin	Anamorsin	[2Fe–2S], [4Fe–4S]	Electron transfer
GLRX3	Glutaredoxin-3	[2Fe–2S], GSH	Fe–S cluster transfer, iron trafficking
BOLA2	Bola-like protein 2	—	Fe–S cluster transfer
Late-acting CIA factors involved in transfer and insertion of Fe–S clusters into the target proteins			
NARFL	Fe–S cluster assembly factor	[4Fe–4S]	Mediates contact between the early and late parts of the CIA machinery
CIAO1	Cytosolic iron–sulfur protein assembly	—	CIA targeting complex
CIA2B	Mitotic spindle-associated MMXD complex subunit MIP18	—	CIA targeting complex
MMS19	Nucleotide excision repair protein homolog	—	CIA targeting complex
CIA2A	MIP18 family protein FAM96A	—	Specific maturation factor of IRP1
YAE1D1	Yae1 domain-containing protein 1	—	Specific maturation factor of the cytosolic ABCE1 protein
ORAOV1	Oral cancer-overexpressed protein 1	—	Specific maturation factor of the cytosolic ABCE1 protein




Fig. 4 Structural overview of the human cytosolic iron-sulfur assembly (CIA) machinery. The tetrameric [NUBP1 NUBP2] complex might receive electrons from the NADPH-NDOR1-Anamorsin electron transfer chain to assemble a $[4\text{Fe}-4\text{S}]^{2+}$ cluster on the C-terminus by reductively coupling two $[2\text{Fe}-2\text{S}]^{2+}$ clusters. These might be received by the holo forms of the GLRX3 and/or [BOLA2 GLRX3] complex. The assembled $[4\text{Fe}-4\text{S}]^{2+}$ cluster is transferred to NARFL, which then delivers it to cytosolic and nuclear target proteins with the contribution of several other CIA proteins. Three of them, CIAO1, MMS19 and CIA2B, form a complex (CIA targeting complex) assisting the transfer of the $[4\text{Fe}-4\text{S}]$ cluster from NARFL to the majority of cytosolic and nuclear target proteins. The complex between CIAO1 and CIA2A specifically matures the cytosolic aconitase/IRP1, while a complex between the CIA targeting complex and the [ORAOV1 YAE1D1] complex specifically assists the maturation of the ABCE1 protein. The crystal structures of the single domains of apo-GLRX3 (the Trx domain PDB ID 2WZ9, the GrxA domain PDB ID 3ZYW and the GrxB domain PDB ID 2YAN), of the FMN-binding domain of NDOR1 (PDB ID 4H2D) and of CIAO1 (PDB ID 3FMNO) have been solved. Two different crystal structures of CIA2A are available (PDB ID 3UX2 and 3UX3); only one of them is shown in the figure. A homology model of BOLA2 was generated based on the available solution structure of the BOLA-like protein from *Mus musculus* (PDB ID 1V9J, sequence identity is 87%). A structural model of the $[2\text{Fe}-2\text{S}]$ CIAPIN1 domain was calculated using diamagnetic and paramagnetic restraints,¹⁶⁶ and the solution structure of the N-terminal domain of Anamorsin (PDB ID 2LD4) has been solved.

events in the human CIA machinery, the plant NUBP1 protein has a scaffold function similar to that of the human [NUBP1 NUBP2] complex.¹⁶⁰

NDOR1 is a diflavin reductase consisting of two domains: the first binds FMN (FMN-binding domain, hereafter), and the second binds FAD and NADPH (FAD-binding domain, hereafter) (Fig. 4). On the basis of the well-known electron transfer mechanism occurring in diflavin reductase enzymes,¹⁶¹ the electrons are transferred from NADPH to FAD and then to FMN, which serves as a donor for one-electron terminal acceptors, *i.e.* the Fe-S cluster(s) of Anamorsin in this case. Anamorsin contains two domains: an N-terminal domain of 172 residues and a C-terminal domain of 90 residues, named cytokine-induced apoptosis inhibitor 1 (CIAPIN1 hereafter), containing two highly conserved, cysteine-rich motifs, $\text{CX}_8\text{CX}_2\text{CXC}$ and $\text{CX}_2\text{CX}_7\text{CX}_2\text{C}$ (Fig. 4). An unstructured and flexible linker of 51 residues connects the two domains (Fig. 4). While the N-terminal domain is well-structured, the C-terminal CIAPIN1 domain is largely unstructured (Fig. 4).¹⁶² The CIAPIN1 domain of Anamorsin binds a $[2\text{Fe}-2\text{S}]$ cluster in the first motif and a $[2\text{Fe}-2\text{S}]$ or $[4\text{Fe}-4\text{S}]$ cluster on the second motif depending on how the preparation of the protein sample has been conducted *in vitro*.¹⁶³⁻¹⁶⁵ To definitively define which is the physiologically relevant cluster bound to the second motif of Anamorsin, an *in vivo* approach would be required to define the cluster bound

state of Anamorsin in the cellular context. The crystal structure of the FMN-binding domain of NDOR1 showed the classical fold of FMN-binding domains of diflavin reductases (Fig. 4).¹⁶⁶ The crystal¹⁶⁷ and solution¹⁶² structures of the N-terminal domain of Anamorsin (Fig. 4) showed an overall structure resembling a typical *S*-adenosylmethionine (SAM)-dependent methyltransferase fold but lacking the glycine-rich motif, which is typically responsible for the binding of the SAM cofactor in methyltransferases. As a result, Anamorsin does not show SAM-dependent methyltransferase activity.¹⁶² On this basis, it has been hypothesized that the N-terminal domain of Anamorsin acts as bait for protein-protein interactions. A structural model of the CIAPIN1 domain shows that the $\text{CX}_8\text{CX}_2\text{CXC}$ and $\text{CX}_2\text{CX}_7\text{CX}_2\text{C}$ regions are connected by a stably formed α -helix and sample a restricted range of conformations that bring the two cluster-binding motifs close to each other (Fig. 4).¹⁶⁶ Anamorsin and NDOR1 form a stable complex, thanks to the specific protein-protein recognition between the FMN-binding domain of NDOR1 and the unstructured linker region of Anamorsin separating the N-terminal domain from the C-terminal CIAPIN1 domain (specifically, residues 185-223).¹⁶⁶ The N-terminal domain of Anamorsin is not involved in such a recognition process. The two protein partners interact permanently forming also a stable interaction in the cell,¹⁵⁷ and no dissociation occurs during the electron transfer process.¹⁶⁶ One electron transfer reaction was observed to occur from the hydroquinone

state of the FMN moiety of NDOR1 to the oxidized $[2\text{Fe}-2\text{S}]^{2+}$ cluster bound to the first motif of Anamorsin (N-terminal $[2\text{Fe}-2\text{S}]$ cluster), while the Fe-S cluster bound to the second motif of Anamorsin does not receive electrons from FMN.¹⁶³

The [Anamorsin NDOR1] complex might therefore supply electrons to the [NUBP1 NUBP2] complex, working similarly to the NADPH-FDX-FDXR electron transfer chain in the mitochondrial ISC assembly machinery.¹⁰⁴ However, at variance with the ISC assembly machinery, the specific molecular targets of the electron transfer flow generated by the [Anamorsin NDOR1] complex are not identified yet. Depending on the molecular nature of the mitochondria-exported X-S compound, the complex might reduce the sulfur moiety of the X-S compound to sulfide, similar to the function of the NADPH-FDX-FDXR electron transfer chain in the mitochondrial ISC system,^{155,156} with the function of building a $[2\text{Fe}-2\text{S}]$ cluster, and/or it could facilitate the fusion of two $[2\text{Fe}-2\text{S}]$ clusters to a $[4\text{Fe}-4\text{S}]$ cluster. The yeast homologues of NDOR1 and Anamorsin (named Tah18 and Dre2, respectively) have been shown not to be involved in the *de novo* formation of a $[2\text{Fe}-2\text{S}]$ cluster.¹⁵⁷ Assuming that the same holds in human cells, we could speculate that the electron transfer chain composed of NADPH-NDOR1-Anamorsin is involved in the fusion of two $[2\text{Fe}-2\text{S}]^{2+}$ clusters to a $[4\text{Fe}-4\text{S}]^{2+}$ cluster on the [NUBP1 NUBP2] complex. The X-S compound, which is needed for the CIA machinery function, should therefore operate upstream of the [NUBP1 NUBP2] complex, to build a $[2\text{Fe}-2\text{S}]$ cluster, and therefore an alternative cytosolic pathway for the *de novo* synthesis of $[2\text{Fe}-2\text{S}]$ clusters might be required. Some ISC components have been detected in small amounts in the cytosol and in the nucleus, being proposed to function in these compartments to assemble a $[2\text{Fe}-2\text{S}]$ cluster,^{55,59} which is then used by the CIA machinery for the maturation of cytosolic and nuclear $[4\text{Fe}-4\text{S}]$ proteins. However, evidence for a direct function of these extra-mitochondrial ISC proteins in Fe-S protein assembly is still under debate.^{50,151,168}

Cytosolic $[2\text{Fe}-2\text{S}]$ cluster trafficking by glutaredoxin-3 and BOLA2 protein

Another $[2\text{Fe}-2\text{S}]$ cluster protein operating in the early stage of the CIA machinery is the cytosolic monothiol glutaredoxin GLRX3 (Fig. 1).¹⁶⁹ GLRX3 consists of an N-terminal thioredoxin (Trx) domain that lacks a redox-active motif and two monothiol glutaredoxin (Grx) domains that both harbor a Cys-Gly-Phe-Ser active site.¹⁷⁰ Crystal structures of the three single domains (Fig. 4) showed that their overall fold is very similar to that of other known monothiol glutaredoxins/thioredoxins. Upon $[2\text{Fe}-2\text{S}]$ cluster binding, GLRX3 forms a dimeric complex bridging two $[2\text{Fe}-2\text{S}]$ clusters between each of the two Grx domains ($[2\text{Fe}-2\text{S}]_2$ GLRX3₂, Fig. 4).^{171,172} Each cluster is shared by the cysteine residue of the active site of each Grx domain and by a GSH molecule per each Grx domain (Fig. 4). Yeast has two homologous proteins of GLRX3, Grx3 and Grx4, which, similar to GLRX3, act in cytosolic/nuclear Fe-S protein maturation.¹⁷³ Yeast Grx3 and Grx4 contain an N-terminal Trx domain, similar to GLRX3, but only one Grx domain, which binds one $[2\text{Fe}-2\text{S}]$ cluster in a dimeric complex similarly to what occurs in GLRX3.¹⁷⁴

In yeast, the binding/assembly of the $[2\text{Fe}-2\text{S}]$ cluster in Grx3 requires the mitochondrial ISC assembly and export machineries, yet occurs independent of the two CIA proteins Nbp35 and Dre2.^{173,174} Conversely, Fe-S cluster binding on both Nar1 and Dre2 strictly depends on Grx3, placing Grx3 early in the CIA pathway. Applying this yeast model to humans, GLRX3 might be the primary target of the X-S compound after its exit from the mitochondria (Fig. 1). This compound might determine the formation of the $[2\text{Fe}-2\text{S}]$ cluster bound state of GLRX3, which might then act as a cytoplasmic $[2\text{Fe}-2\text{S}]$ cluster trafficking protein in the early phase of the CIA pathway (Fig. 1), in analogy to the proposed function of mitochondrial GLRX5. Now, the arising question is how GLRX3 is reconstituted with $[2\text{Fe}-2\text{S}]$ clusters. The crucial point to provide the answer is the identification of the chemical nature of the X-S compound.

The first evidence showing that GLRX3 can work as a $[2\text{Fe}-2\text{S}]$ cluster trafficking protein came from an *in vitro* study.¹⁷⁵ Indeed, it was shown that GLRX3 matures Anamorsin by transferring two $[2\text{Fe}-2\text{S}]$ clusters to the cysteine-rich motif of the CIAPIN1 domain (Fig. 1). Anamorsin and GLRX3 are protein partners in the cellular context as identified by the yeast-two-hybrid assay.¹⁷⁶ It was shown that the transfer mechanism was dependent on the formation of a specific protein-protein complex between the N-terminal domains of GLRX3 and of Anamorsin, with part of the linker of Anamorsin having a further, specific role in stabilizing the interaction.¹⁷⁵ Their interaction is the fundamental requisite to observe the transfer of both $[2\text{Fe}-2\text{S}]$ clusters from Grx domains of GLRX3 to the two cluster-binding motifs of the CIAPIN1 domain of Anamorsin. In conclusion, GLRX3 plays a key role in maturing Anamorsin, specifically working as an intracellular $[2\text{Fe}-2\text{S}]$ cluster-trafficking protein. This indirectly links GLRX3 to all Anamorsin-dependent cellular processes. *In vivo* data showed that silencing of human GLRX3 expression in HeLa cells decreased the activities of the cytosolic Fe-S proteins IRP1 and GPAT.¹⁶⁹ This behavior can be interpreted as an impairment of the GLRX3-dependent Anamorsin maturation process, making the CIA machinery unable to function, since it lacks the electron transfer chain required to form $[4\text{Fe}-4\text{S}]$ clusters on the [NUBP1 NUBP2] complex. Similarly, *in vivo* data in yeast¹⁷³ showed a decrease of Fe-S cluster insertion levels into cytosolic $[4\text{Fe}-4\text{S}]$ target proteins upon Grx4 depletion. Deficiency in yeast Grx3/4 also leads to the impairment of the cytosolic ribonucleotide reductase (RNR) enzyme despite cytosolic iron overload.¹⁷³ Similar to the CIA machinery, the maturation of the di-iron cofactor of RNR depends on the Dre2-Tah18 electron transfer chain,^{177,178} but iron loading into RNR does not require Dre2, while it requires Grx3/Grx4 proteins.¹⁷⁹ On this basis, an iron-trafficking function of cytosolic monothiol glutaredoxins can also be proposed. According to this view, the depletion of GLRX3 in humans and of Grx3 or Grx4 in yeast specifically also impaired the synthesis of heme.¹⁷³ However, direct evidence showing that monothiol Grx proteins are able to transfer iron ions from the bound $[2\text{Fe}-2\text{S}]$ clusters to target proteins is still missing, and such data would be required to validate the proposed iron-trafficking function of cytosolic Grx proteins.

GLRX3 forms a complex *in vitro* with the cytosolic protein BOLA2.^{171,180} Specifically, the *in vitro* studies showed that an apo-complex is formed, with each Grx domain of apo-GLRX3 binding a BOLA2 molecule to form a heterotrimeric complex (Fig. 1).¹⁸⁰ This apo-complex is able to bind two $[2\text{Fe}-2\text{S}]^{2+}$ clusters by chemical reconstitution, with each cluster being bridged between BOLA2 and a Grx domain of GLRX3 ($[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$, Fig. 4).¹⁷¹ The same cluster binding state (*i.e.* two $[2\text{Fe}-2\text{S}]$ clusters per trimer) is observed once the heterotrimeric complex was isolated by cells co-expressing the two BOLA2 and GLRX3 proteins.^{171,180} The $[\text{GLRX3 BOLA2}]$ complex was also observed in mammalian cells.¹⁸¹ This complex is, however, not as abundant as apo-form in mammalian cells, likely due to the low protein-protein affinity ($K_d = 25 \mu\text{M}$), but it accumulates upon $[2\text{Fe}-2\text{S}]$ cluster binding, which stabilizes its interaction.¹⁸¹ *In vitro* studies¹⁸⁰ showed that the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complex transfers both its $[2\text{Fe}-2\text{S}]^{2+}$ clusters to apo-Anamorsin producing its mature holo state (Fig. 1 and 4). The mechanism of cluster transfer relies on the interaction between the N-terminal domain of Anamorsin with that of GLRX3, similarly to what was observed in the cluster transfer from $[2\text{Fe}-2\text{S}]_2\text{GLRX3}_2$ to apo-Anamorsin. *In vivo* studies confirmed that this direct $[2\text{Fe}-2\text{S}]$ cluster transfer process occurs in mammalian cells upon complex formation between $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}]$ and $[\text{apo-Anamorsin NDOR1}]$ complexes.¹⁸¹ Collectively, the experimental data suggested that the heterotrimeric complex can work as a $[2\text{Fe}-2\text{S}]$ cluster transfer component in the CIA machinery, similarly to what the homodimeric $[2\text{Fe}-2\text{S}]_2\text{GLRX3}_2$ does (Fig. 1). Now, the arising question is whether and how the cells handle homodimeric $[2\text{Fe}-2\text{S}]_2\text{GLRX3}_2$ and heterotrimeric $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complexes, *i.e.* which is the predominant species acting as a $[2\text{Fe}-2\text{S}]$ cluster chaperone at the cellular level. *In vivo* data¹⁸¹ support a model in which the heterotrimeric $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complex is the species transferring the $[2\text{Fe}-2\text{S}]$ clusters under physiological conditions. However, we cannot exclude that, in human cells, the $[2\text{Fe}-2\text{S}]_2\text{GLRX3}_2$ homodimers transfer their own clusters to a number of target proteins larger than the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ hetero-complex and, in doing so, they do not accumulate to a detectable degree in cells. Consistent with this hypothesis, the bridged clusters in the GLRX3 homodimer were labile *in vitro*, much more than in the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complex,^{171,182} which, conversely, accumulates in human cells.¹⁸¹ In the hypothesis that the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complex works as the major species in the cytoplasm of human cells transferring the clusters to several target proteins, we would expect similar effects on the *in vivo* Fe-S cluster loading into several target proteins by deleting BOLA2 or GLRX3, besides Anamorsin. We believe that such *in vivo* studies would help in further elucidating the role of the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ complex as the cytoplasmic distributor of $[2\text{Fe}-2\text{S}]$ clusters. In such studies, alternative sources of Fe-S cluster loading into the target proteins might be operative and need to be taken into account. This is indeed what occurs for Anamorsin, where significant amounts of residual iron are retained in Anamorsin complexes in the cells lacking GLRX3 or BOLA2.¹⁸¹ The mitoNEET/miner1 family of

$[2\text{Fe}-2\text{S}]$ proteins has been proposed to be an alternative source of Anamorsin for acquiring $[2\text{Fe}-2\text{S}]$ clusters, since mitoNEET/miner1 proteins can transfer their $[2\text{Fe}-2\text{S}]$ clusters to apo-Anamorsin *in vitro*,¹⁸³ and human mitoNEET can transfer $[2\text{Fe}-2\text{S}]$ clusters to apo-proteins in the cells exposed to oxidative stress.¹⁸⁴⁻¹⁸⁶ This alternative source of Fe-S clusters could also account for the relatively milder phenotypes associated with the depletion of GLRX3 in mammalian cells *versus* yeast, which does not have mitoNEET homologues. However, the role of the mitoNEET/miner1 family in the CIA machinery is still controversial, since still there is no clear evidence from functional and interactomic data supporting the role of this protein family in cluster transfer under physiological conditions. In conclusion, the $[2\text{Fe}-2\text{S}]^{2+}\text{GLRX3}_2$ homodimer and/or the $[[2\text{Fe}-2\text{S}]_2\text{GLRX3 BOLA2}_2]$ heterotrimer might work as cytoplasmic $[2\text{Fe}-2\text{S}]$ chaperones inserting $[2\text{Fe}-2\text{S}]^{2+}$ clusters into cytosolic $[2\text{Fe}-2\text{S}]$ target proteins, such as Anamorsin, or into the $[\text{NUBP1 NUBP2}]$ heterotetrameric complex to assemble a $[4\text{Fe}-4\text{S}]^{2+}$ cluster (Fig. 1).

Target-selective insertion of the assembled $[4\text{Fe}-4\text{S}]$ cluster into cytosolic/nuclear $[4\text{Fe}-4\text{S}]$ target proteins

The late steps of the CIA machinery are quite complex and involve the majority of the CIA proteins. Several studies in the past few years investigated how these proteins work, all showing that the principal component of this process is a stable complex composed of three proteins, $[\text{CIAO1 MMS19 CIA2B}]$, commonly named the ternary CIA targeting complex (Fig. 1).^{187,188} MMS19 contains several HEAT repeats, which are commonly involved in protein-protein interactions. CIA2B (also named FAM96B or MIP18) and CIAO1 (also named CIA1) interact with the C-terminal HEAT repeats of MMS19 to form a docking site for the majority of cytosolic and nuclear $[4\text{Fe}-4\text{S}]$ target proteins.¹⁸⁹ CIA2B directly interacts with the C-terminal HEAT repeats of MMS19, while CIAO1 requires the presence of CIA2B for a stable interaction with MMS19. The direct interaction of MMS19 with the C-terminus of CIA2B protects CIA2B from proteosomal degradation.¹⁸⁹ A structural analysis of the CIA targeting complex will be of fundamental relevance for understanding whether the interaction between MMS19 and CIAO1 is solely mediated by CIA2B or whether CIAO1 displays a weak affinity for MMS19 that is greatly enhanced by CIA2B. Such structural investigation will also be able to clarify which domains/proteins are in direct physical contact with cytoplasmic and nuclear $[4\text{Fe}-4\text{S}]$ target proteins. The most straightforward model is that MMS19, CIA2B and CIAO1, jointly, form a docking site for the $[4\text{Fe}-4\text{S}]$ target proteins, but we cannot exclude that the formation of the ternary complex induces a conformational change in one of the proteins that exposes an interaction site for the $[4\text{Fe}-4\text{S}]$ target proteins, otherwise not accessible. The majority of the $[4\text{Fe}-4\text{S}]$ target proteins bind the CIA targeting complex only in the presence of all three proteins of the complex, *i.e.* MMS19, CIA2B and CIAO1.¹⁸⁹ The only exception is the transcription factor XPD, which binds to the N-terminal HEAT repeats of MMS19 independent of CIA2B and CIAO1.¹⁸⁹ However, it cannot be excluded that XPD features two

interaction sites with the CIA targeting complex: one with the N-terminus of MMS19 and one with the “canonical” [4Fe–4S] docking site at the C-terminus of MMS19 that depends on CIA2B and CIAO1. The only structure available among the proteins of the CIA targeting complex is the crystal structure of CIAO1, solved at a resolution of 1.7 Å, showing a β-propeller fold with seven pseudo-symmetrically orientated blades around a central axis (Fig. 4).¹⁹⁰ This doughnut-shaped geometry is highly conserved among the large family of WD40 proteins, which serve as docking sites in a multitude of cellular networks and mediate the molecular recognition of partner proteins mainly through the top and side surfaces.¹⁹¹

In addition to the cytosolic and nuclear Fe–S target proteins, the CIA targeting complex interacts with another CIA component, the NARFL protein (also named IOP1) (Fig. 1).^{192,193} NARFL has been proposed to also interact with the [NUBP1 NUBP2] complex because of the specific interaction between NUBP2 and NARFL.¹⁹⁴ NARFL can thus function at the interface between the [NUBP1 NUBP2] complex and the CIA targeting complex in the CIA pathway, but its precise molecular role remains to be elucidated (Fig. 1). The protein NARFL exhibits a sequence homology to bacterial [FeFe] hydrogenases,^{195,196} yet the absence of hydrogenase activity in yeast suggests that, instead of a [4Fe–4S] cluster linked to the Fe₂(CO)₃(CN)₂ active-site cluster, NARFL only retains the [4Fe–4S] cluster.¹⁹⁷ Nevertheless, NARFL coordinates two [4Fe–4S] cofactors that are similar to those in hydrogenases.^{197–199} Studies on the yeast homolog Nar1 showed that these clusters are bound to the N- and C-terminal motifs with four conserved cysteine residues each.^{197,198} Homology modelling and stability studies suggested that the C-terminal cluster is buried in the protein and that the more labile N-terminal cluster is surface-exposed, thus potentially being transferred to protein partners.^{197,198} Both motifs are crucial for the function of NARFL as CIA component.¹⁹⁸ In yeast, the depletion of all four early-acting CIA factors (Cdf1, Nbp35, Tah18, and Dre2) impairs the Fe–S cluster incorporation into Nar1 *in vivo*.^{152,157,197} Conversely, depletion of Nar1 has no influence on the Fe–S cluster assembly on Cdf1, Nbp35, or Dre2, but Nar1 is required for the maturation of cytosolic and nuclear [4Fe–4S] target proteins. In contrast, none of the late-acting CIA factors are required for the [4Fe–4S] cluster assembly on Nar1.^{188,200} Protein interaction studies indicated that Nar1 interacts with the yeast homologues of CIAO1 and CIA2B (Cia1 and Cia2, respectively),^{200,201} *i.e.* with the CIA targeting complex, as observed for the human homologues. Assuming that most of the *in vivo* findings on yeast hold in human cells, the most reliable model is that the CIA targeting complex binds NARFL, and that the full complex (the CIA targeting complex + NARFL) may catalyse the late CIA reactions transferring [4Fe–4S] clusters, received by the [NUBP1 NUBP2] complex, to the majority of cytosolic and nuclear [4Fe–4S] target proteins (Fig. 1 and 4). As a result, it might be possible that the observed NARFL–NUBP2 interaction,¹⁹⁴ discussed above, originates both from the maturation of NARFL processed by the early-acting CIA components, and from the [4Fe–4S] cluster transfer from the [NUBP1 NUBP2] complex to an already matured (*i.e.* containing

the C-terminal cluster only) NARFL protein bound to the CIA targeting complex.

NARFL has also been found to be a protein partner of another late-acting CIA component, *i.e.* the CIA2A protein (also named FAM96A), which, in turn, forms a complex with CIAO1 (Fig. 1 and 4).²⁰² The [CIAO1 CIA2A] complex has been proposed to be implicated in the [4Fe–4S] cluster maturation of exclusively IRP1, although no direct interaction with IRP1 was detected, possibly because of a rather labile interaction between apo-IRP1 and the [CIAO1 CIA2A] complex.²⁰² An NMR solution structure of a monomeric form of human CIA2A²⁰³ and two different crystal structures of dimeric CIA2A forms²⁰⁴ have been reported (Fig. 4). So far, the available structural information has not provided any decisive insights into the molecular function of CIA2A. A cysteine residue appears to perform a crucial role in the CIA2A function for maturing Fe–S target proteins.²⁰⁵ Elucidation of the molecular role of this reactive cysteine residue might clarify not only the role of CIA2A but also that of the entire CIA targeting complex in Fe–S protein maturation.

In conclusion, the proposed model assumes NARFL as the CIA component distributing [4Fe–4S] clusters assembled on the [NUBP1 NUBP2] complex to [CIAO1 CIA2A] and CIA targeting complexes. However, since NARFL cannot replace the cellular function of Nar1 in yeast,¹⁹⁷ more robust data need to be provided to prove the existence of a physical interaction between NARFL and the [NUBP1 NUBP2] complex.

Recently, two newly identified CIA factors, termed ORAOV1 and YAE1D1, were shown to function as specific adaptors connecting the CIA targeting complex with a specific [4Fe–4S] target protein, a cytosolic ABCE1 protein (Fig. 4). YAE1D1 and ORAOV1 form a complex that, in turn, interacts with the CIA targeting complex.¹⁹³ Both proteins contain an evolutionary conserved deca-GX₃ motif of 40 residues that is not found in any other eukaryotic protein.¹⁹³ ORAOV1 additionally carries a conserved C-terminal tryptophan (phenylalanine in some organisms).¹⁹³ A chain of binding events facilitates efficient ABCE1 maturation: (i) the CIA targeting complex interacts with ORAOV1 in the [YAE1D1 ORAOV1] complex *via* its C-terminal tryptophan; (ii) the two deca-GX₃ motifs are crucial for [YAE1D1 ORAOV1] complex formation; (iii) YAE1D1 in the [YAE1D1 ORAOV1] complex mediates contact with the ABCE1 protein.^{193,206} Interestingly, a characteristically different but related maturation strategy is followed by the radical-SAM Fe–S enzyme, viperin, an interferon-induced antiviral defense component.²⁰⁷ This protein also uses its conserved C-terminal tryptophan to directly associate with the CIA targeting complex. The removal of this residue abolishes complex formation with the CIA targeting complex, the assembly of viperin's Fe–S cluster, and, consequently, the antiviral function.²⁰⁷ Although it has been reported that CIAO1 forms mutually exclusive complexes with either CIA2A or CIA2B,²⁰² in a very recent work investigating the maturation pathway of the [4Fe–4S] cluster protein viperin, it has been found that both CIA targeting complex and CIA2A interact with viperin.²⁰⁸ Specifically, the CIA targeting complex interacts with the C-terminus of viperin, while CIA2A with the N-terminus. Despite this multitude of viperin interactors, only

CIAO1 is required for the Fe–S cluster incorporation into vipherin. The role of all the other interacting components is still unclear, but it might be possible that a concerted mechanism of cluster insertion is operative, in which CIAO1 is the essential component as it drives the association of all the other interactors.

The mitochondrial iron–sulfur cluster (ISC) export machinery

The main component of the mitochondrial ISC export machinery is an inner membrane ABCB transporter belonging to the ABCB7 group (Fig. 1).^{48,209} This group includes the human ABCB7, yeast Atm1 and *Arabidopsis thaliana* ATM3 transporters and can be found in virtually all eukaryotic species and in some bacterial organisms. *In vivo* data indicate that the members of this subfamily have the same function.^{209–211} In a variety of organisms, this transporter specifically supports the maturation of cytosolic and nuclear Fe–S proteins.^{48,209,211,212} Its depletion is also associated with defects in cellular iron homeostasis resulting in cytosolic iron deficiency and a concomitant iron accumulation in the mitochondria, that is, a similar phenotype observed for the proteins involved in the early stage of ISC protein deficiencies.^{48,213}

The other well-established component of the ISC export machinery is glutathione, which has been found to be essential for the maturation of extra-mitochondrial Fe–S proteins and for iron regulation.^{52,214} It was observed that GSH molecules can form an *in vitro* tetra-GSH-coordinated [2Fe–2S] cluster (defined as (GS)₄–[2Fe–2S]).^{215–217} The (GS)₄–[2Fe–2S] complex is shown to significantly stimulate the ATPase activity of an ABCB7-type transporter in both solution and proteoliposome-bound forms. On the basis of this *in vitro* evidence, it has been suggested that the (GS)₄–[2Fe–2S] complex is the molecule transiting from the mitochondria to the cytosol being the physiological substrate of the membrane transporter.^{218,219} On the other hand, the ATPase activity of the purified, liposome-integrated membrane transporter can be stimulated by many thiol-containing compounds including GSH and GSSG (the oxidized form of GSH),⁵³ questioning the physiological relevance of such an *in vitro* approach. In particular, the stimulatory effect of (GS)₄–[2Fe–2S] was 30-fold weaker than that observed with Cys-containing small peptides. Moreover, the affinity of (GS)₄–[2Fe–2S] binding to Atm1 was in the same range as that of GSH alone, questioning again the specificity of the (GS)₄–[2Fe–2S] interaction with Atm1 *in vitro*.^{218–220} Furthermore, it still needs to be documented that the (GS)₄–[2Fe–2S] complex remains intact under physiological transport conditions, and to verify the existence of a (GS)₄–[2Fe–2S] complex in a living cell. In conclusion, it is possible and reasonable that the (GS)₄–[2Fe–2S] complex is the substrate of the membrane transporter, but further experimental data are required to definitively confirm this conclusion.

The crystal structures of free and glutathione-bound Atm1 were solved and they are virtually identical.²¹⁹ The main outcome of the structure is that a V-shaped molecule enclosing a

6900 Å³ positively charged cavity is present. The cavity is close to the hydrophilic phase of the matrix side of the inner membrane, and likely serves as the substrate-binding site. The fact that Atm1 and a related bacterial ABC transporter²¹⁹ bind GSH or its derivatives suggests that these transporters may exhibit a similar substrate specificity. Both proteins bind GSH or its derivatives in the putative substrate-binding pocket comprised by the large, positively charged cavity close to the internal membrane surface and near to the putative exit channel formed by the twelve trans-membrane helices of Atm1. The cavity features much more space than that occupied by GSH alone, and therefore it is reasonable to suggest that GSH is a part of a larger Atm1 substrate and is required for substrate binding. In agreement with GSH being directly involved in the substrate export process, the Atm1-deficient mitochondria show an increased content of GSH,²²¹ and the ATPase activity of Atm1 is stimulated by GSH.⁵³ In particular, the largest stimulation of ATP hydrolysis by Atm1 from *Novosphingobium aromaticivorans* was observed for the Ag and Hg complexes of GSH.²²² In the crystal structure of Atm1 from *Novosphingobium aromaticivorans*, an oxidized glutathione molecule, GSSG, is bound into the transmembrane region in a location similar to GSH, suggesting that this interaction is specific and that this region of Atm1 family members may allow the binding of a GSH-containing substrate. Another study has suggested the molecule GSSSG, *i.e.* formed by the oxidized form of GSSH (glutathione persulfide) and GSH, as the Atm1 substrate and a potential sulfur carrier.²²³ In conclusion, the putative Atm1-like substrate is currently under debate. The identification on the nature of the molecule exported from the mitochondria to the cytosol by the ABCB transporter will be crucial to define whether a *de novo* synthesis of [2Fe–2S] clusters is required in the cytosol.

In yeast, another proposed component of the ISC export machinery is the sulfhydryl oxidase Erv1 located in the mitochondrial intermembrane space. This proposal is based on a similar phenotype to that observed for Atm1 deficiency.²²⁴ However, it has been recently shown that the observed phenotype was a consequence of strongly decreased glutathione levels, ruling out significant roles for Erv1 in cytosolic Fe–S protein biogenesis and iron regulation.²²⁵

Concluding remarks

Recent research has greatly advanced our understanding of the assembly and insertion of Fe–S clusters into mitochondrial, cytosolic and nuclear target proteins in human cells. The majority of the proteins involved in such processes have been identified, but a complete picture of how they interact and work at a molecular level is still ill-defined. The former steps of the mitochondrial ISC assembly machinery devoted to the assembly of a [2Fe–2S] cluster are best characterized at a molecular level. Nevertheless, the high molecular complexity of the process still does not provide a conclusive model on how [2Fe–2S] clusters are assembled and on the specific role of all components that are part of this complex mechanism. Even more “molecular chaos”

exists in the late stages of the ISC assembly machinery. In particular, the role of IBA57 is still not defined, and how the [4Fe-4S] cluster assembled in mitochondria is inserted into the final target proteins has to be understood. The most relevant CIA machinery components in humans have been identified, but a molecular picture of their function is still largely missing. The available functional data do not provide a clear view on how the CIA machinery components cooperate at the molecular level. Even the numerous interactomic data provided over the past few years are not conclusive in this respect. The investigation of the interactions of the CIA components at a molecular level and the associated structural data are required to make a step forward on defining how the CIA machinery works. We believe that this information will be fundamental to understand the degree of dependence of the CIA machinery on the ISC assembly and export machineries in humans. Finally, the identification of the nature of the molecule exported from mitochondria to the cytosol is a fundamental step to be attained in order to define the role of the ISC isoforms that are present in the cytosol in a *de novo* cytosolic synthesis of [2Fe-2S] clusters. In conclusion, only a combination of structural, biochemical and cell biological approaches can move forward the understanding of the molecular mechanisms responsible for the assembly and insertion of Fe-S clusters into target proteins in humans.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We acknowledge the financial support from the iNEXT Project 653706 "Infrastructure for NMR, EM and X-ray Crystallography for Translational Research". The authors acknowledge the support and the use of resources of Instruct-ERIC, a Landmark ESFRI project, and specifically CERM/CIRMMP Italy Centre. This article is also based upon work from COST Action CA15133, supported by COST (European Cooperation in Science and Technology).

References

- 1 K. J. Waldron, J. C. Rutherford, D. Ford and N. J. Robinson, Metalloproteins and metal sensing, *Nature*, 2009, **460**, 823–830.
- 2 J. J. R. Frausto da Silva and R. J. P. Williams, *The biological chemistry of the elements: the inorganic chemistry of life*, Oxford University Press, New York, 2001.
- 3 I. Bertini, H. B. Gray, E. I. Stiefel and J. S. Valentine, *Biological Inorganic Chemistry*, University Science Books, Sausalito, California, 2006.
- 4 L. A. Finney and T. V. O'Halloran, Transition Metal Speciation in the Cell: Insights from the Chemistry of Metal Ion Receptors, *Science*, 2003, **300**, 931–936.
- 5 B. E. Kim, T. Nevitt and D. J. Thiele, Mechanisms for copper acquisition, distribution and regulation, *Nat. Chem. Biol.*, 2008, **4**, 176–185.
- 6 A. Atkinson and D. R. Winge, Metal acquisition and availability in the mitochondria, *Chem. Rev.*, 2009, **109**, 4708–4721.
- 7 G. J. Brewer, Copper in medicine, *Curr. Opin. Chem. Biol.*, 2003, **7**, 207–212.
- 8 M. Valko, H. Morris and M. T. Cronin, Metals, toxicity and oxidative stress, *Curr. Med. Chem.*, 2005, **12**, 1161–1208.
- 9 R. Lill and U. Muhlenhoff, Maturation of iron-sulfur proteins in eukaryotes: mechanisms, connected processes, and diseases, *Annu. Rev. Biochem.*, 2008, **77**, 669–700.
- 10 T. A. Rouault, Iron metabolism in the CNS: implications for neurodegenerative diseases, *Nat. Rev. Neurosci.*, 2013, **14**, 551–564.
- 11 M. R. Bleackley and R. T. Macgillivray, Transition metal homeostasis: from yeast to human disease, *Biometals*, 2011, **24**, 785–809.
- 12 A. Sheftel, O. Stehling and R. Lill, Iron-sulfur proteins in health and disease, *Trends Endocrinol. Metab.*, 2010, **21**, 302–314.
- 13 E. Madsen and J. D. Gitlin, Copper and iron disorders of the brain, *Annu. Rev. Neurosci.*, 2007, **30**, 317–337.
- 14 K. J. Barnham and A. I. Bush, Metals in Alzheimer's and Parkinson's diseases, *Curr. Opin. Chem. Biol.*, 2008, **12**, 222–228.
- 15 A. V. Davis and T. V. O'Halloran, A place for thioether chemistry in cellular copper ion recognition and trafficking, *Nat. Chem. Biol.*, 2008, **4**, 148–151.
- 16 D. L. Huffman and T. V. O'Halloran, Function, structure, and mechanism of intracellular copper trafficking proteins, *Annu. Rev. Biochem.*, 2001, **70**, 677–701.
- 17 A. K. Boal and A. C. Rosenzweig, Structural biology of copper trafficking, *Chem. Rev.*, 2009, **109**, 4760–4779.
- 18 J. M. Leitch, P. J. Yick and V. C. Culotta, The right to choose: multiple pathways for activating copper, zinc superoxide dismutase, *J. Biol. Chem.*, 2009, **284**, 24679–24683.
- 19 L. Banci, I. Bertini, F. Cantini and S. Ciofi-Baffoni, Cellular copper distribution: a mechanistic systems biology approach, *Cell. Mol. Life Sci.*, 2010, **67**, 2563–2589.
- 20 F. S. La, R. Burke and D. P. Giedroc, Mammalian copper biology: hitting the pause button in celebration of three pioneers and four decades of discovery, *Metallomics*, 2016, **8**, 810–812.
- 21 L. Banci and A. Rosato, Structural genomics of proteins involved in copper homeostasis, *Acc. Chem. Res.*, 2003, **36**, 215–221.
- 22 L. Banci, I. Bertini, S. Ciofi-Baffoni, T. Kozyreva, K. Zovo and P. Palumaa, Affinity gradients drive copper to cellular destinations, *Nature*, 2010, **465**, 645–648.
- 23 C. Andreini, L. Banci, I. Bertini, S. Elmi and A. Rosato, Non-heme iron through the three domains of life, *Proteins: Struct., Funct., Bioinf.*, 2007, **67**, 317–324.
- 24 C. Andreini, L. Banci, I. Bertini and A. Rosato, Occurrence of copper through the three domains of life: a bioinformatic approach, *J. Proteome Res.*, 2008, **1**, 209–216.

- 25 A. C. Dlouhy and C. E. Outten, The iron metallome in eukaryotic organisms, *Met. Ions Life Sci.*, 2013, **12**, 241–278.
- 26 D. C. Johnson, D. R. Dean, A. D. Smith and M. K. Johnson, Structure, function, and formation of biological iron-sulfur clusters, *Annu. Rev. Biochem.*, 2005, **74**, 247–281.
- 27 H. Beinert, R. H. Holm and E. Munck, Iron-sulfur clusters: nature's modular, multipurpose structures, *Science*, 1997, **277**, 653–659.
- 28 P. Venkateswara Rao and R. H. Holm, Synthetic Analogues of the Active Sites of Iron–Sulfur Proteins, *Chem. Rev.*, 2004, **104**, 527–559.
- 29 P. J. Kiley and H. Beinert, The role of Fe–S proteins in sensing and regulation in bacteria, *Curr. Opin. Microbiol.*, 2003, **6**, 181–185.
- 30 Y. Hu and M. W. Ribbe, A journey into the active center of nitrogenase, *J. Biol. Inorg. Chem.*, 2014, **19**, 731–736.
- 31 E. M. Shepard, E. S. Boyd, J. B. Broderick and J. W. Peters, Biosynthesis of complex iron–sulfur enzymes, *Curr. Opin. Chem. Biol.*, 2011, **15**, 319–327.
- 32 R. A. Marcus and N. Sutin, Electron transfer in chemistry and biology, *Biochim. Biophys. Acta*, 1985, **811**, 265–275.
- 33 J. Hirst, Mitochondrial complex I, *Annu. Rev. Biochem.*, 2013, **82**, 551–575.
- 34 S. J. Booker, R. M. Cicchillo and T. L. Grove, Self-sacrifice in radical *S*-adenosylmethionine proteins, *Curr. Opin. Chem. Biol.*, 2007, **11**, 543–552.
- 35 J. Meyer, Iron-sulfur protein folds, iron-sulfur chemistry, and evolution, *J. Biol. Inorg. Chem.*, 2008, **13**, 157–170.
- 36 E. L. Mettert and P. J. Kiley, Fe–S proteins that regulate gene expression, *Biochim. Biophys. Acta*, 2015, **1853**, 1284–1293.
- 37 T. A. Rouault, The role of iron regulatory proteins in mammalian iron homeostasis and disease, *Nat. Chem. Biol.*, 2006, **2**, 406–414.
- 38 C. Andreini, I. Bertini and A. Rosato, Metalloproteomes: a bioinformatic approach, *Acc. Chem. Res.*, 2009, **42**, 1471–1479.
- 39 C. Andreini, L. Banci and A. Rosato, Exploiting bacterial operons to illuminate human iron-sulfur proteins, *J. Proteome Res.*, 2016, **15**, 1308–1322.
- 40 C. Andreini, A. Rosato and L. Banci, The Relationship between Environmental Dioxygen and Iron-Sulfur Proteins Explored at the Genome Level, *PLoS One*, 2017, **12**, e0171279.
- 41 L. Zheng, V. L. Cash, D. H. Flint and D. R. Dean, Assembly of iron-sulfur clusters. Identification of an *iscSUA-hscBA-fdx* gene cluster from *Azotobacter vinelandii*, *J. Biol. Chem.*, 1998, **273**, 13264–13272.
- 42 B. Roche, L. Aussel, B. Ezraty, P. Mandin, B. Py and F. Barras, Iron/sulfur proteins biogenesis in prokaryotes: formation, regulation and diversity, *Biochim. Biophys. Acta*, 2013, **1827**, 455–469.
- 43 O. Stehling, C. Wilbrecht and R. Lill, Mitochondrial iron-sulfur protein biogenesis and human disease, *Biochimie*, 2014, **100**, 61–77.
- 44 N. Maio and T. A. Rouault, Iron-sulfur cluster biogenesis in mammalian cells: new insights into the molecular mechanisms of cluster delivery, *Biochim. Biophys. Acta*, 2015, **1853**, 1493–1512.
- 45 J. O. Fuss, C. L. Tsai, J. P. Ishida and J. A. Tainer, Emerging critical roles of Fe–S clusters in DNA replication and repair, *Biochim. Biophys. Acta*, 2015, **1853**, 1253–1271.
- 46 O. A. Lukianova and S. S. David, A role for iron-sulfur clusters in DNA repair, *Curr. Opin. Chem. Biol.*, 2005, **9**, 145–151.
- 47 D. J. Netz, J. Mascarenhas, O. Stehling, A. J. Pierik and R. Lill, Maturation of cytosolic and nuclear iron-sulfur proteins, *Trends Cell Biol.*, 2014, **24**, 303–312.
- 48 G. Kispal, P. Csere, C. Prohl and R. Lill, The mitochondrial proteins Atm1p and Nfs1p are essential for biogenesis of cytosolic Fe/S proteins, *EMBO J.*, 1999, **18**, 3981–3989.
- 49 A. Biederbick, O. Stehling, R. Rosser, B. Niggemeyer, Y. Nakai, H. P. Elsasser and R. Lill, Role of human mitochondrial Nfs1 in cytosolic iron-sulfur protein biogenesis and iron regulation, *Mol. Cell. Biol.*, 2006, **26**, 5675–5687.
- 50 J. Gerber, K. Neumann, C. Prohl, U. Muhlenhoff and R. Lill, The yeast scaffold proteins Isu1p and Isu2p are required inside mitochondria for maturation of cytosolic Fe/S proteins, *Mol. Cell. Biol.*, 2004, **24**, 4848–4857.
- 51 R. Lill, R. Dutkiewicz, S. A. Freibert, T. Heidenreich, J. Mascarenhas, D. J. Netz, V. D. Paul, A. J. Pierik, N. Richter, M. Stumpfig, V. Srinivasan, O. Stehling and U. Muhlenhoff, The role of mitochondria and the CIA machinery in the maturation of cytosolic and nuclear iron-sulfur proteins, *Eur. J. Cell Biol.*, 2015, **94**, 280–291.
- 52 K. Sipos, H. Lange, Z. Fekete, P. Ullmann, R. Lill and G. Kispal, Maturation of cytosolic iron-sulfur proteins requires glutathione, *J. Biol. Chem.*, 2002, **277**, 26944–26949.
- 53 G. Kuhnke, K. Neumann, U. Muhlenhoff and R. Lill, Stimulation of the ATPase activity of the yeast mitochondrial ABC transporter Atm1p by thiol compounds, *Mol. Membr. Biol.*, 2006, **23**, 173–184.
- 54 C. A. Chen and J. A. Cowan, Characterization of *Saccharomyces cerevisiae* Atm1p: functional studies of an ABC7 type transporter, *Biochim. Biophys. Acta*, 2006, **1760**, 1857–1865.
- 55 T. A. Rouault, Biogenesis of iron-sulfur clusters in mammalian cells: new insights and relevance to human disease, *Dis. Models & Mech.*, 2012, **5**, 155–164.
- 56 N. Maio and T. A. Rouault, Iron-sulfur cluster biogenesis in mammalian cells: new insights into the molecular mechanisms of cluster delivery, *Biochim. Biophys. Acta*, 2015, **1853**, 1493–1512.
- 57 Y. Shi, M. C. Ghosh, W. H. Tong and T. A. Rouault, Human ISD11 is essential for both iron-sulfur cluster assembly and maintenance of normal cellular iron homeostasis, *Hum. Mol. Genet.*, 2009, **18**, 3014–3025.
- 58 W. H. Tong, G. N. Jameson, B. H. Huynh and T. A. Rouault, Subcellular compartmentalization of human Nfu, an iron-sulfur cluster scaffold protein, and its ability to assemble a [4Fe–4S] cluster, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 9762–9767.
- 59 W. H. Tong and T. A. Rouault, Functions of mitochondrial ISCU and cytosolic ISCU in mammalian iron-sulfur cluster biogenesis and iron homeostasis, *Cell Metab.*, 2006, **3**, 199–210.

- 60 F. Acquaviva, I. De Biase, L. Nezi, G. Ruggiero, F. Tatangelo, C. Pisano, A. Monticelli, C. Garbi, A. M. Acquaviva and S. Cocozza, Extra-mitochondrial localisation of frataxin and its association with IscU1 during enterocyte-like differentiation of the human colon adenocarcinoma cell line Caco-2, *J. Cell Sci.*, 2005, **118**, 3917–3924.
- 61 T. Land and T. A. Rouault, Targeting of a human iron-sulfur cluster assembly enzyme, nifs, to different sub-cellular compartments is regulated through alternative AUG utilization, *Mol. Cell*, 1998, **2**, 807–815.
- 62 M. W. Gray, G. Burger and B. F. Lang, Mitochondrial evolution, *Science*, 1999, **283**, 1476–1481.
- 63 C. G. Kurland and S. G. Andersson, Origin and evolution of the mitochondrial proteome, *Microbiol. Mol. Biol. Rev.*, 2000, **64**, 786–820.
- 64 T. A. Rouault and W. H. Tong, Iron-sulfur cluster biogenesis and human disease, *Trends Genet.*, 2008, **24**, 398–407.
- 65 X. M. Xu and S. G. Moller, Iron-sulfur clusters: biogenesis, molecular mechanisms, and their functional significance, *Antioxid. Redox Signaling*, 2011, **15**, 271–307.
- 66 F. Pierrel, P. A. Cobine and D. R. Winge, Metal Ion availability in mitochondria, *Biomaterials*, 2007, **20**, 675–682.
- 67 P. A. Lindahl and M. J. Moore, Labile Low-Molecular-Mass Metal Complexes in Mitochondria: Trials and Tribulations of a Burgeoning Field, *Biochemistry*, 2016, **55**, 4140–4153.
- 68 S. P. McCormick, M. J. Moore and P. A. Lindahl, Detection of Labile Low-Molecular-Mass Transition Metal Complexes in Mitochondria, *Biochemistry*, 2015, **54**, 3442–3453.
- 69 J. Gerber, U. Muhlenhoff and R. Lill, An interaction between frataxin and Isu1/Nfs1 that is crucial for Fe/S cluster synthesis on Isu1, *EMBO Rep.*, 2003, **4**, 906–911.
- 70 D. J. Lane, A. M. Merlot, M. L. Huang, D. H. Bae, P. J. Jansson, S. Sahni, D. S. Kalinowski and D. R. Richardson, Cellular iron uptake, trafficking and metabolism: key molecules and mechanisms and their roles in disease, *Biochim. Biophys. Acta*, 2015, **1853**, 1130–1144.
- 71 A. Pastore and H. Puccio, Frataxin: a protein in search for a function, *J. Neurochem.*, 2013, **126**(suppl 1), 43–52.
- 72 R. Lill, B. Hoffmann, S. Molik, A. J. Pierik, N. Rietzschel, O. Stehling, M. A. Uzarska, H. Webert, C. Wilbrecht and U. Muhlenhoff, The role of mitochondria in cellular iron-sulfur protein biogenesis and iron metabolism, *Biochim. Biophys. Acta*, 2012, **1823**, 1491–1508.
- 73 S. A. Cory, J. G. Van Vranken, E. J. Brignole, S. Patra, D. R. Winge, C. L. Drennan, J. Rutter and D. P. Barondeau, Structure of human Fe–S assembly subcomplex reveals unexpected cysteine desulfurase architecture and acyl-ACP-ISD11 interactions, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E5325–E5334.
- 74 L. Zheng, R. H. White, V. L. Cash and D. R. Dean, Mechanism for the desulfurization of L-cysteine catalyzed by the nifs gene product, *Biochemistry*, 1994, **33**, 4714–4720.
- 75 J. T. Kaiser, T. Clausen, G. P. Bourenkow, H. D. Bartunik, S. Steinbacher and R. Huber, Crystal structure of a Nifs-like protein from *Thermotoga maritima*: implications for iron sulphur cluster assembly, *J. Mol. Biol.*, 2000, **297**, 451–464.
- 76 P. P. Saha, S. Srivastava, S. K. P. Kumar, D. Sinha and P. D'Silva, Mapping Key Residues of ISD11 Critical for NFS1-ISD11 Subcomplex Stability: Implications in the Development of Mitochondrial Disorder, COXPD19, *J. Biol. Chem.*, 2015, **290**, 25876–25890.
- 77 J. G. Van Vranken, M. Y. Jeong, P. Wei, Y. C. Chen, S. P. Gygi, D. R. Winge and J. Rutter, The mitochondrial acyl carrier protein (ACP) coordinates mitochondrial fatty acid synthesis with iron sulfur cluster biogenesis, *Elife*, 2016, **5**, e17828.
- 78 M. Friemel, Z. Marelja, K. Li and S. Leimkuhler, The N-Terminus of Iron-Sulfur Cluster Assembly Factor ISD11 Is Crucial for Subcellular Targeting and Interaction with L-Cysteine Desulfurase NFS1, *Biochemistry*, 2017, **56**, 1797–1808.
- 79 K. Cai, R. O. Frederick, M. Tonelli and J. L. Markley, Mitochondrial Cysteine Desulfurase and ISD11 Coexpressed in *Escherichia coli* Yield Complex Containing Acyl Carrier Protein, *ACS Chem. Biol.*, 2017, **12**, 918–921.
- 80 K. Cai, M. Tonelli, R. O. Frederick and J. L. Markley, Human Mitochondrial Ferredoxin 1 (FDX1) and Ferredoxin 2 (FDX2) Both Bind Cysteine Desulfurase and Donate Electrons for Iron-Sulfur Cluster Biosynthesis, *Biochemistry*, 2017, **56**, 487–499.
- 81 C. L. Tsai and D. P. Barondeau, Human frataxin is an allosteric switch that activates the Fe–S cluster biosynthetic complex, *Biochemistry*, 2010, **49**, 9132–9139.
- 82 S. Schmucker, A. Martelli, F. Colin, A. Page, M. Wattenhofer-Donze, L. Reutenauer and H. Puccio, Mammalian frataxin: an essential function for cellular viability through an interaction with a preformed ISCU/NFS1/ISD11 iron-sulfur assembly complex, *PLoS One*, 2011, **6**, e16199.
- 83 J. Bridwell-Rabb, N. G. Fox, C. L. Tsai, A. M. Winn and D. P. Barondeau, Human frataxin activates Fe–S cluster biosynthesis by facilitating sulfur transfer chemistry, *Biochemistry*, 2014, **53**, 4904–4913.
- 84 N. G. Fox, D. Das, M. Chakrabarti, P. A. Lindahl and D. P. Barondeau, Frataxin Accelerates [2Fe–2S] Cluster Formation on the Human Fe–S Assembly Complex, *Biochemistry*, 2015, **54**, 3880–3889.
- 85 A. Parent, X. Elduque, D. Cornu, L. Belot, J. P. Le Caer, A. Grandas, M. B. Toledano and B. d'Autreaux, Mammalian frataxin directly enhances sulfur transfer of NFS1 persulfide to both ISCU and free thiols, *Nat. Commun.*, 2015, **6**, 5686.
- 86 R. Shi, A. Proteau, M. Villarroja, I. Moukadiri, L. Zhang, J. F. Trempe, A. Matte, M. E. Armengod and M. Cygler, Structural basis for Fe–S cluster assembly and tRNA thiolation mediated by IscS protein-protein interactions, *PLoS Biol.*, 2010, **8**, e1000354.
- 87 E. N. Marinoni, J. S. de Oliveira, Y. Nicolet, E. C. Raulfs, P. Amara, D. R. Dean and J. C. Fontecilla-Camps, (IscS-IscU)₂ complex structures provide insights into Fe₂S₂ biogenesis and transfer, *Angew. Chem., Int. Ed.*, 2012, **51**, 5439–5442.
- 88 W. H. Tong and T. Rouault, Distinct iron-sulfur cluster assembly complexes exist in the cytosol and mitochondria of human cells, *EMBO J.*, 2000, **19**, 5692–5700.

- 89 S. Schmucker, M. Argentini, N. Carelle-Calmels, A. Martelli and H. Puccio, The in vivo mitochondrial two-step maturation of human frataxin, *Hum. Mol. Genet.*, 2008, **17**, 3521–3531.
- 90 R. A. Vaubel and G. Isaya, Iron-sulfur cluster synthesis, iron homeostasis and oxidative stress in Friedreich ataxia, *Mol. Cell. Neurosci.*, 2013, **55**, 50–61.
- 91 P. Cavadini, J. Adamec, F. Taroni, O. Gakh and G. Isaya, Two-step processing of human frataxin by mitochondrial processing peptidase. Precursor and intermediate forms are cleaved at different rates, *J. Biol. Chem.*, 2000, **275**, 41469–41475.
- 92 S. Schmucker, M. Argentini, N. Carelle-Calmels, A. Martelli and H. Puccio, The in vivo mitochondrial two-step maturation of human frataxin, *Hum. Mol. Genet.*, 2008, **17**, 3521–3531.
- 93 R. K. Jobling, M. Assoum, O. Gakh, S. Blaser, J. A. Raiman, C. Mignot, E. Roze, A. Durr, A. Brice, N. Levy, C. Prasad, T. Paton, A. D. Paterson, N. M. Roslin, C. R. Marshall, J. P. Desvignes, N. Roeckel-Trevisiol, S. W. Scherer, G. A. Rouleau, A. Megarbane, G. Isaya, V. Delague and G. Yoon, PMPCA mutations cause abnormal mitochondrial protein processing in patients with non-progressive cerebellar ataxia, *Brain*, 2015, **138**, 1505–1517.
- 94 H. Li, O. Gakh, D. Y. Smith, W. K. Ranatunga and G. Isaya, Missense mutations linked to friedreich ataxia have different but synergistic effects on mitochondrial frataxin isoforms, *J. Biol. Chem.*, 2013, **288**, 4116–4127.
- 95 A. J. Andrew, J. Y. Song, B. Schilke and E. A. Craig, Post-translational regulation of the scaffold for Fe–S cluster biogenesis, *Isu, Mol. Biol. Cell*, 2008, **19**, 5259–5266.
- 96 O. Gakh, T. Bedekovics, S. F. Duncan, D. Y. Smith, D. S. Berkholz and G. Isaya, Normal and Friedreich ataxia cells express different isoforms of frataxin with complementary roles in iron-sulfur cluster assembly, *J. Biol. Chem.*, 2010, **285**, 38486–38501.
- 97 F. Colin, A. Martelli, M. Clemancey, J. M. Latour, S. Gambarelli, L. Zeppieri, C. Birck, A. Page, H. Puccio and C. S. Ollagnier de, Mammalian frataxin controls sulfur production and iron entry during de novo Fe4S4 cluster assembly, *J. Am. Chem. Soc.*, 2013, **135**, 733–740.
- 98 O. Gakh, W. Ranatunga, D. Y. Smith, E. C. Ahlgren, S. Al-Karadaghi, J. R. Thompson and G. Isaya, Architecture of the Human Mitochondrial Iron-Sulfur Cluster Assembly Machinery, *J. Biol. Chem.*, 2016, **291**, 21296–21321.
- 99 T. L. Stemmler, E. Lesuisse, D. Pain and A. Dancis, Frataxin and mitochondrial FeS cluster biogenesis, *J. Biol. Chem.*, 2010, **285**, 26737–26743.
- 100 J. D. Cook, K. C. Kondapalli, S. Rawat, W. C. Childs, Y. Murugesan, A. Dancis and T. L. Stemmler, Molecular details of the yeast frataxin-Isu1 interaction during mitochondrial Fe–S cluster assembly, *Biochemistry*, 2010, **49**, 8756–8765.
- 101 H. Li, O. Gakh, D. Y. Smith and G. Isaya, Oligomeric yeast frataxin drives assembly of core machinery for mitochondrial iron-sulfur cluster synthesis, *J. Biol. Chem.*, 2009, **284**, 21971–21980.
- 102 A. D. Sheftel, O. Stehling, A. J. Pierik, H. P. Elsasser, U. Muhlenhoff, H. Webert, A. Hobler, F. Hannemann, R. Bernhardt and R. Lill, Humans possess two mitochondrial ferredoxins, Fdx1 and Fdx2, with distinct roles in steroidogenesis, heme, and Fe/S cluster biosynthesis, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 11775–11780.
- 103 Y. Shi, M. Ghosh, G. Kovtunovych, D. R. Crooks and T. A. Rouault, Both human ferredoxins 1 and 2 and ferredoxin reductase are important for iron-sulfur cluster biogenesis, *Biochim. Biophys. Acta*, 2012, **1823**, 484–492.
- 104 H. Webert, S. A. Freibert, A. Gallo, T. Heidenreich, U. Linne, S. Amlacher, E. Hurt, U. Muhlenhoff, L. Banci and R. Lill, Functional reconstitution of de novo mitochondrial Fe/S cluster synthesis reveals an essential role of ferredoxin, *Nat. Commun.*, 2014, **5**, 5013–5025.
- 105 H. Uhrigshardt, A. Singh, G. Kovtunovych, M. Ghosh and T. A. Rouault, Characterization of the human HSC20, an unusual DnaJ type III protein, involved in iron-sulfur cluster biogenesis, *Hum. Mol. Genet.*, 2010, **19**, 3816–3834.
- 106 N. Maio, A. Singh, H. Uhrigshardt, N. Saxena, W. H. Tong and T. A. Rouault, Cochaperone binding to LYR motifs confers specificity of iron sulfur cluster delivery, *Cell Metab.*, 2014, **19**, 445–457.
- 107 Y. Shan and G. Cortopassi, HSC20 interacts with frataxin and is involved in iron-sulfur cluster biogenesis and iron homeostasis, *Hum. Mol. Genet.*, 2012, **21**, 1457–1469.
- 108 K. Cai, R. O. Frederick, J. H. Kim, N. M. Reinen, M. Tonelli and J. L. Markley, Human Mitochondrial Chaperone (mtHSP70) and Cysteine Desulfurase (NFS1) Bind Preferentially to the Disordered Conformation, Whereas Co-chaperone (HSC20) Binds to the Structured Conformation of the Iron-Sulfur Cluster Scaffold Protein (ISCU), *J. Biol. Chem.*, 2013, **288**, 28755–28770.
- 109 A. K. Fuzery, J. J. Oh, D. T. Ta, L. E. Vickery and J. L. Markley, Three hydrophobic amino acids in Escherichia coli HscB make the greatest contribution to the stability of the HscB-Isu complex, *BMC Biochem.*, 2011, **12**, 3.
- 110 S. J. Ciesielski, B. A. Schilke, J. Osipiuk, L. Bigelow, R. Mulligan, J. Majewska, A. Joachimiak, J. Marszalek, E. A. Craig and R. Dutkiewicz, Interaction of J-protein co-chaperone Jac1 with Fe–S scaffold Isu is indispensable in vivo and conserved in evolution, *J. Mol. Biol.*, 2012, **417**, 1–12.
- 111 J. R. Cupp-Vickery and L. E. Vickery, Crystal structure of Hsc20, a J-type Co-chaperone from Escherichia coli, *J. Mol. Biol.*, 2000, **304**, 835–845.
- 112 E. Bitto, C. A. Bingman, L. Bittova, D. A. Kondrashov, R. M. Bannen, B. G. Fox, J. L. Markley and G. N. Phillips, Jr., Structure of human J-type co-chaperone HscB reveals a tetracysteine metal-binding domain, *J. Biol. Chem.*, 2008, **283**, 30184–30192.
- 113 J. Amick, S. E. Schlanger, C. Wachnowsky, M. A. Moseng, C. C. Emerson, M. Dare, W. I. Luo, S. S. Ithychanda, J. C. Nix, J. A. Cowan, R. C. Page and S. Misra, Crystal structure of the nucleotide-binding domain of mortalin, the mitochondrial Hsp70 chaperone, *Protein Sci.*, 2014, **23**, 833–842.

- 114 H. Schroder, T. Langer, F. U. Hartl and B. Bukau, DnaK, DnaJ and GrpE form a cellular chaperone machinery capable of repairing heat-induced protein damage, *EMBO J.*, 1993, **12**, 4137–4144.
- 115 U. Muhlenhoff, J. Gerber, N. Richhardt and R. Lill, Components involved in assembly and dislocation of iron-sulfur clusters on the scaffold protein Isu1p, *EMBO J.*, 2003, **22**, 4815–4825.
- 116 M. A. Uzarska, R. Dutkiewicz, S. A. Freibert, R. Lill and U. Muhlenhoff, The mitochondrial Hsp70 chaperone Ssq1 facilitates Fe/S cluster transfer from Isu1 to Grx5 by complex formation, *Mol. Biol. Cell*, 2013, **24**, 1830–1841.
- 117 P. Shakamuri, B. Zhang and M. K. Johnson, Monothiol glutaredoxins function in storing and transporting [Fe2S2] clusters assembled on IscU scaffold proteins, *J. Am. Chem. Soc.*, 2012, **134**, 15213–15216.
- 118 M. M. Molina-Navarro, C. Casas, L. Piedrafita, G. Belli and E. Herrero, Prokaryotic and eukaryotic monothiol glutaredoxins are able to perform the functions of Grx5 in the biogenesis of Fe/S clusters in yeast mitochondria, *FEBS Lett.*, 2006, **580**, 2273–2280.
- 119 R. A. Wingert, J. L. Galloway, B. Barut, H. Foott, P. Fraenkel, J. L. Axe, G. J. Weber, K. Dooley, A. J. Davidson, B. Schmid, B. H. Paw, G. C. Shaw, P. Kingsley, J. Palis, H. Schubert, O. Chen, J. Kaplan and L. I. Zon, Deficiency of glutaredoxin 5 reveals Fe–S clusters are required for vertebrate haem synthesis, *Nature*, 2005, **436**, 1035–1039.
- 120 M. T. Rodriguez-Manzanique, J. Tamarit, G. Belli, J. Ros and E. Herrero, Grx5 is a mitochondrial glutaredoxin required for the activity of iron/sulfur enzymes, *Mol. Biol. Cell*, 2002, **13**, 1109–1121.
- 121 L. Banci, D. Brancaccio, S. Ciofi-Baffoni, R. Del Conte, R. Gadepalli, M. Mikolajczyk, S. Neri, M. Piccioli and J. Winkelmann, [2Fe–2S] cluster transfer in iron-sulfur protein biogenesis, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 6203–6208.
- 122 N. Maio and T. A. Rouault, Mammalian Fe–S proteins: definition of a consensus motif recognized by the co-chaperone HSC20, *Metalloomics*, 2016, **8**, 1032–1046.
- 123 H. Angerer, Eukaryotic LYR Proteins Interact with Mitochondrial Protein Complexes, *Biology*, 2015, **4**, 133–150.
- 124 D. Ghezzi, P. Goffrini, G. Uziel, R. Horvath, T. Klopstock, H. Lochmuller, P. D'Adamo, P. Gasparini, T. M. Strom, H. Prokisch, F. Invernizzi, I. Ferrero and M. Zeviani, SDHAF1, encoding a LYR complex-II specific assembly factor, is mutated in SDH-defective infantile leukoencephalopathy, *Nat. Genet.*, 2009, **41**, 654–656.
- 125 N. Maio, D. Ghezzi, D. Verrigni, T. Rizza, E. Bertini, D. Martinelli, M. Zeviani, A. Singh, R. Carrozzo and T. A. Rouault, Disease-Causing SDHAF1 Mutations Impair Transfer of Fe–S Clusters to SDHB, *Cell Metab.*, 2016, **23**, 292–302.
- 126 J. L. Faria, M. Berberan-Santos and M. J. Prieto, A comment on the localization of cyanine dye binding to brush-border membranes by the fluorescence quenching of n-(9-anthroyloxy) fatty acid probes, *Biochim. Biophys. Acta*, 1990, **1026**, 133–134.
- 127 N. G. Fox, M. Chakrabarti, S. P. McCormick, P. A. Lindahl and D. P. Barondeau, The Human Iron-Sulfur Assembly Complex Catalyzes the Synthesis of [2Fe–2S] Clusters on ISCU2 That Can Be Transferred to Acceptor Molecules, *Biochemistry*, 2015, **54**, 3871–3879.
- 128 L. K. Beilschmidt, C. S. Ollagnier de, M. Fournier, I. Sanakis, M. A. Hograindleur, M. Clemancey, G. Blondin, S. Schmucker, A. Eisenmann, A. Weiss, P. Koebel, N. Messaddeq, H. Puccio and A. Martelli, ISCA1 is essential for mitochondrial Fe4S4 biogenesis in vivo, *Nat. Commun.*, 2017, **8**, 15124.
- 129 K. D. Kim, W. H. Chung, H. J. Kim, K. C. Lee and J. H. Roe, Monothiol glutaredoxin Grx5 interacts with Fe–S scaffold proteins Isa1 and Isa2 and supports Fe–S assembly and DNA integrity in mitochondria of fission yeast, *Biochem. Biophys. Res. Commun.*, 2010, **392**, 467–472.
- 130 A. D. Sheftel, C. Wilbrecht, O. Stehling, B. Niggemeyer, H. P. Elsasser, U. Muhlenhoff and R. Lill, The human mitochondrial ISCA1, ISCA2, and IBA57 proteins are required for [4Fe–4S] protein maturation, *Mol. Biol. Cell*, 2012, **23**, 1157–1166.
- 131 I. Cozar-Castellano, M. M. del Valle, E. Trujillo, M. F. Arteaga, T. Gonzalez, P. Martin-Vasallo and J. Avila, hIsca: a protein implicated in the biogenesis of iron-sulfur clusters, *Biochim. Biophys. Acta*, 2004, **1700**, 179–188.
- 132 U. Muhlenhoff, N. Richter, O. Pines, A. J. Pierik and R. Lill, Specialized function of yeast Isa1 and Isa2 proteins in the maturation of mitochondrial [4Fe–4S] proteins, *J. Biol. Chem.*, 2011, **286**, 41205–41216.
- 133 D. Brancaccio, A. Gallo, M. Mikolajczyk, K. Zovo, P. Palumaa, E. Novellino, M. Piccioli, S. Ciofi-Baffoni and L. Banci, Formation of [4Fe–4S] clusters in the mitochondrial iron-sulfur cluster assembly machinery, *J. Am. Chem. Soc.*, 2014, **136**, 16240–16250.
- 134 D. Brancaccio, A. Gallo, M. Piccioli, E. Novellino, S. Ciofi-Baffoni and L. Banci, [4Fe–4S] Cluster Assembly in Mitochondria and Its Impairment by Copper, *J. Am. Chem. Soc.*, 2017, **139**, 719–730.
- 135 L. T. Jensen and V. C. Culotta, Role of *Saccharomyces cerevisiae* ISA1 and ISA2 in iron homeostasis, *Mol. Cell. Biol.*, 2000, **20**, 3918–3927.
- 136 A. Kaut, H. Lange, K. Diekert, G. Kispal and R. Lill, Isa1p is a component of the mitochondrial machinery for maturation of cellular iron-sulfur proteins and requires conserved cysteine residues for function, *J. Biol. Chem.*, 2000, **275**, 15955–15961.
- 137 A. Navarro-Sastre, F. Tort, O. Stehling, M. A. Uzarska, J. A. Arranz, T. M. Del, M. T. Labayru, J. Landa, A. Font, J. Garcia-Villoria, B. Merinero, M. Ugarte, L. G. Gutierrez-Solana, J. Campistol, A. Garcia-Cazorla, J. Vaquerizo, E. Riudor, P. Briones, O. Elpeleg, A. Ribes and R. Lill, A fatal mitochondrial disease is associated with defective NFU1 function in the maturation of a subset of mitochondrial Fe–S proteins, *Am. J. Hum. Genet.*, 2011, **89**, 656–667.
- 138 J. M. Cameron, A. Janer, V. Levandovskiy, N. MacKay, T. A. Rouault, W. H. Tong, I. Ogilvie, E. A. Shoubridge and B. H. Robinson, Mutations in iron-sulfur cluster

- scaffold genes NFU1 and BOLA3 cause a fatal deficiency of multiple respiratory chain and 2-oxoacid dehydrogenase enzymes, *Am. J. Hum. Genet.*, 2011, **89**, 486–495.
- 139 K. Bych, S. Kerscher, D. J. Netz, A. J. Pierik, K. Zwicker, M. A. Huynen, R. Lill, U. Brandt and J. Balk, The iron-sulphur protein Ind1 is required for effective complex I assembly, *EMBO J.*, 2008, **27**, 1736–1746.
- 140 A. D. Sheftel, O. Stehling, A. J. Pierik, D. J. Netz, S. Kerscher, H. P. Elsasser, I. Wittig, J. Balk, U. Brandt and R. Lill, Humanind1, an iron-sulfur cluster assembly factor for respiratory complex I, *Mol. Cell. Biol.*, 2009, **29**, 6059–6073.
- 141 A. Melber, U. Na, A. Vashisht, B. D. Weiler, R. Lill, J. A. Wohlschlegel and D. R. Winge, Role of Nfu1 and Bol3 in iron-sulfur cluster transfer to mitochondrial clients, *Elife*, 2016, **5**, e15991.
- 142 K. Cai, G. Liu, R. O. Frederick, R. Xiao, G. T. Montelione and J. L. Markley, Structural/Functional Properties of Human NFU1, an Intermediate [4Fe–4S] Carrier in Human Mitochondrial Iron-Sulfur Cluster Biogenesis, *Structure*, 2016, **24**, 2080–2091.
- 143 M. A. Uzarska, V. Nasta, B. D. Weiler, F. Spantgar, S. Ciofi-Baffoni, M. R. Saviello, L. Gonnelli, U. Muhlenhoff, L. Banci and R. Lill, Mitochondrial Bol1 and Bol3 function as assembly factors for specific iron-sulfur proteins, *Elife*, 2016, **5**, e16673.
- 144 H. Li and C. E. Outten, Monothiol CGFS glutaredoxins and BolA-like proteins: [2Fe–2S] binding partners in iron homeostasis, *Biochemistry*, 2012, **51**, 4377–4389.
- 145 V. Nasta, A. Giachetti, S. Ciofi-Baffoni and L. Banci, Structural insights into the molecular function of human [2Fe–2S] BOLA1-GRX5 and [2Fe–2S] BOLA3-GRX5 complexes, *Biochim. Biophys. Acta*, 2017, **1861**, 2119–2131.
- 146 V. D. Paul and R. Lill, Biogenesis of cytosolic and nuclear iron-sulfur proteins and their role in genome stability, *Biochim. Biophys. Acta*, 2015, **1853**, 1528–1539.
- 147 A. K. Sharma, L. J. Pallesen, R. J. Spang and W. E. Walden, Cytosolic iron-sulfur cluster assembly (CIA) system: factors, mechanism, and relevance to cellular iron regulation, *J. Biol. Chem.*, 2010, **285**, 26745–26751.
- 148 A. D. Tsaousis, E. Gentekaki, L. Eme, D. Gaston and A. J. Roger, Evolution of the cytosolic iron-sulfur cluster assembly machinery in Blastocystis species and other microbial eukaryotes, *Eukaryotic Cell*, 2014, **13**, 143–153.
- 149 O. Stehling, D. J. Netz, B. Niggemeyer, R. Rosser, R. S. Eisenstein, H. Puccio, A. J. Pierik and R. Lill, Human Nbp35 is essential for both cytosolic iron-sulfur protein assembly and iron homeostasis, *Mol. Cell. Biol.*, 2008, **28**, 5517–5528.
- 150 T. Okuno, H. Yamabayashi and K. Kogure, Comparison of intracellular localization of Nubp1 and Nubp2 using GFP fusion proteins, *Mol. Biol. Rep.*, 2010, **37**, 1165–1168.
- 151 A. Hausmann, D. J. Aguilar Netz, J. Balk, A. J. Pierik, U. Muhlenhoff and R. Lill, The eukaryotic P loop NTPase Nbp35: an essential component of the cytosolic and nuclear iron-sulfur protein assembly machinery, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 3266–3271.
- 152 D. J. Netz, A. J. Pierik, M. Stumpfig, U. Muhlenhoff and R. Lill, The Cfd1-Nbp35 complex acts as a scaffold for iron-sulfur protein assembly in the yeast cytosol, *Nat. Chem. Biol.*, 2007, **3**, 278–286.
- 153 L. J. Pallesen, N. Solodovnikova, A. K. Sharma and W. E. Walden, Interaction with Cfd1 increases the kinetic lability of FeS on the Nbp35 scaffold, *J. Biol. Chem.*, 2013, **288**, 23358–23367.
- 154 E. J. Camire, J. D. Grossman, G. J. Thole, N. M. Fleischman and D. L. Perlstein, The Yeast Nbp35-Cfd1 Cytosolic Iron-Sulfur Cluster Scaffold Is an ATPase, *J. Biol. Chem.*, 2015, **290**, 23793–23802.
- 155 H. Lange, A. Kaut, G. Kispal and R. Lill, A mitochondrial ferredoxin is essential for biogenesis of cellular iron-sulfur proteins, *Proc. Natl. Acad. Sci. U. S. A.*, 2000, **97**, 1050–1055.
- 156 J. Li, S. Saxena, D. Pain and A. Dancis, Adrenodoxin reductase homolog (Arh1p) of yeast mitochondria required for iron homeostasis, *J. Biol. Chem.*, 2001, **276**, 1503–1509.
- 157 D. J. Netz, M. Stumpfig, C. Dore, U. Muhlenhoff, A. J. Pierik and R. Lill, Tah18 transfers electrons to Dre2 in cytosolic iron-sulfur protein biogenesis, *Nat. Chem. Biol.*, 2010, **6**, 758–765.
- 158 L. Vernis, C. Facca, E. Delagoutte, N. Soler, R. Chanet, B. Guiard, G. Faye and G. Baldacci, A newly identified essential complex, Dre2-Tah18, controls mitochondria integrity and cell death after oxidative stress in yeast, *PLoS One*, 2009, **4**, e4376.
- 159 E. L. Bastow, K. Bych, J. C. Crack, N. E. Le Brun and J. Balk, NBP35 interacts with DRE2 in the maturation of cytosolic iron-sulphur proteins in Arabidopsis thaliana, *Plant J.*, 2017, **89**, 590–600.
- 160 K. Bych, D. J. Netz, G. Vigani, E. Bill, R. Lill, A. J. Pierik and J. Balk, The essential cytosolic iron-sulfur protein Nbp35 acts without Cfd1 partner in the green lineage, *J. Biol. Chem.*, 2008, **283**, 35797–35804.
- 161 M. B. Murataliev, R. Feyereisen and F. A. Walker, Electron transfer by diflavin reductases, *Biochim. Biophys. Acta*, 2004, **1698**, 1–26.
- 162 L. Banci, I. Bertini, S. Ciofi-Baffoni, F. Boscaro, A. Chatzi, M. Mikolajczyk, K. Tokatlidis and J. Winkelmann, Anamorsin is a 2Fe2S cluster-containing substrate of the Mia40-dependent mitochondrial protein trapping machinery, *Chem. Biol.*, 2011, **18**, 794–804.
- 163 L. Banci, S. Ciofi-Baffoni, M. Mikolajczyk, J. Winkelmann, E. Bill and M. E. Pandelia, Human anamorsin binds [2Fe–2S] clusters with unique electronic properties, *J. Biol. Inorg. Chem.*, 2013, **18**, 883–893.
- 164 Y. Zhang, C. Yang, A. Dancis and E. Nakamaru-Ogiso, EPR studies of wild type and mutant Dre2 identify essential [2Fe–2S] and [4Fe–4S] clusters and their cysteine ligands, *J. Biochem.*, 2017, **161**, 67–78.
- 165 D. J. Netz, H. M. Genau, B. D. Weiler, E. Bill, A. J. Pierik and R. Lill, The conserved protein Dre2 uses essential [2Fe–2S] and [4Fe–4S] clusters for its function in cytosolic iron-sulfur protein assembly, *Biochem. J.*, 2016, **473**, 2073–2085.
- 166 L. Banci, I. Bertini, V. Calderone, S. Ciofi-Baffoni, A. Giachetti, D. Jaiswal, M. Mikolajczyk, M. Piccioli and J. Winkelmann,

- Molecular view of an electron transfer process essential for iron-sulfur protein biogenesis, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 7136–7141.
- 167 G. Song, C. Cheng, Y. Li, N. Shaw, Z. C. Xiao and Z. J. Liu, Crystal structure of the N-terminal methyltransferase-like domain of anamorsin, *Proteins*, 2014, **82**, 1066–1071.
- 168 U. Muhlenhoff, J. Balk, N. Richhardt, J. T. Kaiser, K. Sipos, G. Kispal and R. Lill, Functional characterization of the eukaryotic cysteine desulfurase Nfs1p from *Saccharomyces cerevisiae*, *J. Biol. Chem.*, 2004, **279**, 36906–36915.
- 169 P. Haunhorst, E. M. Hanschmann, L. Brautigam, O. Stehling, B. Hoffmann, U. Muhlenhoff, R. Lill, C. Berndt and C. H. Lillig, Crucial function of vertebrate glutaredoxin 3 (PICOT) in iron homeostasis and hemoglobin maturation, *Mol. Biol. Cell*, 2013, **24**, 1895–1903.
- 170 E. Herrero and M. A. de la Torre-Ruiz, Monothiol glutaredoxins: a common domain for multiple functions, *Cell. Mol. Life Sci.*, 2007, **64**, 1518–1530.
- 171 H. Li, D. T. Mapolelo, S. Randeniya, M. K. Johnson and C. E. Outten, Human glutaredoxin 3 forms [2Fe–2S]-bridged complexes with human BOLA2, *Biochemistry*, 2012, **51**, 1687–1696.
- 172 P. Haunhorst, C. Berndt, S. Eitner, J. R. Godoy and C. H. Lillig, Characterization of the human monothiol glutaredoxin 3 (PICOT) as iron-sulfur protein, *Biochem. Biophys. Res. Commun.*, 2010, **394**, 372–376.
- 173 U. Muhlenhoff, S. Molik, J. R. Godoy, M. A. Uzarska, N. Richter, A. Seubert, Y. Zhang, J. Stubbe, F. Pierrel, E. Herrero, C. H. Lillig and R. Lill, Cytosolic monothiol glutaredoxins function in intracellular iron sensing and trafficking *via* their bound iron-sulfur cluster, *Cell Metab.*, 2010, **12**, 373–385.
- 174 H. Li, D. T. Mapolelo, N. N. Dingra, S. G. Naik, N. S. Lees, B. M. Hoffman, P. J. Riggs-Gelasco, B. H. Huynh, M. K. Johnson and C. E. Outten, The yeast iron regulatory proteins Grx3/4 and Fra2 form heterodimeric complexes containing a [2Fe–2S] cluster with cysteinyl and histidyl ligation, *Biochemistry*, 2009, **48**, 9569–9581.
- 175 L. Banci, S. Ciofi-Baffoni, K. Gajda, R. Muzzioli, R. Peruzzini and J. Winkelmann, N-terminal domains mediate [2Fe–2S] cluster transfer from glutaredoxin-3 to anamorsin, *Nat. Chem. Biol.*, 2015, **11**, 772–778.
- 176 Y. Saito, H. Shibayama, H. Tanaka, A. Tanimura, I. Matsumura and Y. Kanakura, PICOT is a molecule which binds to anamorsin, *Biochem. Biophys. Res. Commun.*, 2011, **408**, 329–333.
- 177 Y. Zhang, H. Li, C. Zhang, X. An, L. Liu, J. Stubbe and M. Huang, Conserved electron donor complex Dre2-Tah18 is required for ribonucleotide reductase metallocofactor assembly and DNA synthesis, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, E1695–E1704.
- 178 Y. Zhang, L. Liu, X. Wu, X. An, J. Stubbe and M. Huang, Investigation of *in vivo* diferric tyrosyl radical formation in *Saccharomyces cerevisiae* Rnr2 protein: requirement of Rnr4 and contribution of Grx3/4 AND Dre2 proteins, *J. Biol. Chem.*, 2011, **286**, 41499–41509.
- 179 H. Li, M. Stumpfig, C. Zhang, X. An, J. Stubbe, R. Lill and M. Huang, The diferric-tyrosyl radical cluster of ribonucleotide reductase and cytosolic iron-sulfur clusters have distinct and similar biogenesis requirements, *J. Biol. Chem.*, 2017, **292**, 11445–11451.
- 180 L. Banci, F. Camponeschi, S. Ciofi-Baffoni and R. Muzzioli, Elucidating the molecular function of human BOLA2 in GRX3-Dependent anamorsin maturation pathway, *J. Am. Chem. Soc.*, 2015, **137**, 16133–16134.
- 181 A. G. Frey, D. J. Palenchar, J. D. Wildemann and C. C. Philpott, A Glutaredoxin-BOLA Complex Serves as an Iron-Sulfur Cluster Chaperone for the Cytosolic Cluster Assembly Machinery, *J. Biol. Chem.*, 2016, **291**, 22344–22356.
- 182 X. Nuttle, G. Giannuzzi, M. H. Duyzend, J. G. Schraiber, I. Narvaiza, F. Camponeschi, S. Ciofi-Baffoni, P. H. Sudmant, O. Penn, G. Chiatante, M. Malig, J. Huddleston, C. Benner, H. A. F. Stessman, M. C. N. Marchetto, L. Denman, L. Harshman, C. Baker, A. Raja, K. Penewit, W. J. Tang, M. Ventura, F. Antonacci, J. M. Akey, C. T. Amemiya, L. Banci, F. H. Gage, A. Reymond and E. E. Eichler, Emergence of a *Homo sapiens*-specific gene family and the evolution of autism risk at chromosome 16p11.2, *Nature*, 2016, **536**, 205–209.
- 183 C. H. Lipper, M. L. Paddock, J. N. Onuchic, R. Mittler, R. Nechushtai and P. A. Jennings, Cancer-Related NEET Proteins Transfer 2Fe–2S Clusters to Anamorsin, a Protein Required for Cytosolic Iron-Sulfur Cluster Biogenesis, *PLoS One*, 2015, **10**, e0139699.
- 184 I. Ferecatu, S. Goncalves, M. P. Golinelli-Cohen, M. Clemancey, A. Martelli, S. Riquier, E. Guittet, J. M. Latour, H. Puccio, J. C. Drapier, E. Lescop and C. Bouton, The diabetes drug target MitoNEET governs a novel trafficking pathway to rebuild an Fe–S cluster into cytosolic aconitase/iron regulatory protein 1, *J. Biol. Chem.*, 2014, **289**, 28070–28086.
- 185 M. P. Golinelli-Cohen, E. Lescop, C. Mons, S. Goncalves, M. Clemancey, J. Santolini, E. Guittet, G. Blondin, J. M. Latour and C. Bouton, Redox Control of the Human Iron-Sulfur Repair Protein MitoNEET Activity via Its Iron-Sulfur Cluster, *J. Biol. Chem.*, 2016, **291**, 7583–7593.
- 186 F. Camponeschi, S. Ciofi-Baffoni and L. Banci, Anamorsin/Ndor1 Complex Reduces [2Fe–2S]-MitoNEET via a Transient Protein-Protein Interaction, *J. Am. Chem. Soc.*, 2017, **139**, 9479–9482.
- 187 K. Gari, A. M. Leon Ortiz, V. Borel, H. Flynn, J. M. Skehel and S. J. Boulton, MMS19 links cytoplasmic iron-sulfur cluster assembly to DNA metabolism, *Science*, 2012, **337**, 243–245.
- 188 O. Stehling, A. A. Vashisht, J. Mascarenhas, Z. O. Jonsson, T. Sharma, D. J. Netz, A. J. Pierik, J. A. Wohlschlegel and R. Lill, MMS19 assembles iron-sulfur proteins required for DNA metabolism and genomic integrity, *Science*, 2012, **337**, 195–199.
- 189 D. C. Odermatt and K. Gari, The CIA Targeting Complex Is Highly Regulated and Provides Two Distinct Binding Sites for Client Iron-Sulfur Proteins, *Cell Rep.*, 2017, **18**, 1434–1443.
- 190 C. Xu and J. Min, Structure and function of WD40 domain proteins, *Protein Cell*, 2011, **2**, 202–214.

- 191 V. Srinivasan, D. J. Netz, H. Webert, J. Mascarenhas, A. J. Pierik, H. Michel and R. Lill, Structure of the yeast WD40 domain protein Cia1, a component acting late in iron-sulfur protein biogenesis, *Structure*, 2007, **15**, 1246–1257.
- 192 M. Seki, Y. Takeda, K. Iwai and K. Tanaka, IOP1 protein is an external component of the human cytosolic iron-sulfur cluster assembly (CIA) machinery and functions in the MMS19 protein-dependent CIA pathway, *J. Biol. Chem.*, 2013, **288**, 16680–16689.
- 193 V. D. Paul, U. Muhlenhoff, M. Stumpfig, J. Seebacher, K. G. Kugler, C. Renicke, C. Taxis, A. C. Gavin, A. J. Pierik and R. Lill, The deca-GX3 proteins Yae1-Lto1 function as adaptors recruiting the ABC protein Rli1 for iron-sulfur cluster insertion, *Elife*, 2015, **4**, e08231.
- 194 E. L. Huttlin, L. Ting, R. J. Bruckner, F. Gebreab, M. P. Gygi, J. Szpyt, S. Tam, G. Zarraga, G. Colby, K. Baltier, R. Dong, V. Guarani, L. P. Vaites, A. Ordureau, R. Rad, B. K. Erickson, M. Wuhr, J. Chick, B. Zhai, D. Kolippakkam, J. Mintseris, R. A. Obar, T. Harris, S. Artavanis-Tsakonas, M. E. Sowa, C. P. De, J. A. Paulo, J. W. Harper and S. P. Gygi, The BioPlex Network: A Systematic Exploration of the Human Interactome, *Cell*, 2015, **162**, 425–440.
- 195 Y. Nicolet and J. C. Fontecilla-Camps, Structure-function relationships in [FeFe]-hydrogenase active site maturation, *J. Biol. Chem.*, 2012, **287**, 13532–13540.
- 196 D. W. Mulder, E. M. Shepard, J. E. Meuser, N. Joshi, P. W. King, M. C. Posewitz, J. B. Broderick and J. W. Peters, Insights into [FeFe]-hydrogenase structure, mechanism, and maturation, *Structure*, 2011, **19**, 1038–1052.
- 197 J. Balk, A. J. Pierik, D. J. Netz, U. Muhlenhoff and R. Lill, The hydrogenase-like Nar1p is essential for maturation of cytosolic and nuclear iron-sulphur proteins, *EMBO J.*, 2004, **23**, 2105–2115.
- 198 E. Urzica, A. J. Pierik, U. Muhlenhoff and R. Lill, Crucial role of conserved cysteine residues in the assembly of two iron-sulfur clusters on the CIA protein Nar1, *Biochemistry*, 2009, **48**, 4946–4958.
- 199 C. Cavazza, L. Martin, S. Mondy, J. Gaillard, P. Ratet and J. C. Fontecilla-Camps, The possible role of an [FeFe]-hydrogenase-like protein in the plant responses to changing atmospheric oxygen levels, *J. Inorg. Biochem.*, 2008, **102**, 1359–1365.
- 200 J. Balk, D. J. Aguilar Netz, K. Tepper, A. J. Pierik and R. Lill, The essential WD40 protein Cia1 is involved in a late step of cytosolic and nuclear iron-sulfur protein assembly, *Mol. Cell. Biol.*, 2005, **25**, 10833–10841.
- 201 M. Costanzo, B. VanderSluis, E. N. Koch, A. Baryshnikova, C. Pons, G. Tan, W. Wang, M. Usaj, J. Hanchard, S. D. Lee, V. Pelechano, E. B. Styles, M. Billmann, L. J. van Leeuwen, D. N. van Dyk, Z. Y. Lin, E. Kuzmin, J. Nelson, J. S. Piotrowski, T. Srikumar, S. Bahr, Y. Chen, R. Deshpande, C. F. Kurat, S. C. Li, Z. Li, M. M. Usaj, H. Okada, N. Pascoe, B. J. San Luis, S. Sharifpoor, E. Shuteriqi, S. W. Simpkins, J. Snider, H. G. Suresh, Y. Tan, H. Zhu, N. Malod-Dognin, V. Janjic, N. Przulj, O. G. Troyanskaya, I. Stagljar, T. Xia, Y. Ohya, A. C. Gingras, B. Raught, M. Boutros, L. M. Steinmetz, C. L. Moore, A. P. Rosebrock, A. A. Caudy, C. L. Myers, B. Andrews and C. Boone, A global genetic interaction network maps a wiring diagram of cellular function, *Science*, 2016, 353.
- 202 O. Stehling, J. Mascarenhas, A. A. Vashisht, A. D. Sheftel, B. Niggemeyer, R. Rosser, A. J. Pierik, J. A. Wohlschlegel and R. Lill, Human CIA2A-FAM96A and CIA2B-FAM96B integrate iron homeostasis and maturation of different subsets of cytosolic-nuclear iron-sulfur proteins, *Cell Metab.*, 2013, **18**, 187–198.
- 203 B. OuYang, L. Wang, S. Wan, Y. Luo, L. Wang, J. Lin and B. Xia, Solution structure of monomeric human FAM96A, *J. Biomol. NMR*, 2013, **56**, 387–392.
- 204 K. E. Chen, A. A. Richards, J. K. Ariffin, I. L. Ross, M. J. Sweet, S. Kellie, B. Kobe and J. L. Martin, The mammalian DUF59 protein Fam96a forms two distinct types of domain-swapped dimer, *Acta Crystallogr., Sect. D: Biol. Crystallogr.*, 2012, **68**, 637–648.
- 205 E. Weerapana, C. Wang, G. M. Simon, F. Richter, S. Khare, M. B. Dillon, D. A. Bachovchin, K. Mowen, D. Baker and B. F. Cravatt, Quantitative reactivity profiling predicts functional cysteines in proteomes, *Nature*, 2010, **468**, 790–795.
- 206 C. Zhai, Y. Li, C. Mascarenhas, Q. Lin, K. Li, I. Vyrides, C. M. Grant and B. Panaretou, The function of ORAOV1/LTO1, a gene that is overexpressed frequently in cancer: essential roles in the function and biogenesis of the ribosome, *Oncogene*, 2014, **33**, 484–494.
- 207 A. S. Upadhyay, K. Vonderstein, A. Pichlmair, O. Stehling, K. L. Bennett, G. Dobler, J. T. Guo, G. Superti-Furga, R. Lill, A. K. Overby and F. Weber, Viperin is an iron-sulfur protein that inhibits genome synthesis of tick-borne encephalitis virus via radical SAM domain activity, *Cell. Microbiol.*, 2014, **16**, 834–848.
- 208 A. S. Upadhyay, O. Stehling, C. Panayiotou, R. Rosser, R. Lill and A. K. Overby, Cellular requirements for iron-sulfur cluster insertion into the antiviral radical SAM protein viperin, *J. Biol. Chem.*, 2017, **292**, 13879–13889.
- 209 P. Csere, R. Lill and G. Kispal, Identification of a human mitochondrial ABC transporter, the functional orthologue of yeast Atm1p, *FEBS Lett.*, 1998, **441**, 266–270.
- 210 S. Kushnir, E. Babychuk, S. Storozhenko, M. W. Davey, J. Papenbrock, R. R. De, G. Engler, U. W. Stephan, H. Lange, G. Kispal, R. Lill and M. Van Montagu, A mutation of the mitochondrial ABC transporter Sta1 leads to dwarfism and chlorosis in the Arabidopsis mutant starik, *Plant Cell*, 2001, **13**, 89–100.
- 211 S. Chen, R. Sanchez-Fernandez, E. R. Lyver, A. Dancis and P. A. Rea, Functional characterization of AtATM1, AtATM2, and AtATM3, a subfamily of Arabidopsis half-molecule ATP-binding cassette transporters implicated in iron homeostasis, *J. Biol. Chem.*, 2007, **282**, 21561–21571.
- 212 M. Martinez-Garcia, J. Campos-Salinas, M. Cabello-Donayre, E. Pineda-Molina, F. J. Galvez, L. M. Orrego, M. P. Sanchez-Canete, S. Malagarie-Cazenave, D. M. Koeller and J. M. Perez-Victoria, LmABC3, an atypical mitochondrial ABC transporter essential for Leishmania major virulence,

- acts in heme and cytosolic iron/sulfur clusters biogenesis, *Parasites Vectors*, 2016, **9**, 7.
- 213 N. Mitsuhashi, T. Miki, H. Senbongi, N. Yokoi, H. Yano, M. Miyazaki, N. Nakajima, T. Iwanaga, Y. Yokoyama, T. Shibata and S. Seino, MTABC3, a novel mitochondrial ATP-binding cassette protein involved in iron homeostasis, *J. Biol. Chem.*, 2000, **275**, 17536–17540.
- 214 C. Kumar, A. Igarria, B. d'Autreaux, A. G. Planson, C. Junot, E. Godat, A. K. Bachhawat, A. Delaunay-Moisan and M. B. Toledano, Glutathione revisited: a vital function in iron metabolism and ancillary role in thiol-redox control, *EMBO J.*, 2011, **30**, 2044–2056.
- 215 W. Qi, J. Li, C. Y. Chain, G. A. Pasquevich, A. F. Pasquevich and J. A. Cowan, Glutathione complexed Fe–S centers, *J. Am. Chem. Soc.*, 2012, **134**, 10745–10748.
- 216 J. Li, S. A. Pearson, K. D. Fenk and J. A. Cowan, Glutathione-coordinated [2Fe–2S] cluster is stabilized by intramolecular salt bridges, *J. Biol. Inorg. Chem.*, 2015, **20**, 1221–1227.
- 217 W. Qi, J. Li, C. Y. Chain, G. A. Pasquevich, A. F. Pasquevich and J. A. Cowan, Glutathione-complexed iron-sulfur clusters. Reaction intermediates and evidence for a template effect promoting assembly and stability, *Chem. Commun.*, 2013, **49**, 6313–6315.
- 218 J. Li and J. A. Cowan, Glutathione-coordinated [2Fe–2S] cluster: a viable physiological substrate for mitochondrial ABCB7 transport, *Chem. Commun.*, 2015, **51**, 2253–2255.
- 219 W. Qi, J. Li and J. A. Cowan, A structural model for glutathione-complexed iron-sulfur cluster as a substrate for ABCB7-type transporters, *Chem. Commun.*, 2014, **50**, 3795–3798.
- 220 V. Srinivasan, A. J. Pierik and R. Lill, Crystal structures of nucleotide-free and glutathione-bound mitochondrial ABC transporter Atm1, *Science*, 2014, **343**, 1137–1140.
- 221 G. Kispal, P. Csere, B. Guiard and R. Lill, The ABC transporter Atm1p is required for mitochondrial iron homeostasis, *FEBS Lett.*, 1997, **418**, 346–350.
- 222 J. Y. Lee, J. G. Yang, D. Zhitnitsky, O. Lewinson and D. C. Rees, Structural basis for heavy metal detoxification by an Atm1-type ABC exporter, *Science*, 2014, **343**, 1133–1136.
- 223 T. A. Schaedler, J. D. Thornton, I. Kruse, M. Schwarzlander, A. J. Meyer, H. W. van Veen and J. Balk, A conserved mitochondrial ATP-binding cassette transporter exports glutathione polysulfide for cytosolic metal cofactor assembly, *J. Biol. Chem.*, 2014, **289**, 23264–23274.
- 224 H. Lange, T. Lisowsky, J. Gerber, U. Muhlenhoff, G. Kispal and R. Lill, An essential function of the mitochondrial sulfhydryl oxidase Erv1p/ALR in the maturation of cytosolic Fe/S proteins, *EMBO Rep.*, 2001, **2**, 715–720.
- 225 H. K. Ozer, A. C. Dlouhy, J. D. Thornton, J. Hu, Y. Liu, J. J. Barycki, J. Balk and C. E. Outten, Cytosolic Fe–S Cluster Protein Maturation and Iron Regulation Are Independent of the Mitochondrial Erv1/Mia40 Import System, *J. Biol. Chem.*, 2015, **290**, 27829–27840.