

Proteomic Quantification of Human Blood-Brain Barrier SLC and ABC Transporters in Healthy Individuals and Dementia Patients

DOI:

[10.1021/acs.molpharmaceut.8b01189](https://doi.org/10.1021/acs.molpharmaceut.8b01189)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Al-Majdoub, Z., Al Feteisi, H., Achour, B., Warwood, S., Neuhoff, S., Rostami-Hodjegan, A., & Barber, J. (2019). Proteomic Quantification of Human Blood-Brain Barrier SLC and ABC Transporters in Healthy Individuals and Dementia Patients. *Molecular Pharmaceutics*, 16(3), 1220-1233. <https://doi.org/10.1021/acs.molpharmaceut.8b01189>

Published in:

Molecular Pharmaceutics

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Proteomic Quantification of Human Blood-Brain Barrier SLC and ABC Transporters in Healthy Individuals and Dementia Patients

Journal:	<i>Molecular Pharmaceutics</i>
Manuscript ID	mp-2018-01189c.R1
Manuscript Type:	Article
Date Submitted by the Author:	02-Jan-2019
Complete List of Authors:	Al-Majdoub, Zubida; University of Manchester, Centre for Applied Pharmacokinetic Research; Division of Pharmacy and Optometry Al Feteisi, Hajar; University of Manchester, Centre for Applied Pharmacokinetic Research; Division of Pharmacy and Optometry Achour, Brahim; University of Manchester, Centre for Applied Pharmacokinetic Research; Division of Pharmacy and Optometry Warwood, Stacey; University of Manchester, Biological Mass Spectrometry Core Facility Neuhoff, Sibylle; Simcyp Ltd. , (a Certara company) Rostami-Hodjegan, Amin; University of Manchester, Centre for Applied Pharmacokinetic Research; Division of Pharmacy and Optometry; Simcyp Ltd. , (a Certara company) Barber, Jill; University of Manchester, Centre for Applied Pharmacokinetic Research; Division of Pharmacy and Optometry

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Proteomic Quantification of Human Blood-Brain**
6
7
8
9 **Barrier SLC and ABC Transporters in Healthy**
10
11
12
13 **Individuals and Dementia Patients**
14
15
16
17
18
19

20 **Zubida M. Al-Majdoub^{1*}, Hajar Al Feteisi¹, Brahim Achour¹,**
21 **Stacey Warwood², Sibylle Neuhoff³, Amin Rostami-Hodjegan^{1,3},**
22 **Jill Barber^{1*}**
23
24
25
26
27
28

29 ¹ Centre for Applied Pharmacokinetic Research (CAPKR), University of Manchester, UK.

30 ² Biological Mass Spectrometry Core Facility, University of Manchester, UK.

31 ³ Certara UK Limited, Simcyp Division, Level 2-Acero, 1 Concourse Way, Sheffield, S1 2BJ,
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
UK.

*To whom correspondence should be addressed: Zubida.al-majdoub@manchester.ac.uk;
Jill.barber@manchester.ac.uk

KEYWORDS: Blood-Brain Barrier (BBB), Alzheimer's Disease (AD), Dementia with
Lewy Bodies (DLB), Solute carrier (SLC) and ATP-binding cassette (ABC) transporters.

ABSTRACT

The Blood-Brain Barrier (BBB) maintains brain homeostasis by controlling traffic of molecules from the circulation into the brain. This function is predominantly dependent on proteins expressed at the BBB, especially transporters and tight junction proteins. Alterations to the level and function of BBB proteins can impact on the susceptibility of the central nervous system to exposure to xenobiotics in the systemic circulation with potential consequent effects on brain function. In this study, expression profiles of drug transporters and solute carriers in the BBB were assessed in tissues from healthy individuals (n=12), Alzheimer's patients (n=5) and Dementia with Lewy Bodies patients (n=5), using targeted, AMRT (Accurate Mass Retention Time) and global proteomic methods. A total of 66 transporters were quantified, 19 for the first time in the BBB. A further 20 novel transporters were identified but not quantified. The global proteomic method identified another 3333 BBB proteins. Transporter abundances, taken together with the scaling factor microvessel protein content per unit tissue (BMvPGB also measured here) can be used in quantitative systems pharmacology models predicting drug disposition in the brain and permitting dose adjustment (precision dosing) in special populations of patients, such as those with dementia. Even in this small study, we see differences in transporter profile between healthy and diseased brain tissue.

INTRODUCTION

The two main barrier interfaces between the blood and the brain are the blood–cerebral spinal fluid barrier (BCSFB) across the choroid plexus epithelial cells and the blood–brain barrier (BBB) across the cerebral endothelial cells. These barriers play vital roles in controlling transport between systemic blood and brain tissue; this means that they control the concentration of drugs in the central nervous system (CNS) and this in turn affects their efficacy and toxicity.^{1,2} The contribution of proteins regulating traffic of molecules across the BBB to the maintenance of brain homeostasis, and their effect on the disposition of substrate drugs targeted at CNS conditions is widely appreciated.^{3,4} These transporters at the BBB act as gatekeepers to the CNS and play a critical role in drug and xenobiotic brain disposition.⁵

Membrane transporters can be categorized molecularly into two major superfamilies: solute carriers (SLCs) and ABC-type efflux transporters. Based on function, transporters serve two main roles: efflux and uptake (influx) transport. All ATP-dependent transporters are efflux transporters from the point of view of the cell. However, SLCs can work as uptake, efflux, as well as equilibrative transporters. These transporters are a key protective element of the BBB, actively transporting a wide variety of xenobiotics back into the circulation. Uptake transporters, such as several SLCs, can facilitate uptake into the cerebrospinal fluid from the blood.⁶

One of the major challenges of modern day medicine is the understanding, diagnosis, and treatment of complex and heterogeneous age-related disorders, such as dementia. Despite the significant commercial opportunities that exist, attempts to discover effective drugs to treat neurological diseases have proved so difficult that many companies have withdrawn

1
2 from the field.⁷ Compared with other disease areas, failure rates in clinical trials are high for
3
4 neurological disease largely due to the complexity of the human brain.⁸
5
6

7 The most common cause of dementia is Alzheimer's disease (AD), which is
8 characterized by a progressive decline in memory and cognitive functions.⁹ The classical
9
10 neuropathological hallmarks of AD are extracellular deposits of amyloid-beta and tau
11
12 proteins.^{10,11} Recent evidence suggests that the neurodegenerative process that leads to the
13
14 development of AD is highly associated with a significant dysfunction of the BBB.¹² Unlike
15
16 the genome, which is relatively static, the proteome is extremely dynamic and constantly
17
18 changing in response to internal (e.g. ageing, disease status) and external (e.g. toxin
19
20 exposure) stimuli. Possibly, alterations in the expression of proteins have functional
21
22 consequences and lead to a variety of pathophysiological conditions. Indeed, proteome
23
24 analysis has been suggested to hold the key to improving understanding, diagnosis, and
25
26 treatment of AD, as well as monitoring the progression of the disease.^{13,14} It is therefore
27
28 crucial to understand changes in transporter expression and evaluate inter-individual variation
29
30 in protein levels between healthy and diseased brains.
31
32
33
34
35

36
37 *In vitro* disposition data obtained from cellular brain models, such as neurovascular
38
39 endothelial cells, are commonly scaled using *in vitro-in vivo* extrapolation (IVIVE) to
40
41 ultimately predict drug/xenobiotic disposition *in vivo*. Helms *et al.* evaluated the advantages
42
43 and disadvantages of available brain cell-culture models and commented that, to date, there is
44
45 no ideal BBB *in vitro* model.¹⁵ IVIVE relies upon robust estimates of physiologically-
46
47 relevant scaling factors, including protein content of the microvascular fraction (brain
48
49 microvessels protein per gram of brain, BMvPGB) and transporter expression at the BBB.
50
51 These data are essential for filling the existing gaps in models for human brain disease and
52
53 prediction of appropriate drug dosing for different disease populations.¹⁶ It is notable that
54
55 there are limited data in the literature for the protein content scaling factor (BMvPGB), which
56
57
58
59
60

1
2 has only been reported in healthy individuals,¹⁷ and therefore, there is a need to assess both
3
4 protein content and transporter expression in health and dementia.
5
6

7 The aim of this study was to investigate alterations in the proteomic composition of
8
9 the blood-brain barrier between healthy and dementia brains, focusing on drug and
10
11 endogenous transporters, using a proteomic strategy that combines both targeted and global
12
13 analyses. In addition, the total protein content of the microvascular fraction (BMvPGB) was
14
15 assessed in healthy and dementia brains. Brain samples were from 22 donors, consisting of
16
17 healthy individuals and patients with either AD or DLB.
18
19
20
21
22

23 **MATERIALS AND METHODS**

24
25 Materials and chemicals were purchased from Sigma-Aldrich (Poole, UK) unless
26
27 otherwise indicated. The unlabeled non-naturally occurring peptide (NNOP) calibrator (Glu-
28
29 fibrinopeptide B (EGVNDNEEGFFSAR, purity 95%) was purchased from Severn Biotech
30
31 (Worcestershire, UK). Lysyl Endopeptidase (Lys-C) was purchased from Wako (Osaka,
32
33 Japan). CComplete Mini, EDTA-free protease inhibitor cocktail, and recombinant proteomic-
34
35 grade trypsin were supplied by Roche Applied Sciences (Mannheim, Germany). Two
36
37 TransCATs (artificial proteins, see below) were used; the Liver TransCAT protein was
38
39 produced in-house as previously described.¹⁸ The Brain TransCAT was designed and
40
41 expressed in-house. Both genes were synthesized by Entelechon GmbH (Regensburg,
42
43 Germany) and the QconCAT proteins were used for quantification of brain transporters.
44
45 Bicinchoninic acid (BCA) protein assay kit was purchased from Pierce (Rockford, IL, USA).
46
47 PiColorLock™ phosphate detection system was purchased from Innova Biosciences
48
49 (Cambridge, UK). Solvents used in this study were of HPLC Chromasolv grade (Thermo
50
51 Fisher, Waltham, MA).
52
53
54
55
56
57
58
59
60

QconCAT standards “TransCATs”

Two transporter QconCAT “TransCAT” standards were used in this study: named Liver TransCAT and Brain TransCAT. The “Liver TransCAT” standard has previously been described.¹⁸ The “Brain TransCAT” is a novel QconCAT for the quantification of human brain drug transporter proteins that have a crucial role in disposition of drugs and endogenous compounds across the BBB. The unique tryptic peptides making up the Brain TransCAT were two each from nine transporters (ABCA1, ABCA2, ABCC5, SLC7A5, SLC16A1, SLC22A4, SLC29A2, SLCO2B1, SLCO1C1) and three each from seven transporters (ABCA7, ABCC1, SLC22A2, SLC22A6, SLC22A8, SLC47A2, SLC29A1). Two peptides unique to each neurovascular unit cell marker protein of endothelial cells (GLUT1, PECAM1), pericytes (CSPG4, NG2), astrocytes (GFAP) and neurons (SYP) were also included to assess quantitatively the enrichment and the level of contamination of the isolated microvessels. In addition, two receptor proteins, vascular endothelial growth factor receptor 2 (KDR) and nuclear receptor subfamily 1 group I member 2 (NR1I2), represented by two peptides each, were included.

The candidate peptide list was generated by combining sequences from SRMATlas (www.srmatlas.org) with those predicted to ionize efficiently in addition to peptides identified experimentally by global LC–MS/MS analysis of brain microvessel proteins. Peptides containing methionine or cysteine residues and non-unique or membrane embedded sequences were excluded for the final list. Peptides were checked against the Uniprot database for known polymorphisms and post-translational modifications. Finally, a Glu-fibrinopeptide B (GFB) peptide was included to enable quantification of the TransCAT itself. The full list of peptides of the Brain TransCAT is included in Table ES1.

Brain TransCAT gene synthesis and expression

The DNA construct of the TransCAT was produced by PolyQuant GmbH (Regensburg, Germany) using the expression vector pET21a with codons optimized for both expression and minimized secondary structure formation in mRNA. His-tag was added to the C- terminus to enable purification. The TransCAT sequence was fused to a sequence comprising peptides from *E. coli* core ribosomal proteins that had previously served as a QconCAT for those proteins¹⁹ and sacrificial sequences were added to the N-terminus to protect the proteotypic peptides from potential N-terminal degradation. TransCAT transformation, expression in ¹³C heavy-isotope enriched medium and purification were performed as previously described.¹⁸ TransCAT (77 kDa) was then added in isotopically labelled form to the brain tissue prior to protease digestion. Proteolysis allows release of standard peptides in 1:1 stoichiometry, leading to the estimation of the concentrations of the target proteins using the QconCAT peptides as standards.

The Brain TransCAT was quantified using a label-free AQUA peptide “Glu-fibrinopeptide B (GFB)”. A known amount of GFB peptide was added to the TransCAT preparation and analyzed simultaneously with sample proteins. LC-MS/MS analysis of the TransCAT-analyte mixture yields the relative intensity of the TransCAT and analyte ions, and since the TransCAT is present at a known concentration, the quantity of analyte can be calculated directly.

Human brain samples

Frozen brain frontal cortexes of 22 donors were supplied by the Manchester Brain Bank (University of Manchester) with approval for collection, transfer, storage and handling (Newcastle and North Tyneside REC) under Manchester Brain Bank generic tissue bankethics (H09/0906). The samples were supplied from healthy (n=12), DLB (n=5) and AD (n=5). Donor demographic and clinical details are presented in Table S1. These samples were

1
2 used for global and targeted proteomic analysis of isolated brain microvessels as described
3
4 below.

7 **Tissue preparation**

9 Microvessels were isolated using optimized tissue processing methods, as previously
10 described,²⁰ with some modifications. Briefly, 3 grams of frozen human brain tissue were
11 thawed on ice and all subsequent steps were performed on ice. Brain cortex was finely
12 minced and homogenized using 20 up-and-down non-rotated strokes in lysis buffer (Buffer
13 A) using volumes of 4-5 mL/g tissue (8 g/L NaCl, 400 mg/L KCl, 185.4 mg/L CaCl₂·2H₂O,
14 60 mg/L KH₂PO₄, 200 mg/L MgSO₄·7H₂O, 350 mg/L NaHCO₃, 1 g/L dextrose anhydrous,
15 90 mg/L Na₂HPO₄·7H₂O, pH 7.4) in a 50 mL tube. Protease inhibitor was added to the
16 homogenate to prevent proteolytic degradation in subsequent sample preparation steps,
17 according to the manufacturer's instructions (1 tablet per 10 mL). The homogenates were
18 centrifuged at 2000 g for 10 min at 4°C. The resulting pellet was suspended in 16% dextran
19 and subsequently centrifuged at 5,500 g for 15 min at 4°C. The supernatant was transferred to
20 a fresh tube and centrifuged again in a similar way before suspending the two pellets in
21 10 mL of Buffer B (lysis buffer containing 5 mg/mL bovine serum albumin [BSA]). The
22 suspension was then passed through a 20 µm nylon mesh, using <20 mL Buffer B to rinse the
23 tube and mesh of residual homogenate. The retained fraction was transferred to centrifuge
24 tubes and centrifuged at 2000 g for 5 min at 4°C. The suspended pellet was then dissolved in
25 1 mL Buffer B and further centrifuged at 1,000 g for 5 min at 4°C. Finally, the microvessels
26 pellet was resuspended in 1 mL isotonic buffer (1 M Tris-HCl, 10 mM NaCl, 1.5 mM MgCl₂,
27 pH 7.4) and sonicated using a S220 Focused-ultrasonicator (COVARIS, UK), then stored at -
28 80°C until further analysis. Protein content in the microvessels and homogenates was
29 determined using Bicinchoninic acid (BCA) protein assay kit according to the manufacturer's
30 instructions.

Measurement of ATPase enzymatic activity

To assess the level of enrichment of microvessel proteins, adenosine triphosphatase (ATPase) activity was determined in isolated microvessels and homogenates by a colorimetric method.

The assay was performed according to manufacturer's protocol.

Sample preparation of brain microvessels for global and targeted proteomics

The microvessels protein was digested on filter units. The filter-aided sample preparation (FASP) protocol used here is an optimized version of the previously reported method^{19,21} and the detailed procedure is described in our previous study.²⁰ Briefly, the protocol involved sodium deoxycholate incubation for solubilisation. We chose FASP over in-solution digestion for its simplicity, time-efficiency, efficient removal of contaminants, and its applicability to clinically-relevant enzymes and transporters.²⁰

Three technical replicates for each of the 22 human brain microvessel samples were prepared. The first technical replicate was spiked with Liver TransCAT only, the second replicate was spiked with Liver and Brain TransCATs and the third replicate was spiked with Brain TransCAT only. A mixture of protein extracts (20-50 μ g) and TransCATs (250-500 fmol) were solubilized with 10% (w/v) sodium deoxycholate at room temperature for 10 min, and cysteine disulfide bonds were reduced by adding 0.1 M Tris-HCl buffer, pH 8.5, containing 60 mM dithiothreitol and 8 M urea followed by incubation with mixing at 56°C for 30 min. Ultracel 10,000 MWCO (molecular weight cut-off of 10 kDa) centrifugal filter units (Amicon Ultra, Merck) were washed using two steps of centrifugation at 14,000 g for 20 minutes with 200 μ L of 8 M urea in 0.1 M Tris-HCl, pH 8.5. Solubilized and reduced protein samples were then added to the filter units, followed by a centrifugation step at 14,000 g for 30 min. Protein extracts were buffer-exchanged using 8 M urea in 0.1 M Tris-HCl, pH 8.5, by centrifugation at 14,000 g for 20 min. Alkylation of reduced disulfide bonds was carried out by incubation with 200 μ L of 50 mM iodoacetamide in 8 M urea and 0.1 M

1
2 Tris-HCl, pH 8.5, in the dark for 30 min followed by centrifugation as above. A 200 μ L
3
4 volume of 8 M urea in 0.1 M Tris-HCl, pH 8.5, was added and centrifugation was carried out
5
6 to remove the alkylating agent, followed by three buffer exchanges with 200 μ L of 1 M urea
7
8 in 50 mM ammonium bicarbonate buffer, pH 8.0. A volume of 40 μ L of 50 mM ammonium
9
10 bicarbonate buffer, pH 8.0, was added to the filter to prevent evaporation of the digestion
11
12 mix. Lys-C (10 μ L, 0.1 mg/ml), at mass ratio 1:50 enzyme:protein, was added and samples
13
14 incubated at 30°C for 3 h, followed by a second dose of Lys-C and incubation for another 3
15
16 h. Trypsin (10 μ L, 0.2 mg/ml), at mass ratio 1:25 enzyme:protein was added and incubated
17
18 overnight (16 h) at 37°C. The Lys-C treatment step was repeated to increase digestion
19
20 efficiency and minimize missed cleavages. The samples were then centrifuged at 14000 g for
21
22 20 min and 100 μ L of 0.5 M NaCl solution was added to the filter followed by centrifugation
23
24 at 14,000 g for 20 min to collect the remaining peptides. Samples were then vacuum dried
25
26 and desalted on a C18 spin-column, followed by analysis by targeted LC-MRM (multi
27
28 reaction monitoring) MS/MS on a QTrap 6500 mass spectrometer and global LC-MS/MS on
29
30 a Q Exactive Orbitrap. Amount of total protein between 1.2-1.5 μ g was loaded on a QTrap
31
32 6500 mass spectrometer for MRM analysis.
33
34
35
36
37
38

39 **Global LC-MS/MS proteomic analysis**

41 For LC-MS/MS analysis using the Q Exactive platform, desalted samples were
42
43 resuspended in 10 μ L of loading buffer (5% acetonitrile in 0.1% formic acid [FA]) and 1 μ L
44
45 (2.5 μ g total protein) was loaded onto an UltiMate® 3000 (Dionex, Surrey, UK) liquid
46
47 chromatography (LC) system coupled on-line to a Q Exactive™ HF Hybrid Quadrupole-
48
49 Orbitrap™ mass spectrometer (Thermo Fisher, Bremen, Germany). Peptides were reversed-
50
51 phase separated on a CSH C18 analytical column (250 mm \times 75 μ m i.d., 1.7 μ m) (Waters,
52
53 UK). A multi-step gradient was used from 4% Buffer B (100% acetonitrile, 0.1% FA) and
54
55 96% Buffer A (0.1% FA in HPLC water) to 60% Buffer B over 165 min at a flow rate of 300
56
57
58
59
60

1
2 nL min⁻¹. The LC eluent was injected into the mass spectrometer. All m/z values of eluting
3
4 ions were measured in an Orbitrap MS, set at a resolution of 120,000. Data dependent scans
5
6 were employed to automatically select the 12 most abundant ions for fragmentation by
7
8 higher-energy collisional dissociation (HCD) in the quadrupole mass analyser. Only peptide
9
10 ions with charge states of 2+, 3+, and 4+ were selected for fragmentation. Finally, the
11
12 measurement of the resulting fragment ions was performed in the Orbitrap analyser, set at a
13
14 resolution of 30,000.
15
16
17

18 **MRM assays and targeted LC–MS/MS analysis**

19
20 Peptide samples were analysed by LC-MRM-MS/MS using an UltiMate® 3000 Rapid
21
22 Separation LC (RSLC, Thermo Fisher Scientific, Waltham, MA) coupled to a 6500 QTRAP
23
24 (SCIEX, Framingham, MA) mass spectrometer. Peptide mixtures were separated using a
25
26 gradient from 95% Buffer A (0.1% FA in HPLC water) and 5% Buffer B (0.1% FA in
27
28 acetonitrile) to 30% B, over 46 min at 300 nL min⁻¹, using a M-Class CSH C18 analytical
29
30 column (250 mm x 75 µm i.d., 1.7 µm) (Waters, Milford, MA).
31
32
33

34
35 Signature peptide sequences from both TransCATs were concatenated *in silico*, along
36
37 with the sequence of Glu-fibrinopeptide B (GFB), used as an internal standard for the
38
39 QconCATs. Skyline version 4.1

40
41 (<https://skyline.ms/project/home/software/Skyline/begin.view>)²² was used to generate
42
43 theoretical transitions for each peptide in the sequence. The peptide settings were as follows:
44
45 digestion enzyme was trypsin, no missed cleavages were allowed. Peptide lengths were 6-20
46
47 residues, and carbamidomethylation (C) was included as a fixed modification, oxidation (M)
48
49 as a variable modification. The transition settings were as follows: precursor charge states of
50
51 2+ and 3+ were included, with y product ions included as 1+ only. Products ions (2-5) were
52
53 selected from each precursor ion. In addition, any product ions resulting from fragmentation
54
55
56
57
58
59
60

1
2 N-terminal to proline and C-terminal to glutamic acid or aspartic acid were included. The m/z
3
4 range was set to a minimum of 350 Th and a maximum of 1250 Th.
5
6

7 **Data analysis and protein quantification**

8
9 For label-free quantification, raw Orbitrap MS files were imported into Progenesis software
10
11 (Nonlinear Dynamics, version 4.0). Data in Mascot generic format (mgf) were exported
12
13 directly from Progenesis, and MS/MS spectra were searched using MASCOT against a
14
15 human proteome database containing 71,599 entries (UniProt, May 2017) and a decoy
16
17 database of the predicted proteome for Homo sapiens. The search criteria were set as follows:
18
19 Full tryptic specificity was required (cleavage after lysine or arginine residues); one missed
20
21 cleavage was allowed; carbamidomethylation (C) was set as a fixed modification; and
22
23 oxidation (M), deamidation (NQ) were set as variable modifications. The mass tolerance was
24
25 set to 5 ppm for precursor ions and 0.5 Da for fragment ions. Results from the database
26
27 search were imported into Progenesis, with mapping of peptide identifications to MS1
28
29 features. Quantitative data analysis of global proteomics was carried out as previously
30
31 described.²³ Briefly, label-free protein quantification was performed using the Hi3/Hi2
32
33 method.²⁴ This approach uses the sum of signal intensities of the three (Hi3) or two (Hi2) best
34
35 ionizing unique peptides of any given protein relative to the sum of signal intensities of the
36
37 three (Hi3) or two (Hi2) best ionizing unique peptides of a reference protein (in this case
38
39 ATP1A1 or ABCB1) quantified using MRM. Percentage identical peptides (PIP) and
40
41 percentage identical proteins (PIPr) were estimated as previously described.²⁰
42
43
44
45
46
47

48 For MRM analysis, all peak integration and review were performed using Skyline and
49
50 results were subsequently exported to Microsoft Excel 2010 for quantitative calculations.
51
52 Protein quantification data are provided in Table ES3 and ES6.
53
54

55 Using LC-Orbitrap MS data, Accurate Mass and Retention Time (AMRT) method
56
57 was performed as a second targeted proteomic approach to quantify transporters within
58
59
60

1
2 human brain microvessels. This approach is based on the peptide intensity in the MS
3 spectrum by using the accurate mass measurement and corresponding retention time for a
4 peptide signal. The ion peak intensity of the heavy and light versions of the same peptide are
5 compared across samples and the relative abundance of the parent protein is inferred from the
6 intensity ratios.²⁵

13 **Sub-cellular localization and functional analysis**

16 Protein data annotation for sub-cellular localization was done by database search
17 against UniProtKB (<http://www.uniprot.org/>), the Gene Ontology Project (GO,
18 <http://geneontology.org/>) and the Human Protein Atlas (HPA, <https://www.proteinatlas.org/>).
19 The PANTHER (Protein ANalysis THrough Evolutionary Relationships) classification
20 system version 12.0 (<http://www.pantherdb.org/>) was used for bioinformatics analysis of all
21 identified proteins to assign protein class and molecular function.²⁶

29 **Statistical data analysis**

32 For the three sample sets, the abundance data were described as mean \pm SD,
33 coefficients of variation (CV), and fold difference (maximum/minimum). Normality of
34 distribution was assessed and outliers were identified using iterative Grubbs' test using a cut-
35 off *p*-value of 0.01. Statistical difference between mean values of healthy and disease sets
36 was determined by ANOVA analysis followed by post-hoc unpaired *t*-test. The *p*-value for
37 significance was set at 0.05, and Bonferroni-corrected for multiple tests (0.025). Principal
38 coordinates/components analysis (PCA) and hierarchical cluster analysis (HCA) were
39 performed using combined data from each sample and relationships were computed using
40 Euclidian distance between samples and Spearman correlation (*R*_s) between genes. For inter-
41 correlations, *R*_s > 0.60 and *R*² > 0.30 and a Bonferroni-corrected *p*-value of < 0.004 were
42 considered for significant correlations taking into account the effect of the units of
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 measurement²³ Graphs were generated using GraphPad Prism version 7.04/7.0c for Windows
3
4 (GraphPad Software, San Diego, California, USA) and R version 3.1.0.
5
6

7 **RESULTS**

8 **Isolation of human brain microvessels**

9
10
11 Microvessels isolation was performed according to our previous published method²⁰
12
13 with additional new steps to optimize for human (as opposed to rat) tissue; the use of a 20 μm
14
15 nylon mesh instead of 100 μm and 20 μm mesh sizes for crude microvessel extraction was
16
17 adopted to remove small contaminants. We also introduced a focused-ultrasonicator step to
18
19 shear the membrane and liberate proteins; this sonicator provides additional energy and
20
21 temperature control over traditional low frequency laboratory sonicators. We measured
22
23 protein concentration following sonication using the BCA assay and compared the results to
24
25 those obtained using a traditional laboratory sonicator. We found the protein concentration
26
27 was higher and more reproducible when using the focused sonicator compared with the
28
29 traditional laboratory sonicator. Our optimized steps may overcome some of the difficulties
30
31 of isolation of microvessels from human tissue. Figure 1A shows the experimental workflow
32
33 followed in this study.
34
35
36
37
38

39 **Characterisation of the quality of isolated microvessels by LC-MS/MS**

40
41 To confirm the enrichment and purity of the isolated microvessels, the absolute
42
43 protein expression of cell-specific markers of endothelial cells (GLUT1, PECAM1 and
44
45 VWF), astrocytes (GFAP), neurons (SYP) and pericytes (CSPG4/NG2) were measured using
46
47 quantitative global and MRM approaches. Based on the quantification values, the expression
48
49 of endothelial markers ranged from high (GLUT1 between 4.05-57.8 pmol/mg protein) to
50
51 medium/low abundance (VWF, 0.39-9.65 pmol/mg protein; PECAM1, 0.30-1.67 pmol/mg
52
53 protein). GLUT1 protein was highly expressed in the microvessels, and levels of the pericyte
54
55 marker (CSPG4/NG2) and neuron marker (SYP) were very low in all the 22 samples. The
56
57
58
59
60

1
2 second highest marker abundance (after GLUT1) was recorded for the GFAP marker,
3
4 representing contamination by astrocyte endfeet, which form an integral part of the
5
6 neurovascular unit and are extremely difficult to remove without extensive loss of endothelial
7
8 cells (Table ES3, ES6 and Table 1).
9
10

11
12 Upon assessing the fold enrichment based on ATPase activity (commonly used as a
13
14 marker for membranes) in microvessels and homogenates, the average enrichment was
15
16 approximately 7.9-fold (Table ES7), suggesting efficient extraction of microvascular proteins
17
18 from brain tissue homogenates.
19

20 21 **Untargeted quantification of BBB transporters**

22
23 To our knowledge, this is the first global proteomic study aimed at analysing human
24
25 SLCs and ABC transporters in order to obtain a global view of the proteome dynamics, and a
26
27 comparison between healthy and disease datasets. Following this strategy, approximately
28
29 5,000 proteins were identified from an average of 23,043 peptide matches per nanoLC-
30
31 MS/MS run. The initial output from Progenesis included many duplicated entries because
32
33 several databases were used. A ‘razor’ was applied within Excel until each protein remaining
34
35 contained at least one peptide unique to the data sheet. A ‘razor’ in proteomic terms assigns
36
37 shared peptides to the most likely proteins in order to avoid or minimize assumptions in
38
39 identification.²⁷ After applying the razor, the average number of proteins was found to be
40
41 4,697, which was refined by exclusion of proteins with no unique peptide sequences against
42
43 the proteome, truncated proteins and cDNA sequences, to 3,390 proteins. This was to ensure
44
45 that only functional proteins with confirmed identification were used in the analysis. The
46
47 global analysis demonstrated no significant difference in the quality of profiles of healthy and
48
49 disease tissue. The PIPr (percentage identical proteins) and PIP (percentage identical
50
51 peptides) between the samples were calculated and were found to range within 51-78% and
52
53 32-60%, respectively. In this global study, the Hi3/Hi2 peptide ion intensities (calculated as a
54
55
56
57
58
59
60

1
2 mean) were used for quantification of endogenous and drug transporters. The criteria used for
3
4 protein inclusion in the study included selection of only peptides identified as unique with
5
6 high scores (>15), with proteins identified with only one unique peptide being considered as
7
8 supporting evidence for transporter expression. Several transporters were found in fewer than
9
10 10 samples (Table ES2), these transporters were only considered for identification and
11
12 quantification if their identified peptides met the applied criteria. In Table ES3, proteins were
13
14 quantified based on the plasma membrane marker, ATP1A1, as a reference for normalization
15
16 of data generated using the Mascot search engine. A second reference protein, ABCB1
17
18 (MDR1), was then used to confirm the results. The global analysis was able to generate
19
20 abundances similar to MRM targeted quantification (Tables ES3 and ES6) in this study
21
22 except for a few low abundance transporters. Others have observed the same trend.^{17,28}
23
24
25
26

27 Agreement between methods used in this study is confirmed by Figure 1B and C.
28
29

30 31 **Novel drug and endogenous transporters identified and quantified using the global** 32 33 **approach** 34

35
36 Out of the 77 proteins (14 ABC, 52 SLC and 11 cell markers, i.e. receptors and tight
37
38 junction proteins) (Table ES3) identified and quantified using the global approach, 37
39
40 proteins (ABCA2, ABCA8, ABCA9, ABCB1, ABCG2, ABCC3, ABCC4, ABCC9,
41
42 SLC1A2, SLC1A3, SLC1A4, SLC1A5, SLC3A2, SLC6A6, SLC6A12, SLC6A13, SLC7A1,
43
44 SLC7A5, SLC16A1, SLC19A1, SLC27A4, SLC47A1, SLC51A, SLC51B, SLCO1A2,
45
46 SLCO2B1, SLCO3A1, SLC2A1, PECAM1, VWF, GFAP, GSPG4, SYP, LRP1, TFR1,
47
48 INSR and CLDN5) were identified and quantified in earlier studies based on mRNA
49
50 expression profiling and MRM analysis in human brain microvessels.^{17,28} We have identified
51
52 an additional 39 human ABC/SLC transporters. 20 of these (ABCB9, ABCE1, ABCF1,
53
54 ABCF2, SLC1A6, SLC1A7, SLC2A13, SLC4A10, SLC8A3, SLC9A2, SLC9A6, SLC12A4,
55
56 SLC12A6, SLC14A1, SLC16A2, SLC27A1, SLC30A1, SLC33A1, SLC38A5, SLC41A3)
57
58
59
60 16

1
2 were identified only, and 19 (ABCD3, ABCF3, SLC4A1, SLC4A4, SLC6A1, SLC6A7,
3
4 SLC6A11, SLC8A1, SLC8A2, SLC9A3R1, SLC9A3R2, SLC12A2, SLC12A7, SLC17A7,
5
6 SLC30A3, SLC38A3, SLC39A10, SLC44A1, SLC44A2) were also quantified for the first
7
8 time in this study by label-free analysis in human brain microvessels. Table ES3 lists novel
9
10 and previously identified and quantified SLC/ABC transporters, cell markers, receptors and
11
12 tight junction proteins, with relevant references to other complementary data from targeted
13
14 studies.
15
16
17

18 **Accurate Mass and Retention Time (AMRT) methodology**

19
20
21 The AMRT-based targeted method ²⁵ for protein quantification was used to measure
22
23 five transporters (ABCB1, ABCG2, SLC22A3, SLC51A, SLCO1A2) and one plasma protein
24
25 marker ATP1A1 (Table S2) using one peptide each for the transporters and two peptides for
26
27 ATP1A1. Importantly, we found that the average abundance values of five transporters and
28
29 plasma membrane marker levels showed a strong and statistically significant correlation
30
31 (Figure S2A) between AMRT and MRM approaches with $R^2 \geq 0.99$. Supplementary Table
32
33 ES4 shows AMRT data for individual donors.
34
35
36

37 Technical reproducibility was evaluated by measuring the abundances of ABCB1,
38
39 ABCG2, SLC22A3, SLC51A, SLCO1A2 and ATP1A1 in three samples (one dementia and
40
41 two healthy individuals) using the AMRT approach with QconCAT peptides used as
42
43 standard. The technical variability represented as a relative error was $\%RE < 30\%$, with fold
44
45 errors of less than 1.33 for the transporters abundances between all the 3 samples analysed
46
47 (Figure S2B, Supplementary Table ES5); this suggests that there is a very low variability
48
49 between the two technical replicates.
50
51
52

53 **Gene ontology and functional enrichment of proteins in human brain microvessels**

54
55 The identified proteins (3,390 in total) were annotated for sub-cellular location and
56
57 protein class by analysing gene names against GO, UniProt, HPA and PANTHER databases
58
59
60

1
2
3 (Figure 2A, B). Based on these results, 19% of proteins were assigned to the plasma
4 membrane; 36% and 22% were identified as belonging to cytoplasm and nucleus,
5 respectively. Proteins from endoplasmic reticulum, Golgi body and mitochondria represented
6 11%, 3% and 10% of identified proteins, respectively. It is worth noting that several proteins
7 can be found in multiple subcellular locations. Figure 2B shows assignment of proteins based
8 on protein class, in which the majority of proteins were classified as nucleic acid binding
9 proteins (321 proteins), hydrolases (284 proteins), enzyme modulators (282 proteins), and
10 cytoskeletal proteins (191 proteins). Importantly, 131 proteins were identified as transporters,
11 out of which ABC transporters and SLCs represented 13% and 46%, respectively (Figure
12 2C).

23 24 25 **Microvessel protein content and its variability**

26
27
28 The values of BMvPGB in the 22 samples formed a normal distribution ($p = 0.05$),
29 with minimum and maximum values of 0.19 and 0.76 mg microvessel protein per gram brain
30 tissue, respectively, representing a 4-fold variation. The mean BMvPGB content \pm SD was
31 0.44 ± 0.15 mg/g brain (Table ES7). Variation in BMvPGB content was assessed by brain
32 donor gender and age and analyzed by unpaired t -test and Spearman correlation, respectively.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

51 52 53 **Membrane expression levels of transporters and markers in isolated human brain** 54 **microvessels**

55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2 membrane transporter proteins and 6 cell markers in isolated microvessels fractions. The
3
4 quantification was performed by means of multiplexed MRM/SRM with the same specific
5
6 peptides from the Liver TransCAT used in our previous report ¹⁸ and with new peptides from
7
8 the Brain TransCAT as shown in Table ES1. One membrane protein (ABCC1) was below the
9
10 lower limit of quantification (Table ES6) estimated at 0.01 pmol/mg protein and determined
11
12 as the protein amount that would produce less than 5000 count of peak area in the
13
14 chromatogram when sample protein is measured by LC-MS/MS.
15
16
17

18
19 For MRM analysis, technical replication was based on three analyses with Liver
20
21 TransCAT, Brain TransCAT or both TransCATs used as standards. These preparations were
22
23 analyzed only once each. Several quality controls were applied at the data analysis stage to
24
25 ensure that only reliably quantified transporters are included. For transporters quantified in
26
27 duplicate, the technical variability measured as a relative error (%RE) was shown to be
28
29 within 40%, except OATP1B3 which was within 42%. Three transporters (SLC7A5,
30
31 SLC22A1 and SLCO2B1) were quantified using two peptides, and the results were within 1.2
32
33 fold (Supplementary Table ES6).
34
35

36
37 ABCB1, ABCG2 and the ABCC family are considered to be the most important
38
39 efflux transporters implicated in drug resistance.²⁹ ABCB1/MDR1 and ABCG2/BCRP were
40
41 the most abundant ABC transporters (Figure 3B). ABCC6/MRP6 was also present, but at a
42
43 much lower concentration than ABCB1/MDR1 (5.4-fold) and ABCG2/BCRP (4.6-fold). For
44
45 SLCs, endothelial cell marker (SLC2A1/GLUT1) showed the most abundant expression
46
47 indicating the enrichment of endothelial cells (Table 1, Figure 3A), while organic anion
48
49 transporter 2 (SLC22A7/OAT2) and monocarboxylate transporter 1 (SLC16A1/MCT1)
50
51 showed high variability between donors and were expressed in the range of 2.50-16.20 and
52
53 1.05-14.10 pmol/mg protein (Figure 3B). The quantities of the other SLC22 members
54
55 (SLC22A1/OCT1, SLC22A3/OCT3, SLC22A6/OAT1, SLC22A8/OAT3 and
56
57
58
59
60

1
2 SLC22A9/UST3) were in the range of 0.23-0.80 pmol/mg protein. SLC22A4/OCTN1 was
3
4 expressed at 0.04 pmol/mg protein and this transporter was at the limit of quantification. The
5
6 expression of OATPs transporters in brain samples was in the order of OATP1C1 >
7
8 OATP2B1 > OATP1B3 > OATP1A2. Inter-individual variability was assessed based on fold
9
10 change in transporter abundance. Transporter protein expression showed considerable
11
12 biological variability with a CV in a range of 19%-100% except for ABCC6/MRP6,
13
14 SLC22A3/OCT3, SLC22A8/OAT3, SLCO2B1/OATP2B1 and SLCO1C1/OATP1C1, which
15
16 were less variable with CV values of 11.5%-14.8%. The protein expression levels of specific
17
18 cell markers such as PECAM1 (endothelial cells), NG2 (pericytes), GFAP (astrocytes), and
19
20 SYP (neurons) and the plasma membrane marker Na⁺/K⁺-ATPase were measured in the
21
22 ranges 0.30-1.67, 0.08-0.24, 2.88-65.3, 0.40-6.56 and 3.77-78.7 pmol/mg total protein,
23
24 respectively. The GFAP and Na⁺/K⁺-ATPase cell markers exhibited the greatest inter-
25
26 individual differences in protein expression, with a 22.7 and 20.9-fold difference for GFAP
27
28 and Na⁺/K⁺-ATPase between the donors with the highest and lowest levels, respectively.
29
30 Detailed MRM quantification data are provided in Table ES6.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Protein expression levels of plasma membrane marker, SLCs, ABC transporters, and cell markers for endothelial cells, pericytes, neurons, and astrocytes (pmol/mg total protein) in Healthy and disease brain microvessel samples. Data are expressed as mean \pm SD; ULQ, under limit of quantification; NQ, not quantified; n, the number of brain samples with measurements above the limit of quantification. * represents abundance from two replicates

Absolute protein expression level (means \pm SD, pmol/mg total protein)									
Alias/Gene name	Surrogate peptide used	The present study						Uchida <i>et al.</i> , 2011 (n=7)	Shawhana <i>et al.</i> , 2011 (n=5)
		Healthy	n	DLB	n	AD	n	Healthy	Healthy
Transporters									
<i>ABC transporters</i>									
ABC2/ABCA2	LGGNPLFR	0.08 \pm 0.03	12	0.08 \pm 0.03	5	0.09 \pm 0.03	5	2.86 \pm 0.58	2.11 \pm 0.78
MRP1/ABCC1	TPSGNLVNR	<0.05 \pm 0.05 (ULQ)	9	<0.04 \pm 0.01 (ULQ)	3	<0.06 \pm 0.03 (ULQ)	5	<0.211(ULQ)	<0.21 (ULQ)
MRP5/ABCC5	FFDTPPTGR	<0.01 \pm 0.002 (ULQ)	12	<0.02 \pm 0.002 (ULQ)	5	<0.01 \pm 0.002 (ULQ)	5	<0.497 (ULQ)	<0.5 (ULQ)
MRP6/ABCC6	SSLPSALLGELSK	0.48 \pm 0.06*	12	0.48 \pm 0.04*	5	0.47 \pm 0.07*	5	<0.174 (ULQ)	<0.17 (ULQ)
MDR1/ABCB1	AGAVAEVLAAIR	2.58 \pm 0.93	12	3.18 \pm 0.91	5	2.25 \pm 1.48	5	6.06 \pm 1.69	3.98 \pm 0.88
BCRP/ABCG2	VIQELGLDK	2.22 \pm 0.61*	12	2.31 \pm 0.63*	5	1.91 \pm 0.84*	5	8.14 \pm 2.26	6.15 \pm 1.41
<i>SLC transporters</i>									
OCT1/SLC22A1	MLSLEEDVTEK	0.58 \pm 0.11*	12	0.55 \pm 0.10*	5	0.44 \pm 0.09*	5	<0.288 (ULQ)	<0.29 (ULQ)
OCT1/SLC22A1	GVALPETMK	0.54 \pm 0.06*	12	0.56 \pm 0.07*	5	0.45 \pm 0.17*	5		
OCT3/SLC22A3	GIALPETVDDVEK	0.62 \pm 0.08*	12	0.66 \pm 0.08*	5	0.59 \pm 0.11*	5	<0.207 (ULQ)	<0.21 (ULQ)
OCTN1/SLC22A4	AFILDIFR	<0.04 \pm 0.01 (ULQ)	12	<0.02 \pm 0.02 (ULQ)	5	<0.02 \pm 0.02 (ULQ)	5	<0.123 (ULQ)	<0.12 (ULQ)
OAT1/SLC22A6	TSLAVLGGK	0.48 \pm 0.11	12	0.48 \pm 0.07	5	0.64 \pm 0.07	5	<0.909 (ULQ)	<0.91 (ULQ)
OAT2/SLC22A7	WLLTQGHVK	7.90 \pm 3.80	8	9.92 \pm 3.70	5	6.95 \pm 2.53	4	<0.153 (ULQ)	<0.15 (ULQ)
OAT3/SLC22A8	LSLEELK	0.27 \pm 0.03	12	0.26 \pm 0.02	5	0.24 \pm 0.01	5	<0.348 (ULQ)	<0.35 (ULQ)
OAT7/SLC22A9	DTLTLEILK	0.51 \pm 0.10*	12	0.48 \pm 0.03*	5	0.46 \pm 0.11*	5	<0.326 (ULQ)	<0.33 (ULQ)
ENT1/SLC29A1	WLPSVLAR	0.27 \pm 0.10	12	0.22 \pm 0.09	5	0.24 \pm 0.04	5	0.568 \pm 0.134	0.86 \pm 0.13
MCT1/SLC16A1	SITVFFK	5.37 \pm 3.73*	12	4.48 \pm 2.60*	5	3.14 \pm 1.30*	5	2.27 \pm 0.85	1.46 \pm 0.39
LAT1/SLC7A5	GDVSNLDPNFSFEGTK	0.59 \pm 0.15	12	0.74 \pm 0.09	5	0.56 \pm 0.14	5	0.431 \pm 0.091	0.80 \pm 0.25
LAT1/SLC7A5	SADGSAPAGEGEGVTLQR	0.71 \pm 0.25	9	0.90 \pm 0.29	5	0.55 \pm 0.01	3		
OATP1/SLCO1A2	EGLETNADIHK	0.54 \pm 0.10*	12	0.50 \pm 0.04*	5	0.47 \pm 0.11*	5	<0.695 (ULQ)	<0.69 (ULQ)
OST- α /SLC51A	YTADLLEVLK	0.45 \pm 0.13	12	0.50 \pm 0.09	5	0.43 \pm 0.14	5	<0.292 (ULQ)	<0.29 (ULQ)
OATP8/SLCO1B3	NVTGFFQSLK	0.46 \pm 0.15	12	0.58 \pm 0.13	5	0.52 \pm 0.20	5	<0.572 (ULQ)	<0.57 (ULQ)
OATP2B1/SLCO2B1	IGPAGEVPQVPDK	0.40 \pm 0.04*	12	0.36 \pm 0.03	5	0.35 \pm 0.02	5	<0.337 (ULQ)	<0.34 (NQ)
OATP2B1/SLCO2B1	SSPAVEQQLLVSGPGK	0.48 \pm 0.11*	12	0.36 \pm 0.03	5	0.38 \pm 0.02	5		
OATP1C1/SLCO1C1	DFLPSLK	0.27 \pm 0.03	12	0.28 \pm 0.02	5	0.26 \pm 0.04	5	<0.208 (ULQ)	<0.21 (ULQ)
Markers									
<i>Endothelial Markers</i>									
GLUT1/SLC2A1	TFDEIASGFR	21.9 \pm 9.80	11	36.2 \pm 14.9	5	17.6 \pm 13.4	5	139 \pm 46	78.5 \pm 23
PECAM1	IYDSGTYK	0.89 \pm 0.46	12	0.82 \pm 0.15	5	0.52 \pm 0.13	5	NQ	1.57 \pm 0.21
<i>Plasma membrane marker</i>									
Na ⁺ /K ⁺ ATPase	IVEIPFNSTNK	9.02 \pm 5.64*	12	8.06 \pm 3.27*	5	10.5 \pm 9.34*	5	35.1 \pm 12.6	20.8 \pm 6.5
	NPNTSEPPHLLVMK	17.5 \pm 16.2	12	16.8 \pm 5.92	5	22.1 \pm 28.4	5		
<i>Astrocyte marker</i>									
GFAP	FADLTDAAR	19.2 \pm 18.4	11	11.8 \pm 8.45	5	9.12 \pm 6.66	4	NQ	503.2 \pm 174.09
<i>Pericyte marker</i>									
NG2	APLEVQALGR	0.15 \pm 0.06	12	0.12 \pm 0.06	5	0.12 \pm 0.03	5	NQ	1.07 \pm 0.41
	LVLGQEELR	0.11 \pm 0.05	12	0.18 \pm 0.05	5	0.14 \pm 0.06	5		
<i>Neuron marker</i>									
SYP	ETGWAAPFLR	3.01 \pm 2.45	12	1.95 \pm 1.20	5	3.11 \pm 2.04	4	NQ	1.45 \pm 0.37

1
2
3 **Comparison of membrane protein expression levels in isolated human brain microvessels**
4
5 **with previous studies**
6

7
8 Table 1 shows a comparison of abundance means of 22 transporters and 6 markers in the
9
10 present 22 samples with those in a targeted analysis of 12 individual samples as previously
11
12 reported by Terasaki and co-workers.^{17,28} In the MRM analysis, one additional membrane
13
14 protein (CDH17) was quantified for the first time in this study. Only 5 transporters and 6
15
16 markers were successfully quantified by both groups, and the mean abundances were in the same
17
18 range with typically a factor of 2-4 difference. Working in 2011, Terasaki and co-workers^{17,28}
19
20 found many of the transporter peptides to be below the limit of quantification. In general, they
21
22 reported higher levels of cell markers than in the present study, reflecting different sample
23
24 preparations. At this stage, it is not possible to determine whether the differences in the two
25
26 studies were due to biological factors, tissue handling and storage, or sample preparation.
27
28
29

30
31 Using different proteotypic peptides as well as different sample preparation methods can
32
33 affect protein quantification, and this can explain some of the differences in reported values
34
35 (Table ES3).
36

37 **ABC and SLC transporter–transporter correlation of blood-brain barrier protein**
38
39 **expression**
40

41
42 Transporter-transporter correlation analysis (intercorrelation analysis) was performed on
43
44 abundance data generated by targeted proteomics. In particular, the transporter pairs
45
46 ABCB1/ABCG2, SLC22A9/SLCO1A2, SLC22A9/SLC51A, SLCO1A2/SLC51A and
47
48 SLC2A1/SLC7A5 showed very strong correlation ($R_s = 0.81-0.94$, $p < 0.0001$, $R^2 = 0.58-0.93$).
49
50 ABCB1 and ABCG2 showed a significant and strong correlation with endothelial cell marker
51
52 SLC2A1 ($R_s = 0.80$, $p < 0.0001$, $R^2 = 0.55$; $R_s = 0.71$, $p = 0.0003$, $R^2 = 0.50$, respectively). Amino
53
54
55
56
57
58
59
60

1
2
3 acid transporter SLC7A5 expression was significantly correlated with several markers; such
4 correlations included PECAM1 ($R_s = 0.64$, $p = 0.001$, $R^2 = 0.31$). We also tested a potential
5 association between ATP1A1 expression and transporters. We found expression levels of
6 ATP1A1 to be associated with ABCC6, SLC22A1, SLC22A9, SLCO1A2 and SLC51A
7 expression. The matrix in Figure 4 shows the most significant correlations between SLCs, ABC
8 transporters and cell markers quantified.

16 **Covariates of proteome expression at the blood-brain barrier**

17
18
19 Trends of similarity of proteome expression profiles at the BBB between samples were
20 assessed using percentage identical peptides (PIP) and percentage identical proteins (PIPr) data
21 for the 22 samples. Principal coordinates analysis (PCoA) and hierarchical cluster analysis
22 (HCA) were performed both at the protein and peptide levels and the results were very similar at
23 the two levels (Figure 5) with samples from relatively young healthy donors clustering mainly in
24 the middle with samples from patients with DLB. (Note that the blue ellipse refers to DLB
25 donors only – the cluster is assessed visually, and can be deemed to include patient 3.) Samples
26 from AD patients tended to scatter around the cluster (only patient 12 is within the cluster). Four
27 normal donors are also outside the cluster; three of these were very elderly (87, 89 and 90) at the
28 time of death, but patient 2 was only 54. Note, however, that there is no clustering of these
29 outlier individuals; they are as different from one another as from the main cluster. This suggests
30 that the main covariates of protein expression at the BBB in this dataset were the disease state
31 and age, with some healthy donors of very old age behaving like AD patients.

48 **Covariates of transporter protein expression at the blood-brain barrier**

49
50
51 Transporter expression profiles at the BBB were assessed using principal components
52 analysis (PCA) and hierarchical cluster analysis (HCA) applied to a heatmap of abundance data
53
54
55
56
57
58
59
60

1
2
3 (Figure 6). The PCA analysis data were confirmatory of global proteome profiles; older healthy
4 individuals (donors 2, 9, 14, 17, 18 and 22) with average age 82.3 ± 14.1 years, behaving like
5
6 AD patients, whereas younger healthy donors tended to cluster in the middle (75.2 ± 15.0 years).
7
8 With the exception of sample 12, none of the AD samples clustered with the healthy ones, which
9
10 is a similar trend to the global proteomic analysis. HCA confirmed the cluster and outliers in the
11
12 PCA and the heatmap showed correlations of expression between drug transporters (MDR1 and
13
14 BCRP) and SLCs of endogenous compounds, including glucose and organic cations/anions
15
16 (SLC2A1/SLC7A5 and SLCO1A2/SLC22A9) (Figure 6C). PCA analyses using data from
17
18 different replicates and different peptides (where available) were confirmatory. This reflects that
19
20 disease, age and inter-correlation of gene expression represent significant factors that can affect
21
22 the transporter make-up of the BBB.
23
24
25
26
27

28 **Inter-species differences between rat and human BBB transporter/receptor expression** 29 **levels**

30
31 In-house data for human (Table 1 and Table ES3) and rat²⁰ were used to determine comparative
32
33 levels of transporters and receptors between rat and human brain capillaries. Figure 7A shows
34
35 similarities and differences in expression levels, where several transporters (NKCC1, AE1,
36
37 MCT1, BCRP, EAAT2), receptors (LRP1, TfR1), tight junction and marker proteins (CLDN5,
38
39 ATP1A1) were within 2-fold. Higher expression levels in rat microvessels were observed for
40
41 MDR1, EAAT1, LAT1, CAT1, PMP70 and the glucose transporter GLUT1. This is in overall
42
43 agreement with literature on comparative data between rats and humans³⁰ as shown in Figure 7B.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

Precision dosing,³¹ in which the amount and frequency of drug dosing is tailored to individual patients, can be critical to the outcome of therapy. Precision dosing depends upon understanding the efficacy and toxicity of any drug targeting the individual brain, which is dependent on the concentration at the site of action, and in turn, on the abundance and activity of drug transporters at the BBB.³² We describe the first systematic study, using a quantitative strategy combining both targeted and global methodology, of the proteomic make-up of the BBB, focussing on drug and endogenous transporters. The 22 brains analysed (15 males, 7 females) had an average age of 78 years (range: 53-91 years), and were classified into three different sub-groups: healthy brains (12 samples), brains from Alzheimer's patients (5 samples) and brains from patients with dementia with Lewy bodies (5 samples).

To isolate the microvessels of the BBB, we used an optimized methodology previously performed and validated in a rat brain model²⁰ with minor modifications to sample preparation from frozen brain cortical samples. Differences may be expected when different types of tissue samples (i.e. fresh compared to frozen), storage conditions or sample preparation methods are used.³³ The average enrichment of microvessels was very high (>7.9 fold), and the purity of microvessels was assessed using brain cell-specific markers. Astrocyte endfeet tend to be tightly bound to the neurovascular unit, making it very difficult to remove them entirely. The neuron marker SYP and pericyte marker NG2 were also detected but consistently at very low levels.

The quality of the sample preparation was key to our being able to detect and quantify 19 previously undetected transporters and to detect (but not quantify) a further 20. In total, 22 transporters were quantified using a targeted strategy and 31 using the global, untargeted strategy showing the value of using both methods. 16 of the proteins detected by the targeted approach fell below the limit of quantification of the global proteomic experiment, but 32 additional

1
2
3 transporters were quantifiable by untargeted methods. For the more abundant transporters, the
4 targeted and global methods returned very consistent values (Figure 1C). In the one case where
5
6 targeted and global methods returned very consistent values (Figure 1C). In the one case where
7
8 the difference between targeted and global analysis was considerable, SLC7A5 (which is
9
10 involved in high-grade gliomas), it seems probable that signal overlap gave rise to an
11
12 overestimate in the label-free experiment,³⁴ consistent with previous reports.^{17,28} Disease-related
13
14 transporters quantified for the first time include choline transporter-like proteins 1 and 2,
15
16 SLC44A1 and SLC44A2, reported to be involved in abnormal choline disposition associated
17
18 with papillary glioneuronal tumors.³⁵
19
20

21
22 In drug development, several scaling factors are used for the extrapolation of *in vitro* data
23
24 to *in vivo* parameters to predict drug disposition across the BBB¹⁶, an approach which is starting
25
26 to be accepted for supporting evidence in clinical trials by medicines regulatory agencies.³⁶ One
27
28 important scaling factor is the protein content of brain microvessels per gram tissue
29
30 (BMvPGB).^{16, 17} The corresponding liver and kidney scaling factors are well-established^{37,38} but
31
32 to date, modeling of drug disposition across the human BBB relies on a single study of six
33
34 histologically healthy brains for this parameter.^{16,17} Our data for healthy brains shows limited
35
36 overall variability (4-fold) with an average of 0.44 mg protein/g brain. The previously reported
37
38 values¹⁷ show BMvPGB (0.244 mg protein/g brain) and 7-fold variability. The present study
39
40 therefore trebles healthy brain data that can contribute to BBB models, allowing better
41
42 description of population variability. The mean BMvPGB levels of AD and DLB donors (0.40
43
44 and 0.48 mg/g brain, respectively) were very similar to the levels of healthy donors. BMvPGB
45
46 levels showed no correlation with age, sex or disease state.
47
48
49
50

51
52 Transporter expression data revealed a large degree of inter-individual variability across
53
54 ABC (3- to 11-fold) and SLC (2- to 14-fold) transporters. The analyzed set of samples was from
55
56
57
58
59
60

1
2
3 a small number of donors, which should be considered when using the generated data. Extending
4
5 this study to cover a large number of samples from both health and disease populations should
6
7 confirm these conclusions.
8
9

10 There were no significant univariate sex or age-related differences in the abundances of
11
12 transporters, in line with recent data on liver transporters.³⁹ The expression of four transporters,
13
14 however, correlates significantly with disease (Figure 3). The most significant correlation
15
16 ($p < 0.01$) is the upregulation of SLC22A6 in Alzheimer's disease. This is a drug transporter
17
18 responsible for influx of many drugs, including anti-virals and anti-inflammatory drugs to the
19
20 brain. Multivariate analysis revealed general differences between patient sub-groups, with a set
21
22 of donors, including all the DLB patients and most of the younger healthy individuals forming a
23
24 defined cluster in the principal components analysis. Alzheimer's patients and several of the
25
26 older healthy donors were outside this cluster. This suggests that age in healthy individuals and
27
28 Alzheimer's disease in the disease population are the main contributors to differences in
29
30 expression. This trend was identified in the global proteomic profiling of the blood-brain barrier
31
32 at both the peptide and protein levels (Figure 5), as well as in the more specific analysis of
33
34 transporter levels in brain endothelial cells (Figure 6). The expression trends shown for the first
35
36 time in this study support recent guidelines advocating the definition of Alzheimer's disease by
37
38 its biochemical fingerprint, compared to predominantly anatomically or clinically defined types
39
40 of dementia, such as dementia with Lewy bodies.⁴⁰
41
42
43
44
45
46

47 We also examined inter-species differences in expression between human and rat ^{30,20}
48
49 (Figure 7). The rat is an important pre-clinical species and transport data based on rat need to be
50
51 scalable to human. While some transporter abundances (e.g. BCRP) are clearly useful for
52
53
54
55
56
57
58
59
60

1
2
3 translation (rat to human), others (such as MDR1) show large discrepancies throwing doubt on
4
5 the suitability of the rat as a pre-clinical species where this transporter is critical.
6

7
8 The importance of considering relationships between the expression patterns of proteins
9
10 involved in drug metabolism and disposition has been highlighted in clinical and simulated
11
12 trials.^{37,41} Inter-correlations between individual transporters in the brain were assessed in this
13
14 dataset for the first time (Figure 4), revealing associations that can be attributed to common
15
16 genetic regulatory mechanisms, supported by transcriptomic evidence.⁴² The present dataset
17
18 uncovered several correlations, not only between drug transporters but also between transporters
19
20 involved in brain homeostasis (e.g. glucose and amino acid carriers). One important example of
21
22 expression co-regulation is the correlation between ABCB1 and ABCG2, and patients with
23
24 Alzheimer's disease showed relatively lower expression levels of both transporters, although
25
26 neither reached statistical significance in the univariate analysis (Figure 3).
27
28
29

30 31 **CONCLUSION**

32
33 This study described the protein composition of the blood-brain barrier and will be used
34
35 to inform Quantitative System Pharmacology models of health and disease.
36
37
38
39

40 41 **ASSOCIATED CONTENT**

42 43 **Supplementary Information**

44
45 **Table S1:** Demographic and clinical information of brain donors
46
47

48
49 **Table S2:** Absolute protein expression levels in microvessels isolated from 22 brains
50

51
52 **Table S3:** Summary matrix of correlation analysis of ABC, SLC transporters and cell markers
53
54 expression data
55

56
57 **Table ES1:** Peptide sequences included in brain TransCAT
58
59
60

1
2
3 **Table ES2:** Summary of peptide and protein analysis by Progenesis after application of a Razor

4
5 **Table ES3:** Protein quantification using label free in human brain microvessels

6
7
8 **Table ES4:** Summary of transporters and membrane marker quantified using accurate mass
9
10 retention time (AMRT) approach

11
12 **Table ES5:** Technical variability between 3 brains for ABCB1, ABCG2, SLC22A3 transporters
13
14 and ATP1A1 membrane marker using AMRT

15
16
17 **Table ES6:** Protein abundances in individual human brain microvessels determined using MRM
18
19 approach

20
21 **Table ES7:** Protein activity and BMvPGB protein content estimate data in individual human
22
23 brain donors

24
25
26 **Figure S1:** Assessment of correlation between ATPase activity and protein concentration in
27
28 microvessels and homogenates from human brain tissue

29
30
31 **Figure S2:** Quality control assessment of accurate mass and retention time (AMRT)
32
33 methodology with reference to multi-reaction monitoring (MRM)

34
35
36 **Figure S3:** Percentage identical peptides (PIP) and percentage identical proteins (PIPr) between
37
38 the proteomes of the 22 brains

39 40 **AUTHOR INFORMATION**

41 42 43 **Corresponding Authors**

44
45
46 *Email: Zubida.al-majdoub@manchester.ac.uk

47
48 *Email: Jill.barber@manchester.ac.uk

49 **ORCID**

50
51 **Zubida M. Al-Majdoub:** 0000-0002-1497-3140

52 **Jill Barber:** 0000-0002-5424-0291

CONFLICT OF INTEREST

Sibylle Neuhoff and Amin Rostami-Hodjegan are employee of Certara UK Limited (Simcyp Division).

AUTHOR CONTRIBUTIONS

This study was designed by Zubida M. Al-Majdoub, Hajar Al Feteisi, Brahim Achour, Amin Rostami-Hodjegan and Jill Barber. Zubida M. Al-Majdoub, Hajar Al Feteisi and Stacey Warwood performed the experiments. Data analysis was performed by Zubida M. Al-Majdoub, Brahim Achour and Jill Barber. Zubida M. Al-Majdoub contributed new reagents. Zubida M. Al-Majdoub, Brahim Achour, Sibylle Neuhoff, Amin Rostami-Hodjegan and Jill Barber contributed to the writing of the manuscript.

FUNDING SOURECES

This study was supported by the Simecyp (Certara) Grant and Partnership Scheme (GPS 2014/15).

ACKNOWLEDGMENTS

Tissue samples were supplied by The Manchester Brain Bank, which is part of the Brains for Dementia Research programme, jointly funded by Alzheimer's Research UK and Alzheimer's Society. The authors thank the Biological Mass Spectrometry Core Facility (Bio-MS), University of Manchester, for access to LC-MS/MS instrumentation and analysis software.

ABBREVIATIONS

ABC, ATP-binding cassette; BBB, blood-brain barrier; AD, alzheimer's disease; DLB, dementia with Lewy bodies; DCA, bicinchoninic acid; BSA, bovine serum albumin; CNS, central nervous system; FASP, filter-aided sample preparation; IVIVE, *in vitro-in vivo* extrapolation; LC-MS/MS, liquid chromatography-tandem mass spectrometry; NNOP, non-naturally occurring peptide; PBPK, physiologically-based pharmacokinetics, QconCAT, quantification concatemer; SLC, solute carrier; TransCAT, transporters QconCAT.

References

- (1) Abbott, N. J.; Romero, I. A. Transporting Therapeutics across the Blood-Brain Barrier. *Molecular Medicine Today*. 1996.
- (2) Oby, E.; Janigro, D. The Blood-Brain Barrier and Epilepsy. *Epilepsia*. November 2006, pp 1761–1774.
- (3) Abbott, N. J. Dynamics of CNS Barriers: Evolution, Differentiation, and Modulation. *Cellular and Molecular Neurobiology* **2005**, *25* (1), 5–23.
- (4) Ballabh, P.; Braun, A.; Nedergaard, M. The Blood-Brain Barrier: An Overview: Structure, Regulation, and Clinical Implications. *Neurobiology of Disease* **2004**, *16* (1), 1–13.
- (5) Ito, K.; Suzuki, H.; Horie, T.; Sugiyama, Y. Apical/Basolateral Surface Expression of Drug Transporters and Its Role in Vectorial Drug Transport. *Pharmaceutical Research* **2005**, *22* (10), 1559–1577.
- (6) Ohtsuki, S.; Terasaki, T. Contribution of Carrier-Mediated Transport Systems to the Blood-Brain Barrier as a Supporting and Protecting Interface for the Brain; Importance for CNS Drug Discovery and Development. *Pharmaceutical Research*. August 1, 2007, pp 1745–1758.
- (7) Insel, T. R. Next-Generation Treatments for Mental Disorders. *Science Translational Medicine* **2012**, *4* (155), 1–10.
- (8) Ringel, M.; Tollman, P.; Hersch, G.; Schulze, U. Does Size Matter in R&D Productivity? If Not, What Does? *Nature Reviews Drug Discovery*. December 18, 2013, pp 901–902.
- (9) Blennow, K.; de Leon, M. J.; Zetterberg, H. Alzheimer's Disease. *The Lancet* **2006**, *368*

- (9533), 387–403.
- (10) Tomlinson, B. E.; Blessed, G.; Roth, M. Observations on the Brains of Non Demented Old People. *Journal of Neurological Sciences* **1968**, *7*, 331–356.
- (11) Jensen, M.; Schroder, J.; Blomberg, M.; Engvall, B.; Pantel, J.; Ida, N.; Basun, H.; Wahlund, L.-O.; Werle, E.; Jauss, M.; et al. Cerebrospinal Fluid Abeta42 Is Increased Early in Sporadic Alzheimer’s Disease and Declines with Disease Progression. *Annals of Neurology* **1999**, *45* (4), 504–511.
- (12) Montagne, A.; Zhao, Z.; Zlokovic, B. V. Alzheimer’s Disease : A Matter of Blood – Brain Barrier Dysfunction ? The Journal of Experimental Medicine. *Journal of Experimental Medicine* **2017**, *214* (11), 3151–3169.
- (13) Korolainen, M. A.; Goldsteins, G.; Alafuzoff, I.; Koistinaho, J.; Pirttilä, T. Proteomic Analysis of Protein Oxidation in Alzheimer’s Disease Brain. *Electrophoresis* **2002**, *23* (19), 3428–3433.
- (14) Korolainen, M. A.; Goldsteins, G.; Nyman, T. A.; Alafuzoff, I.; Koistinaho, J.; Pirttilä, T. Oxidative Modification of Proteins in the Frontal Cortex of Alzheimer’s Disease Brain. *Neurobiology of Aging* **2006**, *27* (1), 42–53.
- (15) Helms, H. C.; Abbott, N. J.; Burek, M.; Cecchelli, R.; Couraud, P.-O.; Deli, M. A.; Förster, C.; Galla, H. J.; Romero, I. A.; Shusta, E. V.; et al. In Vitro Models of the Blood–brain Barrier: An Overview of Commonly Used Brain Endothelial Cell Culture Models and Guidelines for Their Use. *Journal of Cerebral Blood Flow & Metabolism* **2016**, *36* (5), 862–890.

- 1
2
3 (16) Gaohua, L.; Neuhoff, S.; Johnson, T. N.; Rostami-Hodjegan, A.; Jamei, M. Development
4 of a Permeability-Limited Model of the Human Brain and Cerebrospinal Fluid (CSF) to
5 Integrate Known Physiological and Biological Knowledge: Estimating Time Varying CSF
6 Drug Concentrations and Their Variability Using in Vitro Data. *Drug Metabolism and*
7 *Pharmacokinetics* **2016**, *31* (3), 224–233.
8
9
10
11
12
13
14
15 (17) Uchida, Y.; Ohtsuki, S.; Katsukura, Y.; Ikeda, C.; Suzuki, T.; Kamiie, J.; Terasaki, T.
16 Quantitative Targeted Absolute Proteomics of Human Blood-Brain Barrier Transporters
17 and Receptors. *Journal of Neurochemistry* **2011**, *117* (2), 333–345.
18
19
20
21
22
23 (18) Russell, M. R.; Achour, B.; Mckenzie, E. A.; Lopez, R.; Harwood, M. D.; Rostami-
24 Hodjegan, A.; Barber, J. Alternative Fusion Protein Strategies to Express Recalcitrant
25 QconCAT Proteins for Quantitative Proteomics of Human Drug Metabolizing Enzymes
26 and Transporters. *Journal of Proteome Research* **2013**, *12* (12), 5934–5942.
27
28
29
30
31
32
33 (19) Al-Majdoub, Z. M.; Carroll, K. M.; Gaskell, S. J.; Barber, J. Quantification of the Proteins
34 of the Bacterial Ribosome Using QconCAT Technology. *Journal of Proteome Research*
35 **2014**.
36
37
38
39
40
41 (20) Al Feteisi, H.; Al-Majdoub, Z. M.; Achour, B.; Couto, N.; Rostami-Hodjegan, A.; Barber,
42 J. Identification and Quantification of Blood-Brain Barrier Transporters in Isolated Rat
43 Brain Microvessels. *Journal of Neurochemistry* **2018**, *146* (6), 670–685.
44
45
46
47
48 (21) Wis, J. R.; Zougman, A.; Nagaraj, N.; Mann, M. Universal Sample Preparation Method
49 for Proteome Analysis. **2009**.
50
51
52
53
54 (22) MacLean, B.; Tomazela, D. M.; Shulman, N.; Chambers, M.; Finney, G. L.; Frewen, B.;

- 1
2
3 Kern, R.; Tabb, D. L.; Liebler, D. C.; MacCoss, M. J. Skyline: An Open Source Document
4 Editor for Creating and Analyzing Targeted Proteomics Experiments. *Bioinformatics*
5
6 **2010**, *26* (7), 966–968.
7
8
9
10
11 (23) Achour, B.; Al Feteisi, H.; Lanucara, F.; Rostami-Hodjegan, A.; Barber, J. Global
12 Proteomic Analysis of Human Liver Microsomes: Rapid Characterization and
13 Quantification of Hepatic Drug-Metabolizing Enzymes. *Drug Metabolism and Disposition*
14
15 **2017**, *45* (6), 666–675.
16
17
18
19
20
21 (24) Silva, J. C. Absolute Quantification of Proteins by LCMSE: A Virtue of Parallel Ms
22 Acquisition. *Molecular & Cellular Proteomics* **2005**, *5* (1), 144–156.
23
24
25
26 (25) Silva, J. C.; Denny, R.; Dorschel, C. a; Gorenstein, M.; Kass, I. J.; Li, G.; Mckenna, T.;
27 Nold, M. J.; Richardson, K.; Young, P.; et al. Quantitative Proteomic Analysis by
28 Accurate Mass Retention Time Pairs. *Analytical Chemistry* **2005**, *77* (7), 2187–2200.
29
30
31
32
33
34 (26) Mi, H.; Muruganujan, A.; Thomas, P. D. PANTHER in 2013: Modeling the Evolution of
35 Gene Function, and Other Gene Attributes, in the Context of Phylogenetic Trees. *Nucleic*
36
37 *Acids Research* **2013**, *41* (D1), 377–386.
38
39
40
41
42 (27) Cox, J.; Mann, M. MaxQuant Enables High Peptide Identification Rates, Individualized
43 p.p.b.-Range Mass Accuracies and Proteome-Wide Protein Quantification. *Nature*
44
45 *Biotechnology* **2008**, *26* (12), 1367–1372.
46
47
48
49
50 (28) Shawahna, R.; Uchida, Y.; Declèves, X.; Ohtsuki, S.; Yousif, S.; Dauchy, S.; Jacob, A.;
51 Chassoux, F.; Dumas-Duport, C.; Couraud, P.-O.; et al. Transcriptomic and Quantitative
52 Proteomic Analysis of Transporters and Drug Metabolizing Enzymes in Freshly Isolated
53
54
55
56
57
58
59
60

- 1
2
3 Human Brain Microvessels. *Molecular Pharmaceutics* **2011**, 8 (4), 1332–1341.
4
5
6 (29) Giacomini, K. M.; Huang, S.-M.; Tweedie, D. J.; Benet, L. Z.; Brouwer, K. L. R. R.; Chu,
7
8 X.; Dahlin, A.; Evers, R.; Fischer, V.; Hillgren, K. M.; et al. Membrane Transporters in
9
10 Drug Development. *Nature Reviews Drug Discovery* **2010**, 9 (3), 215–236.
11
12
13
14 (30) Hoshi, Y.; Uchida, Y.; Tachikawa, M.; Inoue, T.; Ohtsuki, S.; Terasaki, T. Quantitative
15
16 Atlas of Blood-Brain Barrier Transporters, Receptors, and Tight Junction Proteins in Rats
17
18 and Common Marmoset. *Journal of Pharmaceutical Sciences* **2013**, 102 (9), 3343–3355.
19
20
21
22 (31) Polasek, T. M.; Shakib, S.; Rostami-Hodjegan, A. Precision Dosing in Clinical Medicine:
23
24 Present and Future. *Expert Review of Clinical Pharmacology*. August 3, 2018, pp 743–
25
26 746.
27
28
29
30 (32) Rostami-Hodjegan, A.; Tucker, G. T. Simulation and Prediction of in Vivo Drug
31
32 Metabolism in Human Populations from in Vitro Data. *Nature reviews. Drug discovery*
33
34 **2007**, 6 (2), 140–148.
35
36
37
38 (33) Yousif, S.; Marie-Claire, C.; Roux, F.; Scherrmann, J. M.; Declèves, X. Expression of
39
40 Drug Transporters at the Blood-Brain Barrier Using an Optimized Isolated Rat Brain
41
42 Microvessel Strategy. *Brain Research* **2007**, 1134 (1), 1–11.
43
44
45
46 (34) Achour, B.; Dantonio, A.; Niosi, M.; Novak, J. J.; Al-Majdoub, Z. M.; Goosen, T. C.;
47
48 Rostami-Hodjegan, A.; Barber, J. Data Generated by Quantitative Liquid
49
50 Chromatography–Mass Spectrometry Proteomics Are Only the Start and Not the Endpoint:
51
52 Optimization of Quantitative Concatemer-Based Measurement of Hepatic Uridine-5'-
53
54 Diphosphate–Glucuronosyltransferase Enzymes with Ref. *Drug Metabolism and*
55
56
57
58
59
60

- 1
2
3 *Disposition* **2018**, *46* (6), 805–812.
- 4
5
6 (35) Soffietti, R.; Rudà, R.; Reardon, D. Rare Glial Tumors. In *Handbook of Clinical*
7
8 *Neurology*; 2016; Vol. 134, pp 399–415.
- 9
10
11 (36) Shebley, M.; Sandhu, P.; Emami Riedmaier, A.; Jamei, M.; Narayanan, R.; Patel, A.;
12
13 Peters, S. A.; Reddy, V. P.; Zheng, M.; de Zwart, L.; et al. Physiologically Based
14
15 Pharmacokinetic Model Qualification and Reporting Procedures for Regulatory
16
17 Submissions: A Consortium Perspective. *Clinical Pharmacology and Therapeutics* **2018**,
18
19 *104* (1), 88–110.
- 20
21
22 (37) Barter, Z. E.; Perrett, H. F.; Yeo, K. R.; Allorge, D.; Lennard, M. S.; Rostami-Hodjegan,
23
24 A. Determination of a Quantitative Relationship between Hepatic CYP3A5*1/*3 and
25
26 CYP3A4 Expression for Use in the Prediction of Metabolic Clearance in Virtual
27
28 Populations. *Biopharmaceutics and Drug Disposition* **2010**, *31* (8–9), 516–532.
- 29
30
31 (38) Scotcher, D.; Billington, S.; Brown, J.; Jones, C. R.; Brown, C. D. A. A.; Rostami-
32
33 Hodjegan, A.; Galetin, A. Microsomal and Cytosolic Scaling Factors in Dog and Human
34
35 Kidney Cortex and Application for In Vitro-In Vivo Extrapolation of Renal Metabolic
36
37 Clearance. *Drug Metabolism and Disposition* **2017**, *45* (5), 556–568.
- 38
39
40 (39) Prasad, B.; Evers, R.; Gupta, A.; Hop, C. E. C. A. C. a; Salphati, L.; Shukla, S.;
41
42 Ambudkar, S. V.; Unadkat, J. D. Interindividual Variability in Hepatic Organic Anion-
43
44 Transporting Polypeptides and P-Glycoprotein (ABCB1) Protein Expression:
45
46 Quantification by Liquid Chromatography Tandem Mass Spectroscopy and Influence of
47
48 Genotype, Age, and Sex. *Drug Metabolism and Disposition* **2013**, *42* (1), 78–88.
- 49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (40) Jack, C. R.; Bennett, D. A.; Blennow, K.; Carrillo, M. C.; Dunn, B.; Haeberlein, S. B.;
4 Holtzman, D. M.; Jagust, W.; Jessen, F.; Karlawish, J.; et al. *NIA-AA Research*
5 *Framework: Toward a Biological Definition of Alzheimer's Disease*; Elsevier Inc., 2018;
6 Vol. 14, pp 535–562.
7
8
9
10
11
12
13 (41) Doki, K.; Darwich, A. S.; Achour, B.; Tornio, A.; Backman, J. T.; Rostami-Hodjegan, A.
14 Implications of Intercorrelation between Hepatic CYP3A4-CYP2C8 Enzymes for the
15 Evaluation of Drug-Drug Interactions: A Case Study with Repaglinide. *British Journal of*
16 *Clinical Pharmacology* **2018**, *84* (5), 972–986.
17
18
19
20
21
22
23 (42) Wortham, M.; Czerwinski, M.; He, L.; Parkinson, A.; Wan, Y. J. Y. Expression of
24 Constitutive Androstane Receptor, Hepatic Nuclear Factor 4 α , and P450 Oxidoreductase
25 Genes Determines Interindividual Variability in Basal Expression and Activity of a Broad
26 Scope of Xenobiotic Metabolism Genes in the Human Liver. *Drug Metabolism and*
27 *Disposition* **2007**, *35* (9), 1700–1710.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Brief experimental workflow and assessment of quantitative methods used to analyse the proteins expressed at the human blood-brain barrier. Several optimization steps (A) were introduced to isolate the microvessel fraction and extract membrane proteins, which were then proteolytically-digested and analysed using three LC-MS/MS proteomic methods: targeted multi-reaction monitoring (MRM), targeted accurate mass and retention time (AMRT) methodology, and label-free global proteomics using the two/three most abundant peptide peaks per protein. The assessment of transporter and marker proteins using the three methods showed limited differences quantitative results (B) and excellent overall correlation between data (C), with the label-free (LF) method being relatively less precise and accurate than the targeted MRM and AMRT methods. AMRT was used to measure five transporters (ABCB1, ABCG2, SLC22A3, SLC51A and SLCO1A2) and one plasma protein marker ATP1A1 in 22 brains.

261x159mm (300 x 300 DPI)




Figure 2: Global proteomic analysis of brain microvascular proteins showing the localization of the proteins (A), their functional class (B), and membrane and transporter protein classes. Other transporters (C, lower pie) refer to non-drug transporters, including, for example, ion transporters (C). Brain microvessel protein per gram brain was assessed in the microvascular fraction and ATPase activity allowed assessment of the level of enrichment of each of the 22 brains microvessel samples relative to their homogenates (D); ABC, ATP-binding cassette transporters; AD, Alzheimer's disease; BMvPGB, brain microvessel protein per gram brain; DLB, dementia with Lewy bodies; ER, endoplasmic reticulum; PM, plasma membrane; SLC, solute carriers. In A, percentages add to more than 100% due to expression of proteins in more than one subcellular location.

152x202mm (300 x 300 DPI)




Figure 3: Expression of blood-brain barrier (BBB) markers and transporters using MRM targeted proteomics. The abundances of the marker proteins of the neurovascular unit (A) showing markers for endothelial cells (GLUT1, PECAM1), neurons (SYP), astrocytes (GFAP), and pericytes (NG2); ATP1A1 is used as a plasma membrane marker for quality control and also as a reference for global proteomic quantification. After assessment of the purity of the microvascular fraction, the abundance of ABC and SLC transporters was quantified at the blood-brain barrier (B). Three sets of samples were assessed: healthy, AD and DLB, and ratios of average expression in AD and DLB sets relative the healthy set were calculated to assess differences (C). Only ratios >1.5 or <0.67 (horizontal lines) with statistical significance (stars) were considered meaningful; ABC, ATP-binding cassette transporters; AD, Alzheimer's disease; DLB, dementia with Lewy bodies; SLC, solute carrier transporters; *, $p < 0.05$; **, $p < 0.01$.

203x186mm (300 x 300 DPI)




Figure 4: Summary matrix of correlation analysis of ABC, SLC transporters and cell markers expression data. R_2 , linear regression correlation coefficient; R_s , Spearman Rank order correlation coefficient; only significant correlations are shown; transporters with strong correlation ($R_s > 0.75$) are marked with dark red; moderate correlations ($R_s > 0.60$) are marked with faint red; correlations with ($R_s < 0.60$) are marked with faint blue. $R_2 > 0.30$ is considered for limited scatter. Number of samples $n=21-22$.

181x177mm (300 x 300 DPI)




Figure 5: (A) Hierarchical cluster analysis (HCA) and (B) principal coordinates analysis (PCoA) of peptide and protein data for samples from healthy donors (H, green symbols) and patients with Alzheimer's disease (AD, red symbols) or dementia with Lewy bodies (DLB, blue symbols). The ellipses represent 95% confidence intervals. The ages of healthy donors outside the cluster were 54 (donor 2), 87 (donor 14), 89 (donor 21) and 90 (donor 22). Analysis of peptide and protein data returned confirmatory results.

256x159mm (300 x 300 DPI)




Figure 6: Analysis of the expression of transporters at the blood-brain barrier (BBB): (A) the number of ATP-binding cassette (ABC) and solute carrier (SLC) transporters identified using global and targeted proteomics. The total number of transporters is shown in brackets. (B) Principal components analysis (PCA) of targeted transporter data for samples from healthy donors (H, green symbols) and patients with Alzheimer's disease (AD, red symbols) or dementia with Lewy bodies (DLB, blue symbols). The ellipses represent 95% confidence intervals. The data were confirmatory of global protein and peptide cluster analysis. (C) HCA and heat map with correlation analysis of transporters protein expression. The associations between the genes (rows) were assessed using correlation analysis, whereas the hierarchical cluster analysis of the samples (columns) was based on Euclidian distance. The heat map is based on log-transformed abundance data.

224x146mm (300 x 300 DPI)





Figure 7: A comparison of protein expression levels of transporters in brain capillaries between Sprague Dawley rat and human using (A) in-house data 20 and (B) literature data.^{30,17}

149x77mm (600 x 600 DPI)



TOC
Graphical abstract

307x188mm (300 x 300 DPI)