

Brief Communication
Psychiatry & Psychology



Psychological Consequences of Survivors of COVID-19 Pneumonia 1 Month after Discharge

Hye Yoon Park ,¹ Jongtak Jung ,² Hye Youn Park ,³ So Hee Lee ,⁴
Eu Suk Kim ,² Hong Bin Kim ,² and Kyoung-Ho Song

¹Department of Psychiatry, Seoul National University Hospital, Seoul, Korea

²Department of Internal Medicine, Seoul National University Bundang Hospital, Seoul National University College of Medicine, Seongnam, Korea

³Department of Neuropsychiatry, Seoul National University Bundang Hospital, Seongnam, Korea

⁴Department of Psychiatry, National Medical Center, Seoul, Korea



Received: Aug 14, 2020

Accepted: Nov 17, 2020

Address for Correspondence:

Kyoung-Ho Song, MD, PhD

Department of Internal Medicine, Seoul National University Bundang Hospital, Seoul National University College of Medicine, 82 Gumi-ro 173-beon-gil, Bundang-gu, Seongnam 13620, Republic of Korea.
E-mail: khsongmd@snu.ac.kr

© 2020 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Hye Yoon Park

<https://orcid.org/0000-0003-4114-5102>

Jongtak Jung

<https://orcid.org/0000-0003-3497-0796>

Hye Youn Park

<https://orcid.org/0000-0002-5097-532X>

So Hee Lee

<https://orcid.org/0000-0002-9005-3207>

Eu Suk Kim

<https://orcid.org/0000-0001-7132-0157>

Hong Bin Kim

<https://orcid.org/0000-0001-6262-372X>

Kyoung-Ho Song

<https://orcid.org/0000-0002-4517-3840>

ABSTRACT

As the coronavirus disease 2019 (COVID-19) has rapidly spread worldwide, there are growing concerns about patients' mental health. We investigated psychological problems in COVID-19 patients assessed with self-reported questionnaires including the Patient Health Questionnaire-9, Generalized Anxiety Disorder-7 scale, and Impact of Event Scale-Revised Korean version. Ten patients who recovered from COVID-19 pneumonia without complications underwent self-reported questionnaires about 1 month after discharge. Of them, 10% reported depression and posttraumatic stress disorder (PTSD) while 50% had depression during the treatment. Perceived stigma and history of psychiatric treatment affected PTSD symptom severity, consistent with previous emerging infectious diseases. Survivors also reported that they were concerned about infecting others and being discriminated and that they chose to avoid others after discharge. Further support and strategy to minimize their psychosocial difficulties after discharge should be considered.

Keywords: COVID-19; SARS-CoV-2; Depression; PTSD; Survivor; Pneumonia

The coronavirus disease 2019 (COVID-19) has rapidly spread worldwide¹; accordingly, there are growing concerns in various fields about COVID-19 patients' mental health. Patients with severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) have been shown to experience psychological distress after cure as well as during their illness; in 2003, while 35% SARS survivors in Hong Kong reported significant anxiety and/or depressive symptoms at four weeks or more after discharge,² 43% of the MERS survivors had posttraumatic stress disorder (PTSD) and 27% had depression at 12 months after the outbreak.³ COVID-19, as a new type of emerging infectious disease (EID), shows different characteristics when compared with past EIDs, like high infectivity, prevalence, and public impact.¹ However, knowledge on COVID-19 patients' psychological distress after cure remains meager; thus, we retrospectively investigated the psychological problems in COVID-19 patients about 1 month after discharge from hospital.

We searched for all patients who were confirmed to have experienced a severe acute respiratory syndrome coronavirus 2 infection through reverse transcription polymerase chain

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Park HY,¹ Lee SH, Song KH; Data curation: Park HY,¹ Jung J, Kim ES, Song KH; Formal analysis: Park HY,¹ Song KH; Investigation: Kim ES, Kim HB, Song KH; Methodology: Park HY,¹ Jung J, Park HY,² Lee SH, Kim HB; Project administration: Song KH; Supervision: Park HY,² Lee SH, Kim HB; Writing - original draft: Park HY,¹ Song KH; Writing - review & editing: Jung J, Park HY,² Lee SH, Kim ES, Kim HB.
Park HY,¹ Park Hye Yoon; Park HY,² Park Hye Yoon.

reaction using respiratory specimen, and who received isolation treatment in the nationally designated high-level isolation unit in Seoul National University Bundang Hospital from January to 31 May 2020. Among these, we reviewed patients who responded to the following self-reported questionnaires to evaluate their depression, anxiety, and PTSD symptoms: the Patient Health Questionnaire-9 (PHQ-9),⁴ Generalized Anxiety Disorder-7 scale,⁵ and Impact of Event Scale-Revised Korean version (IES-R-K),⁶ respectively. The patients we reviewed also responded to an 8-item tool that was used in a previous study for MERS survivors,³ and that was utilized at this time to assess perceived stigma regarding COVID-19. This perceived stigma questionnaire was adopted from the 40-item human immunodeficiency virus (HIV) stigma scale, which was developed by Berger et al.⁷ and the 8-item short version of the HIV stigma scale.⁸ Responses were rated on a 4-point Likert scale, where higher scores meant higher levels of perceived stigma; we defined a high level of perceived stigma as any score that was two standard deviations from the mean. To compare mental health outcomes according to the sociodemographic characteristics and clinical features, we applied the Mann-Whitney U test with the significance set at $P < 0.05$.

Ten patients completed the self-reported questionnaires, and this was done about 1 month after discharge. Descriptive statistics are presented in **Table 1**. All patients were diagnosed with pneumonia through radiologic examination (i.e., chest and/or chest computed tomography [CT]); additionally, five patients required oxygen supplement therapy during hospitalization. The median duration of isolation treatment was 20 days (range, 18–38 days). All patients successfully recovered from COVID-19 pneumonia without complications or sequelae in the lung; this was assessed using chest CT \pm pulmonary function test. One month after discharge, all patients visited the hospital for a follow-up consultation (median, 25 days; range, 13–50 days).

Among the survivors, at 1 month post-discharge, 10% reported depression (i.e., PHQ-9 \geq 10) and PTSD (IES-R-K \geq 25); additionally, 50% reported depression during the treatment. No significant anxiety scores were observed after discharge. Interestingly, patients with high perceived stigma ($n = 4$) tended to have higher scores for PTSD symptoms (their IES-R-K mean score = 16.3 vs. 3.5 in patients with low perceived stigma; $P = 0.067$), especially in the hyperarousal (mean score = 4.8 vs. 0.2, $P = 0.010$) and numbness (mean score = 3.3 vs. 0.5, $P = 0.019$) domains. Survivors with a previous history of psychiatric treatment ($n = 3$) also had higher scores for PTSD symptoms (mean score = 20.3 vs. 3.6 in patients without a previous history of psychiatric treatment, $P = 0.017$); still, they did not show scores for depression and anxiety that differed significantly from survivors without such a history. There were no significant differences in the scores for depression, anxiety, or PTSD by pneumonia severity and whether their tracing routes were disclosed to the public. Notwithstanding, 40% of the survivors worried about infecting others and being discriminated against by neighbors owing to their history of COVID-19. They wanted to receive support from neighbors (60%), from their families (50%), and to undergo regular hospital check-ups after discharge (50%). Moreover, they needed accurate information from the government (40%) and access to psychiatric service (30%).

Our results indicated that survivors who get completely cured of COVID-19 pneumonia, without experiencing significant complications, may be at low risk for post-discharge depression or anxiety; still, they could develop or express depressive symptoms/depression during treatment. Moreover, our findings suggested that perceived stigma regarding COVID-19 and a previous history of psychiatric treatment may affect the severity of PTSD

Table 1. Participants' demographic and clinical characteristics and mental health status at 1 month after discharge from COVID-19 pneumonia in Korea

Characteristics	Variables	Total (n = 10)	Case No.									
			1	2	3	4	5	6	7	8	9	10
Sex	Male/Female	8/2	M	M	M	F	M	M	M	M	M	F
Age, yr	Mean (SD)	62.6 (14.9)	67	81	56	63	82	60	29	55	65	68
Marriage status	Married = yes	7	Yes	Yes	Yes	Yes		Yes		Yes	Yes	
Income	> 3,500 USD/month = yes	6	Yes		Yes			Yes	Yes	Yes	Yes	NA
Job	Employed = yes	6			Yes	Yes		Yes	Yes	Yes	Yes	
Religion	Having = yes	5		Yes		Yes	Yes				Yes	Yes
Underlying physical illness	Type of disease	6	HTN	HTN, DL	HTN, LC		HTN, CVD	CVD, DM			HTN	
Previous history of psychiatric treatment	Yes	3			Yes			Yes				Yes
Duration of hospitalization, day	Median (range)	20.9 (13-38)	13	22	19	17	15	26	15	21	23	38
Interval between follow-up visit and discharge, day	Median (range)	25.0 (13-50)	35	16	16	18	20	50	36	30	47	13
Pneumonia	Yes	10	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Required oxygen supplement therapy	Yes	5		Yes				Yes		Yes	Yes	Yes
Mechanical ventilation	Yes	0										
Having a family member or an acquaintance who was infected by COVID-19	Yes	4			Yes	Yes			Yes	Yes		
Having a family member or an acquaintance who died from COVID-19	Yes	2	Yes		Yes							
Mental health status												
During admission ^a	Depression (PHQ-9)	5 ^b	NA	2	20	19	17	11	11	4	4	6
	Anxiety (GAD-7)	1 ^c	NA	0	10	5	8	2	6	0	0	1
Post-discharge at 1 month	Depression (PHQ-9)	1 ^b	0	2	0	7	2	0	0	0	0	15
	Anxiety (GAD-7)	0 ^c	0	0	0	2	0	0	0	0	0	2
	PTSD (IES-R-K)	1 ^d	5	7	12	5	0	15	4	2	2	34
	Intrusion score		1	3	3	3	0	5	0	2	2	10
	Avoidance score		3	1	4	2	0	4	0	0	0	7
	Hyperarousal score		0	1	2	0	0	3	3	0	0	11
	Numbness score		1	2	3	0	0	3	1	0	0	6
Psychosocial experiences & patients' needs												
Perceived stigma regarding COVID-19		4 ^e	2	6	14	8	7	14	17	9	2	21
Worry about infecting others		4		Yes	Yes			Yes		Yes		
Being discriminated against by neighbors owing to their history of COVID-19 diagnosis		4		Yes	Yes			Yes				Yes
Worry about invasion of privacy		3				Yes		Yes				Yes
Avoiding other people		3		Yes				Yes				Yes
Need for support from family members		5			Yes	Yes	Yes	Yes			Yes	
Need for support from the neighborhood		6			Yes	Yes	Yes	Yes		Yes	Yes	
Need for regular medical check-ups after discharge		5		Yes	Yes		Yes	Yes				Yes
Need for psychiatric service		3	Yes				Yes	Yes				
Need for accurate information from the government		4	Yes				Yes	Yes			Yes	

COVID-19 = coronavirus disease 2019, SD = standard deviation, PHQ-9 = Patient Health Questionnaire-9, GAD-7 = Generalized Anxiety Disorder-7, PTSD = posttraumatic stress disorder, IES-R-K = Impact of Event Scale-Revised Korean version, HTN = hypertension, DL = dyslipidemia, LC = liver cirrhosis, CVD = cardiovascular disease, DM = diabetes mellitus, NA = not available.

^aThe data were acquired by recall; ^bNumber of patients with PHQ-9 ≥ 10; ^cNumber of patients with GAD-7 ≥ 10; ^dNumber of patients with IES-R-K ≥ 25; ^eNumber of patients with the perceived stigma score > 11.1. Perceived stigma regarding COVID-19 was assessed with an 8-item questionnaire which was used in a previous study.³ The range of the score was 0–24. High level of perceived stigma was defined as scores above > 11.1, which was two SD = 2.7 from the mean = 5.7.

symptoms; these were consistent with the highlights from a past study on MERS survivors, which was conducted in Korea in 2015.³ In 2015, Korea experienced an outbreak of MERS, and a study on survivors from this period showed that they experienced discrimination; specifically, people regarded them as perpetrators of the infection, even after being cured.⁹ Moreover, previous studies on SARS survivors in Hong Kong also remarked that they suffered from SARS-associated stigma after they returned to their community and it did not decrease over time.^{10,11} Perceived stigma that may be reshaped by survivors' experiences in

their community after discharge, for example, being discriminated against by neighbors could modify their response to—the potentially traumatic COVID-19 situation. Thus, although COVID-19 patients in our sample seemed to be able to recover from depression if they experienced physical improvements, dealing with their PTSD symptoms may prove more complicated; these symptoms depend on the attitudes of the community/society toward the survivors along the course of the pandemic, and they can persist for a long time if the attitudes are negative. Our findings also suggested that survivors could have been experiencing psychosocial difficulties after discharge, namely, when they interacted with other people. Some survivors still worried about infecting others as a perpetrator and exposure of their privacy that may be a source of being blamed by people regarding COVID-19 infection, which may lead to social avoidance.

This research presented only the preliminary results of an assessment of the mental health status post-infection of a small sample of people who were diagnosed with COVID-19 pneumonia, were subsequently cured, and who did not need ventilator treatment—namely, did not incur in major complications. Accordingly, studies with larger samples and that also include patients who received intensive care are warranted.

Nevertheless, we could infer from our results that some of the survivors might have experienced mental health problems according to their psychiatric history or their perception of COVID-19 associated stigma and could have incurred in psychosocial difficulties after discharge, especially in worry about discrimination. Thus, to enhance the psychosocial recovery of patients after COVID-19, stakeholders should utilize strategies aimed at managing their mental health and facilitating their reintegration into the community.

Ethics statement

The Institutional Review Board (IRB) of Seoul National University Bundang Hospital approved the study (IRB No. B-2006-618-104). The IRB waived the requirement for informed consent.

REFERENCES

1. World Health Organization (WHO). Novel coronavirus (2019-nCoV) situation reports. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. Updated 2020. Accessed June 9, 2020.
2. Cheng SK, Wong CW, Tsang J, Wong KC. Psychological distress and negative appraisals in survivors of severe acute respiratory syndrome (SARS). *Psychol Med* 2004;34(7):1187-95.
[PUBMED](#) | [CROSSREF](#)
3. Park HY, Park WB, Lee SH, Kim JL, Lee JJ, Lee H, et al. Posttraumatic stress disorder and depression of survivors 12 months after the outbreak of Middle East respiratory syndrome in South Korea. *BMC Public Health* 2020;20(1):605.
[PUBMED](#) | [CROSSREF](#)
4. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16(9):606-13.
[PUBMED](#) | [CROSSREF](#)
5. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166(10):1092-7.
[PUBMED](#) | [CROSSREF](#)
6. Lim HK, Woo JM, Kim TS, Kim TH, Choi KS, Chung SK, et al. Reliability and validity of the Korean version of the Impact of Event Scale-Revised. *Compr Psychiatry* 2009;50(4):385-90.
[PUBMED](#) | [CROSSREF](#)

7. Berger BE, Ferrans CE, Lashley FR. Measuring stigma in people with HIV: psychometric assessment of the HIV stigma scale. *Res Nurs Health* 2001;24(6):518-29.
[PUBMED](#) | [CROSSREF](#)
8. Wiklander M, Rydström LL, Ygge BM, Navér L, Wettergren L, Eriksson LE. Psychometric properties of a short version of the HIV stigma scale, adapted for children with HIV infection. *Health Qual Life Outcomes* 2013;11:195.
[PUBMED](#) | [CROSSREF](#)
9. Sim M. Psychological trauma of Middle East respiratory syndrome victims and bereaved families. *Epidemiol Health* 2016;38:e2016054.
[PUBMED](#) | [CROSSREF](#)
10. Lee S, Chan LY, Chau AM, Kwok KP, Kleinman A. The experience of SARS-related stigma at Amoy Gardens. *Soc Sci Med* 2005;61(9):2038-46.
[PUBMED](#) | [CROSSREF](#)
11. Siu JY. The SARS-associated stigma of SARS victims in the post-SARS era of Hong Kong. *Qual Health Res* 2008;18(6):729-38.
[PUBMED](#) | [CROSSREF](#)