

PTSD Symptoms across Pregnancy and Early Postpartum

Among Women with Lifetime PTSD Diagnosis

Maria Muzik, M.D., M.S.¹;

Ellen W. McGinnis, M.S.¹;

Erika Bocknek, PhD, LMFT²;

Diana Morelen, Ph.D.¹

Katherine Rosenblum Ph.D.¹;

Israel Liberzon M.D.¹,

Julia Seng PhD, CNM, FAAN¹

& James L. Abelson, MD, PhD¹

Affiliated Universities: ¹University of Michigan, Ann Arbor, MI, USA ²Wayne State University, Detroit, MI, USA

Acknowledgements

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/da.22465](#).

This article is protected by copyright. All rights reserved.

The research presented was supported through funds from the Department of National Institute of Health (NR008767; PI: Seng), the Department of National Institute of Health-Michigan Mentored Clinical Scholars Program awarded to MM (K12 RR017607-04, PI: D. Steingart), the National Institute of Mental Health -Career Development Award K23 (K23 MH080147-01, PI: Muzik), and the Michigan Institute for Clinical and Health Research (MICHHR, UL1TR000433, PI: Muzik).

Corresponding author: Maria Muzik, MD, M.S. Department of Psychiatry, University of Michigan muzik@med.umich.edu 4250 Plymouth Rd Room 2736 Ann Arbor MI 48109-2700 [734-232-0206](tel:734-232-0206) Fax: (734) 998-7268

Short Title: Peripartum PTSD Trajectories in Women At-Risk

Keywords: PTSD, Trauma, Pregnancy and Postpartum, Depression, maternal-child

Conflict of interest: None of the authors have any conflict of interest or financial disclosures related to this work.

Abstract

Background: Little is known about trajectories of PTSD symptoms across the peripartum period in women with trauma histories, specifically those who met lifetime PTSD diagnoses prior to pregnancy. The present study seeks to identify factors that influence PTSD symptom load across pregnancy and early postpartum, and study its impact on postpartum adaptation. **Method:** The current study is a secondary analysis on pregnant women with a Lifetime PTSD diagnosis ($N=319$) derived from a larger community sample who were interviewed twice across pregnancy (28 and 35 weeks) and again at 6 weeks postpartum, assessing socio-economic risks, mental health, past and ongoing trauma exposure, and adaptation to postpartum. **Results:** Using trajectory analysis, first we examined the natural course of PTSD symptoms based on patterns across peripartum, and found 4 distinct trajectory groups. Secondly, we explored factors (demographic, historical, and gestational)

that shape the PTSD symptom trajectories, and examined the impact of trajectory membership on maternal postpartum adaptation. We found that child abuse history, demographic risk, and lifetime PTSD symptom count increased pregnancy-onset PTSD risk, whereas gestational PTSD symptom trajectory was best predicted by interim trauma and labor anxiety. Women with the greatest PTSD symptom rise during pregnancy were most likely to suffer postpartum depression and reported greatest bonding impairment with their infants at 6 weeks postpartum. **Conclusions:** Screening for modifiable risks (interpersonal trauma exposure and labor anxiety) and /or PTSD symptom load during pregnancy appears critical to promote maternal wellbeing.

Introduction

Post-traumatic stress disorder (PTSD) is common following trauma, and low-income and minority women are at particularly high risk for both exposure to trauma^{1,2} and development of PTSD³⁻⁵. Prevalence rates among women ranges from 10-20% for lifetime PTSD and around 5%^{6,7} for current PTSD. With growing awareness of cross-generational transmission of risk for psychopathology there is increasing attention to PTSD during the peripartum (i.e., pregnancy and first year postpartum). PTSD rates are up to 24% during pregnancy for high-risk women who are racial minorities, teens, less educated or poor⁸.

Maternal PTSD in pregnancy, even after controlling for demographic factors, increases the odds for poor pregnancy and birthing outcomes, such as ectopic pregnancy, miscarriage, preterm labor⁸, and low birth weight⁹. Furthermore, maternal gestational stress, trauma, and PTSD adversely impact fetal functional and structural brain development. Maternal stress hormones during pregnancy modulate fetal stress hormone reactivity¹⁰ and babies of “stressed” mothers are likely to show altered HPA axis functioning of basal cortisol^{11,12} and stress reactivity^{13,14}. Additionally, animal work demonstrates a link between prenatal stress exposure and alterations in specific brain structures¹⁵. Although evidence from humans is still sparse in this area¹⁶, there is initial evidence that prenatal maternal stress, trauma or PTSD might be associated with child difficult temperament¹⁷⁻¹⁹ attention problems^{20,21}, mood and anxiety symptoms²²⁻²⁴, cognitive delay^{25,26}, and adult disease²⁷.

The peripartum period may be experienced as stressful and/or traumatic, particularly for women with past trauma histories²⁸. PTSD often exhibits chronic course with fluctuating symptom severity that can be exacerbated by re-exposure to traumatic/stressful events^{29,30}. Women with prior reproductive trauma³¹ or prior interpersonal trauma history, especially childhood maltreatment, may have increased risk for re-emergence of PTSD symptoms^{29,32,33}. Onoye and colleagues³⁴ reported that community-recruited women peaked with PTSD symptoms in late gestation if they were exposed to trauma during pregnancy.

Thus, reducing the prevalence of maternal trauma exposure and the incidence or severity of PTSD during pregnancy has important public health implications, for both the mother and child. Systematic screening for trauma exposure and signs of PTSD relapse early in gestation may reduce risk for relapse to clinical PTSD diagnosis during the peripartum, and subsequently reduce risks for adverse outcomes in both mother and baby^{11,35,36}. Women with PTSD symptoms in pregnancy might be in critical needs for services³⁷ in order to facilitate a healthy course of pregnancy and to promote adequate fetal growth and development.

Although cumulatively prior research has demonstrated clinically significant impact of PTSD on peripartum women³⁷ and their infants⁹⁻¹², there are no longitudinal studies of PTSD across peripartum in women at greatest risk. Therefore, the primary goal of this study is to examine the natural course of PTSD symptoms across the peripartum period in women with heightened vulnerability for illness or relapse (i.e., current/lifetime PTSD diagnoses due to interpersonal violence)³⁰. Based on prior works, we hypothesize that the majority of such vulnerable women will experience an increase in PTSD symptoms during the peripartum period^{28,31-34}. Additionally, although research shows that past trauma²⁸, poverty⁸, concurrent stress/trauma³⁴, history of

PTSD³⁰, and pregnancy²⁸ itself can increase risk for PTSD, we have no actual data to model the course of PTSD symptoms across pregnancy and early postpartum in these vulnerable women. Therefore, our secondary research question is to identify risk factors (demographic, historical, gestational) that shape the natural course of peripartum PTSD symptom trajectories. We hypothesize that more risk factors, especially experiencing trauma, will be associated with a greater increase in PTSD symptoms. Finally, our third research question will examine the impact of peripartum PTSD symptom trajectories on women's postpartum functioning in other domains relevant to motherhood (i.e., depression, subjective bonding with infant, breastfeeding).

This study is the first to longitudinally follow women with PTSD risk across gestation and early postpartum with the goals of describing naturalistic PTSD symptom patterns, identifying risk factors, and examining consequences of these PTSD patterns on postpartum adaptation. We aim to shed light on those vulnerable for and resilient to symptom exacerbation with the hope that findings can guide development of more targeted interventions to reduce illness burden and reduce transmission of risk to the child.

Method

The current study is a secondary analysis from a larger study focused on determining the effects of PTSD on childbearing outcomes (i.e., pregnancy health, birth, and postpartum adjustment) in a community sample of pregnant women (NIH NR008767; "the Stress, Trauma, Anxiety, and the Childbearing Year, STACY project"). Eligible women presenting to antenatal care (≥ 18 years, English speaking, expecting their first infant, < 28 weeks completed gestation) were recruited from 2005 to 2008. The study oversampled for low SES by recruiting from prenatal clinics of three academic health centers, two of which were in an urban area, with one serving 90% women who were publically insured. Women initiating prenatal care in these clinics were invited to participate by obstetric nurses and those interested in participating gave contact information and assent to be called up by a research survey organization (DataStat, Inc.) for more interviewing over the phone. DataStat team was comprised of 13 diverse women who specialized in health-related standardized telephone interviews, who conducted all participant interviews averaging 32 minutes each. Ten percent of interviews were monitored live by a supervisor. The phone contacts were as follows: at or before 28 weeks gestation (wave 1); at 35 weeks gestation (wave 2); and at 6 weeks postpartum (wave 3). Of the 2,689 interested women, 758 were unreachable, 284 were ineligible, 58 declined at the start of the first interview, and 8 interviews were interrupted and not completed, leaving a total of 1,581 women enrolled in the study. Prior published work shows detailed screening and enrollment flow chart on these 1,581 eligible women with complete first interviews at wave 1³⁸. Among these 1,581 women, 319 (20.2%) had met PTSD diagnosis (lifetime or current), 532 (33.6%) were subsyndromal (i.e., had few PTSD symptoms but never had met diagnostic criteria), 380 (24%) were trauma-exposed resilient without PTSD, and finally, 350 (22.1%) were healthy controls. For the current study, we selected the subsample of women with heightened risk for PTSD, those who either at present or in lifetime had met diagnostic criteria for PTSD ($n=319$, 20.2% of total sample), with the aim to study the course of symptoms among these risk-women across gestation and early postpartum (see Table 1).

Measures

Demographics. Demographics were assessed using the Perinatal Risk Assessment Monitoring Survey (PRAMS)³⁹. A score of “1” was assigned for each “risk” (under 21 years, non-White, no partner, high school diploma or less, income less than \$15,000); scores were summed as a cumulative risk index for analyses⁴⁰.

Historical and Gestational Characteristics. The National Women's Study PTSD Module (NWS-PTSD)⁴¹ was utilized for the assessment of both continuous PTSD 17 symptom count and a diagnosis based on DSM-IV criteria⁴¹. Depression was assessed at wave 1 with the Composite International Diagnostic Interview short form (CIDI)⁴². The Dissociative Experience Scale-Taxonomic Version (DES-T)⁴³ assessed lifetime status of dissociative experiences commonly associated with PTSD and childhood abuse⁴⁴. The Life Stressor Checklist-Revised (LSC-R)⁴⁵ assessed lifetime trauma for 29 potential trauma exposures. Three subscales were derived from this checklist including: child abuse (5 items), adult abuse (3 items), and non-abuse trauma (21 items). In wave 2, 13 of the items assessed the occurrence of concurrent stress/trauma, the most frequent of which included ‘severe financial problems’, ‘family member put in jail’ and ‘care for an ill family member’. Also at wave 2, labor anxiety was assessed using a 5-item subscale of the Childbirth Self-Efficacy Inventory⁴⁶, social support was assessed using the Family APGAR⁴⁷, a five-item 5-point scale that measures family functioning, and the client’s appraisal of the quality of her maternity care providers was assessed using the Health Care Alliance Questionnaire, a 16-item scale adapted from therapeutic alliance measurements⁴⁸.

Postpartum Adaptation. Using the Postpartum Bonding Questionnaire⁴⁹, participants self-rated problematic bonding with their infant and indicated whether they were breastfeeding at wave 3. Depression was assessed using the Postpartum Depression Screening Scale (PDSS)⁵⁰ which is more detailed and valid in obstetric settings compared to other common screening measures⁵¹. It provides a continuous score (35-175), with a validated cut-off >80 indicating probable depressive diagnosis⁵⁰.

Plan for Analyses

Missingness. Prior to analyses addressing the primary research aim, we performed imputation of missing data. First, item-level imputation was done to compensate for a single item omission error of a PTSD symptom at wave 2 and 3 using the dissociation score and trauma items (Receptive Operative Characteristics curve of .91). Following guidelines⁵², we also imputed for attrition because 42.9% had missing data at wave 2 and 51.1% at wave 3. These rates are consistent with methodological studies showing imputation should not yield bias in parameter estimates^{53–55}. Attrition analyses found those who completed all three waves of data collection had significantly lower cumulative sociodemographic risk than those lost to follow up, but there were no significant

differences in PTSD status or outcome variables of interest. We proceeded with the SAS PROC TRAJ procedure with a Poisson distribution, an approach that handles random wave nonresponse. Results describing the complete dataset ($N=319$) and a smaller dataset without missing values ($n=156$) were similar.

Hypothesis testing. To examine the course of PTSD symptoms across peripartum, we conducted semiparametric, group-based trajectory analysis⁵⁶. Conclusions about the best number and shape of the trajectory groups are based on the Bayesian Information Criterion (BIC)⁵⁶, seeking the most parsimonious models⁵⁶. To examine demographic, historical, and gestational predictors ($N=319$) and postpartum adaptation ($n=156$) that might explain the distinct PTSD trajectories, we utilized two-tailed ANOVAs and chi-square tests, and General Linear Model regressions with manual backward steps.

Results

PTSD symptom trajectory groups

Posterior probability scores $\geq .80$ were calculated to assign each participant to a trajectory yielding the optimal number of distinct PTSD trajectories of four, with quadratic shapes (BIC=-2431; Figure 1). This means that PTSD symptoms over time were likely to follow a non-linear path for all groups; however, the groups meaningfully differed from each other. We identified two groups with rapid symptom changes across gestation: a rapidly increasing group (Increase), and a rapidly decreasing group (Decrease). These groups had significantly different starting points in early pregnancy, with Increase intercept significantly lower than Decrease ($t(173)=-15.54, p < .001$). Across gestation the trajectories of these 2 groups cross— Increase shows significant rise from T1 to T2 ($t(84)=-10.00, p=.000$), whereas Decrease shows significant decline ($t(98)=15.85, p=.000$). At 6 weeks postpartum Increase endpoint is significantly higher than Decrease ($t(181)=5.41, p < .001$).

In contrast, the other two trajectory groups show more stable patterns. One group shows constant low PTSD symptoms across pregnancy to 6 weeks postpartum (Low; T1 to T2 $t(49)=-.49, p=ns$, and T2 to T3 ($t(49)=-2.19, p=.033$). Finally, the remaining trajectory group shows a slow significant decline of PTSD symptoms across this time period (T1 to T2 ($t(86)=3.93, p=.000$); and T2 to T3 ($t(86)=6.45, p=.000$), yet remains overall significantly above all other PTSD trajectory groups (High).

PTSD diagnostic rates highlight the clinical significance of these PTSD symptom trajectories throughout peripartum (see bottom of Figure 1).

Demographic, historical, and gestational factors by trajectory group

First we analyzed demographic and historical predictors to PTSD trajectory groups (Table 2). Women in the trajectory groups that showed *low* PTSD symptom load in early gestation (i.e., Low and Increase) were *less* likely non-White ($\chi^2(3, N=318)=44.05, p=.000$), poor ($F(3, 306)=20.42, p=.000$), single ($\chi^2(3, N=319)=45.30, p=.000$), and very young ($F(3, 315)=29.75, p=.000$), and overall had *less cumulative risk* ($F(3, 315)=39.59, p=.000$) (< 21 years, < \$15,000, <=HS education, non-White, and not having a partner) compared to women in the other 2 groups (i.e., Decrease and High). Not having a high school education ($\chi^2(3, N=319)=72.38, p=.000$) followed a dose effect in the expected direction (Low > Increase > Decrease > High).

As historical predictors, we tested lifetime PTSD and dissociation symptoms, and lifetime trauma exposure. We found that Low and Increase had fewer lifetime PTSD symptoms than Decrease and High ($F(3, 315)=28.87, p=.000$). Moreover, Low experienced significantly less child abuse than High ($F(3, 315)=3.77, p=.011$). Finally, lifetime dissociation scores also differed by group ($F(3, 315)=4.27, p=.006$); Low and Decrease were similar in dissociation scores, and both had significantly lower scores than High. Lifetime adult abuse and lifetime non-abuse trauma did not differentiate group membership.

We next explored gestational factors and found anticipatory labor anxiety, social support, comorbid depression, and trauma exposure during pregnancy as relevant for symptom course. Specifically, we found that Low experienced less labor anxiety compared to all other groups ($F(3, 147)=6.09, p=.001$); and had significantly more social support than High ($F(3, 178)=3.48, p=.017$). Low and Increase also had less comorbid depression compared to Decrease and High ($\chi^2(3, N=319)=13.43, p=.004$). Trauma-exposure differed by group status ($F(3, 178)=3.48, p=.017$), but this time Low and Decrease were less likely to experience trauma compared to Increase and High. Health care alliance did not differentiate group membership.

Post-Hoc Analyses Predicting PTSD symptom crossing patterns

Intrigued by the crossing pregnancy PTSD trajectory patterns (Increase vs Decrease), we investigated predictors of these patterns, using for each as contrast the group with complimentary starting point. We found as predictors for Increase (compared to Low) having more stressors during pregnancy ($\beta=1.448, p=.013$) and more labor anxiety ($\beta=.274, p=.015$) ($\chi^2(3, N=72)=16.49, p=.000$; Nagelkerke $R^2=.281$). By contrast, predictors for Decrease (compared to High) were less pregnancy stressors ($\beta=1.307, p=.016$) and less lifetime PTSD symptoms ($\beta=.376, p=.001$) ($\chi^2(3, N=79)=25.18, p=.000$; Nagelkerke $R^2=.364$).

Trajectory group impact on postpartum adaptation

PTSD trajectory group membership, rather than initial baseline PTSD status, was significantly associated with postpartum adaptation (Table 3). Women across the PTSD trajectory groups differed on 6 weeks postpartum depression ($\chi^2(3, N=156)=12.17, p=.007$), with Increase group

showing greatest prevalence of major depression (52.3%), followed by Decrease and High (48.4% and 41.7%, respectively), and then Low (15.2%). Results for bonding impairment paralleled depression findings ($F(3, 152)=6.08, p=.001$), such that Increase mothers reported highest bonding problems with their infants compared to all other groups. Low rates of breastfeeding at 6 weeks postpartum were associated with mothers' PTSD symptoms in early pregnancy, but not concurrent PTSD ($F(3, 155)=15.843, p=.001$).

Discussion

In this secondary data analysis paper utilizing a sample of pregnant community women with PTSD risk, we aimed to examine naturalistically occurring patterns of PTSD symptoms across gestation and early postpartum (i.e., peripartum), and relate symptom occurrence to presence or absence of various demographic, historical, and gestational factors. We found four distinct patterns of PTSD symptoms across peripartum. We had proposed, that women with PTSD risk based on lifetime/current PTSD diagnoses such as those in our sample, would all increase in PTSD symptoms across gestation. However, in contrast to our primary hypothesis, only one group (26% of sample) significantly increased in PTSD symptoms over peripartum. This was surprising and unexpected. Moreover, the majority of women in our sample (58%) started with high PTSD rates, and these women all significantly decreased in symptom load throughout peripartum. We identified, not surprisingly, several risk factors (i.e., current/lifetime PTSD symptoms, history of child abuse, demographic risk) associated with higher PTSD symptoms at the early pregnancy starting point. Furthermore, variation in women's PTSD trajectories across peripartum was best predicted by their experience of pregnancy stress/trauma exposure, labor anxiety, and lifetime PTSD symptoms; confirming our second hypothesis that stress/trauma during pregnancy would be a particularly salient predictor for symptom exacerbation. Finally, PTSD trajectory was significantly associated with postpartum adjustment such that women whose PTSD symptoms rose during pregnancy were most impacted by postpartum depression and subjective bonding impairment with their infant at 6 weeks postpartum.

In contrast with our first hypothesis and despite substantial impairment during early pregnancy, the majority of vulnerable women in our cohort did not increase in PTSD symptoms during peripartum. In fact, the two highest-risk groups of women who had started pregnancy with high PTSD symptoms (High and Decrease) both dropped in PTSD diagnosis across peripartum by almost half, confirming prior work³⁴ that showed PTSD symptom reduction in postpartum. We speculate that increases in estradiol and progesterone concentrations at the end of pregnancy⁵⁷ may be associated with lower perceived distress and more positive perception of peripartum to promote reproduction and offspring care⁵⁸. This was pronounced in High mothers who reported low subjective bonding impairment. These significant improvements in the High group occurred despite these women carrying great historical and psychosocial burden: 50-70% of these High women had entered pregnancy with a current PTSD diagnosis, the majority had high lifetime PTSD symptoms, childhood abuse exposure, were younger, more likely poor, uneducated, single and of ethnic

minority status. All these risks are in fact associated with greater illness burden in the literature (^{28,59,60,61}), thus, it is interesting that PTSD women actually improve in symptom load across pregnancy. More research with larger samples and a wider range of potential predictors (e.g., physical health complications, factors related to maternity care, mental health treatment, biological markers etc) is needed to shed light on this phenomenon.

Women classified in Decrease dropped dramatically in PTSD diagnoses across pregnancy (56.6 % to only 1.8%). Interestingly, Decrease members shared one similar feature with the Low group—only a few experienced stress/trauma during pregnancy. Based on these findings, we propose that excessive stress/trauma exposure during pregnancy is the most salient factor for mental health in pregnancy and immediate postpartum, confirming our second hypothesis and prior work on the detrimental effects of trauma/stress on maternal perinatal outcomes ^{8,10,62} ..

The women showing the greatest change in risk constellation across gestation were mothers in the Increase group; starting with low PTSD symptoms and increasing to high rates of active illness in postpartum (PTSD: 23%; MDD: 52%). The predictors for this striking deterioration were 2-fold more exposure to stress/trauma during pregnancy, and higher labor-related anxiety compared to members in the Low group. This is consistent with past studies finding that interim stress or trauma leads to a 2.4-fold increase in PTSD symptoms across pregnancy ³⁴.

Increase mothers showed highest rates of postpartum mental illness (>50% depression and 22% PTSD) and reported the highest levels of subjective bonding impairment with their infants. This finding is consistent with prior research showing link between postpartum mental illness and poor bonding⁶³. While bonding impairment is a mother's self-perception and open to distortions, prior research has show associations to observed parenting⁶³. Because bonding impairment was associated with PTSD trajectory across pregnancy, and not simply the postpartum PTSD symptom load alone, our data seem to validate the clinical utility of longitudinal data collection over and above cross-sectional data. Interestingly, breastfeeding frequencies were not associated with group trajectories, even in Increase mothers, who had high rates of postpartum psychopathology. Perhaps breastfeeding may be more a product of education and other demographic factors⁶⁴, than driven by psychopathology.

Our findings are a novel contribution to the field as no known past research has considered factors related to PTSD symptom trajectories across the perinatal period. Put together, our results highlight the essential need for repeated assessment of symptomology during this time given the substantial fluctuation in symptoms. This is relevant for several reasons. First, it gives a message of hope that a current/lifetime diagnosis of PTSD is by no means deterministic of worsening PTSD in the perinatal period. In fact, results indicate that it is more likely than not that PTSD symptoms will remain the same or get better during the perinatal period. Second, it highlights a vulnerable group (Increase) and describes factors that providers can identify as heightening risk for symptom exacerbation (e.g., pregnancy stressors/trauma, labor-related anxiety, lifetime PTSD load). Preventing symptom exacerbation and manifest postpartum illness protects the offspring from the detrimental impact of postpartum mental illness on parental capacity⁶⁵. Treatment providers could

quickly identify (via a few questions/brief screening measures) mothers who are experiencing such risk factors and heighten awareness for support, case management and referrals for treatment. Lastly, our results identify a modifiable protective factor that promotes resilience in risk women with lifetime-PTSD, namely social support (more so than healthcare alliance).

Limitations

This study selected participants with heightened risk for PTSD who at baseline met either lifetime or current diagnostic criteria, thus women's symptom levels at baseline were high. As such, generalizability of the findings from this population to other obstetrics populations and women with less baseline risk may be limited. In order to maximize our sample size, we imputed data; this is a limitation, which is mitigated by the use of a sophisticated approach and by validation of results of the total dataset using the non-imputed data ($n=156$). It would have benefitted the study if measures assessed at only wave 2 were assessed at multiple time points to better understand their roles as group determinants. Although we may infer negative child outcomes of mothers with subjective impaired bonding and postpartum psychopathology based on the literature, we remain cautious in this implication and suggest future research directly investigate this link. As our study was a secondary analysis with limited range of collected variables potential confounding factors (e.g., physical health complications etc.) that may influence PTSD risk were unavailable. Lastly, we focused on the progression of PTSD, and although the overlap of PTSD with depression was high for some groups, for others the link was more distinct. Future studies could better understand the interaction of these two psychopathologies over time. Please note, PTSD diagnosis is based on DSM-IV⁶⁶ criteria.

References

1. Gillespie CF, Bradley B, Mercer K, et al. Trauma exposure and stress-related disorders in inner city primary care patients. *Gen Hosp Psychiatry*. 2009;31(6):505-514. doi:10.1016/j.genhosppsy.2009.05.003.
2. Dailey DE, Humphreys JC, Rankin SH, Lee KA. An exploration of lifetime trauma exposure in pregnant low-income African American women. *Matern Child Health J*. 2011;15(3):410-418. doi:10.1007/s10995-008-0315-7.
3. Alim TN, Graves E, Mellman TA, et al. Trauma exposure, posttraumatic stress disorder and depression in an African-American primary care population. *J Natl Med Assoc*. 2006;98(10):1630-1636.
4. Schwartz AC, Bradley RL, Sexton M, Sherry A, Ressler KJ. Posttraumatic stress disorder among African Americans in an inner city mental health clinic. *Psychiatr Serv*. 2005;56(2):212-215. doi:10.1176/appi.ps.56.2.212.
5. Norris F, Foster J, Weishaar D. The epidemiology of gender differences in PTSD across developmental, societal, and research contexts. In: *Gender and PTSD*. Kimerling, Ouimette, & Wofle. New York: The Guilford Press; 2002.
6. Resnick HS, Kilpatrick DG, Dansky BS, Saunders BE, Best CL. Prevalence of civilian trauma and posttraumatic stress disorder in a representative national sample of women. *J Consult Clin Psychol*. 1993;61(6):984-991.
7. Seng JS, Low LK, Sperlich M, Ronis DL, Liberzon I. Prevalence, trauma history, and risk for posttraumatic stress disorder among nulliparous women in maternity care. *Obstet Gynecol*. 2009;114(4):839-847. doi:10.1097/AOG.0b013e3181b8f8a2.
8. Seng JS, Oakley DJ, Sampsel CM, Killion C, Graham-Bermann S, Liberzon I. Posttraumatic stress disorder and pregnancy complications. *Obstet Gynecol*. 2001;97(1):17-22.
9. Seng JS, Low LK, Sperlich M, Ronis D, Liberzon I. Post-traumatic stress disorder and birthweight: methodological challenges. *An International Journal of Obstetrics & Gynaecology*. 2012;119(111). doi:10.1111/j.1471-0528.2011.03200.x.
10. Waffarn F, Davis EP. Effects of antenatal corticosteroids on the hypothalamic-pituitary-adrenocortical axis of the fetus and newborn: experimental findings and clinical considerations. *Am J Obstet Gynecol*. 2012;207(6):446-454. doi:10.1016/j.ajog.2012.06.012.
11. Yehuda R, Engel SM, Brand SR, Seckl J, Marcus SM, Berkowitz GS. Transgenerational effects of posttraumatic stress disorder in babies of mothers exposed to the World Trade Center attacks during pregnancy. *J Clin Endocrinol Metab*. 2005;90(7):4115-4118. doi:10.1210/jc.2005-0550.

12. Brennan PA, Pargas R, Walker EF, Green P, Newport DJ, Stowe Z. Maternal depression and infant cortisol: influences of timing, comorbidity and treatment. *J Child Psychol Psychiatry*. 2008;49(10):1099-1107. doi:10.1111/j.1469-7610.2008.01914.x.
13. Davis EP, Snidman N, Wadhwa PD, Glynn LM, Schetter CD, Sandman CA. Prenatal Maternal Anxiety and Depression Predict Negative Behavioral Reactivity in Infancy. *Infancy*. 2004;6(3):319-331. doi:10.1207/s15327078in0603_1.
14. Grant K-A, McMahon C, Austin M-P, Reilly N, Leader L, Ali S. Maternal prenatal anxiety, postnatal caregiving and infants' cortisol responses to the still-face procedure. *Dev Psychobiol*. 2009;51(8):625-637. doi:10.1002/dev.20397.
15. Charil A, Laplante D, Vaillancourt C, King S. Prenatal Stress and Brain Development. *Brain Research Reviews*. 2010;65(1):56-79.
16. Buss C, Davis EP, Muftuler LT, Head K, Sandman CA. High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6-9-year-old children. *Psychoneuroendocrinology*. 2010;35(1):141-153. doi:10.1016/j.psyneuen.2009.07.010.
17. Van den Bergh BRH, Mennes M, Stevens V, et al. ADHD deficit as measured in adolescent boys with a continuous performance task is related to antenatal maternal anxiety. *Pediatr Res*. 2006;59(1):78-82. doi:10.1203/01.pdr.0000191143.75673.52.
18. Van Os J, Selten JP. Prenatal exposure to maternal stress and subsequent schizophrenia. The May 1940 invasion of The Netherlands. *Br J Psychiatry*. 1998;172:324-326.
19. Wadhwa PD, Porto M, Garite TJ, Chicx-DeMet A, Sandman CA. Maternal corticotropin-releasing hormone levels in the early third trimester predict length of gestation in human pregnancy. *Am J Obstet Gynecol*. 1998;179(4):1079-1085.
20. Weaver ICG, Cervoni N, Champagne FA, et al. Epigenetic programming by maternal behavior. *Nat Neurosci*. 2004;7(8):847-854. doi:10.1038/nn1276.
21. Weaver ICG, Champagne FA, Brown SE, et al. Reversal of maternal programming of stress responses in adult offspring through methyl supplementation: altering epigenetic marking later in life. *J Neurosci*. 2005;25(47):11045-11054. doi:10.1523/JNEUROSCI.3652-05.2005.
22. Weaver ICG, D'Alessio AC, Brown SE, et al. The transcription factor nerve growth factor-inducible protein a mediates epigenetic programming: altering epigenetic marks by immediate-early genes. *J Neurosci*. 2007;27(7):1756-1768. doi:10.1523/JNEUROSCI.4164-06.2007.
23. Wei Q, Lu X-Y, Liu L, et al. Glucocorticoid receptor overexpression in forebrain: a mouse model of increased emotional lability. *Proc Natl Acad Sci USA*. 2004;101(32):11851-11856. doi:10.1073/pnas.0402208101.

24. Welberg LA, Seckl JR, Holmes MC. Inhibition of 11 β -hydroxysteroid dehydrogenase, the foeto-placental barrier to maternal glucocorticoids, permanently programs amygdala GR mRNA expression and anxiety-like behaviour in the offspring. *Eur J Neurosci*. 2000;12(3):1047-1054.
25. Whincup PH, Kaye SJ, Owen CG, et al. Birth weight and risk of type 2 diabetes: a systematic review. *JAMA*. 2008;300(24):2886-2897. doi:10.1001/jama.2008.886.
26. Kapoor A, Petropoulos S, Matthews SG. Fetal programming of hypothalamic-pituitary-adrenal (HPA) axis function and behavior by synthetic glucocorticoids. *Brain Res Rev*. 2008;57(2):586-595. doi:10.1016/j.brainresrev.2007.06.013.
27. Cottrell EC, Seckl JR. Prenatal stress, glucocorticoids and the programming of adult disease. *Front Behav Neurosci*. 2009;3:19. doi:10.3389/neuro.08.019.2009.
28. Seng JS, Rauch SAM, Resnick H, et al. Exploring posttraumatic stress disorder symptom profile among pregnant women. *J Psychosom Obstet Gynaecol*. 2010;31(3):176-187. doi:10.3109/0167482X.2010.486453.
29. Breslau N, Peterson EL. Assaultive violence and the risk of posttraumatic stress disorder following a subsequent trauma. *Behaviour Research and Therapy*. 2010;48(10):1063-1066. doi:10.1016/j.brat.2010.07.001.
30. Breslau N, Peterson EL, Schultz LR. A second look at prior trauma and the posttraumatic stress disorder effects of subsequent trauma: a prospective epidemiological study. *Arch Gen Psychiatry*. 2008;65(4):431-437. doi:10.1001/archpsyc.65.4.431.
31. Born L, Soares CN, Phillips S-D, Jung M, Steiner M. Women and reproductive-related trauma. *Ann N Y Acad Sci*. 2006;1071:491-494. doi:10.1196/annals.1364.049.
32. Banyard VL, Williams LM, Siegel JA. The long-term mental health consequences of child sexual abuse: an exploratory study of the impact of multiple traumas in a sample of women. *J Trauma Stress*. 2001;14(4):697-715. doi:10.1023/A:1013085904337.
33. Roodman AA, Clum GA. Revictimization rates and method variance: a meta-analysis. *Clin Psychol Rev*. 2001;21(2):183-204.
34. Onoye JM, Shafer LA, Goebert DA, Morland LA, Matsu CR, Hamagami F. Changes in PTSD symptomatology and mental health during pregnancy and postpartum. *Arch Womens Ment Health*. 2013;16(6):453-463. doi:10.1007/s00737-013-0365-8.
35. Famularo R, Fenton T, Kinscherff R, Ayoub C, Barnum R. Maternal and child posttraumatic stress disorder in cases of child maltreatment. *Child Abuse Negl*. 1994;18(1):27-36.
36. Spell AW, Kelley ML, Wang J, et al. The moderating effects of maternal psychopathology on children's adjustment post-Hurricane Katrina. *J Clin Child Adolesc Psychol*. 2008;37(3):553-563. doi:10.1080/15374410802148210.

37. Smith MV, Poschman K, Cavaleri MA, Howell HB, Yonkers KA. Symptoms of posttraumatic stress disorder in a community sample of low-income pregnant women. *Am J Psychiatry*. 2006;163(5):881-884. doi:10.1176/appi.ajp.163.5.881.
38. Seng JS, Kane Low LM, Sperlich MI, Ronis DL, Liberzon I. Prevalence of PTSD in relation to antecedent trauma exposures in a community sample of pregnant women. *Obstet Gynecol*. 2009;114:839-847.
39. Beck L, Lipscomb L, Johnson C, Gaffield ME, Rogers M, Gilvert B. Prevalence of selected maternal behaviors and experiences, Pregnancy Risk Assessment Monitoring System (PRAMS). *Morbidity and Mortality Weekly Report Surveillance Summaries (Washington, DC: 2002)*. 1999;51(SS02):1-26.
40. Sameroff AJ, Seifer R, Baldwin A, Baldwin C. Stability of intelligence from preschool to adolescence: the influence of social and family risk factors. *Child Dev*. 1993;64(1):80-97.
41. Kilpatrick D, Resnick H, Saunders B, Best C. The National Women's Study PTSD Module. *Medical University of South Carolina, Dept of Psychiatry and Behavioral Sciences*. 1989.
42. Wittchen HU. Reliability and validity studies of the WHO--Composite International Diagnostic Interview (CIDI): a critical review. *J Psychiatr Res*. 1994;28(1):57-84.
43. Waller N, Putnam FW, Carlson EB. Types of dissociation and dissociative types: A taxometric analysis of dissociative experiences. *Psychological Methods*. 1996;1(3):300-321. doi:10.1037/1082-989X.1.3.300.
44. Roth S, Newman E, Pelcovitz D, van der Kolk B, Mandel FS. Complex PTSD in victims exposed to sexual and physical abuse: results from the DSM-IV Field Trial for Posttraumatic Stress Disorder. *J Trauma Stress*. 1997;10(4):539-555.
45. Wolfe J, Kimerling R. Gender issues in the assessment of Posttraumatic Stress Disorder. In: *Assessing Psychological Trauma and PTSD*. New York: Guilford Press; 1997:192-238.
46. Drummond J, Rickwood D. Childbirth confidence: validating the Childbirth Self-Efficacy Inventory (CBSEI) in an Australian sample. *J Adv Nurs*. 1997;26(3):613-622.
47. Smilkstein G. The family APGAR: a proposal for a family function test and its use by physicians. *J Fam Pract*. 1978;6(6):1231-1239.
48. Roosevelt LK, Holland KJ, Hiser J, Seng JS. Psychometric assessment of the Health Care Alliance Questionnaire with women in prenatal care. *J Health Psychol*. October 2013. doi:10.1177/1359105313506027.
49. Brockington IF, Fraser C, Wilson D. The Postpartum Bonding Questionnaire: a validation. *Arch Womens Ment Health*. 2006;9(5):233-242. doi:10.1007/s00737-006-0132-1.

50. Beck CT, Gable RK. Postpartum Depression Screening Scale: development and psychometric testing. *Nurs Res*. 2000;49(5):272-282.
51. Beck CT, Gable RK. Comparative analysis of the performance of the Postpartum Depression Screening Scale with two other depression instruments. *Nurs Res*. 2001;50(4):242-250.
52. Acock AC. Working with missing values. *J Marriage Fam*. 2005;67(4):1012-1028. doi:10.1111/j.1741-3737.2005.00191.x.
53. Graham JW. Analysis of Missing Data. *Missing Data*. 2012:47-69.
54. Jelčić H, Phelps E, Lerner RM. Why missing data matter in the longitudinal study of adolescent development: using the 4-H Study to understand the uses of different missing data methods. *J Youth Adolesc*. 2010;39(7):816-835. doi:10.1007/s10964-010-9542-5.
55. Newman DA. Longitudinal modeling with randomly and systematically missing data: A simulation of ad hoc, maximum likelihood, and multiple imputation techniques. *Organizational Research Methods*. 2003;6(3):328-362.
56. Nagin D, Tremblay RE. Trajectories of boys' physical aggression, opposition, and hyperactivity on the path to physically violent and nonviolent juvenile delinquency. *Child Dev*. 1999;70(5):1181-1196.
57. Martin M, Hoffman P. The endocrinology of pregnancy. In: *Basic and Clinical Endocrinology*. 2nd ed. Los Altos, CA: Lange; 1986:476-500.
58. Stern D. *The Motherhood Constellation: A Unified View of Parent–infant Psychotherapy*. New York, NY US: Basic Books; 1995.
59. Morland L, Goebert D, Onoye J, et al. Posttraumatic stress disorder and pregnancy health: preliminary update and implications. *Psychosomatics*. 2007;48(4):304-308. doi:10.1176/appi.psy.48.4.304.
60. Hobfoll SE, Ritter C, Lavin J, Hulsizer MR, Cameron RP. Depression prevalence and incidence among inner-city pregnant and postpartum women. *J Consult Clin Psychol*. 1995;63(3):445-453.
61. Spinelli MG, Goetz RR. Symptoms of depression in a minority population of women from an urban obstetrics and gynecology clinic. *Gend Med*. 2011;8(5):320-324. doi:10.1016/j.genm.2011.06.001.
62. Seng JS, Low LK, Sperlich M, Ronis DL, Liberzon I. Post-traumatic stress disorder, child abuse history, birthweight and gestational age: a prospective cohort study. *BJOG*. 2011;118(11):1329-1339. doi:10.1111/j.1471-0528.2011.03071.x.

63. Muzik M, Bocknek EL, Broderick A, et al. Mother–infant bonding impairment across the first 6 months postpartum: the primacy of psychopathology in women with childhood abuse and neglect histories. *Archives of Women's Mental Health*. 2012:1-10.
64. Flacking R, Nyqvist KH, Ewald U. Effects of socioeconomic status on breastfeeding duration in mothers of preterm and term infants. *Eur J Public Health*. 2007;17(6):579-584. doi:10.1093/eurpub/ckm019.
65. Waxler E, Thelen K, Muzik M. Maternal Perinatal Depression: Impact on Infant and Child Development. 2011.
66. American Psychiatric Association. *Diagnostic and Statistical Manual for Mental Disorders*. 4th ed. text revision (DSM-IV-TR). Washington, DC US: American Psychiatric Press; 2000.

Table 1. Sample Characteristics (N=319)

| Characteristic | Category | n or M | SD or % |
|----------------|------------------------|--------|---------|
| Maternal Age | | 26 | 6 |
| Race | Caucasian | 107 | 34 |
| | African American | 155 | 49 |
| | Asian | 17 | 5 |
| | Hispanic | 4 | 1 |
| | Native American | 8 | 3 |
| Marital Status | Married/partnered | 164 | 51 |
| | Never married/divorced | 155 | 49 |

| | | | |
|------------------|-------------------------|------|------|
| Education | College or greater | 144 | 45 |
| | High school/GED or less | 175 | 55 |
| Household Income | <\$15,000 | 100 | 32 |
| | \$15,000-\$25,000 | 56 | 18 |
| | \$25,000-\$50,000 | 60 | 19 |
| | >\$50,000 | 94 | 30 |
| Traumatic Events | Child Abuse | .83 | 1.05 |
| | Adult Abuse | .50 | .76 |
| | Non-Abuse Trauma | 3.79 | 2.14 |

Table 2. Demographic, and Historical Characteristics by Trajectory Group (N=319)

| | | <i>Low (N=49)</i> | | <i>Increase (N=84)</i> | | <i>Decrease (N=99)</i> | | <i>High (N=87)</i> | |
|---------------------|-------------|--------------------|-------------|------------------------|-------------|------------------------|-------------|--------------------|-------------|
| | | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> |
| Demographics | | | | | | | | | |
| Ethnicity | | | | | | | | | |
| | % Non-White | 44.90 ^a | | 46.40 ^a | | 77.80 ^b | | 84.90 ^b | |
| Education | | | | | | | | | |
| | % < HS | 18.40 ^a | | 34.50 ^b | | 65.70 ^c | | 82.80 ^d | |

Marital Status

| | | | | | | | | |
|-----------------|--------------------|--------|--------------------|--------|--------------------|--------|--------------------|--------|
| % No Partner | 8.20 ^a | | 21.40 ^a | | 36.40 ^b | | 48.30 ^b | |
| Income (1-4) | 3.22 ^a | (0.96) | 2.93 ^a | (1.19) | 2.20 ^b | (1.15) | 1.92 ^b | (1.23) |
| Age | 29.55 ^a | (5.96) | 28.38 ^a | (6.02) | 23.40 ^b | (4.43) | 22.91 ^b | (5.13) |
| Cumulative Risk | 1.08 ^a | (1.53) | 1.52 ^a | (1.75) | 2.91 ^b | (1.54) | 3.51 ^b | (1.31) |

Historical Factors

| | | | | | | | | |
|------------------|-------------------|--------|--------------------|--------|--------------------|--------|--------------------|--------|
| Lifetime PTSD sx | 9.78 ^a | (2.53) | 9.89 ^a | (2.48) | 10.35 ^a | (2.48) | 12.81 ^b | (2.09) |
| Lifetime | 1.80 ^a | (3.67) | 2.56 ^{ab} | (3.24) | 2.15 ^a | (3.15) | 3.76 ^b | (4.37) |
| Dissociation | | | | | | | | |
| Child Abuse | 0.55 ^a | (0.89) | 0.77 ^{ab} | (1.03) | 0.76 ^{ab} | (0.98) | 1.13 ^b | (1.17) |
| Adult Abuse | 0.47 ^a | (0.79) | 0.39 ^a | (0.74) | 0.52 ^a | (0.69) | 0.61 ^a | (0.85) |
| Non Abuse | 4.00 ^a | (2.05) | 3.60 ^a | (2.08) | 3.58 ^a | (2.15) | 4.10 ^a | (2.22) |
| Trauma | | | | | | | | |

Gestational Factors

| | | | | | | | | |
|-------------------------|--------------------|---------|---------------------|---------|---------------------|---------|--------------------|--------|
| % W1 MDD dx | 12.2 ^a | | 59.6 ^b | | 36.6 ^a | | 73.7 ^b | |
| W2 Labor Anxiety | 12.11 ^a | (2.30) | 14.15 ^b | (2.72) | 13.93 ^b | (3.60) | 15.26 ^b | (2.63) |
| W2 Social Support | 21.54 ^a | (3.23) | 20.24 ^{ab} | (3.11) | 19.81 ^{ab} | (5.22) | 18.44 ^b | (4.72) |
| W2 Health Care Alliance | 68.62 ^a | (10.26) | 62.52 ^a | (12.74) | 64.07 ^a | (13.09) | 64.62 ^a | (9.52) |

Note: A different superscript letter indicates groups are significantly different;

Income categories are as follows: (1) <15k, (2) 15k-25k, (3) 25k-50k, (4) >50k


Wave 1 <=28weeks

Cumulative Risk was the sum of risk status from the following: non-White, <HS education, no partner, <22years, <15k annually.

Note: A different superscript letter indicates groups are significantly different.

Table 3. Six Week Postpartum Adaptation by Trajectory Group (n=156)

| | <i>Low (N=33)</i> | | <i>Increase (N=44)</i> | | <i>Decrease (N=48)</i> | | <i>High (N=31)</i> | |
|--------------------|---------------------|-------------|------------------------|-------------|------------------------|-------------|--------------------|-------------|
| | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> | <i>M or %</i> | <i>(SD)</i> |
| % MDD dx | 15.20 ^a | | 52.30 ^b | | 41.70 ^b | | 48.40 ^b | |
| % Breastfeeding | 75.80 ^a | | 72.70 ^a | | 47.90 ^b | | 36.70 ^b | |
| Bonding Impairment | 19.61 ^{ab} | (6.34) | 23.86 ^b | (9.68) | 17.06 ^a | (7.07) | 17.87 ^a | (8.85) |



Author