



Published in final edited form as:

J Cell Biochem. 2008 March 1; 103(4): 1231–1245. doi:10.1002/jcb.21503.

Pur α as a Cellular Co-Factor of Rev/RRE-Mediated Expression of HIV-1 Intron-Containing mRNA

Rafal Kaminski¹, Nune Darbinian¹, Bassel E. Sawaya¹, Dorota Slonina¹, Shohreh Amini^{1,2}, Edward M. Johnson³, Jay Rappaport¹, Kamel Khalili^{1,*}, and Armine Darbinyan¹

¹Department of Neuroscience, Center for Neurovirology, Temple University School of Medicine, Philadelphia, Pennsylvania 19122

²Department of Biology, College of Science and Technology, Temple University, Philadelphia, Pennsylvania 19122

³Department of Microbiology and Molecular Cell Biology, Eastern Virginia Medical School, Norfolk, Virginia 23501

Abstract

To ensure successful replication, HIV-1 has developed a Rev-mediated RNA transport system that promotes the export of unspliced genomic RNA from nuclei to cytoplasm. This process requires the Rev responsive element (RRE) that is positioned in the viral transcript encoding Env protein, as well as in unspliced and singly spliced viral transcripts. We identified Pur α , a single-stranded nucleic acid binding protein as a cellular partner for Rev that augments the appearance of unspliced viral RNAs in the cytoplasm. A decrease in the level of Pur α expression by siRNA diminishes the level of Rev-dependent expression of viral RNA. Through its nucleic acid binding domain, Pur α exhibits the ability to interact with the multimerization and RBD domains of Rev. Similar to Rev, Pur α associates with RRE and in the presence of Rev forms a complex with slower electrophoretic mobility than those from Rev:RRE and Pur α :RRE. The interaction of Pur α with RRE occurs in the cytoplasm where enhanced association of Rev with RRE is observed. Our data indicate that the partnership of Pur α with Rev is beneficial for Rev-mediated expression of the HIV-1 genome.

Keywords

Pur α ; HIV-1; Rev

During the course of HIV-1 infection, three classes of viral RNAs are produced. These include 9 kb unspliced RNAs which are packaged into progeny virions as genomic RNA that can also serve as a transcript for protein products of gag/pol genes [Schwartz et al., 1990; Purcell and Martin, 1993]. The second class is singly spliced mRNAs of approximately 4 kb that encode the Vif, Vpr, and Vpu/Env proteins. Finally, the third group is doubly spliced 2 kb transcripts that are responsible for the production of Tat, Rev, and Nef [Purcell and Martin, 1993]. To facilitate transport of unspliced or partially spliced pre-mRNA into nuclei, HIV-1 has developed an efficient mechanism that is orchestrated by the viral protein Rev and its association with a 351 nucleotide RNA sequence named Rev Responsive Element (RRE) that spans the Env gene and is present in unspliced and singly spliced RNAs [Malim et al., 1989; Zapp and Green, 1989; Pollard and Malim, 1998]. The association of Rev and RRE in the

*Correspondence to: Kamel Khalili, 1900 North 12th Street, 015-96, Room 203, Philadelphia, PA 19122. E-mail: kamel.khalili@temple.edu

nucleus promotes engagement of several other cellular proteins including CRM1/Exportin-1 and RanGTP that eventually leads to the exit of Rev-associated RNA cages from the nucleus [Fornerod et al., 1997; Henderson and Per-cipalle, 1997; Neville et al., 1997; Stade et al., 1997; Nakielny and Dreyfuss, 1999; Kjems and Askjaer, 2000; Daelemans et al., 2005].

The cellular protein, Pura α , which has a strong affinity for associating with single-stranded nucleic acid containing the GGC(A)GGA(C) sequence, exhibits diverse biological activities on RNA transcription, cell cycle, neuronal cell differentiation, and DNA replication [Gallia et al., 2000; Khalili et al., 2003; Zhang et al., 2005; Johnson et al., 2006; Knapp et al., 2006; Shimotai et al., 2006]. Earlier studies revealed that some of the activity of Pura α is regulated upon its association with small RNA molecules [Hereault et al., 1995; Tretiakova et al., 1998; Gupta et al., 2003]. The interaction of Pura α with the mRNA sequence from exon 3 of the mouse VSM α -actin which is placed in the 5' untranslated region of a reporter mRNA suppresses its translation [Kelm et al., 1999].

Recent studies indicate that Pura α participates in dendritic transport of mRNAs and associates with ribosomes [Ohashi et al., 2000, 2002; Li et al., 2001; Johnson et al., 2006]. Pura α was isolated from RNA containing granules in dendrites together with a group of RNA binding proteins including hnRNP-U, Pur β , PSF, DDX1, DDX3, SYNCRIP, TLS, NonO, HSPC117, ALY, CGI-99, Staufen (Tang et al., 2001) three FMRPs, EF-1 α , mRNAs for CaMKII α and Arc. These RNP complexes were associated with kinesin [Kanai et al., 2004]. Pura α was also identified in mRNA/Protein complexes containing mStaufen, Fragile X mental retardation protein (FMRP) and myosin Va and associated with rough endoplasmic reticulum [Ohashi et al., 2002]. Unique features and functions of Pura α raise the possibility that Pura α may be involved in post- transcriptional regulation of HIV-1 gene expression.

Here we provide evidence that Pura α potentiates HIV-1 gene expression from unspliced RNAs and augments Rev/RRE-mediated stimulation of gene expression from viral mRNAs containing intron, probably facilitating their translocation, and targeting them to sites of translation or virion assembly.

MATERIALS AND METHODS

Cell Culture and Transfection

U-87MG (ATCC HTB14), a human glioblastoma cell line, was grown in Dulbecco's modified Eagle's medium supplemented with 10% heat-inactivated fetal bovine serum. Transfection of U-87MG cells was performed using Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA) or the calcium phosphate precipitation method [Graham and van der Eb, 1973]. Leptomycin B (LMB, Sigma-Aldrich, St. Louis, MO) treatment was performed 24 h after transfection at 2 nM concentration for 16 h.

Plasmids

pIIIAR was previously described [Rosen et al., 1988]. pcRev was a generous gift from Dr. Roger Pomerantz (Tibotec, Inc., Yardley, PA). The standard PCR cloning technique was used to construct the following expression plasmids, with the templates, primers, and cloning backbone vectors shown in parentheses (restriction enzyme sites underlined): pcDNA6/*myc*-His-A-Rev 1-116 and deletion mutants, pGEX-5X-1-Rev 1-116 and deletion mutants (pcRev; forward for Rev 1-116, Rev 1-83, Rev 1-67, Rev 1-50, Rev 1-33, Rev 1-18: 5'-TAGGCAGAATTCATGGCAGGA AGA-3'; forward for Rev 86-116 5'-GCAGAATTCGCCATGGAG GAT TGTGGA-3', forward for Rev 68-116 5'-TCAGAATTCGCCATGGAGCCT GTG CCT-3'; reverse for Rev 1-116, Rev 86-116, Rev 68-116: 5'-CTAACTCGAGTTCTTT AGCTCC-3'; reverse for Rev 1-83: 5'-

CTAACTCGAGAAGAGTAAGTCT-3'; reverse for Rev 1-67: 5'-CTAACTCGAGAGATCGTCCCAG-3'; reverse for Rev 1-50: 5'-CTAACTCGAGTCTCTGTCTCTC-3'; reverse for Rev 1-33: 5'-CTAACTCGAGCCCCTCGGGGTT-3'; reverse for Rev 1-18: 5'-CTAACA CTGAGTCTGA CTGCCTT-3'; pcDNA6/myc-His-A (Invitrogen) and pGEX-5X-1, pDs-Rev-Red1 (pcDNA6/myc-His-A-Rev, forward: 5'-TAGGCAGGATCC GAATTCATGGCAGGAAGA-3', reverse: 5'-CCTAACAGGATCCCTGAGTTCTTTAGCTCC-3', pDsRed1 (CLONTECH, Mountain View, CA); pBluescript II KS(+)-RRE 1-234 (pIIIAR, forward 5'-CGCCAAGCTTGAATAGGAGCTTTGTTCC-3', reverse 5'-CTAGGATCCAGGAGCTGTTGATCCTTTAGG-3', pBlue-script II KS(+)); pBluescript II KS(+)-RRE 34-110 (pIIIAR, 5'-TCACAAGCTTAGGAAGCACTAT GGGCGC-3', reverse 5'-CTAGGATCCGCTGCTGCACTATACCAG-3', pBluescript II KS(+)).

Construction of pcDNA3.1-SD4*-luciferase-RRE-SA7 and pcDNA3.1-SD4*-luciferase-SA7 was performed in few steps: First, pIIIAR was digested with *Bgl*II and *Xho*I and DNA fragment (1,276 bp) that contains the RRE (full-length) and splice acceptor 7 (SA7) sequences (corresponding to 7,650-8,926 nt of HIV-1 NCBI X01762) was removed and subcloned into pcDNA3.1 vector (Invitrogen) using *Bam*HI and *Xho*I enzyme recognition sites producing pcDNA3.1-RRE-SA7 vector. Next, DNA fragment (562 bp), which contains sequences of the first intron of Rev and splicing donor 4 (SD4) (corresponding to 5,945-6,380 nt of HIV-1 NCBI X01762) was obtained by digestion of pIIIAR with *Sal*I and *Kpn*I restriction enzymes followed with blunting of overhangs. This fragment was subcloned into pGL3-Basic vector (Promega, Madison, WI), digested with *Hind*III and blunted, generating pGL3AR vector. Site-directed mutagenesis was then performed to mutate ATG of Rev present in pIIIAR and pGL3AR, using QuikChange II XL Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA), pGL3AR vector as a template and following primers: forward 5'-CTTAGGCATCTCCTTAAGCAGGAAGAAGCGGAGAC-3', reverse 5'-GTCTCCGCTTCTTCTGCTTAAGGAGATGCCTAAG-3'. Site-directed mutagenesis resulted in pGL3AR* plasmid. Next, pGL3AR* was digested with *Xba*I, blunted then digested with *Nhe*I and fragment containing SD4* and luciferase gene was obtained (2,266 bp). Next, pcDNA3.1-RRE-SA7 was digested with *Kpn*I, blunted and then digested with *Nhe*I. The 2,266 bp fragment flanked by the *Nhe*I and blunt end was subcloned into digested pcDNA3.1-RRE-SA7, resulting in pcDNA3.1-SD4*-luciferase-RRE-SA7. Finally, for pcDNA3.1-SD4*-luciferase-SA7, pcDNA3.1-RRE-SA7 was digested with *Hind*III, blunted and then digested with *Nhe*I restriction enzyme removing RRE sequence and keeping 756 nt fragment from pIIIAR (corresponding to 8,170-8,926 nt of HIV-1 NCBI X01762), encompassing SA7. Subcloning of the 2,266 bp fragment, flanked by the *Nhe*I and blunt end, generated pcDNA3.1-SD4*-luciferase-SA7. PLEGFP-Purα [Darbinian et al., 2006], pEBV-Purα and deletion mutants and pGEX1λT-Purα and deletion mutants were previously described [Johnson et al., 1995; Gallia et al., 1998]. All recombinant plasmids were verified by sequencing.

RNAi and siRNA Transfections

Purα-specific siRNA, ACAAGTACGGCGTGTATTAT, was derived from sequences corresponding to 772-790 of Purα mRNA (Dharamcon, Lafayette, CO). U-87MG cells were transfected with 50 nM Purα/siRNA using Oligofectamine kit (Invitrogen).

Luciferase and CAT Assay

For luciferase assay, cells were harvested at designated time points, and protein extracts (20 μg) were used to examine the level of luciferase activity with the dual-luciferase reporter assay system (Promega, Madison, WI). The pRLTK plasmid was used as an internal control for transfection efficiency. Chloramphenicol acetyltransferase (CAT) activity was determined by the established method [Gorman et al., 1982].

Protein Extracts and Western Blot Analysis

For whole cell extract preparation, cells were lysed for 30 min on ice in LB1 (50 mM Hepes, pH 7.5/150 mM NaCl/1.5 mM MgCl₂/1 mM EGTA/10% glycerol/1% Triton X-100) buffer containing protease inhibitor cocktail (Sigma, P8340) and 0.2 mM Na-orthovanadate. Cell debris was removed by centrifugation at 14,000 rpm for 15 min at 4°C. Nuclear and cytoplasmic fractions were prepared using NE-PER nuclear and cytoplasmic extraction reagents (Pierce Biotechnology, Rockford, IL #78833). Western blot analysis was performed as described [Kaniowska et al., 2006].

Co-immunoprecipitation

U-87MG cells were transfected with pcDNA3.1-SD4*-luciferase-RRE-SA7 along with Pura and/or Rev expressing plasmids. Nuclear and cytoplasmic extracts were prepared and 100 µg of nuclear or 300 µg of cytoplasmic lysates were incubated with rabbit polyclonal antibody against Pura or normal rabbit serum as a negative control for immunoprecipitation in 500 µl HNTG buffer overnight at 4°C. Immunocomplexes were precipitated by the addition of protein A-Sepharose beads, washed four times with rocking at 4°C in 1 ml of HNTG buffer, and resolved by SDS-PAGE followed by Western blotting using an anti-myc antibody for detection of Rev.

GST Fusion Proteins and GST Pull-Down Assay

For preparation of bacterially produced GST fusion proteins, we followed the methods previously described. Overnight cultures (100 ml) of *Escherichia coli* DH5α, were transformed with pGEX-5X1-Rev, pGEX1λT-Pura [Gallia et al., 1998]. GST pull-down assay was performed according to the procedures described previously [Kaniowska et al., 2006].

RNA Preparation and Northern Blot Analysis

Total RNA was isolated using RNeasy kit (Qiagen, Valencia, CA, #74104) according to the manufacturer's directions. Preparation of cytoplasmic RNA was based on the protocol provided with RNeasy kit and using RLN buffer containing 50 mM Tris-HCl, pH 8.0, 140 mM NaCl, 1.5 mM MgCl₂, 0.5% (v/v) Nonidet P-40 (1.06 g/ml), 1,000 U/ml RNase inhibitor, 1 mM DTT. Ten micrograms of RNA were fractionated on 1.2% agarose, 0.4% formaldehyde, 1 × morpholinepropanesulfonic acid (Mops) gel and transferred to a Hybond-N nylon membrane (Amersham, Piscataway, NJ). For the detection of CAT RNAs, the membranes were hybridized with [³²P]-labeled CAT DNA probe obtained by digestion of pIIIAR plasmid with *Bgl*III and *Nco*I (850 bp). For the detection of RRE containing RNAs, the filters were probed with a PCR-amplified and [³²P]-labeled DNA fragment consisting of nucleotides 1-234 of the RRE. Radiolabeled DNA probes were prepared with Random Primed DNA Labeling Kit (Roche Molecular Biochemicals, Indianapolis, IN), followed by removal of unincorporated radionucleotides with MicroSpinTMG-50 columns (Amersham). Relative levels of mRNA were determined by using the housekeeping gene GAPDH as an internal standard.

RNA Immunoprecipitation and RT-PCR

Nuclear and cytoplasmic protein extracts from cells transfected with pcDNA3.1-SD4*-luciferase-RRE-SA7 vector along with Pura and/or Rev expressing plasmids were prepared and precleared with protein A sepharose beads by 1 h incubation at 4°C. The resultant lysates were incubated with protein A sepharose beads and anti-T7 antibody against Pura. In control samples, normal mouse serum was added instead of anti-T7 antibody. For RT control Rev expressing extracts were immunoprecipitated with anti-myc antibody. After overnight incubation at 4°C, the beads were washed four times (3 min each) with HNTG buffer (20 mM HEPES, pH 7.5/150 mM NaCl/0.1% Triton X-100/10% Glycerol, protease inhibitor cocktail (Sigma, P8340) and 0.2 mM Na-orthovanadate) and resuspended in 150 µl of buffer 2 (50 mM

Tris-HCl, pH 7.0/5 mM EDTA/1% SDS/10 mM DTT). Then RNA was extracted using phenol TE buffer pH 4.5 saturated/chloroform/isoamyl (25/24/1) and chloroform/isoamyl (24/1). Extracted RNA was precipitated by incubation with 7.5 M ammonium acetate/ethanol (0.5/2.5, V/V) and GlycoBlue (Ambion, Austin, TX) for 30 min at -70°C followed with washing in 70% ethanol, drying and rehydration in 20 µl of nuclease free water. Samples were treated with DNase1 in the presence of RNase inhibitor (Roche) for 1 h at 37°C followed with phenol/chloroform extraction and precipitation as described above. RT reaction was performed using random primers (p(dN)₆; Roche) and M-MuLV RT enzyme. For PCR, RRE specific primers and FailSafe™ PCR reagents (Epicentre, Madison, WI) were used. DNA products generated as a result of RT-PCR were analyzed by 2% agarose gel electrophoresis.

Fluorescent Technique

U-87MG cells (1×10^5) were transfected with 5 µg of GFP-Purα or Rev-Red plasmids, alone or in combination, then seeded in poly-L-lysinecoated glass chamber slides, and after 16 h incubation, cells were fixed in 4% paraformaldehyde in 1 × PBS. Cells were then washed in PBS, and proteins were visualized for green or red fluorescence. Fluorescent images were captured using an inverted fluorescent Nikon microscope with deconvolution software (Slide-Book 4.0.1.34; Intelligent Imaging, Denver, CO).

RNA Electrophoretic Mobility Shift Assay

The RRE-containing RNA probe was in vitro transcribed using pBluescript IKS(+) RRE 1-234 and pBluescript IKS(+) RRE 34-110 plasmids as DNA template. In vitro transcription reaction was performed by incubating the above described template DNAs, T3 RNA polymerase, transcription optimized 5 × buffer, DTT 100 mM, recombinant RNasin ribonuclease inhibitor, rATP, rGTP, rCTP (2.5 mM each) and [α -³²P]-UTP for 1 h at 37-40°C. For RNA-protein interaction studies 0.2 µM of total recombinant proteins GST, GST-Rev, GST-Purα were incubated with 100,000 cpm of [α -³²P]-UTP RNA probe for 15 min at room temperature in 20 µl binding buffer containing 12 mM HEPES (pH 7.9), 4 mM Tris-HCl (pH 7.5), 60 mM KCl, 5 mM MgCl₂, 0.8 mM dithiothreitol (DTT), 0.5 µg of poly[dI-dC] as a nonspecific competitor, 10% glycerol and RNase inhibitor. Reaction products were analyzed on a 1% agarose gel followed by autoradiography.

Antibodies

Anti-α-tubulin, clone B512 was obtained from Sigma-Aldrich. Anti-myc antibody was purchased from Invitrogen and anti-T7 tag antibody was from Novagen (Madison, WI). Rabbit polyconal anti-Purα antibody was raised against full-length Purα, cleaved from GST-Purα and injected into rabbits (Lampire Biological Laboratories, Pipersville, PA).

RESULTS

To investigate the effect of Purα on Rev activity, we constructed and utilized a luciferase-based reporter plasmid where the luciferase gene was placed within the HIV-1 intronic sequence flanked by splicing donor 4 and splicing acceptor 7 (Fig. 1A top). The translation start site of Rev, which is located 76 nucleotides upstream from the splice donor, was disrupted and the RRE was placed at the 3' end of the luciferase gene within the intron. The control plasmid had no RRE sequence, yet encompassed all the other features of the reporter plasmid (Fig. 1A, bottom). As shown in Figure 1B, Purα enhanced the ability of Rev to stimulate luciferase activity directed by the RRE-containing plasmid and exhibited no stimulatory effect on the expression from the control RRE negative plasmid. Leptomycin, which targets the CRM1/Exportin-1 and inhibits Rev nuclear export [Nishi et al., 1994; Wolff et al., 1997] suppressed cooperativity of Rev and Purα, suggesting that Purα is important for Rev:RRE-mediated stimulation of HIV-1 expression. A decrease of approximately 40% in the level of endogenous

Pur α by siRNA (Fig. 1C) caused comparable levels of decline in the level of Rev-mediated activation of luciferase expression (Fig. 1D).

To demonstrate the effect of Pur α on Rev-mediated RNA accumulation in the cytoplasm, total and cytoplasmic RNAs from cells transfected with pIIIAR [Rosen et al., 1988] along with pEBV-Pur α and pCMV-Rev were analyzed by Northern blot using DNA probes derived from RRE or CAT sequences. The RRE probe detects 4.4 kb unspliced RNA whereas the CAT DNA probe identifies both 2.37 kb spliced and 4.4 kb unspliced RNA species. As seen in Figure 2A, a significant increase in the level of the unspliced form of cytoplasmic RNA was observed upon Rev expression. Interestingly, Pur α also increased the level of unspliced RNA in the cytoplasm. Co-expression of both proteins resulted in an increase in the amount of unspliced RNA in the cytoplasm, more than that observed with Rev alone. Note that unspliced RNA in total RNA extract was also maximal when Rev and Pur α were co-expressed. A similar pattern for 4.4 kb RNA was observed when DNA sequence from the CAT gene was used as a probe (Fig. 2B). The integrity of RNA preparation in various samples is shown in panel C. The level of hGAPDH in each sample served as an internal control (Fig. 2, panel D).

Earlier studies have shown that Rev predominantly accumulates in the nucleolus with a fraction shuttling between the nucleus and cytoplasm [Cullen et al., 1988; Cochrane et al., 1990; Wolff et al., 1995]. To investigate the subcellular localization of Pur α and Rev in our cell culture system, we constructed plasmids expressing Pur α and Rev in fusion with green and red fluorescent proteins, respectively. Activity of Rev-Red fusion protein was verified by transfection assay using a reporter plasmid that is responsive to Rev activity. As seen in Figure 3A, similar to previous observations, Rev fusion protein had a stimulatory effect on reporter plasmid containing RRE, and this effect was augmented by expression of GFP-Pur α . Co-transfection of cells with GFP Pur α and/or Rev-Red expressing plasmids allowed visualization of the corresponding proteins in the cells. As shown in Figure 3B, Rev was detected within nucleoli with some accumulation in the perinuclear region (Fig. 3B). Pur α was predominantly localized in the perinuclear region and cytoplasm of cells (Fig. 3C), although, a trace of Pur α was also detected in the nucleoli. In cells co-transfected with both plasmids, some levels of Pur α and Rev were found in the perinuclear compartment (Fig. 3D).

Co-localization of Pur α and Rev prompted us to investigate the possible interaction of these two proteins. Results from the GST-based protein binding assay revealed the ability of Pur α to interact with full-length Rev protein and highlighted the importance of the region of Rev that spans aa 18-50, which overlaps nuclear localization signal (NLS) and RNA binding domain (RBD) and multimerization domain (MD), in this interaction. Figure 4A (top panel) illustrates results from GST pull-down assay and the various mutants of Rev that were utilized in this experiment. The integrity of GST-Rev and its mutant variants were examined by SDS-PAGE (Fig. 4A, middle panel). Figure 4A (bottom panel) illustrates the linear organization of full-length Rev, its deletion mutants, and their binding ability to Pur α . In a reciprocal study, we utilized a similar approach and identified a region of Pur α located between amino acids 73-123 as a minimum domain that is recognized by Rev (Fig. 4B, top panel). Note that Rev, due to its modification, appears as a double band. Treatment with alkaline phosphatase or lambda protein phosphatase results in the disappearance of the top band (unpublished observations). The integrity of GST-Pur α and its various mutants is shown in Figure 4B (middle panel). Figure 4B (bottom panel) also illustrates the structural organization of the full-length Pur α , its mutant variants, and summarizes their binding abilities to Rev. The interaction of Pur α and Rev was also detected by immunoprecipitation/Western blot analysis of the protein extracts expressing both proteins (Fig. 4C,D). Nuclear (100 μ g) or cytoplasmic (300 μ g) fractions were prepared from U-87MG cells transfected with pcDNA3.1-SD4*-luciferase-RRE-SA7 along with Pur α and/or Rev expressing plasmids and the presence of proteins was verified by Western blot analysis using antibodies against Rev (Fig. 4C, top panel) and Pur α (Fig. 4C, middle panel).

Rabbit polyclonal antibody against Pur α detects endogenous and overexpressed Pur α (middle panel). Fractionation and equal loading was controlled by α -Tubulin detection (Fig. 4C, bottom panel). We immunoprecipitated protein complexes bound to Pur α using anti-Pur α antibody (Fig. 4D, lanes 1-10) in the presence of protein A Sepharose. After washing, protein complexes were separated by SDS-PAGE and analyzed by Western blot using anti-Myc antibody for detection of Rev. As shown in panel D, Rev was associated with endogenous Pur α (lane 2) and overexpressed Pur α (lane 4) in cytoplasmic fractions. Although, a faint band that points to the interaction of overexpressed Pur α and Rev was also detected in nuclear fraction (lane 9).

To investigate the relevance of Pur α :Rev interaction to the observed functional cooperativity of these two proteins, we examined the effect of Rev on the RRE-containing reporter in the absence and presence of full-length and mutant Pur α with no binding ability to Rev. As shown in Figure 5A, the activity of Rev was enhanced in the presence of full-length Pur α , but not mutant Pur α encompassing the C-terminus (216-332) of this protein. Of note, full-length Pur α , but not its mutant 216-322, showed a modest increase in the level of luciferase gene expression (Fig. 5B). While full-length Pur α was able to cooperate with full-length Rev (aa 1-116), it exhibited no cooperativity with the Rev mutant 1-33 which showed reduced binding activity to Rev (data not shown). These observations provided evidence that physical association between Pur α and Rev may be important, at least in part, for their coordinated effect on the expression of the viral genome.

As Pur α has a great affinity for binding to single-stranded RNAs containing GGCG sequence, a feature that is found in HIV-1 RRE, in the next series of experiments we evaluated the ability of Pur α to bind to RRE and its impact on the interaction of Rev with RRE. We performed RNA electrophoretic gel mobility shift assays using RRE RNA probe and recombinant mutant GST-Rev, GST-Pur α , and GST proteins. Two regions of the RRE spanning nucleotides 1-234 and 34-110 (numbering according to Malim et al., 1989) were synthesized *in vitro* and used as a probe. As shown in Figure 6A, incubation of both labeled RNAs with GST-Pur α or GST-Rev resulted in the formation of complexes with slower electrophoretic mobilities. Co-incubation of the probes with both Rev and Pur α resulted in the formation of a higher molecular weight complex than those seen by either protein alone (Fig. 6A, compare lane 5 to lanes 3 and 4, and lane 10 with lanes 8 and 9). The specificity of *in vitro* RNA binding was tested in competition assay with excess unlabeled probes. The addition of unlabeled RRE-containing RNAs 1-234 and 34-110 to the reaction mixture abrogated binding of both proteins with the labeled full-length RNA probe (data not shown).

In attempts to assess the ability of Pur α to associate with pre-existing Rev:RRE complex, RNA probe was first incubated with Rev for 10 min prior to the addition of Pur α to the binding reaction. As shown in Figure 6 (panel B), the addition of Pur α resulted in the formation of higher molecular weight complexes with RRE. Similar results were obtained in reciprocal experiments where Rev was added to the reaction after the incubation of RRE probe with Pur α . These observations suggest that Pur α binds to RNA sequences that are not bound to Rev and that Pur α interacts with RRE RNA that is associated with Rev.

Next, we performed RNA immunoprecipitation assay to determine whether the association of Pur α and Rev with RRE RNA occurs in the cytoplasm or in nuclei of the cells. The human astrocytic cell lines, U-87MG, was co-transfected with pcDNA3.1SD4**lucRRE-SA7* vector and plasmids expressing T7-tagged Pur α and myc-tagged Rev, either alone or in combination. Cytoplasmic and nuclear fractions were prepared, and expression of Rev and Pur α was determined by Western blot analysis (Fig. 7A and B, respectively). The quality of each fraction was tested by measuring the presence of α -tubulin (Fig. 7C). Next, RNA:protein complexes were immunoprecipitated using antibody that recognizes Pur α (anti-T7) (Fig. 8A,B, lanes 1-5), Rev (anti-myc) (Fig. 8A,B, top, lane 7) or control normal mouse serum (nms). The

immunocomplexes were extensively washed and their associated RNAs were prepared and subjected to reverse transcription (RT) reaction using random primers (Fig. 8A). RRE was subsequently amplified by polymerase chain reaction (PCR) using primers specific for RRE. The RRE fragment which was directly amplified from pcDNA3.1SD4**lucRRE-SA7* plasmid served as a positive control. Reactions lacking reverse transcriptase (-RT) served as negative controls in this study (Fig. 8B). A high intensity band, corresponding to RRE fragment was detected in cytoplasmic extracts from cells expressing both Pur α and Rev proteins (Fig. 8A, lane 5, cytoplasmic fraction). In cells expressing only Pur α , a weak signal pointing to the association of Pur α with RRE was detected in nuclear fractions. RNA input was verified by extraction of RNA directly from nuclear and cytoplasmic fractions (without prior immunoprecipitation) following RT-PCR (Fig. 8C). Rev:Pur α protein complexes were also detected in cytoplasmic fraction using IP-Western (data not shown). In addition, we repeated RNA immunoprecipitation experiments using anti-Pur α antibody instead of anti-T7 antibody and detected association of endogenous Pur α with RRE RNA and Rev in the cytoplasm (data not shown). These observations point to the *in vivo* association of Pur α with RRE in the presence of Rev. Moreover, Pur α :RRE and Pur α :RRE:Rev complexes are present in the cytoplasmic compartment of cells.

DISCUSSION

Replication of HIV-1 is a highly regulated process and is dependent on cooperation between viral and host proteins. In an earlier study, we identified Pur α as a potential partner for HIV-1 Tat that may facilitate Tat activation of the responsive promoters [Krachmarov et al., 1996; Chepenik et al., 1998; Gallia et al., 1999]. In this report we show that Pur α enhances the action of Rev protein, the main regulator of export of unspliced viral mRNAs, and that depletion of endogenous Pur α by siRNA diminishes Rev:RRE-mediated reporter activation. Similarly, silencing of Pur α by siRNA during the course of HIV-1 infection of human primary culture of microglial cells caused nearly 25% suppression in viral replication pointing to the importance of Pur α in HIV-1 gene expression and replication (data not shown).

Using reporter constructs containing luciferase gene sequence placed within intron alone or in combination with RRE, we demonstrated that Pur α can function as a cofactor for Rev in activating gene expression from intronic sequences. Co-localization of Pur α and Rev was observed within cytoplasm of cells expressing both proteins. Rev/RRE/Pur α complexes were detected *in vivo* in the cytoplasmic compartment of cells also in RNA immunoprecipitation experiments. Thus Pur α , which can also directly bind to RRE may act in post-transcriptional processing steps involving Rev function. In support of this notion, recent studies demonstrated that Pur α is involved in dendritic transport of a subset of mRNAs and translation. Pur α has been identified in mRNA/protein complexes containing mStaufen, FMRP, proteins that have been implicated to play a key role in targeting mRNAs to polyribosomes [Feng et al., 1997; Marion et al., 1999; Brown et al., 2001; Krichevsky and Kosik, 2001; Tang et al., 2001; Ohashi et al., 2002]. In addition, Staufen 1, which was found in HIV-1 Gag RNP complex, was implicated in regulating retroviral genomic encapsidation [Mouland et al., 2000; Chatel-Chaix et al., 2004]. Immunoprecipitation of these proteins with an anti-Pur α antibody was abolished by RNase A treatment, suggesting an RNA dependence for Pur α in the assembly of mRNPs [Ohashi et al., 2002]. These observations led to the hypothesis that Pur α via binding to rRNA may mediate the association of mRNPs with polyribosomes. Interestingly, our earlier studies have revealed that Pur α binds to RNAs that are homologous to 18S ribosomal RNA and to 7SL RNA [Tretiakova et al., 1998; Gallia et al., 1999, 2001]. Pur α has also been isolated from kinesin-associated granules in dendrites together with a group of RNA binding proteins [Kanai et al., 2004], including DDX1, DDX3, staufen, PSF, EF-1 α , which are involved in RNA splicing, nuclear RNA export and RNA translation and have been linked to retroviral replication [reviewed in Cochrane et al., 2006]. It was shown that Pur α recognizes dendrite-

targeting RNA motifs rich in G and U residues in the 5' portion of BC1 RNA and links BC1 RNA to microtubules [Ohashi et al., 2000] and associates with ribosomes in neuronal cytoplasm [Li et al., 2001]. Non-coding BC1 RNA acts as a molecular scaffold for the formation of BC1 ribonucleoprotein particles (BC1 RNPs). These particles have been shown to be important for the dendritic delivery of mRNAs and factors regulating translation in dendrites. All of these observations are consistent with the notion that Pur α via interaction with Rev and RRE may be involved in directing viral unspliced RNA association with RNP complexes to sites of translation.

Another interesting, although preliminary, observation comes from our experiments with RRE-negative construct showing that the level of cytoplasmic unspliced RNA in cells with expressing Pur α and Rev is higher compared to that from cells with reporter construct alone, yet the level of protein remains comparable. Thus, one may speculate that in the presence of Rev, Pur α and Rev may cooperate to enhance gene expression at the post-transcriptional level from intronic sequences, again supporting a role for Pur α in post-transcription, that is, translation level. Our preliminary data also demonstrate that Pur α can increase, albeit modestly, the level of cytoplasmic unspliced reporter RNAs (with or without RRE) and consequently, expression of the reporter protein in LMB insensitive manner. This observation along with the ability of Pur α to associate with the RRE in the absence of Rev suggests that Pur α may also be involved in CRM-1-independent nuclear export pathway that includes RNA helicase A (RHA), TAP, and Sam68 [Braun et al., 2001; Reddy et al., 2000].

The HIV-1 RRE has double helical, highly looped secondary structure with GG rich motifs. All known RNA molecules interacting with Pur α in vivo contain such structures [Johnson et al., 2006]. It is possible that Pur α through its interaction with Rev and RRE can act in remodeling of RNA cargo assisting DDX3, human RNA helicase, a member of DEAD-box family, which binds CRM1 and restructures cargo HIV-1 RNA to permit translocation [Yedavalli et al., 2004]. This is an interesting notion in light of earlier studies showing local helix-unwinding activity of Pur α [Darbinian et al., 2001; Wortman et al., 2005].

Altogether, our results demonstrate that Pur α participates in cytoplasmic processing of unspliced mRNAs containing intronic viral RNA sequences. Further, we demonstrate that Pur α cooperates with Rev to stimulate expression of HIV-1 genes from unspliced mRNAs. Pur α acts as a cellular factor, which by interaction with Rev alleviates molecular barriers in Rev function [Neumann et al., 1995; Brack-Werner 1999; Gorry et al., 2003]. Moreover, our results show that Pur α is able to promote translocation and expression of unspliced mRNA in the absence of Rev and in LMB insensitive manner.

These observations together with the demonstration of Pur α association with RNA transporting granules imply a new role for Pur α in stabilization and transport of unspliced viral RNA to sites of translation or viral assembly for encapsidation and virion formation. Pur α has been implicated in the transport of Map2 mRNA to sites of translation in neuronal dendrites [Johnson et al., 2006]. In light of earlier reports on the export of intron-containing mRNAs from nucleus and pre-mRNA splicing in neuronal dendrites [Brown et al., 2001; Rodrigues et al., 2001; Jin et al., 2003; Glanzer et al., 2005], it will be of particular interest to determine if Pur α has any role in the intracellular transport, stability and translation of cellular pre-mRNAs.

ACKNOWLEDGMENTS

We wish to thank past and present members of the Department of Neuroscience and Center for Neurovirology for their continued support, insightful discussion, and sharing of reagents and ideas. We also wish to thank C. Schriver for editorial assistance. This work was made possible by grants awarded by NIH to KK and SA.

Grant sponsor: NIH.

REFERENCES

- Brack-Werner R. Astrocytes: HIV cellular reservoirs and important participants in neuropathogenesis. *AIDS* 1999;13:1–22. [PubMed: 10207540]
- Braun IC, Herold A, Rode M, Conti E, Izaurralde E. Overexpression of TAP/p15 heterodimers bypasses nuclear retention and stimulates nuclear mRNA export. *J Biol Chem* 2001;276:20536–20543. [PubMed: 11259411]
- Brown V, Jin P, Ceman S, Darnell JC, O'Donnell WT, Tenenbaum SA, Jin X, Feng Y, Wilkinson KD, Keene JD, Darnell RB, Warren ST. Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell* 2001;107:477–487. [PubMed: 11719188]
- Chatel-Chaix L, Clément JF, Martel C, Bériault V, Gatignol A, DesGroseillers L, Mouland AJ. Staufen is part of the HIV-1 Gag Ribonucleoprotein complex and is involved in the generation of infectious viral particles. *Mol Cell Biol* 2004;24:2637–2648. [PubMed: 15024055]
- Chepenik L, Tretiakova AP, Krachmarov CP, Johnson EM, Khalili K. The single-stranded DNA binding protein, Pur-alpha, binds HIV-1 TAR RNA and activates HIV-1 transcription. *Gene* 1998;210:37–44. [PubMed: 9524214]
- Cochrane AW, Perkins A, Rosen CA. Identification of sequences important in the nucleolar localization of human immunodeficiency virus Rev: Relevance of nucleolar localization to function. *J Virol* 1990;64:881–885. [PubMed: 2404140]
- Cochrane AW, McNally MT, Mouland AJ. The retrovirus RNA trafficking granule: From birth to maturity. *Retrovirology* 2006;3:18. [PubMed: 16545126]
- Cullen BR, Hauber J, Campbell K, Sodroski JG, Haseltine WA, Rosen CA. Subcellular localization of the human immunodeficiency virus trans-acting art gene product. *J Virol* 1988;62:2498–2501. [PubMed: 2836628]
- Daelemans D, Costes SV, Lockett S, Pavlakis GN. Kinetic and molecular analysis of nuclear export factor CRM1 association with its cargo in vivo. *Mol Cell Biol* 2005;25:728–739. [PubMed: 15632073]
- Darbinian N, Gallia GL, Khalili K. Helix-destabilizing properties of the human single-stranded DNA- and RNA-binding protein Puralpha. *J Cell Biochem* 2001;80:589–595. [PubMed: 11169743]
- Darbinian N, White MK, Khalili K. Regulation of the Pur-alpha promoter by E2F-1. *J Cell Biochem* 2006;99:1052–1063. [PubMed: 16741925]
- Feng Y, Absher D, Eberhart DE, Brown V, Malter HE, Warren ST. FMRP associates with polyribosomes as an mRNP, and the I304 N mutation of severe fragile X syndrome abolishes this association. *Mol Cell* 1997;1:109–118. [PubMed: 9659908]
- Fornierod M, Ohno M, Yoshida M, Mattaj IW. CRM1 is an export receptor for leucine-rich nuclear export signals. *Cell* 1997;90:1051–1060. [PubMed: 9323133]
- Gallia GL, Safak M, Khalili K. Interaction of the single-stranded DNA-binding protein Puralpha with the human polyomavirus JC virus early protein T-antigen. *J Biol Chem* 1998;273:32662–32669. [PubMed: 9830007]
- Gallia GL, Darbinian N, Tretiakova A, Ansari SA, Rappaport J, Brady J, Wortman B, Johnson EM, Khalili K. Association of HIV-1 Tat with the cellular protein, Puralpha, is mediated by RNA. *Proc Natl Acad Sci USA* 1999;96:11572–11577. [PubMed: 10500218]
- Gallia GL, Johnson EM, Khalili K. Puralpha: A multifunctional single-stranded DNA- and RNA-binding protein. *Nucleic Acids Res* 2000;28:3197–3205. [PubMed: 10954586]Review
- Gallia GL, Darbinian N, Jaffe N, Khalili K. Single-stranded nucleic acid-binding protein, Pur alpha, interacts with RNA homologous to 18S ribosomal RNA and inhibits translation in vitro. *J Cell Biochem* 2001;83:355–363. [PubMed: 11596104]
- Glanzer J, Miyashiro KY, Sul JY, Barrett L, Belt B, Haydon P, Eberwine J. RNA splicing capability of live neuronal dendrites. *Proc Natl Acad Sci USA* 2005;102:16859–16864. [PubMed: 16275927]
- Gorman CM, Moffat LF, Howard BH. Recombinant genomes which express chloramphenicol acetyltransferase in mammalian cells. *Mol Cell Biol* 1982;2:1044–1051. [PubMed: 6960240]
- Gorry PR, Ong C, Thorpe J, Bannwarth S, Thompson KA, Gatignol A, Wesselingh SL, Purcell DF. Astrocyte infection by HIV-1: Mechanisms of restricted virus replication, and role in the pathogenesis of HIV-1-associated dementia. *Curr HIV Res* 2003;1:463–473. [PubMed: 15049431]

- Graham FL, van der Eb AJ. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* 1973;52:456–467. [PubMed: 4705382]
- Gupta M, Sueblinvong V, Raman J, Jeevanandam V, Gupta MP. Single-stranded DNA-binding proteins PURalpha and PURbeta bind to a purine-rich negative regulatory element of the alpha-myosin heavy chain gene and control transcriptional and translational regulation of the gene expression. Implications in the repression of alpha-myosin heavy chain during heart failure. *J Biol Chem* 2003;278:44935–44948. [PubMed: 12933792]
- Henderson BR, Percipalle P. Interactions between HIV Rev and nuclear import and export factors: The Rev nuclear localisation signal mediates specific binding to human importin-beta. *J Mol Biol* 1997;274:693–707. [PubMed: 9405152]
- Hereault Y, Chatelain G, Brun G, Michel D. RNA-dependent DNA binding activity of the Pur factor, potentially involved in DNA replication and gene transcription. *Gene Expr* 1995;4:85–93. [PubMed: 7537563]
- Jin L, Guzik BW, Bor YC, Rekosh D, Hammarskjold ML. Tap and NXT promote translation of unspliced mRNA. *Genes Dev* 2003;17:3075–3086. [PubMed: 14701875]
- Johnson EM, Chen PL, Krachmarov CP, Barr SM, Kanovsky M, Ma Z-W, Lee WH. Association of human Pur alpha with the retinoblastoma protein, Rb, regulates binding to the single-stranded DNA Pur alpha recognition element. *J Biol Chem* 1995;270:24352–24360. [PubMed: 7592647]
- Johnson EM, Kinoshita Y, Weinreb DB, Wortman MJ, Simon R, Khalili K, Winckler B, Gordon J. Role of Puralpha in targeting mRNA to sites of translation in hippocampal neuronal dendrites. *J Neurosci Res* 2006;83:929–943. [PubMed: 16511857]
- Kanai Y, Dohmae N, Hirokawa N. Kinesin transports RNA: Isolation and characterization of an RNA-transporting granule. *Neuron* 2004;43:513–525. [PubMed: 15312650]
- Kaniowska D, Kaminski R, Amini S, Radhakrishnan S, Rappaport J, Johnson EM, Khalili K, Del Valle L, Darbinyan A. Cross-interaction between JC virus agnoprotein and human immunodeficiency virus type 1 (HIV-1) Tat modulates transcription of the HIV-1 long terminal repeat in glial cells. *J Virol* 2006;80:9288–9299. [PubMed: 16940540]
- Kelm RJ Jr, Elder PK, Getz MJ. The single-stranded DNA-binding proteins, Puralpha, Purbeta, and MSY1 specifically interact with an exon 3-derived mouse vascular smooth muscle alpha-actin messenger RNA sequence. *J Biol Chem* 1999;274:38268–38275. [PubMed: 10608902]
- Khalili K, Del Valle L, Muralidharan M, Gault WJ, Darbinian N, Otte J, Meier E, Johnson EM, Daniel DC, Kinoshita Y, Amini S, Gordon J. Puralpha is essential for postnatal brain development and developmentally coupled cellular proliferation as revealed by genetic inactivation in the mouse. *Mol Cell Biol* 2003;23:6857–6875. [PubMed: 12972605]
- Kjems J, Askjaer P. Rev protein and its cellular partners. *Adv Pharmacol* 2000;48:251–298. [PubMed: 10987094]
- Knapp AM, Ramsey JE, Wang SX, Godburn KE, Strauch AR, Kelm RJ Jr. Nucleoprotein interactions governing cell type-dependent repression of the mouse smooth muscle alpha-actin promoter by single-stranded DNA-binding proteins Pur alpha and Pur beta. *J Biol Chem* 2006;281:7907–7918. [PubMed: 16436378]
- Krachmarov CP, Chepenik LG, Barr-Vagell S, Khalili K, Johnson EM. Activation of the JC virus Tat-responsive transcriptional control element by association of the Tat protein of human immunodeficiency virus 1 with cellular protein Pur alpha. *Proc Natl Acad Sci USA* 1996;93:14112–14117. [PubMed: 8943069]
- Krichevsky AM, Kosik KS. Neuronal RNA granules: A link between RNA localization and stimulation-dependent translation. *Neuron* 2001;32:683–696. [PubMed: 11719208]
- Li Y, Koike K, Ohashi S, Funakoshi T, Tadano M, Kobayashi S, Anzai K, Shibata N, Kobayashi M. Pur alpha protein implicated in dendritic RNA transport interacts with ribosomes in neuronal cytoplasm. *Biol Pharm Bull* 2001;24:231–235. [PubMed: 11256476]
- Malim MH, Hauber J, Le SY, Maizel JV, Cullen BR. The HIV-1 rev trans-activator acts through a structured target sequence to activate nuclear export of unspliced viral mRNA. *Nature* 1989;338:254–257. [PubMed: 2784194]

- Marion RM, Fortes P, Beloso A, Dotti C, Ortin JA. The human sequence homologue of Staufen is an RNA-binding protein that is associated with polysomes and localizes to the rough endoplasmic reticulum. *Mol Cell Biol* 1999;19:2212–2219. [PubMed: 10022908]
- Mouland AJ, Mercier J, Luo M, Bernier L, DesGroseillers L, Cohen EA. The double-stranded RNA-binding protein Staufen is incorporated in human immunodeficiency virus type 1: Evidence for a role in genomic RNA encapsidation. *J Virol* 2000;74:5441–5451. [PubMed: 10823848]
- Nakielnny S, Dreyfuss G. Transport of proteins and RNAs in and out of the nucleus. *Cell* 1999;99:677–690. [PubMed: 10619422]
- Neumann M, Felber BK, Kleinschmidt A, Froese B, Erfle V, Pavlakis GN, Brack-Werner R. Restriction of human immunodeficiency virus type 1 production in a human astrocytoma cell line is associated with a cellular block in Rev function. *J Virol* 1995;69:2159–2167. [PubMed: 7884864]
- Neville M, Stutz F, Lee L, Davis LI, Rosbash M. The importin-beta family member Crm1p bridges the interaction between Rev and the nuclear pore complex during nuclear export. *Curr Biol* 1997;7:767–775. [PubMed: 9368759]
- Nishi K, Yoshida M, Fujiwara D, Nishikawa M, Horinouchi S, Beppu T. Leptomycin B targets a regulatory cascade of crm1, a fission yeast nuclear protein, involved in control of higher order chromosome structure and gene expression. *J Biol Chem* 1994;269:6320–6324. [PubMed: 8119981]
- Ohashi S, Kobayashi S, Omori A, Ohara S, Omae A, Muramatsu T, Li Y, Anzai K. The single-stranded DNA- and RNA-binding proteins pur alpha and pur beta link BC1 RNA to microtubules through binding to the dendrite-targeting RNA motifs. *J Neurochem* 2000;75:1781–1790. [PubMed: 11032866]
- Ohashi S, Koike K, Omori A, Ichinose S, Ohara S, Kobayashi S, Sato TA, Anzai K. Identification of mRNA/protein (mRNP) complexes containing Puralpha, mStaufen, fragile X protein, and myosin Va and their association with rough endoplasmic reticulum equipped with a kinesin motor. *J Biol Chem* 2002;277:37804–37810. [PubMed: 12147688]
- Pollard VW, Malim MH. The HIV-1 Rev protein. *Annu Rev Microbiol* 1998;52:491–532. [PubMed: 9891806]
- Purcell DF, Martin MA. Alternative splicing of human immunodeficiency virus type 1 mRNA modulates viral protein expression, replication, and infectivity. *J Virol* 1993;67:6365–6378. [PubMed: 8411338]
- Reddy TR, Tang H, Xu W, Wong-Staal F. Sam68, RNA helicase A and Tap cooperate in the post-transcriptional regulation of human immunodeficiency virus and type D retroviral mRNA. *Oncogene* 2000;19:3570–3575. [PubMed: 10951562]
- Rodrigues JP, Rode M, Gatfield D, Blencowe BJ, Carmo-Fonseca M, Izaurralde E. REF proteins mediate the export of spliced and unspliced mRNAs from the nucleus. *Proc Natl Acad Sci USA* 2001;98:1030–1035. [PubMed: 11158589]
- Rosen CA, Terwilliger E, Dayton A, Sodroski JG, Haseltine WA. Intragenic cis-acting art gene-responsive sequences of the human immunodeficiency virus. *Proc Natl Acad Sci USA* 1988;85:2071–2075. [PubMed: 2832844]
- Schwartz S, Felber BK, Benko DM, Fenyo EM, Pavlakis GN. Cloning and functional analysis of multiply spliced mRNA species of human immunodeficiency virus type 1. *J Virol* 1990;64:2519–2529. [PubMed: 2335812]
- Shimotai Y, Minami H, Saitoh Y, Onodera Y, Mishima Y, Kelm RJ Jr, Tsutsumi K. A binding site for Pur alpha and Pur beta is structurally unstable and is required for replication in vivo from the rat aldolase B origin. *Biochem Biophys Res Commun* 2006;340:517–525. [PubMed: 16376299]
- Stade K, Ford CS, Guthrie C, Weis K. Exportin 1 (Crm1p) is an essential nuclear export factor. *Cell* 1997;90:1041–1050. [PubMed: 9323132]
- Tang S, Meulemans D, Vazquez L, Colaco N, Schuman E. A role for a rat homolog of Staufen in the transport of RNA to neuronal dendrites. *Neuron* 2001;32:463–475. [PubMed: 11709157]
- Tretiakova A, Gallia GL, Shcherbik N, Jameson B, Johnson EM, Amini S, Khalili K. Association of Pur with RNAs homologous to 7SL determines its binding ability to the myelin basic protein promoter DNA sequence. *J Biol Chem* 1998;273:22241–22247. [PubMed: 9712838]
- Wolff B, Cohen G, Hauber J, Meshcheryakova D, Rabeck C. Nucleocytoplasmic transport of the Rev protein of human immunodeficiency virus type 1 is dependent on the activation domain of the protein. *Exp Cell Res* 1995;217:31–41. [PubMed: 7867718]

- Wolff B, Sanglier JJ, Wang Y. Leptomycin B is an inhibitor of nuclear export: Inhibition of nucleocytoplasmic translocation of the human immunodeficiency virus type 1 (HIV-1) Rev protein and Rev-dependent mRNA. *Chem Biol* 1997;4:139–147. [PubMed: 9190288]
- Wortman MJ, Johnson EM, Bergemann AD. Mechanism of DNA binding and localized strand separation by Pur alpha and comparison with Pur family member, Pur beta. *Biochim Biophys Acta* 2005;1743:64–78. [PubMed: 15777841]
- Yedavalli VS, Neuveut C, Chi YH, Kleiman L, Jeang KT. Requirement of DDX3 DEAD box RNA helicase for HIV-1 Rev-RRE export function. *Cell* 2004;119:381–392. [PubMed: 15507209]
- Zapp ML, Green MR. Sequence-specific RNA binding by the HIV-1 Rev protein. *Nature* 1989;342:714–716. [PubMed: 2556643]
- Zhang Q, Pedigo N, Shenoy S, Khalili K, Kaetzel DM. Puralpha activates PDGF-A gene transcription via interactions with a G-rich, single-stranded region of the promoter. *Gene* 2005;348:25–32. [PubMed: 15777709]




Fig. 1.

A: Structural organization of pcDNA3.1-SD4*-luciferase-RRE-SA7 and pcDNA3.1-SD4*-luciferase-SA7, a luciferase-based reporter construct. **B:** Effect of Purα on Rev-mediated luciferase activity in the presence of Leptomycin B (LMB) treatment. U-87MG cells were transfected with reporter vectors with or without RRE along with Rev and/or Purα expressing plasmids. One set of transfected cells was treated with LMB (2 nM, 16 h). Cells were harvested and luciferase activity was determined. Data represent fold increase relative to basal levels—reporter alone. **C:** Down-regulation of Purα expression in U-87MG cells by siRNA. Cells were transfected with non-target (NT) siRNA (**lane 1**) or Purα directed siRNA (**lane 2**). Western blot shows the levels of Purα and the housekeeping tubulin protein that served as a loading control. Efficiency of silencing is quantified. **D:** Effect of silencing of Purα gene expression on Rev/RRE-mediated luciferase activity. Rev expressing cells were transfected with pcDNA3.1-SD4*-luciferase-RRE-SA7 reporter plasmid and non-target siRNA or siPurα. Data represent fold increase relative to basal levels—Rev and luciferase-RRE. The experiments were repeated five times.




Fig. 2. Effect of Purα on Rev-mediated accumulation of intron-containing unspliced RNA in the cytoplasm of cells. Total (T) and cytoplasmic (C) RNA from U-87MG cells transfected with pIIIAR, pEBV-Purα and pCMV-Rev were analyzed by Northern blot. Probes were RRE (**A**), CAT (**B**) and hGAPDH (**D**). **Panel C** demonstrates the RNA integrity and the ribosomal RNA bands 28S (5.0 kb) and 18S (1.9 kb) after transfer to a membrane.




Fig. 3. Subcellular localization of Rev and Pur α in cells transfected with PLEGFP-Pur α and pDs-Rev-Red1 plasmids. **A:** U-87MG cells were transfected with reporter vectors with or without RRE along with plasmids expressing fusion Red-Rev and GFP-Pur α fusion fluorescent proteins in various combinations as denoted. Cells were fixed and red fluorescence from Rev (**B**) and green fluorescence from Pur α (**C**) were detected by microscopy. Composite of the two colors demonstrates co-localization of Pur α and Rev (**D**). DAPI Blue is for nuclear staining.




Fig. 4. Interaction of Pura with Rev. **A:** Binding of Rev to Pura and localization of the interaction domain of Rev. Protein extract from cells expressing Pura was used in GST pull-down assay. The binding of Pura to purified GST-Rev and its various deletion mutants bound to GST beads (0.5 μM) was assessed by Western blot analysis with antibody against Pura (top panel). GST alone was used for control. The schematic structural organization of Rev is illustrated and binding abilities of GST proteins are summarized (bottom panel). **B:** Binding of Pura to Rev and mapping of the interaction region of Pura. Protein extracts from cells expressing Rev were incubated with GST-Pura or its deletion mutants (0.5 μM) in GST pull-down assay for identification of the region of Pura that binds to Rev (top panel). The schematic representation

of structure of Rev protein and binding activity of GST proteins are illustrated (bottom panel). Integrity of fusion GST proteins used in assays is demonstrated (A,B, **middle panels**). **C**: U-87MG cells were transfected with pcDNA3.1-SD4*-luciferase-RRE-SA7 along with Pura and/or Rev expressing plasmids. Nuclear and cytoplasmic extracts were prepared and the presence of proteins was verified by Western blot analysis using antibodies against Rev (top panel) and Pura (middle panel). Rabbit polyclonal antibody against Pura detects endogenous and overexpressed Pura (middle panel). For loading and fractionation control, α -Tubulin was used (bottom panel). **D**: Rev and Pura were co-immunoprecipitated in cytoplasmic and nuclear lysates. Nuclear (100 μ g) or cytoplasmic (300 μ g) fractions prepared from cells as described in panel C, were immunoprecipitated using anti-Pura antibody (**lanes 1-9**). Immunoprecipitated protein complexes were washed, separated by SDS-PAGE followed by Western blot analysis using anti-Myc antibody for detection of Rev.




Fig. 5. Mutant Pur α with no binding activity to Rev is unable to enhance Rev-mediated RRE directed reporter gene expression. Cells were transfected with luciferase-RRE reporter along with plasmids expressing Rev and full-length Pur α or mutant Pur α which exhibited no binding activity to Rev. Data represent fold increase relative to basal levels-luciferase-RRE and Rev (A). Full-length Pur α demonstrates modest effect on luciferase expression in the absence of Rev (B).




Fig. 6. Interaction of Pur α with RRE in vitro. **A:** Pur α binds to RRE and enhances binding of Rev to RRE. GST, GST-Pur α and GST-Rev proteins (100 nM) were assessed in RNA gel mobility shift assay with labeled RRE 1-234 (spanning stem I, stem loops IIA, IIB, IIC, III, IV, V) and RRE 34-110 (spanning stem loops IIA, IIB, IIC). **B:** Pur α associates with RRE and with pre-existing Rev:RRE complex. Rev was incubated with labeled RRE for 10' prior to the addition of Pur α to the reaction (**lane 7**). Pur α was incubated with labeled RRE 1-234 for 10' prior to the addition of Rev to the reaction (**lane 9**). GST was used as a control in similar reactions (**lanes 6 and 8**).




Fig. 7. Cytoplasmic and nuclear presence of Pur α and Rev. U-87MG cells were transfected with pcDNA3.1-SD4*-Luciferase-RRE-SA7 along with Pur α and/or Rev expressing plasmids. Nuclear and cytoplasmic extracts were prepared and presence of proteins was verified in Western blot analysis using antibodies against Rev (**A**) and Pur α (**B**). For loading and fractionation control, tubulin was used (**C**).




Fig. 8. Interaction of Purα with RRE in vivo. Antibodies against Purα (anti-T7, **lanes 1-5**) and Rev (anti-myc, **lane 7** in “+RT” panels) were used to immunoprecipitate interacting components from cytoplasmic (**A**) and nuclear (**B**) lysates from cells as described in Figure 7. Normal mouse serum (nms) was used as a negative control for immunoprecipitation (**lane 6**). RNA extracted from these complexes was subjected to RT-PCR with (“+RT”) and without (“-RT”) reverse transcriptase (RT) using RRE 1-234 specific primers. For PCR control, pcDNA3.1-SD4*-luciferase-RRE-SA7 and RRE 1-234 specific primers were used (**lanes 8-9** in “+RT” panels and lanes 7 and 8 “-RT” panels). M: DNA marker. RNA input was verified by extraction of

RNA directly from nuclear and cytoplasmic fractions (without prior immunoprecipitation) following RT-PCR (C).