

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Papers in Microbiology

Papers in the Biological Sciences

---

October 1978

## Purification of the Protein Crystal from *Bacillus thuringiensis* by Zonal Gradient Centrifugation

Kenneth W. Nickerson

*University of Nebraska-Lincoln*, knickerson1@unl.edu

Barabara J. Ang

*University of Nebraska-Lincoln*

Follow this and additional works at: <https://digitalcommons.unl.edu/bioscimicro>



Part of the [Microbiology Commons](#)

---

Nickerson, Kenneth W. and Ang, Barabara J., "Purification of the Protein Crystal from *Bacillus thuringiensis* by Zonal Gradient Centrifugation" (1978). *Papers in Microbiology*. 17.

<https://digitalcommons.unl.edu/bioscimicro/17>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Purification of the Protein Crystal from *Bacillus thuringiensis* by Zonal Gradient Centrifugation

BARBARA J. ANG AND KENNETH W. NICKERSON\*

*School of Life Sciences, University of Nebraska, Lincoln, Nebraska 68588*

Received for publication 10 July 1978

A method is described for the large-scale purification of the *Bacillus thuringiensis* protein crystal by zonal gradient centrifugation. NaBr gradients are employed in a Beckman J21-B centrifuge equipped with a JCF-Z rotor.

*Bacillus thuringiensis* is an economically important microbial insecticide which differs from most other spore formers by synthesizing a discrete parasporal protein crystal in addition to the endospore. This protein crystal appears to be the insecticidal agent because crystal preparations which do not contain any viable cells are still highly toxic when ingested by lepidopteran insects (2). However, efficient separation of spores from parasporal crystals is a technical prerequisite of studying the chemistry of the crystal itself. The earliest attempts to obtain purified crystals relied on spontaneous germination and autolysis of the spores, followed by repeated differential centrifugation of the crystals (4, 8). These techniques were followed by several diphasic systems in which organic solvent emulsions were formed; the crystals remained in the aqueous phase. Among the solvents employed were trifluorotrchloroethane (3), tetrabromoethane (8), carbon tetrachloride (10), and chloroform (9). Unfortunately, the extractions had to be repeated several times to achieve acceptable purity, yields were low, and the danger of crystal modification by the organic solvents was ever present. This last objection was overcome by the use of diphasic systems containing dextran sulfate 500 and polyethylene glycol 6000; the spores preferentially entered the phase richer in polyethylene glycol (5, 7).

Fast (6) later described a method of crystal purification employing isopycnic density gradient centrifugation in CsCl, whereas Sharpe et al. (11) recommended the use of Renografin gradients. Both of these methods are convenient and give crystal preparations of high purity. However, the biochemical characterization of the *B. thuringiensis* crystal requires larger quantities of purified crystal than are conveniently available from density gradient centrifugation in a swinging bucket rotor.

One solution to this requirement for greater capacity is zonal gradient centrifugation (1).

Zonal centrifugation employs a hollow bowl instead of tubes. Thus, the entire 360° rotor radius is available for the density gradient, and the consequent larger gradient volume allows a larger sample capacity. We report here an isopycnic zonal centrifugation method for the large-scale purification of the insecticidal *B. thuringiensis* protein crystals employing NaBr gradients in a Beckman J-21B centrifuge equipped with a JCF-Z 40-1900 titanium rotor.

Figure 1 shows the results of a typical separation. Satisfactory resolution of the spores and crystals has been achieved with both *B. thuringiensis* var. *thuringiensis* (NRRL B-4039) and *B. thuringiensis* var. HD-1. A sample size of 100 ml was employed routinely, containing 3.0 to 7.0 g (dry weight) of spores and crystals. Purified crystal free of NaBr could subsequently be prepared either by dialysis or by dilution followed by centrifugation. The final crystal yields ranged between 15 and 20%. No systematic attempts were made to modify gradient shape or to maximize sample loading capacity.

The buoyant densities we have observed for the crystals and spores in NaBr (1.32 and 1.38 g/cm<sup>3</sup>, respectively) are, of course, different from those previously reported in CsCl (6) and Renografin (11-13). These differences are to be expected because buoyant densities are not strictly comparable between gradients of different types due to changes in the extent of hydration and possible specific ion binding. Significantly, the difference between our spore and crystal buoyant densities (0.06 g/cm<sup>3</sup>) in NaBr is of approximately the same magnitude as those reported by other workers (6, 11-13). The two lighter debris peaks (Fig. 1) are similar to those observed previously in tube gradients (6, 11).

Selection of an appropriate gradient type is especially important because of the large gradient volumes employed in zonal centrifugation. The rotor volume is 1,900 ml, and an additional 1,900 ml of heavy-density material (1.42 g/cm<sup>3</sup>




FIG. 1. Particle separation by isopycnic zonal centrifugation on NaBr gradients. The JCF-Z rotor was loaded at 2,000 rpm. The following solutions were pumped sequentially through the edge at 25 ml/min: (i) 100 ml of water, taper volume; (ii) 150 ml of 5% (wt/wt) NaBr; (iii) 150 ml of 10% NaBr; (iv) 150 ml of 15% NaBr; (v) 150 ml of 20% NaBr; (vi) 1,000 ml of a linear NaBr gradient formed on an M 146-C gradient former (1-liter capacity; MRA Corp., Clearwater, Fla.); 500 ml of 25% NaBr was mixed with 500 ml of 40% NaBr; (vii) 200 ml 42% NaBr cushion; the rotor was now full and further loading proceeded through the center port; (viii) 100 ml of sample (3.5 g of homogenized spores and crystals from *B. thuringiensis* var. HD-1 in water) displacing the taper volume; (ix) 100 ml of water overlay to force the sample into the gradient. Subsequently, the rotor was accelerated to 10,000 rpm. After a 4-h centrifugation at 4°C the rotor was decelerated to 2,000 rpm and unloaded by pumping in 1,900 ml of 42% NaBr from the edge at 25 ml/min. A total of 38 fractions of 50 ml each were collected and monitored according to their density, absorbance (Klett<sub>66</sub> colorimeter), and microscopic appearance. Fraction density was measured with two hydrometers (Fisher Scientific Co.) with ranges of 1.000 to 1.200 and 1.200 to 1.400 g/cm<sup>3</sup>. At least 50 ml of solution was required for their proper use. A total of 0.45 g of purified crystal was recovered.

of NaBr) must be pumped in through the edge in order to unload the rotor from the center. Both Renografin and CsCl are too expensive to be used in such large amounts. Thus, the ideal gradient must be of sufficient density (up to 1.42 g/cm<sup>3</sup>), cheap, and of low viscosity because low viscosity allows the use of less expensive gravity-flow gradient makers. NaBr was considered the best candidate. It can be purchased in bulk (350-lb. [ca. 159-kg] drums) from Great Lakes Chemical Corp., West Lafayette, Ind.

This research was supported by grant 516-15-121 from the United States Department of Agriculture CSRS and by Public Health Service Biomed Research Support grant RR-07055 from the Division of Research Resources, National Institutes of Health.

#### LITERATURE CITED

- Anderson, N. G. 1962. The zonal ultracentrifuge. A new instrument for fractionating mixtures of particles. *J. Phys. Chem.* **66**:1984-1989.
- Angus, T. A. 1956. Extraction, purification, and properties of *Bacillus sotto* toxin. *Can. J. Microbiol.* **2**:416-426.
- Angus, T. A. 1959. Separation of bacterial spores and parasporal bodies with a fluorocarbon. *J. Insect Pathol.* **1**:97-98.
- Cooksey, K. E. 1971. The protein crystal toxin of *Bacillus thuringiensis*: biochemistry and mode of action, p. 247-274. In H. D. Burges and N. W. Hussey (ed.), Biological control of insects and mites. Academic Press Inc., London.
- Delaffield, F. P., H. J. Somerville, and S. C. Rittenberg. 1968. Immunological homology between crystal and spore protein of *Bacillus thuringiensis*. *J. Bacteriol.* **96**:713-720.
- Fast, P. G. 1972. The  $\delta$ -endotoxin of *Bacillus thuringiensis* III. A rapid method for separating parasporal bodies from spores. *J. Invertebr. Pathol.* **20**:139-140.
- Goodman, N. S., R. J. Gottfried, and M. H. Rogoff. 1967. Biphasic system for separation of spores and crystals of *Bacillus thuringiensis*. *J. Bacteriol.* **94**:485.
- Lecadet, M. M. 1970. *Bacillus thuringiensis* toxins—the proteinaceous crystal, p. 437-471. In T. C. Montie, S. Kadis, and S. S. Ajl (ed.), *Microbial toxins*, vol. III. Academic Press Inc., New York.
- Murray, E. D., and E. Y. Spencer. 1966. A simplified purification technique for parasporal inclusions from certain varieties of *Bacillus thuringiensis*. *J. Invertebr. Pathol.* **8**:418-420.
- Pendleton, I. R., and R. B. Morrison. 1966. Separation of the spores and crystals of *Bacillus thuringiensis*. *Nature (London)* **212**:728-729.
- Sharpe, E. S., K. W. Nickerson, L. A. Bulla, Jr., and J. N. Aronson. 1975. Separation of spores and parasporal crystals of *Bacillus thuringiensis* in gradients of certain X-ray contrasting agents. *Appl. Microbiol.* **30**:1052-1053.
- Tamir, H., and C. Gilvarg. 1966. Density gradient centrifugation for the separation of sporulating forms of bacteria. *J. Biol. Chem.* **241**:1085-1090.
- Wise, J., A. Swanson, and H. O. Halvorson. 1967. Dipicolinic acid-less mutants of *Bacillus cereus*. *J. Bacteriol.* **94**:2075-2076.