

Research paper

Purification, properties and alternate substrate specificities of arginase from two different sources: *Vigna catjang* cotyledon and buffalo liverSnehal Dabir¹, Pankaj Dabir² and Baburao Somvanshi¹¹ Department of Biochemistry, Dr BA Marathwada University, Aurangabad – 400 001 (MS), India² Department of Biochemistry and JB Tropical Disease Research Centre, Mahatma Gandhi Institute of Medical Sciences, Sevagram - 442 102, (MS), India

Corresponding address: Dr. Snehal Dabir, Phone: (+91)9822562943, E-mail: snehal_214@rediffmail.com

Received: 2005.03.10; Accepted: 2005.07.08; Published: 2005.08.01

Arginase was purified from *Vigna catjang* cotyledons and buffalo liver by chromatographic separations using Bio-Gel P-150, DEAE-cellulose and arginine AH Sepharose 4B affinity columns. The native molecular weight of an enzyme estimated on Bio-Gel P-300 column for *Vigna catjang* was 210 kDa and 120 kDa of buffalo liver, while SDS-PAGE showed a single band of molecular weight 52 kDa for cotyledon and 43 kDa for buffalo liver arginase. The kinetic properties determined for the purified cotyledon and liver arginase showed an optimum pH of 10.0 and pH 9.2 respectively. Optimal cofactor Mn⁺⁺ ion concentration was found to be 0.6 mM for cotyledon and 2 mM for liver arginase. The Michaelis-Menten constant for cotyledon arginase and hepatic arginase were found to be 42 mM and 2 mM respectively. The activity of guanidino compounds as alternate substrates for *Vigna catjang* cotyledon and buffalo liver arginase is critically dependent on the length of the amino acid side chain and the number of carbon atoms. In addition to L-arginine cotyledon arginase showed substrate specificity towards agmatine and L-canavanine, whereas the liver arginase showed substrate specificity towards only L-canavanine.

Key words: Purification, kinetic properties, substrate specificity arginine amidino hydrolase, *Vigna catjang* cotyledon, buffalo liver**1. Introduction**

L-Arginase (L-arginine amidino hydrolase, EC 3.5.3.1.) is a binuclear manganese metalloenzyme that catalyzes the hydrolysis of L-arginine to form L-ornithine and urea. Arginase is the terminal enzyme of the urea cycle among the six enzymes. The enzyme was found to exist in two forms and has a broad tissue distribution. A cytosolic form, AI is highly expressed in the liver or hepatic cells and is important in ureogenesis. Extra-hepatic AII form is thought to be involved in the biosynthesis of polyamines, the amino acids ornithine, proline and glutamate and in the inflammatory process, among others [1]. Recently studies have shown that increased stimulation of arginase expression in animal systems leads to production of polyamines that promote tumor cell proliferation [2] and wound healing [3].

Most of the studies on plant arginase have focused on its role in mobilizing arginine during early seedling germination. Storage proteins are mobilized to provide amino acids for protein synthesis in the expanding axis. Arginase activity increases sharply during germination in several species including soybean [4, 5], arabidopsis and loblolly pines [6, 7]. In higher plants urea and ornithine formed as products of arginase reaction are used in the assimilation of nitrogen into amino acids and the synthesis of polyamines through urease and ornithine decarboxylase respectively [8, 9]. The enzyme has been isolated and characterized from cotyledons as well as axis of soybean [5, 8]. But no properties of the purified enzyme from the cotyledons of *Vigna catjang* and buffalo liver have been reported. In animal's arginase is purified and characterized from a variety of sources having different characteristic properties with regard to molecular weight, subunit structure and its regulation especially in the urea cycle [10].

In contrast far less is known about the purification and properties of the enzyme in higher plants [11, 12]. Hence the present study was undertaken for the purification and properties of arginase from *Vigna catjang* cotyledon and buffalo liver, and also reports physicochemical and kinetic properties of the purified enzyme.

This purified enzyme can be used to study the effect of L-amino acids on the inhibition of arginases, to know the composition of amino acids at the active site of enzyme and may be useful in investigating its role in mobilizing the storage proteins.

2. Materials and methods**Source of enzyme**

Seeds of cow pea *Vigna catjang* were obtained from the local market; surface sterilized by treating with Lysol, soaked in water for 6 hrs and then allowed to germinate for 24 hrs on moist filter paper at 25°C.

Buffalo liver sample was collected from the local slaughterhouse. A portion of liver tissue was removed within 15 min after sacrificing the animal. It was placed in ice bath and brought to the laboratory as early as possible. The liver sample was immediately used for the isolation and purification of arginase.

Cotyledon arginase activity assay

Arginase activity was measured according to the Roman and Ruy's method [13]. Briefly the reaction mixture consisting of 10 mM carbonate bicarbonate buffer (pH 10), 2 mM MnCl₂, 130 mM L-arginine and enzyme solution in a total volume of 1 ml was incubated for 30 minutes at 37 °C. The reaction was terminated by adding 10% TCA. Protein was removed by centrifugation; 1 ml of the supernatant was mixed with 1ml of anhydrous glacial acetic acid and ninhydrin reagent. The mixture was kept

in boiling water bath for 20 min. The intense red colour developed was diluted with 2 mls of glacial acetic acid. The reagent blank was run in a parallel way and the results were compared with control assay. The optical density was read at 515 nm on spectrophotometer (UV 160A Shimadzu, Japan.) against reagent blank. The activity of arginase was expressed in units. A unit of an enzyme activity is defined as that amount of an enzyme, which produces one μ mole of L-ornithine per min at 37°C.

Liver arginase activity assay

Enzyme assay and experimental conditions used for buffalo liver arginase were same as described above for *Vigna catjang* cotyledon arginase, except millimolar concentrations of $MnCl_2$ (6 mM), L-arginine (25 mM) and carbonate bicarbonate buffer (50 mM, pH 9.2).

Protein determination

The protein content of each fraction was determined by the method of Lowry et al. [14] with the bovine serum albumin (Sigma, USA) as standard.

Purification of cotyledon arginase

Purification of the enzyme was carried out at 0-4 °C. The cotyledons (100 gm) were homogenized in a chilled mixer with 300 ml of 50 mM Tris-HCl buffer (pH 7.5) containing 0.6 mM $MnCl_2$, 100 mM KCl and 10% Glycerol. The homogenate was filtered through four layers of gauze and clarified by centrifugation at 13,000 rpm for 30 min at 4 °C.

Ammonium sulphate precipitation

The supernatant was adjusted to 30% saturation with solid ammonium sulphate and stirred at 4°C for 5 hours. The solution was then centrifuged and the pellet was discarded. The supernatant was brought to 70% saturation with solid $(NH_4)_2 SO_4$ and treated as above except that the pellet was retained. The pellet containing the enzyme was resuspended with 30 mls of 10 mM Tris-HCl buffer (pH 7.5) containing 0.6 mM $MnCl_2$ and 10% glycerol. The extract was purified by centrifugation.

Bio-Gel P-150 chromatography

The clear supernatant was applied to a Bio-Gel P-150 column (2 X 80 cm) equilibrated with Tris-HCl buffer (10 mM, pH 7.5) containing 0.6 mM $MnCl_2$ and 10% glycerol. The enzyme was eluted using the same buffer with a flow rate of 12 ml / hour.

DEAE-cellulose chromatography

Active fractions obtained from above step were applied to DEAE-cellulose (Sigma) column (2.5 X 12 cm) previously equilibrated with Tris-HCl buffer (10 mM, pH 7.5) containing 10% glycerol. The column was washed with the same buffer and the enzyme was eluted using a linear gradient of sodium chloride (0-0.5 M) prepared in column-equilibrated buffer.

Affinity chromatography

Active fractions from above step were pooled and applied on a hydroxyapatite column (1.2 X 10 cm) (Bio-Rad, USA) previously equilibrated with Tris-HCl buffer (10 mM, pH 7.5) containing 0.6 mM $MnCl_2$ and 10% glycerol followed by 15 ml of 0.5 M KCl prepared in column equilibrating buffer. The flow rate of column was adjusted to 12 ml / hour, each 1.0 ml fractions were collected and its protein content and arginase activity were assayed.

Purification of liver arginase

Liver sample was weighed (30 gm) homogenized with 120 ml of chilled Tris-HCl buffer (100 mM, pH 7.5) containing 5 mM $MnCl_2$ and 100 mM KCl. The homogenate was centrifuged at 13,000 rpm, 30 minutes and the supernatant was collected.

Heat treatment

The heat treatment was given to the above supernatant at 60 °C, 20 min with swirling. It was then immediately cooled in ice bath and denatured proteins were removed by centrifugation.

Ammonium sulphate precipitation

The resulting supernatant was adjusted to 30% saturation with solid $(NH_4)_2 SO_4$ and stirred at 4 °C for 1 hour. The solution was then centrifuged and the pellet was discarded. The supernatant was brought to 70% saturation with solid ammonium sulphate and treated as above except that the pellet was retained. The pellet was resuspended in 15 mM Tris-HCl buffer pH 8.0 containing 5 mM $MnCl_2$ and kept for 2 hours. The unresolved proteins were removed by centrifugation and dialyzed against the Tris-HCl buffer used for homogenization.

DEAE-cellulose chromatography

The dialyzed material was applied to a 2.5 X 12 cm column of DEAE-cellulose (Sigma) at 4 °C, equilibrated with the Tris-HCl buffer (15 mM, pH 8.0) containing 2 mM $MnCl_2$. Elution was carried out with a linear gradient of NaCl from 0 to 0.5 M in same buffer containing 2 mM $MnCl_2$ at a flow rate of 20 ml / hour. The active fractions containing arginase activity were pooled together.

Bio-Gel P-150 chromatography

Above pooled fractions were dialyzed against Tris-HCl buffer (15 mM, pH 8.0) containing 2 mM $MnCl_2$ and applied as a concentrated zone on Bio-Gel P-150 (2 X 80 cm) column equilibrated with Tris-HCl buffer (15 mM, pH 8.0) containing 2 mM $MnCl_2$. Each 2 mls of fractions were collected at 12 ml / hour and active fractions containing arginase activity were pooled.

Affinity chromatography

The pooled fractions from above step were dialyzed at 4 °C in 3 X 3 L of Tris-HCl buffer (15 mM, pH 8.0) containing 2 mM $MnCl_2$ and passed through the arginine AH-Sepharose 4B (1.2 X 10 cm) column equilibrated with 15 mM Tris-HCl buffer pH 8.0 containing 1 mM $MnCl_2$. The column was washed with 20 mls of equilibrating buffer without $MnCl_2$. The adsorbed proteins were eluted by passing 20 mls of 10 mM Tris-HCl (pH 8.0), 10 mls of 50 mM Tris-HCl (pH 7.5), 10 mls of 50 mM Tris-HCl (pH 8.5) and finally with 1.0 M sodium chloride in column equilibrating buffer. The fractions of 2 ml were collected with a flow rate of 12 ml / hour and arginase activity and protein content were assayed.

SDS-PAGE electrophoresis

SDS-polyacrylamide gel electrophoresis (10%) was carried out according to Laemmli et al (15). The purified enzyme was treated with 1% SDS and β -mercaptoethanol for 10 minutes at 100 °C and loaded in wells. Sample stacking was done at 10 mM and resolution was carried out at 15 mA constant current. Molecular weight standards were bovine serum albumin (66,000), Egg albumin (45,000), Glyceraldehydes triphosphate

dehydrogenase (35,000), Carbonic anhydrase (29,000) and β -lactoglobuline (18,400) (Sigma). After electrophoresis gel was stained in 0.25% w/v Coomassie brilliant blue (R-250) prepared in 50% v/v methanol and 10% v/v acetic acid. The gels were destained by passive diffusion of dye in 50% v/v methanol and 1% acetic acid after changing the destaining solution for 2-3 times.

Native molecular weight determination

The molecular weight of the purified enzyme was estimated by gel filtration with a Bio-Gel P-300 column (2.2 X 78 cm). Blue dextran was used to measure the void volume. Catalase (bovine) (2, 40,000), Aldolase (rabbit muscle) (147,000), Albumin (bovine serum) (67,000), Albumin (egg) (45,000), Chymotrypsinogen A (25,000) and Cytochrome C (12,300) were used as a molecular weight marker. The elution volume of arginase was determined by estimating enzymatic activity whereas the protein content of each fraction was determined by Lowry's method [14]. The position of each marker protein was defined by plotting the elution volume against protein concentration. The molecular weight of an arginase was estimated by plotting log of molecular weight versus K_{av} for the standards. $K_{av} = (V_e - V_o) / (V_t - V_o)$, where V_o is the void volume of the column; V_t is the total volume of the column and V_e is the elution volume of a protein.

Effect of pH on arginase activity

The pH dependence of the enzymatic activity of arginase from cotyledon and liver were determined by using three buffer systems: 10 mM Potassium phosphate (pH 6.0 to 6.5), Tris-HCl (pH 7.0 to 9.0) and carbonate bicarbonate buffer (pH 9.0 to 11.0).

Effect of divalent cations

To know the effect of divalent cations on the activity of arginase, the purified cotyledon arginase was diluted (1:2) with distilled water. The enzyme was dialyzed extensively with distilled water and further with 10 mM Tris-HCl buffer (pH 7.5) in the absence of manganese and then assayed the purified enzyme with the various metals.

The same procedure was followed to study the effect of different cations on the activity of buffalo liver arginase under the suitable enzyme assay conditions.

Estimation of urea

Arginase activity assay of the *Vigna catjang* cotyledon and hepatic tissue was carried out according to the procedure of Marsh WH et al. [16] by estimating urea. The reaction mixture consisted of 10 mM carbonate bicarbonate buffer (pH 10), 2 mM $MnCl_2$, 130 mM L-arginine and enzyme solution in a total of 1 ml was incubated for 30 minutes at room temperature. This enzymatic reaction was terminated by adding 10% TCA. Two mls of reaction mixture was taken for urea estimation. The control tube was run simultaneously by inactivating the enzyme with 10% TCA prior to the addition of compound tested as alternate substrate (s) for arginase. The tubes were centrifuged, 2 ml of the clear supernatant was transferred from each tube and the enzyme activity was recorded by spectrophotometric detection of urea at 520 nm by using diacetyl monoxime as described by Marsh WH et al. Enzyme assay and experimental conditions used for buffalo liver arginase were the same as described above for the *Vigna catjang* arginase, except millimolar concentrations of $MnCl_2$ (6

mM), L-arginine (25 mM) and carbonate bicarbonate buffer (50 mM, pH 9.2).

Substrate specificity

The substrate specificity of *Vigna catjang* cotyledon and buffalo liver arginase was investigated by testing its activity towards structurally similar compounds with guanidino group. The agmatine, d-arginine, L-arginine, L-canavanine, L-argininic acid, β -guanidino propionic acid, γ -guanidino butyric acid, L-homoarginine and streptomycin (Sigma, USA.) tested as a potential alternate substrate for arginase. The solutions of above compounds (30 mM) were prepared in 10 mM carbonate bicarbonate buffer. Enzyme activity was measured by spectrophotometric detection of urea as described above.

3. Results and Discussion

Purification

The purification of arginase from *Vigna catjang* cotyledon and buffalo liver was carried out as described earlier 'Materials and Methods'. The results from the purification of arginase from cotyledons and liver are summarized in Table 1. The fractional precipitation of arginase with chilled acetone (-10 °C) and heat treatment results in reduced stability of the cotyledon arginase, but fractional precipitation with ammonium sulphate yielded good results. It showed 2.2-fold purification corresponding to a specific activity of 70.15. The enzyme fractions were freed of ammonium sulphate and low molecular weight protein by passing the solution through a Bio-Gel P-150 column. Arginase was adsorbed on a DEAE-cellulose column and eluted at about 300 mM sodium chloride in column equilibrating buffer (Tris-HCl, 10 mM, pH 7.5) containing 0.6 mM $MnCl_2$ and 10% glycerol. Inclusion of 10% glycerol in the column buffers has given stability to arginase activity during chromatographic purification. The enzyme was specifically bound on an arginine Sepharose 4B column and was eluted at a high molarity of Tris-HCl buffer (Figure 1). The specific activity of the arginase obtained from affinity column was about five fold higher than DEAE-cellulose chromatography. Probably arginine AH-Sepharose 4B minimized non-specific adsorption. The purification of *Vigna catjang* cotyledon arginase was achieved about 64 fold with 1.64% yield. This is about two and half fold lower than that reported for the axis of Soybean [5].

Arginase isozymes have not been reported in *Iris bulbosus* and *Pisum sativum* and *Jerusalem artichoke tubers* [11, 12, 17]. We could not find any evidence of the presence of arginase isozymes in purification steps.

Heat treatment was given to the crude extract of buffalo liver to enhance the coagulation of thermolabile proteins, which were further removed by centrifugation and supernatant was collected to test the enzymatic activity. It was found that arginase from buffalo liver was stable at 60 °C. Such thermal stability of arginase has also been reported from livers of beef [18] and rabbit [19]. 70% saturation of arginase with solid ammonium sulphate has resulted in increased specific activity of liver arginase. We did not observe any isozyme for buffalo liver arginase on DEAE-cellulose chromatography. The enzyme was eluted in a single symmetrical peak, with linear gradient of sodium chloride (Figure 2). The above active enzyme fractions were passed through Bio-Gel P-150 chromatography and a single homogenous peak of

arginase activity was appeared. We have obtained about 2028.97 fold purification from arginine AH-Sepharose 4B affinity adsorbent (Table 1). The overall recovery was about 18 %. From the purification data it is clear that the protocol which we have used for the purification of *Vigna catjang* cotyledon and buffalo liver arginase is simple and effective.

Molecular weight

The molecular weight of the native enzyme determined by gel filtration was found to be 210 kDa for *Vigna catjang* and 120 kDa for buffalo liver arginase (Figure 3). The molecular weight on SDS-PAGE was found to be 52 kDa and 43 kDa for *Vigna catjang* and buffalo liver arginase respectively (Figure 4).

The molecular weight obtained for cotyledon arginase is smaller, compared with those of soybean axes 220 kDa [5] and *E. prunastri* 330 kDa [20]. Reports obtained for buffalo hepatic arginase agrees with the reported molecular weights of arginases from beef 115 kDa [18] and rabbit liver 110 kDa [19].

Kinetic properties

The effect of incubation time on the activities of cotyledon and liver arginase was studied up to 75 min. There was a direct relationship although not linear, between the product formed and incubation time for the enzymes. From the observations common incubation time selected for the studies was 30 min.

Buffalo liver arginase is stable at 55-60 °C as in rat mammary gland [21] and rabbit liver [19], but in case of cotyledon arginase it is found to be less stable at the same temperature. The maximum activity of cotyledon and liver arginase was found at 35 °C and 42 °C respectively (Figure 5).

Effect of pH

The activity of cotyledon and liver arginase was determined in the buffer pH range from 6.0 to 11 at 37 °C. The optimum pH of cotyledon arginase was found to be 10.0. This is found to be slightly higher than *Evernia prunastri* and Soybean axes [5, 20]. Liver arginase showed high activity at pH 9.2 (Figure 6). Generally mammalian arginases appear to have basic pH optima of 9.5-10.5, although some exceptions have been noted. Gasiorowska [22] have found four different isoenzymes of arginase in rat tissues, one of which had a pH optimum of 7.5. On the other hand, Patil et al [23] noted optimum pH 11.5 for Ox-erythrocyte arginase at 55 °C. The variation of activity with pH suggests that an ionisable group may function at the catalytic site [24].

Michaelis-Menten constant (Km)

The effect of substrate concentration on the activity of cotyledon and liver arginase was determined by Lineweaver-Burk, double reciprocal (1/V Vs 1/S) plots. The Km values for cotyledon and liver arginase were found to be 42 mM and 2 mM respectively as shown in figure 7 (A) and (B). The Km value of *Vigna catjang* cotyledon arginase is in the range of other reported Km (40-83 mM) for plant arginases [5, 11, 25]. The Km value of buffalo liver arginase is in the range reported for type I arginase isolated from human 6.3 mM [26] and rabbit liver [19]. A variation in millimolar concentration of arginine is may be due to the variety of assays used and non-physiological conditions generally used [27, 28]. It is interesting to note that at a higher concentration of

substrate, double reciprocal plot is non-linear and seems to indicate excess substrate inhibition in both *Vigna catjang* cotyledon and buffalo liver arginase.

Effect of metal ions

Arginase is a metallo enzyme in which manganese acts as a cofactor as well as activator in almost all reported arginases [20, 29]. However the amount required for optimal activity of the enzyme varies. About 0.6 mM and 2 mM MnCl₂ in the reaction mixture were found to be optimal for the activity of *Vigna catjang* and buffalo liver arginase respectively.

Replacement of Mn⁺⁺ ion with other metal ions such as Mg⁺⁺ and Co⁺⁺ restored more than 60 % while Ni⁺⁺ and Fe⁺⁺ restored about 50 % activity, whereas Ca⁺⁺ and Zn⁺⁺ restored about 35 to 38 % and Cd⁺⁺ restored only 10 % of the original activity of *Vigna catjang* cotyledon arginase. In case of buffalo liver arginase, Mg⁺⁺ ions restored almost the original activity, whereas Ca⁺⁺, Ni⁺⁺, Co⁺⁺ and Cd⁺⁺ restored about 40-50 % of the original activity of the enzyme and Fe⁺⁺ and Zn⁺⁺ has completely inactivated the arginase activity (Table 2).

The amount of Mn⁺⁺ required for optimal activity of the soybean axes enzyme was reported as 1 mM [5], 1-2 mM for human liver [30], 8-10 mM for human erythrocytes [31], 40 mM for mammalian liver arginase [32] and 50 mM for beef liver arginase [18]. Maggini et al [33] have measured arginase activity by reassessment of assay conditions using near physiological steady-state concentrations of arginine (5-35 mM) and excess Mn⁺⁺ (30 mM). They obtained specific activity values for rat liver arginase of 1.2 U / g of tissue, compared with activity values of 70 U / g from initial velocity experiments. This modified activity values suggest that arginase could play a role in the regulation of urea cycle activity, its activity being regulated by the concentrations of arginine and Mn⁺⁺ ions. Increased streptozotocin-induced in diabetic rat's resulted in activation of hepatic arginase [34]. Similarly, the concentration of manganese in the diet has been shown to regulate the activity of hepatic arginase in the rat [35].

Substrate specificity

Hepatic arginase of ureotelic and urecotelic animals have nearly same substrate specificity towards L-arginine and to some extent L-canavanine [36]. The leupine arginase hydrolyzes L-homoarginine more efficiently than L-arginine [37]. With this background it was interesting to investigate the substrate specificity of *Vigna catjang* cotyledon and buffalo liver arginase. Different substrates containing guanidino group common in their structure were tested as potential substrates for *Vigna catjang* cotyledon among these only agmatine, and L-canavanine were hydrolyzed significantly in addition to L-arginine. The ratio of the reaction product, urea of L-arginine to agmatine and L-arginine to L-canavanine is six and fifteen respectively Table 3. Arginase/canavanine high ratios (>10) have been reported from arginase from rat liver [35] and silk moth [38] Table 3.

Buffalo liver arginase hydrolyzed only L-arginine and L-canavanine under the standard assay conditions and the results are shown in (Table 3). The ratio of L-arginine to L-canavanine is more than ten. The enzyme does not hydrolyze the isomer of L-arginine, the D-arginine. Agmatine has no α -carboxyl group, whereas γ -guanidino butyric acid and β -guanidino propionic acid

lack α -amino group. Streptomycin does not possess α -NH₂ as well as α -COOH groups. Therefore buffalo liver arginase does not utilize these guanidine compounds as its substrate to liberate urea as one of the product.

Conflict of interest

The authors have declared that no conflict of interest exists.

References

- Cederbaum SD, Yu H, Grody WW, et al. Arginase I and II: do their functions overlap? *Med Genet Metab* 2004; 81 (suppl 1):S38-44.
- Chang CI, et al. Macrophage arginase promotes tumor cell growth and suppresses nitric oxide-mediated tumor cytotoxicity. *Cancer Res* 2001; 61:1100-1106.
- Satriano J. Agmatine: At the Crossroads of the Arginine Pathways. *Ann N Y Acad Sci* 2003; 1009:34-43.
- Kang JH, Cho YD. Purification and properties of arginase from soybean (*Glycine max*). *Plant physiol* 1990; 93:1230-1234.
- Matsubara S, Suzuki Y. Arginase activity in the cotyledons of soybean seedlings. *Plant physiol* 1984; 62:309-314.
- Zonia LE, et al. Essential role of arginase in germination of nitrogen limited *Arabidopsis thaliana* seeds. *Plant physiol.* 1995 107 :1097-1103
- King JE, Gifford DJ. Amino acid utilization in seeds of Loblolly pine during germination and early seedlings growth L-Arginine and arginase activity. *Plant physiol.* 1997; 113:1125-1135.
- Dawnun KR, Rosenthal GA, Cohen WS. L-Arginine and L-canavanine metabolism in jack bean, *Canavalia ensiformis* (L) DC and soybean, *Glycine max* (L) mere. *Plant physiol* 1983; 73:965-968.
- Polacco JC, Holland MA. Roles of urease in plant cells. *Int Rev Cytol* 1993; 146:65-103.
- Carvajal N, Ascoria M, Rodriguez JP, et al. Evidence of cooperative effects in human liver arginase. *Biochim Biophys Acta* 1982; 701:146-148.
- Butin JP. Purification, properties and subunit structure of arginase from *Iris bulbosus*. *Eur J Biochem* 1982; 127:237-243.
- Wright LC, Brady GJ, Hinde RW. Purification and properties of arginase from Jerusalem artichoke tubers. *Phytochem* 1981; 20:2641-2645.
- Roman W, Ruys J. Simple method for the estimation of l-ornithine. *J Clin. Enzyme* 1969; 2:121-128.
- Lowry OH, Roserbrough NJ, Farr AL, et al. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; 193:265-275.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 1970; 227:680-685.
- Marsh WH, Fingerhut B, Miller H. Automated and manual direct methods for determination of urea. *Clin Chem* 1965; 11:624-627.
- Talor AA, Stevart GR. Tissue and Subcellular localization of enzymes to arginine metabolism in *Pisum sativum*. *Biochem Biophys Res Commun* 1981; 101:1281-1289.
- Harell D, Sokolovsky M. Beef-liver arginase. Isolation and molecular properties. *Eur J Biochem* 1972; 25:102-108.
- George AK Harold JS. Isolation and characterization of arginase from rabbit liver. *Biochem J* 1973; 133:779-788.
- Angels MF, Legaz ME. Purification and properties of the constitutive arginase of *Evernia prunastri*. *Plant Physiol* 1984; 76:1065-1069.
- Christopher PJ, Murray RG. Rat mammary arginase: Isolation and characterization. *Biochemical Med and Meta Biol* 1994; 51:156-165.
- Gasiorowska I, Porembaska Z, Jochimowicz J, et al. Isozymes of arginase in rat tissue. *Acta Biochim Polon* 1970; 17:19-30.
- Patil NB, Somvanshi BS, Kothari RM. Isolation and characterization of arginase from Ox-erythrocyte. *Biotech Tech* 1990; 4 (2):133-136.
- Kuhn NJ, Talbor J, Ward S. pH-sensitive control of arginase by Mn (II) ions at submicromolar concentrations. *Arch Biochem Biophys* 1991; 286:217-221.
- Kollofel C, Van Dijke HD. Mitochondrial arginase activity form cotyledons of developing and germinating seeds of *Vicia faba*. *Plant physiol* 1975; 55:507-510.
- Buscur L, Cabellow J, Veliz M, et al. Molecular forms of human liver arginase. *Biochem Biophys Acta* 1966; 128:149-154.
- Ganganta CL, Bond JS. Assay and kinetics of arginase. *Anal Biochem* 1986; 154:388-394.
- Pace CN, Buonanno A, Simmons - Hansen J. Steady-state kinetic studies of arginase with an improved direct spectrophotometric assay. *Anal Biochem* 1980; 109:261-265.
- Helga HK, Greenberg DM. Molecular characteristics of rat liver arginase. *J Biol Chem* 1968; 243:6123.
- Carvajal N, Venegas A., Ostreicher G, et al. The effect of manganese on the quaternary structure of human liver arginase. *Biochim Biophys Acta* 1971; 250:437- 442.
- Nishibe H. Isolation and characterization of arginase in human erythrocyte. *Physiol Chem Physics* 1973; 54:53 - 462.
- Hirsch-Kolb H, Kolb HJ, Greenberg DM. Comparative physical and chemical studies of mammalian arginase. *Comp Biochem Physiol* 1970; 37:345-359.
- Maggini S, Stockin-Tschan FB, Morikofer-Zwez S, et al. New kinetic parameters for rat liver arginase measured at near physiological steady-state concentrations of arginine and Mn²⁺. *Biochem J* 1992; 283:53-660.
- Bond JS, Failla ML, Unger DF. Elevated manganese concentration and arginase activity in livers of streptozotocin - induced diabetic rats. *J Biol Chem* 1983; 258:8004-8009.
- Brock AA, Chapman SA, Ulman EA, et al. Dietary manganese deficiency decreases rat hepatic arginase activity. *Nutrition* 1994; 124:340-344.
- Mora J, Tarrab R., Maratuscelli L, et al. Characteristics of arginases from ureotelic and non-ureotelic animals. *Biochem J* 1965; 96:588-594.
- Muszynska G, Reifer I. Purification, properties and inhibition of plant arginase. *Acta Biochim Pol* 1968; 15:55-66.
- Reddy SRR, Campbell JW. Arginine metabolism in insects: properties of insect fat body arginase. *Comp Biochem Physiol* 1969; 28:515-534.

Tables & Figures

Table 1. Purification of Arginase from *Vigna catjang* cotyledon and buffalo liver

(a) <i>Vigna catjang</i>							
Sr. No	Purification step	Volume (ml)	Total protein (mg)	Total activity (units x 10 ³)	Specific activity (units/mg)	Purification (fold)	Yield (%)
1	Homogenate / Supernatant	280	8,620	274	31.78	1.0	100
2	Ammonium sulphate	20	1,910	134	70.15	2.20	48.90
3	Bio- Gel P-150 chromatography	20	462	47	101.73	3.20	17.15
4	DEAE-cellulose chromatography	18	80	32.4	400	12.58	11.67
5	Affinity chromatography	6	2.2	4.5	2045.45	64.36	1.64
(b) Buffalo liver							
1	Homogenate	130	10,200	459	45	1.0	100
2	Homogenate supernatant	90	2,367	450	190	4.22	98.03
3	Heat treatment	69	584	365	625	13.88	79.52
4	Ammonium sulphate	16	248	312	1,258	27.95	67.97
5	DEAE-cellulose chromatography	34	30	253	8,433	187.40	55.11
6	Bio-Gel P-150 chromatography	16	3.8	179	47,105	1046.77	38.99
7	Affinity chromatography	8	0.92	84	91,304	2028.97	18.30

Table 2. Effect of divalent cations on the activity of *Vigna catjang* cotyledon and buffalo liver arginase.

Metal ion 2 mM	Residual activity (%)	
	Cotyledon arginase	Liver arginase
Mn ⁺⁺	100	100
Mg ⁺⁺	61	93
Ca ⁺⁺	35	52
Co ⁺⁺	66	45
Ni ⁺⁺	55	51
Cd ⁺⁺	11	43
Fe ⁺⁺	52	00
Zn ⁺⁺	38	00

Table 3. Substrate specificities of *Vigna catjang* cotyledon and buffalo liver arginase

Compounds (30 mM)	Structure	Urea in $\mu\text{moles / min}$	
		Cotyledon	Liver
Agmatin	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_4\text{NH}_2$ NH	10.0	0.0
D-Arginine	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_3-\text{C}-\text{COOH}$ NH NH ₂ H	0.0	0.0
L-Arginine	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_3-\text{C}-\text{COOH}$ NH NH ₂ H	63.0	125.0
L-Homoarginine	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_4-\text{C}-\text{COOH}$ NH NH ₂ H	0.0	0.0
L-argininic acid	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_3-\text{C}-\text{COOH}$ NH NH ₂ H	0.0	0.0
L-Canavanine	$\text{H}_2\text{N}-\text{C}-\text{NH}-\text{O}(\text{CH}_2)_2-\text{C}-\text{COOH}$ NH NH ₂ H	8.0	12.0
γ -Guanidino butyric acid	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_3-\text{COOH}$ NH	0.0	0.0
β -Guanidopropionic acid	$\text{H}_2\text{N}-\text{C}-\text{NH}-(\text{CH}_2)_2-\text{COOH}$ NH	0.0	0.0
Streptomycin	$\text{H}_2\text{N}-\text{C}-\text{NH}$ R	0.0	0.0

Figure 1: Elution profiles of *Vigna catjang* arginase activity and protein content. A) Bio-Gel P-150 B) DEAE-cellulose and C) Arginine AH-Sepharose 4B chromatography.




Figure 2: Elution profiles of Buffalo liver arginase activity and protein content. A) DEAE-cellulose B) Bio-Gel P-150 and C) Arginine AH-Sepharose 4B chromatography.




Figure 3: Determination of molecular weight of *Vigna catjang* cotyledon (E) and Buffalo liver (E1) arginase by gel filtration technique on Bio-Gel P-300.




Figure 4: SDS-PAGE (10%) analysis of purified *Vigna catjang* cotyledon and buffalo liver arginase. M: Molecular weight standard, size in kilodalton is shown on the left. A: Purified *Vigna catjang* cotyledon arginase and B: Purified buffalo liver arginase.




Figure 5: Determination of optimal temperature for the activity of *Vigna catjang* (●) cotyledon and buffalo liver (■) arginase. The data represents an average of multiple experiments.




Figure 6: Determination of the optimal pH for *Vigna catjang* (●) cotyledon and buffalo liver (■) arginase. The enzymatic activity of arginase was determined by replacing the buffer in the pH range of 6.0-11.0. The data represents the average of multiple experiments.




Figure 7: Lineweaver-Burk plots of initial reaction velocities ($\mu\text{moles}/\text{min}$) for (A) *Vigna catjang* cotyledon and (B) buffalo liver arginase measured at various concentrations of arginine.

