

Pyrazoline Stereochemistry: Use of Nuclear Magnetic Resonance Shift Reagents with Azo-compounds

By STEPHEN R. WILSON* and (the late) RICHARD B. TURNER

(Department of Chemistry, California Institute of Technology, Pasadena, California 91106)

Summary Eu(fod)₃ shifts the n.m.r. spectra of *cis*-azo-compounds, whereas *trans*-azo-compounds remain almost unaffected; shifts obtained from pyrazolines have been related to their stereochemistry.

We report the results of a study on the effect of the shift reagent Eu(fod)₃ on azo-compounds. Although the use of such n.m.r. shift reagents is commonplace, only passing

reference has been made to their use with pyrazolines¹ and several reports have appeared claiming azo-compounds do not complex.^{2,3} In the course of a synthesis of the sesquiterpene marasmic acid,⁴ we determined the stereochemistry of pyrazoline intermediates. In order to determine the effect of Eu(fod)₃ on azo-compounds, a study of *trans*-azobenzene (**1**), *cis*-azobenzene (**2**), azo-2-methylpropane (**3**), and 1,4-dimethyl-2,3-diazabicyclo[2,2,2]oct-2-ene (**4**) was

undertaken. We found that the *cis*-azo-compounds (2) and (4) complex strongly with the reagent, whereas the *trans*-azo-compounds (1) and (3) remain virtually unaffected. The n.m.r. spectrum of the *cis*-azobenzene (2), for example, shows the *ortho*-protons shifting to a multiplet at δ 13.2 with one equivalent of $\text{Eu}(\text{fod})_3$. When three pyrazoline intermediates were studied, the lines for the shifts of the methylene AB quartet adjacent to the azo-linkage possess identical slopes and the greatest shifts. Equivalent environments of the azo-linkage in the three compounds

can be inferred by the similarly identical slopes of other corresponding protons. The *cis*-azo-linkage complexes less strongly than OH but more strongly than CO or CO_2Me .

We thank the Robert A. Welch Foundation, Houston, Texas, for support of this work, and Dr. Paul Engel, Rice University, Houston, Texas, for gifts of compounds (3) and (4).

(Received, 30th April 1973; Com. 616.)

¹ R. G. Bergman and R. A. Keppel, *J. Amer. Chem. Soc.*, 1972, **94**, 1350.

² C. Beate, Z. W. Wolkowski, and N. Thoai, *Chem. Comm.*, 1971, 700.

³ R. E. Rondeau, M. A. Berwick, R. N. Steppel, and M. P. Serve, *J. Amer. Chem. Soc.*, 1972, **94**, 1096, footnote 16.

⁴ S. R. Wilson and R. B. Turner, *J. Org. Chem.*, in the press.