

# QTL hotspots in eggplant (*Solanum melongena*) detected with a high resolution map and CIM analysis

Amy Frary · Anne Frary · Marie-Christine Daunay ·  
Koen Huvenaars · Rolf Mank · Sami Doğanlar

Received: 30 October 2013 / Accepted: 29 December 2013 / Published online: 18 January 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Fifty-eight F<sub>2</sub> individuals derived from an interspecific cross between cultivated eggplant, *Solanum melongena*, and its wild relative, *S. linnaeanum*, were phenotyped for 42 plant, leaf, flower, and fruit traits. Composite interval mapping analysis using genotypic data from 736 molecular markers revealed the positions of 71 statistically significant ( $P \leq 0.05$ ) quantitative trait loci (QTL) influencing 32 of the morphological traits. Although most QTL were location-specific, QTL governing three traits (leaf lobing, leaf prickles and prickle anthocyanin) were detected in both experimental locations. Analysis of three additional traits (stem prickles, fruit calyx prickles and fruit length) in both locations yielded

QTL in similar but non-overlapping map positions. The majority (69 %) of the QTL corresponded closely with those detected in previous analyses of this data set. However the increased resolution of the linkage map combined with advances in QTL mapping permitted more precise localization, such that the average interval length of these QTL was reduced by 93 %. Thirty-one percent of the QTL were novel, suggesting that simple linear regression with a low density linkage map (the method used in previous studies of this population) missed a substantial portion of significant QTL. Hotspots of QTL affecting plant hairiness, prickliness, and pigmentation were identified on chromosomes 3, 6, and 10, respectively, and may reflect the pleiotropic activity of single structural or regulatory genes at these positions. Based on synteny between the eggplant, tomato, potato and pepper genomes, putative orthologs were identified for 35 % of the QTL suggesting strong conservation of gene function within the Solanaceae. These results should make it

The localization of QTL for 32 morphological traits on the high-resolution map of the eggplant genome has allowed hotspots and putative orthologs with other solanaceous species to be identified.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10681-013-1060-6) contains supplementary material, which is available to authorized users.

A. Frary  
Department of Biological Sciences, Mount Holyoke  
College, South Hadley, MA 01075, USA

A. Frary · S. Doğanlar (✉)  
Plant Molecular Genetics Laboratory, Department of  
Molecular Biology and Genetics, Izmir Institute of  
Technology, Gulbahce Campus, Urla, Izmir 35430,  
Turkey  
e-mail: samidoganlar@iyte.edu.tr

M.-C. Daunay  
Unité de Génétique et Amélioration  
des Fruits et Légumes, INRA, Domaine Saint Maurice,  
BP 94, 84143 Montfavéd Cedex, France

K. Huvenaars · R. Mank  
Keygene N.V., Agro Business Park 90,  
6708 PW Wageningen, The Netherlands

easier to target particular loci for map-based cloning and marker-assisted selection studies.

**Keywords** *Solanum melongena* · Quantitative trait loci · Gene conservation · Solanaceae

## Introduction

The Solanaceae family of plants has long been of interest to plant geneticists both for its agricultural importance and its tractability as a model system. Members of the family are grown as vegetable crops (tomato, potato, pepper, eggplant, tomatillo, and pepino) and for ornamental purposes (*Petunia*, *Calibrachoa*, *Datura*, *Brugmansia*, *Schizanthus*, etc.). The cash crop tobacco also belongs to the family. Thus it is not surprising that some of the earliest work in gene and genome mapping as well as comparative genomics was done in the Solanaceae family. The construction of the first high-density molecular linkage map in tomato (*Solanum lycopersicum*) (Tanksley et al. 1992) was essential in establishing that species as a forerunner in the realm of quantitative trait locus (QTL) mapping and cloning (Alpert et al. 1995; Frary et al. 2000). The close relationship of tomato to potato (*S. tuberosum*), eggplant (*S. melongena*), and pepper (*Capsicum* spp.) has facilitated genome and trait mapping efforts in those species (Tanksley et al. 1992; Doganlar et al. 2002a; Livingstone et al. 1999).

Eggplant has lagged behind in the area of quantitative trait analysis in part because, in economic terms, it is a less important crop. Eggplant placed 25th in the FAO's top commodities ranking for 2010, well behind tomato (8th) and potato (13th) (FAO 2013). As a result, the first QTL mapping in eggplant was performed only 15 years ago and was limited to traits of breeder interest. The first such work involved the localization of a QTL for fruit shape on a random amplified polymorphism (RAPD) map for an intraspecific *S. melongena* F<sub>2</sub> population (derived from a cross between eggplant lines 'EPL1' and 'WCGR112-8') (Nunome et al. 1998). The addition of nearly 100 AFLP<sup>®</sup> markers to the map allowed the detection of QTL for several other traits (fruit, stem, and calyx color) (Nunome et al. 2001). More recently, Barchi et al. (2012) developed a linkage map for their intraspecific F<sub>2</sub> population derived from a cross between two breeding lines, '305E40' and '67/3'. The map included over 400 restriction fragment length polymorphism

(RFLP) and single nucleotide polymorphism (SNP) markers and enabled the localization of QTL influencing seven pigmentation traits on eight of eggplant's twelve chromosomes. Thus a survey of the eggplant literature reveals that few quantitative traits have been analyzed in intraspecific populations. This fact highlights a serious hindrance to using such populations for QTL analysis: they display limited phenotypic variability. Although maps developed from intraspecific populations may be easier to exploit for the purposes of marker-assisted selection and breeding, the higher degree of morphological diversity in interspecific populations allows examination of a greater range of plant traits.

A *S. linnaeanum* MM195 × *S. melongena* MM738 F<sub>2</sub> population was used by Doganlar et al. (2002a) to produce a molecular linkage map consisting of 233 RFLP markers at an average interval of 4.8 cM. Because *S. linnaeanum* is a prickly wild relative of eggplant that produces small, round, green, striped fruit while MM738 is a non-spiny commercial cultivar that produces large, oblong, purple fruit without striping, the F<sub>2</sub> population derived from these parents was highly polymorphic. Twenty-two domestication traits (fruit size, shape and color and plant prickliness) (Doganlar et al. 2002b) and 18 morphological traits (leaf, flower and fruit size, shape, appearance, and development) (Frary et al. 2003a) were evaluated in the population and subjected to single-point linear regression analysis. A total of 125 significant QTL were positioned on the interspecific map. Because of the relatively low resolution of the linkage map, the average length of the QTL detected in these companion studies was 35.8 cM, a fairly broad interval (considering the average linkage group length of 128 cM). In the current study, we have re-analyzed the domestication and morphological trait data for the *S. linnaeanum* × *S. melongena* F<sub>2</sub> population. Several advances in QTL mapping and eggplant genomics merited this strategy. Composite interval mapping (CIM) with marker cofactors is now the standard for QTL detection. By controlling for the effects of other markers on the trait, CIM is a more powerful method of QTL detection and provides greater accuracy in QTL localization (Zheng 1994). The development of a high-resolution map of the eggplant genome comprising over 850 AFLP, RFLP, and COSII (conserved ortholog set) markers at an average spacing of 1.8 cM (Doganlar et al. in press) further enhances our ability to refine the positions of the QTL controlling the

domestication and morphological traits. More precise locations of these loci are essential for marker-assisted selection and/or map-based cloning. In addition, they allow more detailed comparisons with QTL mapping studies in other solanaceous species. Such comparisons can help identify putative orthologs within the family, thereby shedding light on the evolutionary conservation and divergence of genomes.

## Materials and methods

### Plant population

The mapping population of 58 F<sub>2</sub> individuals was generated from a cross between *S. linnaeanum* MM195 and *S. melongena* MM738 made by M.-C. Daunay at Institut National de la Recherche Agronomique, France. The female parent, *S. linnaeanum* Hepper & Jaeger ‘MM195’, is a spiny wild relative that produces small, round, reticulate green fruit. The male parent, *S. melongena* L. ‘MM738’, is a non-spiny European commercial type that bears large, oblong, purple fruit. The F<sub>2</sub> plants were grown in the greenhouse in Ithaca, NY. Rooted vegetative cuttings were sent to Montfavet, France (FR), for field evaluations. Two plants of each genotype were planted at a single stake with 1 meter row spacing between genotypes. Replicates of the parental controls were also grown at both locations.

### Phenotype evaluations

Greenhouse-grown F<sub>2</sub> plants and controls were evaluated in Ithaca, New York (NY), during spring 1999. Field-grown plants were scored in Montfavet, FR, during July–October, 2000. A total of 42 plant, leaf, flower, and fruit traits were evaluated as described below (summarized in Online Resource 1).

Plant height (ht) in centimeters was determined in FR at the beginning of August. The number of days to first flowering (dtf) was counted in NY from the date of greenhouse transplanting until the opening of the first flower. The number of flowers per inflorescence (fn) was determined in FR at various times during the growing season and at several locations on the plant. The mean for each genotype was then used in the analysis. The mean number of fruit per infructescence (ftn) was determined in a similar way. Fruit set (fset)

was an overall measure of fertility evaluated on a 0 (no fruit) to 5 (many fruit) scale.

Hairiness of vegetative plant parts was determined on a scale of 0 (no hairs) to 5 (very many hairs) in FR. The hairiness of plant apices (ah), leaves (lh), and stems (sh) was assessed by eye. The presence/absence of ovary hairs (ovh) was determined by microscopic examination of approximately three ovaries per genotype in the NY material. Fruit glossiness (fglo) was measured in FR on a scale of 1 (dull epidermis) to 3 (glossy epidermis).

Prickliness of leaves (lp), stems (sp) and fruit calyxes (ftcp) were assessed on a 1–5 scale in NY (1, no prickles; 5, many prickles) and a 0–5 scale (0, no prickles; 5, many prickles) in FR. Flower calyx (flcp) and petiole (pp) prickliness were evaluated only in FR using the aforementioned scale.

Anthocyanin content of leaf laminae (lla), stems (sa), and prickles (pa) was scored on a 1 (green) to 3 (dark purple) scale in NY and a 0 (green) to 5 (dark purple) scale in FR. Leaf rib (lra) and flower corolla (ca) anthocyanin was assessed only in FR. Three separate fruit color traits were evaluated in both locations. Fruit anthocyanin presence (fap) recorded whether the fruits were green or purple. Fruit anthocyanin intensity (fai) scored the degree of pigmentation in the purple fruits only on a scale of 1 (light purple) to 3 (dark purple). Fruit stripe (fst) measured the secondary color repartition in the fruit as presence/absence in NY and a 1–3 scale in FR (1, no stripes; 2, irregular striping; 3, uniform reticulate striping). Two additional fruit color traits were scored in FR. Fruit chlorophyll netting (fcn) assessed the pattern of chlorophyll distribution in the fruit on a 1–3 scale in FR (1, no reticulation; 2, irregular reticulation; 3, uniform reticulation). Anthocyanin under the calyx (auc) was scored as a presence (1), absence (0) trait and served as an indirect measure of the sensitivity of fruit anthocyanin synthesis to light.

Size and shape parameters were evaluated for leaves, flowers, and fruit. Leaf width (lw) and length (ll) of 12 leaves per genotype were measured (in cm) in the early autumn in FR. The ratio between leaf length and width (ll/lw) was designated as leaf shape (lsh). Two traits described the appearance of leaves. Leaf lobing (llob) was scored on a scale of 1 (very weak lobing) to 5 (very strong lobing) in both FR and NY. Leaf surface appearance (lsur) was evaluated on a similar scale (1 = smooth leaf, 5 = strongly wrinkled

leaf) in FR. For the flower traits, approximately 12 inflorescences per genotype were harvested between July and October in FR; only the main flower of each inflorescence was measured. Flower diameter (fld) was measured in mm, and flower shape (fls) was assessed on a 1 (orbicular) to 5 (star-shaped) scale. Ovary length (ovl), diameter (ovd), shape (ovs) and area (oa) were evaluated in NY by measuring (in mm) transverse sections of ovaries harvested at anthesis. Ovary locule number (oln) was determined using transverse sections. In general, three ovaries were measured from each genotype for these trait analyses. Five representative fruits in NY and FR were harvested just prior to physiological ripeness for the analysis of fruit traits. Fruit weight (fw) was measured in grams. Fruit length (fl) and diameter (fd) were measured in cm. Fruit shape (fs) was the ratio of length to diameter (fl/fd) such that round fruit had a shape index of 1, oblate fruit had an index < 1, and oblong fruit had an index > 1. Fruit calyx size (cs) in FR was scaled according to the proportion of the fruit covered by the calyx (1 = very short calyx, <10 % of the fruit length covered; 5 = very long calyx, >75 % of fruit length covered).

QGene (Nelson 1997) was used to calculate correlation coefficients between traits.

### Genotype evaluations

Molecular marker analysis and construction of the high-density eggplant map are described in Doganlar et al. (in press). A total of 736 AFLP, RFLP and COSII (conserved ortholog set) markers were used for QTL analysis. QGene version 4.0 (Joehanes and Nelson 2008) was used to map QTL. CIM (a method that combines interval mapping with multiple regression analysis) with automatic forward cofactor selection and a scan interval of 0.1 cM was used for QTL detection. A genome-wide critical threshold value for an experiment-wise type I error rate,  $\alpha = 0.05$  and  $\alpha = 0.01$ , was set by 1,000 random permutations of the trait data (Churchill and Doerge 1994). The percentages of phenotypic variance explained (PVE) were obtained from the generalized  $R^2$  values (Nagelkerke 1991) calculated by QGene. Trait means, and gene actions ( $d/a$ ) were determined for each significant QTL using the CIM results. QTL detected in the present study were compared to those identified in the previous two studies conducted in this population

(Doganlar et al. 2002b; Frary et al. 2003a). Because these QTL were localized on a lower resolution version of the interspecific eggplant map (Doganlar et al. 2002a), the map positions and relative lengths of these previously identified QTL on the current molecular map (Doganlar et al. in press) were determined using shared RFLP markers as anchors.

Quantitative trait loci with 95 % confidence intervals were drawn on the molecular linkage map of eggplant using MapChart 2.2 (Voorrips 2002). To identify QTL hotspots (clusters), a 20 cM sliding window was advanced in 5 cM increments across the linkage map. The number of QTL co-localizing within the window at each position in the genome was recorded with regions containing more than three QTL qualifying as hotspots.

## Results

### Phenotypic variation

A total of 42 plant, leaf, flower, and fruit traits were analyzed. The trait means are summarized in Online Resource 2. Fourteen of the traits were evaluated in both locations, 21 in FR only, and seven in NY only (Online Resource 1). The phenotypic distributions of the fourteen traits were compared across the two locations (data not shown). The majority of the traits showed a similar pattern, however some of the anthocyanin traits (sa, lla, fai, fst) tended to skew toward higher values in FR. This is not surprising as anthocyanin synthesis is closely tied to environmental conditions such as light intensity and temperature both of which are expected to be quite different for field-grown plants in FR as compared to greenhouse-grown plants in NY.

### Correlations between traits

Significant ( $P \leq 0.05$ ) positive correlations existed between all of the traits measured in both FR and NY. These traits included fruit size and shape parameters (fruit weight, fruit shape, fruit diameter, fruit length), plant pigmentation (stem, prickle, leaf lamina anthocyanin, fruit anthocyanin presence and intensity, and fruit stripe) and prickle (stem, leaf, fruit calyx) traits as well as leaf lobing. Correlation coefficients ranged from 0.43 for fruit anthocyanin

intensity (fai) to 0.96 for fruit shape (fs), with an average value of 0.79.

A number of significant correlations were observed between traits measured at a single location. Those relationships are described in the following paragraphs and are summarized in Online Resource 2.

All of the prickle traits (sp, lp, pp, flcp, ftcp) were strongly associated with each other ( $r = 0.70\text{--}0.92$ ). Relationships between these prickle traits and several other morphological measures were also detected. Plant organ prickliness was positively correlated with leaf lobing (llob) ( $r = 0.59\text{--}0.83$ ) as well as the anthocyanin content of stems (sa) ( $r = 0.29\text{--}0.38$ ) and leaf laminae (lla) ( $r = 0.35\text{--}0.42$ ). Significant associations were seen between most of the prickliness traits and the hairiness of stems (sh) ( $r = 0.28\text{--}0.29$ ) and between leaf prickles and ovary hairs (ovh) ( $r = 0.39$ ) as well as the size of calyxes (cs) ( $r = 0.32\text{--}0.56$ ). In addition, associations were found between both flower (flcp) and fruit (ftcp) calyx prickliness and corolla pigmentation (ca) ( $r = 0.47\text{--}0.54$ ). A strong negative relationship existed between prickliness and fruit glossiness (fglo) ( $-0.49 \leq r \leq -0.64$ ). Stem prickles (sp) were also negatively correlated with aspects of fruit size, namely fruit diameter (fd) and length (fl) ( $r = -0.36$  to  $0.38$ ).

While the traits assessing the hairiness of vegetative organs (ah, lh, sh) were strongly correlated with each other ( $r = 0.71\text{--}0.85$ ), no association with ovary hairiness (ovh) was found. Similarly, while ah, lh, and sh were negatively correlated with fruit glossiness ( $-0.39 \leq r \leq -0.50$ ), ovh was unrelated to fglo. Negative relationships between fglo and two shape parameters, llob ( $r = -0.41$ ) and fruit shape (fs) ( $r = -0.28$ ), were observed. While fruit diameter (fd), a key determinant of fruit shape (defined as fruit length/fruit diameter) was associated with fglo ( $r = 0.36$ ), fruit length (fl) was not.

Fruit length (fl) and diameter (fd) were strongly correlated ( $r = 0.84\text{--}0.87$ ), however only fl was significantly associated with fruit shape ( $r = 0.44\text{--}0.48$ ). Not surprisingly, a similar relationship was seen in ovaries; ovary length (ovl) and diameter (ovd) were highly correlated with each other ( $r = 0.85$ ) and ovs was significantly related to ovl only ( $r = 0.34$ ). Interestingly, a positive association between calyx size and fs was seen ( $r = 0.40$ ) such that more oblong fruit tended to have larger calyxes.

Most parameters of leaf size were unrelated. And, while a strong negative association between lw and leaf shape (lsh, defined as ll/lw) existed ( $r = -0.70$ ), ll and lsh were not significantly correlated.

All of the traits assessing pigmentation levels in vegetative tissues [stems (sa), leaf ribs (lra), leaf laminae (lla), and prickles (pa)] and flower corollas (ca) were positively correlated ( $r = 0.36\text{--}0.94$ ). Fruit anthocyanin presence (fap) was correlated with the other pigment traits however fruit anthocyanin intensity (fai) was less reliably associated with the other color traits, showing no significant connection with sa, pa or ca. Most of the anthocyanin traits were positively correlated with plant height (ht) ( $r = 0.31\text{--}0.38$ ) however, fruit stripe (fst) was negatively correlated with ht ( $r = -0.32$ ). Negative associations also existed between fst and other aspects of plant growth, namely flower number (fln) ( $r = -0.31$ ) and fruit set (fset) ( $r = -0.38$ ).

Fruit set (fset) was significantly correlated with fruit and flower traits, including fln ( $r = 0.28$ ), ftn ( $r = 0.61$ ) fl ( $r = 0.29$ ) and fw ( $r = 0.29$ ). The correlation coefficient between flower and fruit number was 0.46.

## QTL analysis

A genetic map consisting of 736 AFLP, RFLP, and COSII markers was used for CIM of QTL. Logarithm of odds (LOD) thresholds for QTL declaration were calculated by 1,000 permutations of the data for each trait. The mean experimental LOD thresholds were 4.92 and 5.91 at the 5 and 1 % significance levels, respectively. Seventy-one statistically significant ( $P \leq 0.05$ ) QTL impacting 32 traits mapped to 11 of eggplant's 12 linkage groups (Table 1; Online Resource 3). Seventy-five percent of these QTL met or exceeded the LOD threshold at the 1 % level of significance. The average number of QTL identified per trait was 2.2. Fruit length (fl) and apex hairs (ah) yielded the greatest number of QTL: five each. The average number of QTL on each linkage group was 5.9, with linkage group 6 having the most (13 QTL) and linkage group 8 having the fewest (0 QTL). The size of the QTL ranged from 0 to 19.3 cM, with a mean QTL length of 3.4 cM. The percentage of phenotypic variation explained (PVE) by the QTL varied from a low of 33 % to a high of 100 %; the mean phenotypic trait variance was 55 %.

**Table 1** QTL detected in the *S. melongena* × *S. linnaeanum* F<sub>2</sub> population across two locations

Trait	QTL <sup>a</sup>	Chr.	Location	Position (cM) <sup>b</sup>	Marker interval/nearest marker	Peak LOD	Peak R <sup>2</sup>	Additive effect <sup>c</sup>
Leaf length (ll)	ll11.1	11	FR	122.3 (121.3–123.1)	CT175	4.83*	0.34	Sl
Leaf width (lw)	lw1.1	1	FR	69.8 (69.2–70.9)	TG59-P15/M47-063.34	6.43**	0.43	Sm
Leaf shape (lsh)	lw4.1	4	FR	53.4 (53.1–53.5)	T0877-E38/M60-216.57	5.51*	0.38	Sm
	lsh1.1	1	FR	69.8 (69.2–70.9)	C2A14g00740-P15/M47-063.34	9.84**	0.58	Sl
Leaf lobing (llob)	lsh5.1	5	FR	58.2 (57.9–58.4)	P15/M47-384.89-E35/M60-283.89	9.27**	0.55	Sm
	llob5.1	5	NY	24 (19.7–39)	P15/M47-117.26-CT151	11.88**	0.64	Sl
	llob6.1	6	NY	72.2 (70–72.8)	TG240-P14/M47-370.34	6.22**	0.42	Sm
	llob6.2	6	FR	106 (103.8–106.3)	CT193-TG279	16.64**	0.76	Sl
Ovary length (ovl)	llob6.2	6	NY	106 (103.8–106.3)	CT193-TG279	29.50**	0.92	Sl
	llob7.1	7	NY	62.7 (61.5–63.3)	E32/M59-313.82-C2A15g56940	8.32**	0.52	Sl
	ov1.1	1	NY	81.5 (79.8–82.2)	TG607-TG83	6.51**	0.57	Sl
	ov9.1	9	NY	69.8 (67.5–71.7)	TG390-TG404	5.47**	0.50	Sl
Ovary diameter (ovd)	ovd9.1	9	NY	95.8 (91.3–96.1)	T0793-C2A1g07310	8.56**	0.67	Sl
Ovary area (oa)	oa6.1	6	NY	48 (46.7–48.4)	E39/M59-214.94-E38/M60-255.77	7.16**	0.60	Sm
	oa11.1	11	NY	119.5 (119.1–119.8)	E32/M59-074.02-P15/M47-098.03	8.61**	0.67	Sm
Locule number (oln)	oln5.1	5	NY	40.2 (40–40.6)	E44/M54-097.34	6.35*	0.58	Sm
Fruit length (fl)	fl1.1	1	FR	0	C2A15g51970	4.97*	0.36	Sm
	fl2.1	2	NY	75.4 (75.2–75.6)	DFR-P15/M47-275.63	5.41*	0.38	Sm
Fruit shape index (fs)	fl2.2	2	FR	79.2 (77.9–81.3)	TG140-C2A15g67370	6.58**	0.44	Sm
	fl7.1	7	NY	62.5 (62–62.5)	E32/M59-293.77	6.09**	0.42	Sm
	fl9.1	9	NY	48.8 (46.9–50.1)	E38/M60-099.47-C2A15g58410	7.28**	0.48	Sm
	fs7.1	7	NY	93.6	E38/M60-263.34	4.63*	0.34	Sm
Fruit weight (fw)	fw1.1	1	FR	0 (0–2.5)	C2A15g51970-T0343	7.83**	0.50	Sm
	fw2.1	2	FR	78 (77.5–79.1)	TG140	5.45**	0.38	Sm
Fruit stripe (fst)	fw9.1	9	NY	46.1 (45.4–46.5)	E38/M60-099.47	4.95*	0.53	Sm
	fw9.2	9	NY	57 (56.9–57)	P15/M47-147.62-E35/M60-390.17	5.21*	0.55	Sm
	fst4.1	4	FR	28.6 (25.6–33.7)	E32/M59-181.88-E44/M54-284.84	13.23**	0.69	Sl
	fst10.1	10	FR	60.7 (60.2–60.9)	CT199	6.48*	0.44	Sm
Fruit chlorophyll netting (fcn)	fcn3.1	3	FR	57.4 (57.2–57.4)	E35/M60-567.26	5.81*	0.40	Sl
	fcn4.1	4	FR	28.6 (25.6–34.6)	E32/M59-181.88-E44/M54-284.84	17.14**	0.78	Sl

**Table 1** continued

Trait	QTL <sup>a</sup>	Chr.	Location	Position (cM) <sup>b</sup>	Marker interval/nearest marker	Peak LOD	Peak R <sup>2</sup>	Additive effect <sup>c</sup>
Fruit glossiness (fglo)	fgl01.1	1	FR	47.7 (47–48.8)	C2At3g23590-P14/M47-226.40	8.38**	0.52	Sm
	fgl06.1	6	FR	88.2 (88.2–89.6)	CT204-TG446	8.83**	0.54	Sm
Flowers/fin	fgl09.1	9	FR	89 (88.2–89.7)	TG248	5.13*	0.37	Sm
	fn3.1	3	FR	120.2	TG284	4.48*	0.33	Sm
	fn4.1	4	FR	67.9 (66.4–68)	E39/M59-340.73-TG62	8.78**	0.54	Sm
	fn3.1	3	FR	65.1	TG130	4.47*	0.33	Sm
Fruit/fin	fn9.1	9	FR	33.3 (32.3–33.5)	C2At2g37240	5.11*	0.36	Sm
	ah2.1	2	FR	50.6 (50.5–57.2)	C2At1g11430-P12/M61-118.74	8.81**	0.54	Sm
Apex hairs (ah)	ah3.1	3	FR	23.3 (23.3–38.7)	TG585-	13.89**	0.70	Sl
	ah3.2	3	FR	68.4 (68.1–70.8)	C2At1g72030-E32/M59-401.16	6.28**	0.42	Sl
	ah3.3	3	FR	105.9 (103–109.5)	TG442-C2At5g08050	9.28**	0.55	Sm
	ah7.1	7	FR	62 (60.3–65.2)	E44/M50-357.60-E32/M59-293.77	7.63**	0.49	Sm
Stem hairs (sh)	sh3.1	3	FR	23.3 (23.3–31.3)	TG585	6.21**	0.42	Sl
	sh10.1	10	FR	68	E44/M50-268.47	6.63**	0.44	Sm
Leaf hairs (lh)	lh3.1	3	FR	23.3 (23.3–38.4)	TG585	9.05**	0.54	Sl
	lh9.1	9	FR	16.4 (14.2–19)	TG18	5.52**	0.38	Sm
Ovary hairs (ovh)	lh10.1	10	FR	93.8 (93.8–95.4)	E38/M60-273.03	5.69**	0.39	Sm
	ovh10.1	10	NY	143.3 (142.7–149.3)	TG420-CD72B	8.29**	0.65	Sm
Stem prickles (sp)	sp1.1	1	NY	56.8 (55.8–59)	C2At5g13700-T1241	7.22**	0.47	Sm
	sp3.1	3	NY	23.3 (23.3–40)	TG585	6.87**	0.45	Sl
	sp6.1	6	NY	102.1 (98.8–102.1)	TG292-CT146	12.00**	0.65	Sl
	sp6.2	6	FR	106 (103.8–106.3)	CT193-TG279	15.26**	0.73	Sl
Leaf prickles (lp)	lp2.1	2	FR	25.7 (24.7–26.3)	CD11-C2At2g34560	5.98**	0.41	Sl
	lp3.1	3	NY	23.3 (23.3–36.1)	TG585	10.28**	0.59	Sl
	lp6.1	6	NY	10.9 (9.7–12.3)	CT119	4.84*	0.34	Sm
Petiole prickles (pp)	lp6.2	6	FR	106 (103.8–106.3)	CT193-TG279	23.36**	0.87	Sl
	lp6.2	6	NY	106 (103.8–106.3)	CT193-TG279	22.35**	0.86	Sl
	pp2.1	2	FR	35.5 (33.1–39.4)	TG554-CT277	5.23**	0.37	Sl
	pp6.1	6	FR	106 (103.8–106.3)	CT193-TG279	18.09**	0.79	Sl
Flower calyx prickles (fcp)	fcp6.1	6	FR	103.6 (102.1–106)	CT146-CT106	14.27**	0.72	Sl
Fruit calyx prickles (fcp)	fcp6.1	6	FR	102.1 (98.8–102.1)	TG292-CT146	11.83**	0.66	Sl
	fcp6.2	6	NY	106 (103.8–106.2)	CT193-CT109	8.38**	0.69	Sl

Table 1 continued

Trait	QTL <sup>a</sup>	Chr.	Location	Position (cM) <sup>b</sup>	Marker interval/nearest marker	Peak LOD	Peak R <sup>2</sup>	Additive effect <sup>c</sup>
Stem anthocyanin (sa)	sa6.1	6	NY	72.8 (72.6–73.3)	C2A4g10030-P14/M47-370.34	6.06**	0.41	SI
	sa10.1	10	FR	109.8 (109.6–110.8)	CT240	5.21*	0.36	Sm
	sa10.2	10	NY	121.2	CT124	4.55*	0.33	Sm
Prickle anthocyanin (pa)	pa10.1	10	FR	109.2 (107–109.8)	CT214-CT240	20.61**	0.89	Sm
	pa10.1	10	NY	109.8 (104.9–111.1)	T0838-CT240	13.36**	0.69	Sm
Leaf rib anthocyanin (lra)	lra10.1	10	FR	110.5 (109.3–113.1)	TG233-TG63	16.74**	0.77	Sm
Leaf lamina anthocyanin (lla)	lla10.1	10	FR	109.8 (121.3–123.1)	TG233-TG63	10.53**	0.60	Sm
Corolla anthocyanin (ca)	ca5.1	5	FR	73.3 (72.3–73.4)	T0612	5.02*	0.36	Sm
Fruit anthocyanin intensity (fai)	<i>fai11.1</i>	11	NY	123.7	CT175	6.34**	0.69	SI
	fai12.1	12	NY	0.2 (0–1.8)	CT79-T0076	10.57**	0.86	SI
Fruit anthocyanin presence (fap)	fap10.1	10	FR	106.4	C2A13g08760	22.69**	0.87	SI
	fap10.2	10	FR	109.8 (109.3–112.8)	TG233-CT240	223.54*	1.00	Sm
						2223.54**		

<sup>a</sup> Newly detected QTL are italicized

<sup>b</sup> Peak position of QTL on linkage group and confidence interval given in parenthesis

<sup>c</sup> Whether alleles from *S. melongena* (Sm) or *S. limmaeanum* (SI) parent increase trait

\*\* Significant at 1 %

\* Significant at 5 %



For the traits assessed in both locations, three QTL in corresponding positions were detected in NY and FR: *llob6.2*, *lp6.2*, and *pa10.1*. Analysis of three other traits in both locations yielded QTL in similar but non-overlapping map positions: *fl2.1/fl2.2*, *sp6.1/sp6.2*, *ftcp6.1/ftcp6.2*. However, of the 23 QTL associated with traits measured in both NY and FR, the majority were specific to a single location.

Twenty-two of the QTL (31 %) identified in this study were novel, not reported in previous QTL analyses of this interspecific population. The map locations of the 49 remaining QTL corresponded fairly closely to those of QTL detected by Doganlar et al. (2002b) and Frary et al. (2003a) using simple linear regression analysis. The peak position of each of these QTL shifted an average of  $\pm 6.8$  cM as compared to the map position of its previously identified counterpart. Average interval size changed dramatically with the new analysis: these common QTL averaged 2.8 cM in length as compared to 39.2 cM in the previous study.

#### Leaf size and shape traits

A single significant QTL explaining 34 % of the phenotypic variation in leaf length was identified on chromosome 11 (*ll11.1*). Interestingly, *S. linnaeanum* alleles at this locus contributed toward increasing leaf length, an effect opposite to that expected.

Leaf width was influenced by QTL on chromosomes 1 and 4. *lw1.1* and *lw4.1* accounted for 43 and 38 % of the phenotypic variation in leaf width, respectively. *S. melongena* alleles at both loci increased leaf width as expected. The QTL *lw4.1* was previously undetected in this population.

Quantitative trait loci on chromosomes 1 and 5 affected leaf shape, the ratio between leaf length and width. Individually, *lsh1.1* and *lsh5.1* explained 58 and 55 %, respectively, of the variation in leaf shape. Alleles at *lsh1.1* had effects opposite to those predicted based on the parental phenotypes.

Four QTL on chromosomes 5, 6 and 7 impacted the degree of leaf lobing. Of the two loci identified on chromosome six, *llob6.1* was identified in NY and *llob6.2* was identified in both growing environments (NY and FR). Interestingly, *S. melongena* alleles at *llob6.1* had the unexpected effect of increasing lobing. The largest phenotypic effects on the trait were seen at *llob6.2* (92 % PVE). *llob5.1* and *llob7.1* were

previously undetected in the NY material. The more significant of these loci (*llob5.1*, 64 % PVE) behaved in an additive manner.

No significant QTL were detected for leaf surface appearance (*lsur*).

#### Flower size and shape traits

Ovary length was affected by two QTL, *ov11.1* and *ov19.1*, which accounted for 57 and 50 %, respectively, of the phenotypic variation in the trait. *S. linnaeanum* alleles at both loci had the unexpected effect of increasing ovary length with *ov11.1* alleles behaving in an additive manner. *ov19.1* was previously undetected in this population.

One QTL (*ovd9.1*) was associated with 67 % of the variation in ovary diameter. As expected, *S. linnaeanum* alleles at this locus contributed toward wider ovaries.

Ovary area, as calculated from transverse sections, was controlled by two QTL, *oa6.1* and *oa11.1* which explained 60–67 % of the phenotypic variation in the trait. Parental alleles at both loci behaved in a predictable fashion with *S. melongena* alleles tending to augment ovary size.

*Solanum melongena* alleles at a single QTL increased ovary locule number. This QTL, *oln5.1*, accounted for 58 % of the variation in the trait.

No significant QTL were detected for flower diameter (*fld*), flower shape (*fls*) or ovary shape (*ovs*).

#### Fruit size and shape traits

Five QTL impacting fruit length were identified, two in FR (*fl1.1* and *fl2.2*) and three in NY (*fl2.1*, *fl7.1*, and *fl9.1*). *fl1.1* and *fl7.1* were previously undetected in this population. Each locus accounted for between 36 and 48 % of the variability in the length of the measured fruit. The two QTL on chromosome 2, *fl2.1* and *fl2.2*, lay within 4 cM of one another. Because the cultivated parent produces oblong fruit, it was not surprising that *S. melongena* alleles at all five loci were associated with longer fruit.

Fruit shape in the NY-grown plants was influenced by a QTL located on chromosome 7 (*fs7.1*) which explained 34 % of the variability in the trait. Parental alleles at this locus behaved in a predictable manner. No fruit shape QTL was detected in FR.

A total of four individual fruit weight QTL were identified, two (*fw1.1* and *fw2.1*) in FR and two (*fw9.1*

and *fw9.2*) in NY. With the exception of *fw2.1* (38 % PVE), each locus accounted for 50–55 % of the variability in fruit weight. As predicted, alleles from the cultivated parent increased fruit weight.

No significant QTL were identified for fruit diameter (fd) and calyx size (cs).

### Fruit appearance traits

Fruit stripe in FR was controlled by QTL on chromosomes 4 and 10. *S. linnaeanum* alleles at *fst4.1* increased the degree of striping in fruit whereas *S. melongena* alleles had the effect of increasing fruit stripe at *fst10.1*. *fst4.1* had a larger impact on phenotype, affecting 69 % of the trait while *fst10.1* explained 44 % of the variation in striping. No significant QTL influencing fruit stripe were detected in the NY-grown population. Fruit chlorophyll netting (fcn) was also affected by two QTL. *fcn4.1* overlapped with *fst4.1* and accounted for 78 % of the phenotypic variation in the trait. The locus on chromosome 3, *fcn3.1*, explained 40 % of the variability in the trait. *S. linnaeanum* alleles increased fruit chlorophyll reticulation at both loci.

Quantitative trait loci associated with fruit glossiness were detected on three chromosomes. Each of the QTL on chromosomes 1 and 6, *fgl1.1* and *fgl6.1* explained around 50 % of the total variation in the trait. Alleles at *fgl9.1* behaved in an additive manner and accounted for 39 % of the variability in fruit glossiness. At all three loci, *S. melongena* alleles enhanced fruit glossiness.

### Plant traits

Two QTL influenced the number of flowers per inflorescence, *fln3.1* and *fln4.1*. *fln4.1* accounted for a greater amount of the phenotypic variation in the trait (54 %) than *fln3.1* (33 %). *S. melongena* alleles at both loci increased flower number, an unexpected effect given the parental phenotypes.

Fruit number per infructescence was determined by QTL on chromosomes 3 and 9. These loci, *fnt3.1* and *fnt9.1* explained 33–36 % of the phenotypic variation in fruit number. Alleles from the cultivated parent increased fruit number at both loci.

Five QTL affecting apex hairs were detected. These included three loci on chromosome 3, and one each on

chromosomes 2 and 7. The most significant of the loci, *ah3.1* explained 70 % of phenotypic variation. The remaining loci (*ah2.1*, *ah3.2*, *ah3.3*, and *ah7.1*) were not previously identified in this population and accounted for 42–55 % of the phenotypic variation in degree of hairiness at the apex. *S. melongena* alleles at *ah2.1*, *ah3.3* and *ah7.1* led to hairier apices whereas *S. linnaeanum* alleles had that effect at *ah3.1* and *ah3.2*.

Stem hairiness was controlled by loci on chromosomes 3 and 10. Each locus explained around 40 % of the phenotypic variation in stem hairiness, however, *S. linnaeanum* alleles increased the trait at *sh3.1* but had the opposite effect at *sh10.1*.

Three QTL were associated with leaf hairiness, *lh3.1*, *lh9.1*, and *lh10.1*. The most significant of these loci was *lh3.1* (54 % PVE); the magnitude of effect at *lh9.1* and *lh10.1* was around 38 %. Interestingly, *S. melongena* alleles at the other two loci contributed toward hairier leaves. *lh9.1* was previously undetected in this population.

A single QTL for ovary hairiness was identified on chromosome 10. *ovh10.1* explained 65 % of the phenotypic variation in ovary hairiness. While *S. linnaeanum* ovaries are typically hairier than those of cultivated eggplant, wild species alleles at *ovh10.1* decreased ovary hairiness.

The degree of prickliness of the stem was controlled by four QTL on three chromosomes. Three of these, *sp1.1*, *sp3.1*, *sp6.1*, were identified in the NY-grown plants. While the fourth QTL, *sp6.2*, was specific to FR, its peak LOD position was within 4 cM of *sp6.1*. These two chromosome 6 loci all had a similar magnitude of effect (65–73 % PVE). *sp1.1* and *sp3.1* were previously undetected in this population. Both explained around 46 % of the phenotypic variation, however, *S. melongena* alleles at *sp1.1* increased stem prickliness.

Four QTL influenced leaf prickliness. The most major of these, *lp6.2* (~86 % PVE), was detected in both NY and FR. An additional locus was associated with leaf prickliness in FR: *lp2.1* (41 % PVE) was not previously detected in this population. Two other novel loci, *lp3.1* and *lp6.1*, were identified in the NY material and accounted for 59 and 34 %, respectively, of the phenotypic variability in leaf prickliness. *S. linnaeanum* alleles at all loci but one (*lp6.1*) acted to increase prickliness.

Petiole prickliness in FR was determined by QTL on chromosomes 2 and 6. The locus on chromosome 6,

*pp6.1* explained 79 % of the variability in petiole prickliness. While *pp2.1* had a smaller effect on phenotype (37 % PVE), it represented a novel QTL for this trait. Petiole prickliness at both loci was enhanced by wild parent alleles. No QTL were significantly associated with petiole prickliness in NY.

A single flower calyx prickliness QTL, *flcp6.1*, was identified in the FR-grown population only. Seventy-two percent of the phenotypic variation in flower calyx prickliness was ascribed to this locus. Prickliness of the fruit calyx was also controlled by chromosome 6 loci. While *flcp6.1* was specific to FR and *flcp6.2* to NY, the peak LOD positions of these two loci were within 4 cM of each other and they had a similar magnitude of effect (66 and 69 % PVE, respectively). *S. linnaeanum* alleles were responsible for increasing calyx prickliness at all of these loci.

No significant QTL were detected for plant height (ht), days to flowering (dtf), or fruit set (fset).

#### Anthocyanin traits

Three QTL on chromosomes 6 and 10 influenced stem anthocyanin levels. The effects of these loci on the phenotypic variation in stem pigmentation were similar, ranging from 33 to 41 % PVE. However, *sa6.1* and *sa10.2* were specific to NY and *sa10.1* was detected in FR only. *S. linnaeanum* alleles unexpectedly increased stem anthocyanin levels at *sa6.1*.

Prickle anthocyanin levels were determined by a single chromosome 10 QTL in both NY and FR however the phenotypic effects of *pa10.1* were greater in FR (89 % PVE) than NY (69 % PVE). Cultivated parent alleles at this locus enhanced prickle pigmentation.

Leaf rib and leaf lamina anthocyanin levels in FR were associated with the same region of chromosome 10. The QTL impacting leaf rib pigmentation (*lra10.1*) explained 80 % of the phenotypic variation whereas that for leaf lamina pigmentation (*lla10.1*) explained only 40 % of the variability in that trait. As expected, *S. melongena* alleles increased anthocyanin levels in leaf ribs and laminae.

A QTL on chromosome 5, *ca5.1*, influenced anthocyanin levels in the corolla and explained 36 % of the variability in petal pigmentation. *S. melongena* alleles at the locus increased corolla anthocyanin levels.

Two QTL on chromosomes 11 and 12 controlled fruit anthocyanin intensity in NY. No QTL for this trait were detected in FR. *fai12.1* explained 86 % of the variability in fruit anthocyanin levels. *fai11.1* accounted for 69 % of the variability and was a novel QTL for this trait. Unexpectedly, wild parent alleles at both loci increased pigmentation in the fruit. The presence of anthocyanin in fruit was associated with two adjacent QTL on chromosome 10, *fap10.1* and *fap10.2*. While both are major QTL, explaining 87 and 100 %, respectively, of the variability in the trait, these QTL were detected in FR but not NY. Cultivated and wild parent alleles had opposite effects at these two loci.

No significant QTL were detected for the anthocyanin under the calyx (auc) trait.

#### QTL hotspots

Quantitative trait loci hotspots, defined for the purposes of this study as clusters containing more than three adjacent or overlapping QTL within a 20 cM window, were found on three linkage groups (Online Resource 3). These hotspots largely consisted of QTL for highly correlated traits. On the short arm of chromosome 3, a cluster of five QTL within a 16.7 cM interval (between map positions 23.3 and 40 cM) impacted leaf (lh), stem (sh), and apex (ah) hairiness as well as leaf (lp) and stem (sp) prickliness. Strong positive correlations existed among the three hairiness traits ( $r = 0.71\text{--}0.85$ ). Similarly, the two prickle traits had a correlation coefficient of 0.78. Correlations between the individual hair and prickle traits were weaker ( $r = 0.29$  for the association between stem hairs and leaf and stem prickles). A hotspot of seven QTL in a 18.2 cM region on the long arm of chromosome 10 (between map positions 104.9 and 123.1 cM) consisted of loci linked to aspects of pigment production in stems (sa), prickles (pa), leaves (lla and lra), and fruit (fap). Once again, correlation analysis revealed all of these traits to be significantly related to one another. The largest QTL hotspot, eight loci within a 7.5 cM region on the long arm of chromosome 6 (between map positions 98.8 and 106.3 cM), affected all five of the prickliness traits (leaf (lp), stem (sp), petiole (pp), flower calyx (flcp), and fruit calyx (ftcp) prickles) as well as leaf lobing (llob). As mentioned earlier, all of these traits were strongly associated ( $r = 0.59\text{--}0.92$ ). In addition to the hotspots, nine smaller clusters consisting of three QTL were

identified on chromosomes 1 (two overlapping clusters within the 55–75 and 65–85 cM windows on the genetic map), 2 (65–85 cM window), 6 (70–90 cM window), 7 (60–80 cM window), 9 (three overlapping clusters: 30–50, 40–60 and 50–70 cM windows), and 11 (105–125 cM window). As with the hotspots, QTL controlling correlated traits occupied most of these clusters with the majority of the loci (67 %) influencing some aspect of organ size (ll, lw, lsh, ovl, fl, and fw).

## Discussion

The discussion that follows is not meant to review the QTL exhaustively but rather to accomplish three goals. One goal is to examine QTL hotspots, defined as clusters containing more than three adjacent or overlapping QTL within a 20 cM window. In some instances, we have given separate QTL designations to pairs of loci that affect a single trait and are in close proximity but do not overlap. Thus, *fl2.1* and *fl2.2* are separated by just 2.3 cM. Similarly, *sp6.1/sp6.2* and *ftcp6.1/ftcp6.2* are located within 1.7 cM of their counterparts. In all three cases, one of the pair of QTL was detected in NY and the other in FR. For this reason we have named the QTL separately even though we suspect that more extensive phenotyping in a larger population or the use of a more accurate method of estimating the confidence interval would reveal them to be overlapping and therefore single loci. Colocalization of QTL for correlated traits was common and may suggest the pleiotropic activity of a single structural or regulatory gene. This phenomenon has been previously reported in tomato: *fw2.2* and *ovate* are quantitative regulatory genes (Frery et al. 2000; Liu et al. 2002) that are most likely responsible for the clustering of fruit size and shape QTL on tomato chromosome 2. Alternatively, clusters of genes influencing related phenomena could arise from gene duplication followed by subfunctionalization such that the duplicate copies evolve slightly different but overlapping functions (for example, tissue-specific expression) (Lynch and Force 2000). Under such circumstances, natural selection would maintain the duplicate loci. Of course, in instances where the traits are obviously codependent (shape and size parameters), clustering of QTL is expected. As previously mentioned, this has been reported for fruit shape and weight QTL on tomato chromosome 2 (Eshed and

Zamir 1995; Grandillo et al. 1999; Lippman and Tanksley 2001). Similar QTL clusters are observed on tomato chromosome 4 (Monforte et al. 2001; Yates et al. 2004). Several strategies have been used to discriminate whether traits associated with clustered QTL are controlled by a single pleiotropic gene or two (or more) tightly-linked genes. Substitution mapping in near-isogenic lines revealed that fruit color and soluble solids content are controlled by two separate, linked QTL on chromosome 1 of the wild tomato relative *S. chmielewskii* (Frery et al. 2003b). Similarly, high-resolution linkage mapping of a *S. pennellii* chromosome 9 introgression uncovered two distinct but closely-linked loci influencing soluble solids content (Fridman et al. 2002). Association mapping is another approach and has also proven useful for resolving relationships between QTL and candidate genes (Wilson et al. 2004).

The second goal of this discussion is to identify potential orthologs. In a number of instances, the eggplant QTL detected in this study seem to have counterparts in the tomato and/or potato genome in the form of QTL or morphological markers for analogous traits (Table 2). Such cases indicate conserved gene function within the Solanaceae.

The third aim is to emphasize overlap with the results of other QTL studies in eggplant. Environmental factors contribute in myriad ways to trait phenotypes, therefore QTL detection and estimates of QTL effects (PVE) can be highly dependent upon experimental location. For the traits measured in both NY and FR, the majority of QTL were location-specific. Even for QTL identified in both locations, variable PVE values were obtained. These discrepancies are not surprising given that the two locations (greenhouse in NY, field in FR) represent quite different environments in terms of both biotic and abiotic factors (soil type and nutrients, light availability, temperature, relative humidity). These variable results limit the broad applicability of findings from QTL analyses. However, because of the profound influence that environmental conditions and genetic background can have on the phenotypic expression of quantitative traits, QTL detected in multiple locations and populations are more likely to represent major genes.

A hotspot for vegetative organ (apices, stems and leaves) hairiness QTL (*ah3.1*, *sh3.1*, *lh3.1*) was detected on chromosome 3. These traits were well-correlated ( $P < 0.01$ ),  $r = 0.71$ – $0.85$ . Together these

**Table 2** QTL with putative conservation in the Solanaceae

Trait	QTL	Putative ortholog		Locus type	Reference		
		Locus name	Location (eggplant/other) <sup>a</sup>				
Leaf length	ll11.1	Leaf length QTL	E1/T4	QTL	Paran et al. (1997)		
		lflr4.1	E1/T4	QTL	Frary et al. (2004)		
Leaf shape	lsh1.1	lr1b	E1/T1	QTL	deVicente and Tanksley (1993)		
		lr5	E5/T5	QTL	deVicente and Tanksley (1993)		
	lsh5.1	lfw12.1	E5/T12	QTL	Frary et al. (2004)		
		lfl12.1	E5/T12	QTL	Frary et al. (2004)		
Leaf lobing	llob6.2	Pts	E6/T6	QTL	Tanksley et al. (1992)		
Fruit length	fl2.1, fl2.2	ovate	E2/T2	Known gene	Ku et al. (1999)		
		Frd2.1	E2/Pe2	QTL	Barchi et al. (2009)		
		Frs2.1	E2/Pe2	QTL	Barchi et al. (2009)		
		fs2.1	E2/Pe2	QTL	Zygier et al. (2005)		
Fruit shape index	fs7.1	fs7.b	E7/T7	QTL	Grandillo et al. (1999)		
Fruit weight	fw1.1	fw1.1	E1/T1	QTL	Grandillo and Tanksley (1996)		
		fw2.1	fw2.2	E2/T2	QTL	Frary et al. (2000)	
	fw2.1	fw2.1	E2/Pe2	QTL	Ben Chaim et al. (2001)		
		fw9.1	fw9.1	E9/T9	QTL	Grandillo et al. (1999)	
	Fruit stripe/chlorophyll netting	fst4.1, fcn4.1	fw9.2	fw9.2	E9/T9	QTL	Grandillo et al. (1999)
			Fs	E4/T10	Morphological	Tanksley et al. (1992)	
Apex hairs	ah3.1	u	E4/T10	Known gene	Tanksley et al. (1992)		
		Ln	E3/T3	Morphological	Tanksley et al. (1992)		
Stem and leaf hairs	sh10.1, lh10.1	TriIV-1	E10/T5	QTL	Maliepaard et al. (1995)		
		type B trichome qtl	E10/Po5	QTL	Bonierbale et al. (1994)		
Ovary hairs	ovh10.1	h	E10/T10	Morphological	Tanksley et al. (1992)		
Stem, leaf, prickle anthocyanin	sa10.1, sa10.2, lla10.1, lra10.1, pa10.1	an2a, an2b	E10/T10	Known gene	De Jong et al. (2004)		
		ant1	E10/T10	Known gene	De Jong et al. (2004)		
		chs	E10/T5	Known gene	De Jong et al. (2004)		
		3GT	E10/T10	Known gene	De Jong et al. (2004)		
Corolla anthocyanin	ca5.1	5GT	E5/T12	Known gene	Barchi et al. (2012)		
Fruit anthocyanin presence	fap10.1, fap10.2	an2a, an2b	E10/T10	Known gene	De Jong et al. (2004)		
		ant1	E10/T10	Known gene	De Jong et al. (2004)		
		chs	E10/T5	Known gene	De Jong et al. (2004)		
		3GT	E10/T10	Known gene	De Jong et al. (2004)		

<sup>a</sup> Chromosome location in eggplant (E) and other solanaceous species [tomato (T), potato (Po) or pepper (Pe)]

results suggest that a single structural or regulatory gene may be influencing trichome density on vegetative organs. A putative ortholog, *Ln*, which produces

very hairy stems when mutated, maps in this vicinity in tomato (Tanksley et al. 1992). On chromosome 10, *sh10.1* and *lh10.1* mapped 25 cM apart suggesting the

existence of two hairiness genes in fairly close proximity on this chromosome. Trichome QTL in tomato (*TriIV-1*, type IV trichome production; Malling et al. 1995) and potato (type B trichome density; Bonierbale et al. 1994) are possible orthologs of *sh10.1* and *lh10.1*. Additional QTL influencing leaf and apex hairiness mapped in independent locations on chromosomes 2 (*ah2.1*), 3 (*ah3.2*, *ah3.3*), 7 (*ah7.1*), and 9 (*lh9.1*) indicating that additional unrelated loci of slightly lesser effect are controlling these traits. Of these, *lh9.1* may be orthologous to *TriIV-2*, a QTL controlling type IV trichome density in tomato (Malling et al. 1995). Ovary hairs were controlled by an independent locus on chromosome 10, *ovh10.1*, which is syntenic with the tomato *h* (hairs absent) mutation (Tanksley et al. 1992).

The prickliness of leaves and stems showed a strong positive correlation ( $r = 0.71$ – $0.88$ ) and these prickly traits (*lp3.1* and *sp3.1*) mapped together on chromosome 3, indicating that a single gene may control the prickliness of both vegetative organs at this location. The previously mentioned QTL *ah3.1*, *sh3.1*, and *lh3* are located in this same region of chromosome 3. While *sh* was rather weakly correlated with *lp* and *sp* ( $r = 0.29$ ), neither *ah* nor *lh* was significantly correlated with either prickly trait. This lack of strong correlation between prickliness and hairiness, combined with the fact that prickly and hair QTL did not overlap anywhere else in the genome, suggests that these two traits are under separate genetic control.

The hotspot of prickliness QTL (*sp6.2*, *lp6.2*, *pp6.1*, *flcp6.1*, *ftcp6.1*, *ftcp6.2*) on chromosome 6 is consistent with the high correlation coefficients among these traits ( $r = 0.70$ – $0.92$ ) and suggests that a major structural or regulatory gene (accounting for 65–87 % of phenotypic variance) for prickliness resides in this location. That a locus controlling >75 % of the phenotypic variation in leaf lobing (*llob6.2*) maps in this same region is interesting. Deeply lobed leaves and prickly plant organs are highly correlated ( $r = 0.59$ – $0.83$ ) traits inherited from the wild *S. linnaeanum* parent. Close linkage between two (or more) genes for prickliness and leaf lobing would mean that deliberate selection against spines during eggplant domestication would have been accompanied by changes in leaf shape. Two tomato mutations map in the vicinity of *llob6.2*: *Pts*, Petroselinum and *c*, potato leaf (Liharska et al. 1997; Tanksley et al. 1992). Both mutations alter leaf complexity: *Pts* produces highly serrate, thrice-divided

leaves while *c* has the opposite effect, reducing both leaflet number and lobing (Hareven et al. 1996). Map-based cloning of *Pts* showed it to encode a novel KNOX1 transcription factor lacking a homeodomain (Kimura et al. 2008). Two wild tomato species with thrice-compound leaves (*S. cheesmanii* and *S. galapagense*) were found to overexpress *Pts* (Kimura et al. 2008). Thus a strong connection exists between this particular gene and the natural variation observed in wild *Solanum* species. Characterization of *C* has shown it to be a member of a family of *R2R3 MYB* transcription factors that control shoot branching. Because fully functional copies of *C* have been found in *S. melongena*, the gene has been ruled out as the gene responsible for the differences in leaf dissection between cultivated and wild eggplant (Busch et al. 2011). These authors suggest that the phenotype of the *sf* (solanifolia) mutant of tomato makes the *Sf* gene a more likely candidate for determining the degree of leaf indentation in eggplant. However, our study identified no QTL for leaf lobing on chromosome 3, the location of *Sf* (Tanksley et al. 1992).

Seven QTL for five highly correlated pigmentation traits (*sa*, *pa*, *lla*, *lra*, *fap*) map within a 16 cM region of chromosome 10. In this same vicinity, Barchi et al. (2012) found a cluster of QTL controlling anthocyanin levels in six plant parts (stem, leaf lamina, leaf veins, corolla, calyx, peduncle) in their intraspecific *S. melongena* mapping population. Not surprisingly, several structural and regulatory genes involved in anthocyanin synthesis map to orthologous regions of the tomato genome. These include *CHS* (encoding chalcone synthase), *3GT* (encoding 3-*O*-glucosyltransferase), and the transcription factors *an2a*, *an2b*, and *ant1* (De Jong et al. 2004). Any of these genes are good candidates for those regulating pigment production in eggplant. In tomato, elevated fruit anthocyanins resulted when either the native gene or the *ant1* allele from the purple-fruited wild tomato *S. chilense* was overexpressed in *S. lycopersicum* (Mathews et al. 2003; Schreiber et al. 2012). As a master switch that upregulates nearby genes involved in anthocyanin biosynthesis (*CHS*) and modification (*3GT*) (Mathews et al. 2003), *ANTI* could be an especially valuable target for improving eggplant peel color. While another major cluster of anthocyanin QTL was localized on chromosome 5 in the Barchi et al. (2012) intraspecific population, corolla anthocyanin (*ca5.1*) is the only trait that maps in this region

in the current study. A single candidate gene resides in this region of the tomato genome, *5GT* (encoding 5-O-glucosyltransferase) (Barchi et al. 2012). Of the other pigment QTL detected in the two studies, the only other overlap was seen on chromosome 11, between loci for abaxial leaf lamina anthocyanin (ablanE11.ML; Barchi et al. 2012) and fruit anthocyanin intensity (*fai1.1*).

Fruit stripe (*fst*) and fruit chlorophyll netting (*fcn*) are related traits in that both measure pigmentation patterns in the fruit. Not surprisingly, a single region on chromosome 4 explains 69 % (*fst4.1*) and 78 % (*fcn4.1*) of the variance in each trait. These effects are likely due to the action of the eggplant *Gv* (green variegation) gene that controls chlorophyll reticulation (Daunay et al. 2004). However, interestingly, independent loci of lesser effect (*fst10.1* and *fcn3.1*) were also identified for each character, suggesting that these traits may not be strictly monogenic. Both *fst* and *fcn* were inherited independently of the other color traits, a result that agrees with Daunay et al.'s (2004) finding that chlorophyll distribution and anthocyanin presence in eggplant fruit are controlled separately. Potential tomato orthologs of *fst4.1* and *fcn4.1* are *Fs* (fruit stripe) and *u* (uniform ripening), two linked genes that affect striping (Clayberg 1962) and shoulder color (MacArthur 1934) in unripe fruit. *u* has been recently characterized (Powell et al. 2012). It encodes a *Golden 2-like* transcription factor that regulates chlorophyll development during fruit formation. Thus, by selecting for unripe fruit that are a uniform light green (a product of the *u* mutation), tomato breeders have selected for reduced chlorophyll content, a trait that has negatively impacted sugar levels in ripe fruit (Powell et al. 2012). If *fst4.1* is indeed an ortholog of *u*, it would be a useful target for altering sugar content in eggplant breeding programs.

As expected, loci influencing codependent size and shape parameters of particular organs often localized to the same region of the genome; examples of this are: *lw1.1* and *lsh1.1*; *fl1.1* and *fw1.1*; *fl2.1*, *fl2.2* and *fw2.1*; *fl9.1* and *fw9.1*. Based on syntenic map positions, several of the QTL affecting leaf size and shape appear to be conserved with loci in tomato. Thus, two potential orthologs of *lll1.1* are positioned on chromosome 4 of tomato: a leaf length QTL identified by Paran et al. (1997) and *lflr4.1* (leaflet width to length ratio) (Frery et al. 2004). The leaf shape QTL *lsh1.1* and *lsh5.1* also have counterparts in the tomato

genome. Tomato *lr1b* and *lr5* (deVicente and Tanksley 1993), two QTL associated with leaflet width/length ratio, map near eggplant *lsh1.1* and *lsh5.1*, respectively. The leaflet width QTL (*lfw12.1*) on tomato chromosome 12 (Frery et al. 2004) is another possible ortholog of *lsh5.1*. The fruit length loci on chromosome 2 (*fl2.1* and *fl2.2*) map in the same genomic region as tomato *ovate* (Ku et al. 1999) and several QTL in pepper. These include loci controlling pepper fruit diameter (*Frd2.1*) and shape (*Frs2.1*, *fs2.1*) (Barchi et al. 2009; Zygier et al. 2005). The sole fruit shape QTL identified in this study (*fs7.1*) also has a counterpart in tomato, *fs7.b* (Grandillo et al. 1999). Fruit weight is perhaps the most widely studied quantitative trait in tomato and possible orthologs exist for all four of the loci detected here. The map positions of *fw1.1*, *fw9.1*, and *fw9.2* are syntenic with synonymously named QTL in tomato (Grandillo and Tanksley 1996; Grandillo et al. 1999) while the location of eggplant *fw2.1* corresponds to that of tomato *fw2.2* (Frery et al. 2000), a gene that controls carpel cell number. Pepper *fw2.1* also maps in this same region (Ben Chaim et al. 2001). These results suggest conservation of gene(s) affecting fruit size and shape on the long arm of chromosome 2 in eggplant, tomato, and pepper. One likely candidate is *ovate* which acts as a negative regulator of growth during early fruit development in tomato (Liu et al. 2002).

In a previous analysis of this interspecific eggplant population, simple linear regression analysis found 123 significant ( $P \leq 0.01$ ) QTL for the 40 morphological traits (Doganalr et al. 2002b; Frery et al. 2003a). Over half of those loci (54 %) were not detected in this follow-up study using CIM, with more stringent individual trait LOD thresholds ( $P \leq 0.01$ ) (determined empirically from 1,000 permutations of the data) to reduce the incidence of false positives. Whereas the QTL identified in the previous studies explained, on average, around 33 % of the phenotypic variance in the traits, the average in this study is 55 %. Because each of the QTL identified in this study account for >10 % of PVE, all of the loci qualify as “major” genes, according to the definition suggested by Collard et al. (2005). However, it should be noted that the relatively small size of the mapping population (58  $F_2$  individuals) has several consequences. In such populations, minor QTL are more difficult to detect and major QTL effects are generally overestimated (Vales et al. 2005). Thus the change in the

magnitude of QTL effect may, in part, be explained by the elimination of false positives and minor QTLs. Another factor to consider is the differing densities of the maps used in the analyses (207 markers in the previous map, 736 in the current map). With improved genome coverage we are more likely to detect a greater number of linked markers associated with each trait; therefore the estimated of phenotypic effects should be more accurate than those obtained previously for this population.

In addition, improvements in the resolution of the eggplant linkage map have yielded QTL approximately 1/10<sup>th</sup> the length of the corresponding loci in the previous studies. Thus, the current analysis has tended to identify fewer loci of greater effect that occupy more precise positions. Our re-examination of the trait data has yielded 22 novel QTL not found in the previous studies. The PVE of the newly identified QTL averaged 53 %, suggesting that CIM on the high-density genetic map revealed loci that were completely missed by simple linear regression with the low density linkage map. Counterparts in the tomato genome have been proposed for over one-third (35 %) of the 71 QTL reported here (Table 2). Conserved gene function in 50 % of the traits (for which significant QTL were identified) is thus expected to exist between tomato and eggplant.

In conclusion, the re-analysis of the morphological and domestication trait data for the *S. linnaeanum* × *S. melongena* F<sub>2</sub> population yielded valuable results. Using the genotypic data from a much larger number of eggplant-specific markers allowed the detection of hitherto unrevealed associations with phenotypic traits. Estimates of QTL effects were improved and individual QTL could be placed with greater accuracy on the high-density map. An important next step would be to confirm the positions and phenotypic effects of these QTL in a much larger mapping population. With this information, particular loci could be targeted for map-based cloning and marker-assisted selection studies. In addition these results provide a starting point for identifying putative orthologs between eggplant and tomato based on synteny. Intraspecific populations of eggplant have also been useful for developing high-density linkage maps that explore the synteny between the tomato and eggplant genomes (Barchi et al. 2012; Fukuoka et al. 2012) and provide for the possibility of additional quantitative trait analyses. Such analyses, combined

with the recently released tomato genome sequence (Tomato Genome Consortium 2012), should facilitate further examination of conserved gene function in the Solanaceae, an economically important family of plants.

**Acknowledgments** This work was supported by a Career Project (TUBITAK 104T224) from the Scientific and Technical Research Council of Turkey to Sami Doğanlar. This execution of the AFLP marker work was financially supported by DeRuiterZonen C.V., Rijk Zwaan Zaadteelt and Zaadhandel B.V., and Vilmorin Clause & Cie S.A. The AFLP<sup>®</sup> technology is covered by patents and/or patent applications of Keygene N.V. AFLP and KeyGene are registered trademarks of Keygene N.V. Other trademarks are the property of their respective owners.

**Conflict of interest** The authors declare that they have no conflict of interest.

## References

- Alpert K, Grandillo S, Tanksley SD (1995) *fw2.2*: a major QTL controlling fruit weight is common to both red- and green-fruited tomato species. *Theor Appl Genet* 91:994–1000
- Barchi L, Lefebvre V, Sage-Palloix A-M, Lanteri S, Palloix A (2009) QTL analysis of plant development and fruit traits in pepper and performance of selective phenotyping. *Theor Appl Genet* 118:1157–1171
- Barchi L, Lanteri S, Portis E, Vale G, Volante A, Pulcini L, Ciriaci T, Acciarri N, Barbierato V, Toppino L, Rotino GL (2012) A RAD tag derived marker based eggplant linkage map and location of QTLs determining anthocyanin pigmentation. *PLoS ONE* 7:e43740
- Ben Chaim A, Paran I, Grube RC, Jahn M, van Wijk R, Peleman J (2001) QTL mapping of fruit-related traits in pepper (*Capsicum annuum*). *Theor Appl Genet* 102:1016–1028
- Bonierbale MW, Plaisted RL, Pineda O, Tanksley SD (1994) QTL analysis of trichome-mediated insect resistance in potato. *Theor Appl Genet* 87:973–987
- Busch BL, Schmitz G, Rossmann S, Piron F, Ding J, Bendahmane A, Theres K (2011) Shoot branching and leaf dissection in tomato are regulated by homologous gene modules. *Plant Cell* 23:3595–3609
- Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. *Genetics* 138:963–971
- Clayberg CD (1962) Inheritance and linkage of fruit stripe *Fs*. *Rep Tomato Genet Coop* 12:22–23
- Collard B, Jahufer M, Brouwer J, Pang E (2005) An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica* 142:169–196
- Daunay MC, Aubert S, Frary A, Doganlar S, Lester RN, Barendse G, van der Weerden G, Hennart JW, Haanstra J, Dauphin F, Jullian E (2004) Eggplant (*Solanum melongena*) fruit colour: pigments measurements and genetics. In:



- Proceedings of the XIIth EUCARPIA meeting on genetics and breeding of *Capsicum* and eggplant, 17–19 May 2004, Noordwijkerhout, The Netherlands, pp 108–116
- De Jong WS, Eannetta NT, De Jong DM, Bodis M (2004) Candidate gene analysis of anthocyanin pigmentation loci in the *Solanaceae*. *Theor Appl Genet* 108:423–432
- deVicente MC, Tanksley SD (1993) QTL analysis of transgressive segregation in an interspecific tomato cross. *Genetics* 134:585–596
- Doganlar S, Frary A, Daunay M, Lester R, Tanksley S (2002a) A comparative genetic linkage map of eggplant (*Solanum melongena*) and its implications for genome evolution in the Solanaceae. *Genetics* 161:1697–1711
- Doganlar S, Frary A, Daunay MC, Lester RN, Tanksley SD (2002b) Conservation of gene function in the Solanaceae as revealed by comparative mapping of domestication traits in eggplant. *Genetics* 161:1713–1726
- Doganlar S, Frary A, Daunay MC, Huvenaars K, Mank R, Frary A (in press) High resolution map of eggplant (*Solanum melongena*) reveals extensive chromosome rearrangement in domesticated members of the Solanaceae. *Euphytica*
- Eshed Y, Zamir D (1995) An introgression line population of *Lycopersicon pennellii* in the cultivated tomato enables the identification and fine-mapping of yield-associated QTL. *Genetics* 141:1147–1162
- FAO Statistics (2013) <http://faostat.fao.org>. Accessed 15 Jan 2013
- Frary A, Nesbitt TC, Frary A, Grandillo S, van der Knapp E, Cong B, Liu J, Meller J, Elber R, Alpert KB, Tanksley SD (2000) *fw2.2*: a quantitative trait locus key to the evolution of tomato fruit size. *Science* 289:85–88
- Frary A, Doganlar S, Daunay MC, Tanksley SD (2003a) QTL analysis of morphological traits in eggplant and implications for conservation of gene function during evolution of Solanaceous species. *Theor Appl Genet* 107:359–370
- Frary A, Doganlar S, Frampton A, Fulton T, Uhlig J, Yates H, Tanksley S (2003b) Fine mapping of quantitative trait loci for improved fruit characteristics from *Lycopersicon chmielewskii* chromosome 1. *Genome* 46:235–243
- Frary A, Fritz LA, Tanksley SD (2004) A comparative study of the genetic bases of natural variation in tomato leaf sepal and petal morphology. *Theor Appl Genet* 109:523–533
- Fridman E, Liu YS, Carmel-Goren L, Gur A, Shores M, Pleban T, Eshed Y, Zamir D (2002) Two tightly linked QTLs modify tomato sugar content via different physiological pathways. *Mon Genet Genomics* 266:821–826
- Fukuoka H, Miyatake K, Nunome T, Negoro S, Shirasawa K, Isobe S, Asamizu E, Yamaguchi H, Ohyama A (2012) Development of gene-based markers and construction of an integrated linkage map in eggplant by using *Solanum* orthologous (SOL) gene sets. *Theor Appl Genet* 125:47–56
- Grandillo S, Tanksley SD (1996) Analysis of horticultural traits differentiating the cultivated tomato from the closely related species *Lycopersicon pimpinellifolium*. *Theor Appl Genet* 92:935–951
- Grandillo S, Ku HM, Tanksley SD (1999) Identifying the loci responsible for natural variation in fruit size and shape in tomato. *Theor Appl Genet* 99:978–987
- Hareven D, Gutfinger T, Parnis A, Eshed Y, Lifschitz E (1996) The making of a compound tomato leaf: genetic manipulation of leaf architecture in tomato. *Cell* 84:735–744
- Joehanes R, Nelson JC (2008) QGene 4.0 an extensible Java QTL-analysis platform. *Bioinformatics* 24:2788–2789
- Kimura S, Koenig D, Kang J, Yoong FY, Sinha N (2008) Natural variation in leaf morphology results from mutation of a novel *KNOX* gene. *Curr Biol* 18:672–677
- Ku H-M, Doganlar S, Chen K-Y, Tanksley SD (1999) The genetic basis of pear-shaped tomato fruit. *Theor Appl Genet* 9:844–850
- Liharska TB, Hontelez J, van Kammen A, Zabel P, Koornneef M (1997) Molecular mapping around the centromere of tomato chromosome 6 using irradiation-induced deletions. *Theor Appl Genet* 95:969–974
- Lippman Z, Tanksley SD (2001) Dissecting the genetic pathway to extreme fruit size in tomato using a cross between the small-fruited wild species *Lycopersicon pimpinellifolium* and *L. esculentum* var. Giant Heirloom. *Genetics* 158:413–422
- Liu J, van Eck J, Cong B, Tanksley SD (2002) A new class of regulatory genes underlying the cause of pear-shaped tomato fruit. *Proc Natl Acad Sci USA* 99:13302–13306
- Livingstone KD, Lackney VK, Blauth JR, van Wijk R, Jahn MK (1999) Genome mapping in *Capsicum* and the evolution of genome structure in the Solanaceae. *Genetics* 152:1183–1202
- Lynch M, Force A (2000) The probability of duplicate gene preservation by subfunctionalization. *Genetics* 154:459–473
- MacArthur JW (1934) Linkage groups in the tomato. *J Genet* 29:123–133
- Maliepaard C, Bas N, van Heusden S, Kos J, Pet G, Verkerk R, Vrieling R, Zabel P, Lindhout P (1995) Mapping of QTLs for glandular trichome densities and *Trialeurodes vaporariorum* (greenhouse whitefly) resistance in an F<sub>2</sub> from *Lycopersicon esculentum* × *Lycopersicon hirsutum* f. *glabratum*. *Heredity* 75:425–433
- Mathews H, Clendennen SK, Caldwell CG, Liu XL, Connors K, Matheis N, Schuster DK, Menasco DJ, Wagoner W, Lightnew J, Wagner DR (2003) Activation tagging in tomato identifies a transcriptional regulator of anthocyanin biosynthesis modification and transport. *Plant Cell* 15:1689–1703
- Monforte A, Friedman E, Zamir D, Tanksley SD (2001) Comparison of a set of allelic QTL-NILs for chromosome 4 of tomato: deductions about natural variation and implications for germplasm collection. *Theor Appl Genet* 102:572–590
- Nagelkerke NJD (1991) A note on a general definition of the coefficient of determination. *Biometrika* 78:691–692
- Nelson JC (1997) QGene: software for marker-based genomic analysis and breeding. *Mol Breed* 3:229–235
- Nunome T, Yoshida T, Hirai M (1998) Genetic linkage map of eggplant. In: Proceedings of the 10th Eucarpia meeting on genetics and breeding of *Capsicum* and eggplant, Avignon France, pp 239–242
- Nunome T, Ishiguro K, Yoshida T, Hirai M (2001) Mapping of fruit shape and color development traits in eggplant (*Solanum melongena* L.) based on RAPD and AFLP markers. *Breed Sci* 51:19–26
- Paran I, Goldman I, Zamir D (1997) QTL analysis of morphological traits in a tomato recombinant inbred line population. *Genome* 40:242–248
- Powell ALT, Nguyen CV, Hill T, Cheng KL, Figueroa-Balderas R, Aktas H, Ashrafi H, Pons C, Fernandez-Muñoz R,

- Vicente A, Lopez-Baltazar J, Barry CS, Liu Y, Chetelat R, Granell A, van Deynze A, Giovannoni JJ, Bennett AB (2012) *Uniform ripening* encodes a *Golden 2-like* transcription factor regulating tomato fruit chloroplast development. *Science* 336:1711–1715
- Schreiber G, Reuveni M, Evenor D, Oren-Shamir M, Ovadia R, Sapir-Mir M, Bootbool-Man A, Nahon S, Shlomo H, Chen L, Levin I (2012) ANTHOCYANIN1 from *Solanum chilense* is more efficient in accumulating anthocyanin metabolites than its *Solanum lycopersicum* counterpart in association with the ANTHOCYANIN FRUIT phenotype of tomato. *Theor Appl Genet* 124:295–307
- Tanksley SD, Ganai MW, de Prince JP, Vicente MC, Bonierbale MW, Broun P, Fulton TM, Giovannoni JJ, Grandillo S, Martin GB, Messeguer R, Miller JC, Miller L, Paterson AH, Pineda O, Roder S, Wing RA, Wu W, Young ND (1992) High-density linkage maps of the tomato and potato genomes. *Genetics* 132:1141–1160
- Tomato Genome Consortium (2012) The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* 485:635–641
- Vales MI, Schon CC, Capettini F, Chen XM, Corey AE, Mather DE, Mundt CC, Richardson KL, Sandoval-Islas JS, Utz HF, Hayes PM (2005) Effect of population size on the estimation of QTL: a test using resistance to barley rust stripe. *Theor Appl Genet* 111:1260–1270
- Voorrips RE (2002) MapChart: software for the graphical presentation of linkage maps and QTLs. *J Hered* 93:77–78
- Wilson LM, Whitt SR, Ibanez AM, Rocheford TR, Goodman MM, Buckler ES (2004) Dissection of maize kernel composition and starch production by candidate gene association. *Plant Cell* 16:2719–2733
- Yates HE, Frary A, Doganlar S, Frampton A, Eannetta N, Uhlig J, Tanksley SD (2004) Comparative fine mapping of fruit quality QTLs on chromosome 4 introgressions derived from two wild tomato species. *Euphytica* 135:283–296
- Zheng ZB (1994) Precision mapping of quantitative trait loci. *Genetics* 136:1457–1468
- Zygier S, Chaim A-B, Efrati A, Kaluzky G, Borovsky Y, Paran I (2005) QTLs mapping for fruit size and shape in chromosomes 2 and 4 in pepper and a comparison of the pepper QTL map with that of tomato. *Theor Appl Genet* 111:437–445