

**Quantification of training and match-load distribution
across a season in elite English Premier League soccer
players**

Journal:	<i>Science and Medicine in Football</i>
Manuscript ID	RSMF-2018-0122.R1
Manuscript Type:	Original Papers
Keywords:	soccer training, mesocycle, microcycle, session RPE-TL, periodisation
Abstract:	<p>Abstract Objective: To examine training and match loads undertaken by soccer players competing in the English Premier League. Methods: Using a retrospective design, external (GPS) and internal training loads (sessions ratings of perceived exertion [sRPE-TL]) were examined in twenty-six players across the competition phase of the 2012-2013 English Premier League season. Within-subject linear mixed-models estimated the mean effects (95% confidence interval [CI]) for load data across 6-week mesocycles and 1-week microcycles. Results: Daily sRPE-TL (95% CI range, 15 to 111 AU) and total distance (95% CI range, 179 to 949 AU) were higher during the early stages (mesocycle 1 and 2) of the competition period. Overall, high-speed activity was similar between mesocycles. Across the training week, load was greater on match day and lower pre match-day (G-1) vs. all other days, respectively ($p < 0.001$). sRPE-TL (~70-90 AU per day) and total distance (~700-800 m per day) progressively declined over the three days before a match ($p < 0.001$). High-speed distance was greater three days (G-3) before a game vs. G-1 (95% CI, 140 to 336 m) while very high-speed distance was greater on G-3 and G-2 than G-1 (95% CI range, 8 to 62 m; $p < 0.001$). Conclusion: Periodisation of in-season training load is mainly evident across the weekly microcycle reflecting the recovery and preparation for matches.</p>

SCHOLARONE™
Manuscripts

1
2
3 **Quantification of training and match-load distribution across a season in elite**
4
5 **English Premier League soccer players**
6
7
8
9

10
11
12 **Running Title:** Seasonal training and match-load in elite soccer
13
14

15
16
17 **Submission Type:** Original Research
18
19

20
21 **Keywords:** soccer training, mesocycle, microcycle, session RPE-TL, periodisation
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Abstract

Objective: To examine training and match loads undertaken by soccer players competing in the English Premier League.

Methods: Using a retrospective design, external (GPS) and internal training loads (sessions ratings of perceived exertion [sRPE-TL]) were examined in twenty-six players across the competition phase of the 2012-2013 English Premier League season. Within-subject linear mixed-models estimated the mean effects (95% confidence interval [CI]) for load data across 6-week mesocycles and 1-week microcycles. **Results:** Daily sRPE-TL (95% CI range, 15 to 111 AU) and total distance (95% CI range, 179 to 949 AU) were higher during the early stages (mesocycle 1 and 2) of the competition period. Overall, high-speed activity was similar between mesocycles. Across the training week, load was greater on match day and lower pre match-day (G-1) vs. all other days, respectively ($p < 0.001$). sRPE-TL (~70-90 AU per day) and total distance (~700-800 m per day) progressively declined over the three days before a match ($p < 0.001$). High-speed distance was greater three days (G-3) before a game vs G-1 (95% CI, 140 to 336 m) while very high-speed distance was greater on G-3 and G-2 than G-1 (95% CI range, 8 to 62 m; $p < 0.001$).

Conclusion: Periodisation of in-season training load is mainly evident across the weekly microcycle reflecting the recovery and preparation for matches.

Keywords: soccer training, mesocycle, microcycle, sRPE-TL, periodisation

Introduction

The complex physiological demands of soccer necessitate the implementation of training programmes which are multifactorial in nature (Morgans et al., 2014). Such requirements are further complicated by the stochastic movement profiles observed in elite soccer. The sporadic work bouts associated with soccer training may therefore result in variability between the desired training load and the actual training load the players are exposed to (Malone et al., 2015). Monitoring the individual player's daily training load therefore represents an important component of the effective planning of a soccer-specific training regimen (Weston, 2018).

The volume and intensity of training, collectively referred to as the training load (Impellizzeri et al., 2005), requires manipulation (periodisation) to elicit an optimum training stimulus (Malone et al., 2015). Many clubs therefore employ practitioners to collect, interpret and feedback information to coaches regarding the players daily load and status (Arkenhead and Nassis, 2016; Weston, 2018). To date, studies focused on training load quantification in soccer have largely focused on isolated training drills (Coutts et al., 2009; Casamichana and Castellano, 2010; Buchheit et al., 2015) or mesocycles of up to 10 weeks (Impellizzeri et al., 2004; Gaudino et al., 2013; Scott et al., 2013; Clemente et al., 2019). In contrast, while a plethora of studies have documented the long-term (season long) periodisation models adopted in other football codes (Gabbett and Jenkins, 2011; Moreira et al., 2016; McGahan et al., 2017), little data currently exists in elite soccer.

1
2
3
4
5
6 Recent studies have provided some insight into the seasonal training loads
7
8 encountered by players competing in the Spanish reserve league (Los Arcos et al.,
9
10 2017; Martin-Garcia et al., 2018), Dutch Eredivisie League (Stevens et al., 2017),
11
12 and the English Premier League (Malone et al., 2015; Anderson et al., 2016). Across
13
14 the competitive season there was little variation in training load between mesocycles
15
16 (6-8 week training blocks) (Malone et al., 2015; Anderson et al., 2016). Within
17
18 weekly microcycles, load was also generally similar between training days with the
19
20 exception of a marked reduction in load on the day preceding the game (Malone et
21
22 al., 2015; Martin-Garcia et al., 2018). Whilst these studies provide valuable insights
23
24 into the training loads experienced by elite players, further observations are required
25
26 in order to gain a comprehensive insight into the collective periodisation practices
27
28 adopted by professional teams (Weston, 2018). Furthermore, a more detailed
29
30 analysis of the nature of the loading incurred by players is required. For example,
31
32 internal training load, or the individual physiological response to the external load
33
34 administered by the coach, represents the stimulus for training induced adaptation
35
36 (Virus and Virus, 2000). Valid and reliable indicators of internal training load are
37
38 therefore essential when monitoring the training process. Session RPE-TL (sRPE-
39
40 TL) represents a valid indicator of the global internal training load during
41
42 intermittent team sports such as soccer (Impellizzeri et al., 2004; Casamichana et al.,
43
44 2013; Kelly et al., 2016). Despite the importance of the internal load in indicating the
45
46 training response, observations on elite players have also been largely restricted to
47
48 descriptions of short-term periods of training (Campos-Vazquez et al., 2015) with
49
50 only one research group to date reporting session RPE-TL responses to long-term
51
52 periods of training in elite players (Malone et al., 2015).
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Most of what is currently known about load monitoring derives from personal experiences or remains unpublished, since many elite teams are often unwilling to publish their data in order to retain competitive advantage. The training approaches adopted by elite teams and the degree to which these approaches incorporate periodisation strategies therefore remains largely unexplored in the literature. A recent survey of practitioners and coaches working in elite English soccer perceived that coaches were mostly responsible, and sports scientists/fitness coaches somewhat responsible, for planning training (Weston, 2018). Coaching practice is heavily influenced by tradition, emulation and historical precedence rather than through critical consideration of the latest research (Stoszkowski and Collins, 2016). Given the diverse coaching philosophies inherent in the modern elite game, further studies are needed to enhance our understanding as to how training loads in soccer are programmed across the annual cycle. The aim of the current investigation therefore was to quantify the combined external and internal training and match-load distribution across the competition phase of one full season at an English Premier League club.

Methods

Participants

Twenty-six elite-level soccer players were monitored across a 36-week competition phase of the 2012-2013 English Premier League (League Champions) season (mean \pm SD: age 27 ± 5.4 years, body mass 77 ± 6.6 kg, height 181 ± 7.0 cm). Players were assigned to one of five positional groups: central defender (CD) ($n = 4$), wide defender (WD) ($n = 4$), central midfielder (CM) ($n = 7$), wide midfielder (WM) ($n =$

1
2
3 3), and attacker (A) ($n = 8$). The team competed in four official competitions
4 throughout the season corresponding to 49 competitive matches in total. All of the
5 players were notified as to the aim of the study, requirements, research procedures,
6 benefits and risks before giving written informed consent. The Ethics committee of
7 the relevant School at Liverpool John Moores University approved the study.
8
9
10
11
12
13
14
15
16

17 *Design*

18 For the purpose of the current study, all of the first team field-based training sessions
19 carried out were considered for the analysis. This was inclusive of sessions involving
20 both the starting line-up and non-starting players. Individual training, rehabilitation,
21 recovery and specific fitness sessions were excluded from the analysis. Goalkeepers
22 were not included in the study. Daily training load data was collected using the
23 sRPE-TL method and micro-technology. Training and match data collection was
24 carried out at the soccer club's training ground on the same natural outdoor grass
25 training pitches, and at both home and away grounds in the English Premier League,
26 respectively. A stadium-based tracking system was used to record match-play
27 activities. All training and match load data observed during a 36-week competition
28 phase of the season were categorised into 6-week mesocycle phases, and subsequent
29 weekly calendar blocks (Sunday to Sunday). This enabled a full season's analysis of
30 both the training and match-play load (Figure 1).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Training and match load data were also analysed in relation to the proximity of the
52 forthcoming competitive game (day type). Six-day types in total were identified and
53 analysed in the current study (G-3, G-2, G-1, match day (MD), G+2, G+3). For
54 example, one day before the game was classified as game day minus one (G-1), two
55
56
57
58
59
60

1
2
3 days before was G-2 etc. whereby G+2 and G+3 were the second- and third-days
4 post-match, respectively. The day immediately following a game (i.e. G+1) was not
5
6 included in the analysis as this was classified as a recovery day which involved a
7
8 reduced load non-weight bearing recovery strategy and was therefore not
9
10 representative of a training day. During the season there were one, two, and three
11
12 game weeks. A one-game week consisted of 6 training days leading into the game.
13
14 The two-game week had 1 recovery day following the first game (e.g. G+1) and 4
15
16 training days leading into the next game. A three-game week had 1 recovery session
17
18 and a training day (G-1) between the first and second game and the second and third
19
20 game respectively. In some instances during two and three game weeks, games were
21
22 played in closer proximity (e.g. Saturday and Tuesday), leaving only two days
23
24 between fixtures. In this scenario, 1 recovery session and a training day (G-1) was
25
26 implemented between games.
27
28
29
30
31
32
33
34

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

****Figure 1 near here****

Methodology

Training Load Assessment

Internal Training Load: Internal training load (sRPE-TL, arbitrary units, AU) was estimated for all players by multiplying total training or match session duration (min) with session ratings of perceived exertion (sRPE) (Foster et al., 2001). Player sRPE was collected in isolation where possible, to avoid the potential effects of peer pressure ~20-minutes after each training session or match. All the players were familiarised with the use of the RPE scale during the pre-season training phase.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

External Training Load: Team Training and Matches. The player's external training session load was monitored using portable micro-technology (GPSports SPI Pro X, Canberra, Australia). The SPI Pro X (GPS and accelerometer integrated; size: 48x20x87mm; 76g) was placed inside a specially made vest, inside a mini pocket and positioned on the player's back, which was located centrally between the scapulae. The player wore micro-technology for the whole duration of the session. The unit was activated ~15 min before data collection to allow for the acquisition of satellite signals (Waldron et al., 2011). During every training session observation, the minimum acceptable number of available satellite signals was 8, which is optimal for the measurement of human movement (Jennings et al., 2010). To avoid inter-unit error, each player wore the same micro-technology device for every training session observation (Jennings et al., 2010). The SPI Pro unit provides raw position, velocity and distance data at a rate of 15 samples-per-second (15 Hz). Every 3 raw data points were averaged for the purpose of the current study to provide a sampling frequency of 5 Hz. This type of system has been shown to provide a reliable and valid estimate of the high-speed distance covered during multi-directional sports such as soccer (Portas et al., 2010; Randers et al., 2010; Varley et al., 2012; Waldron et al., 2011).

All training sessions and competitive matches during the 2012-13 season were observed and subsequently recorded. The mean number of training sessions completed, and the average match observations during each month ($n = 5$) are shown in Figure 2. Mean training session duration across all positions was 59 ± 7 min (Figure 3). Matches were inclusive of domestic (Premier League, F.A. Cup, League Cup), and European (Champions League) fixtures. Friendly games were excluded

1
2
3 from the analysis. A total of 49 matches were observed during the 36-week
4
5 competition phase of the season. Individual player's activities were monitored during
6
7 each game using a stadium-based multiple-camera match analysis system (Prozone
8
9 Sports Limited, Leeds, UK). Data from both home and away fixtures were included.
10
11 Only data from completed 90 min matches were used for the analysis. The median
12
13 number of completed matches by individual players was 16 (range: 2-38). All
14
15 Prozone data were processed using the appropriate software package (Prozone 3
16
17 Version 12.0.4.2., Prozone Sports Limited, Leeds, UK). This was carried out post-
18
19 game(s) by the club's performance analyst and exported into a Microsoft Excel
20
21 spreadsheet database (Microsoft Corporation, U.S.) for the analysis.
22
23
24
25
26
27
28

29 ******Figure 2 near here******
30

31
32
33 ******Figure 3 near here******
34
35
36
37

38 The observed training and match-play activities (external load markers) identified for
39
40 subsequent analysis were: total distance (m), distance (m) completed at high-speeds
41
42 >14.4 km/h (m), and distance (m) completed at very high-speeds 19.8-25.2 km/h.
43
44 The current authors acknowledge that some differences in the measures derived from
45
46 the micro-technology devices and Prozone system exist. In particular, it has been
47
48 shown previously that high-intensity running distances are slightly-to-moderately
49
50 greater when tracked using Prozone in comparison to GPSports devices (Buchheit et
51
52 al., 2014). However, for the purpose of the current investigation, both the GPSports
53
54 (training load), and Prozone (match load) data were combined together for the
55
56 analysis (Anderson et al., 2016).
57
58
59
60

Statistical Analysis

[Data are](#) represented as means \pm S.D. A multi-factorial linear mixed model was used to quantify mean differences between mesocycles, day-type and playing position.

[Use of linear mixed-modelling is suitable to examine repeated-measures data and unbalanced observations over time as, for example, in the context of our study where players differ in the number of training sessions and matches \(Cnaan et al., 1997\).](#)

Linear mixed modeling can also cope with the mixture of random and fixed level effects (Cnaan et al., 1997) as well as with missing and ‘nested’ data (hierarchical models). [The main effects for sub-group comparisons](#) of each factor were [summarised using](#) least significance difference (LSD) multiple contrasts (Perneger, 1998).

[Mean differences are presented with 95% confidence intervals \(CI\) as markers of uncertainty in the estimates. In the absence of an established anchor, despite the lack of real-world relevance of standardised effect sizes \(Lenth, 2001\), Cohen’s d was reported as an additional statistic for interpreting the magnitude of the estimated effects \(Cook et al., 2018\).](#) Effect size (ES), estimated from the ratio of the mean difference to the pooled standard deviation were also calculated. The ES magnitude was classified as trivial (<0.2), small ($>0.2-0.6$), moderate ($>0.6-1.2$), large ($>1.2-2.0$) and very large ($>2.0-4.0$) (Hopkins et al., 2009). [Within this particular context and to address the potential inflation of error rates associated with the large number of inferences in the present study, effects were declared meaningful if the point estimate for the mean difference expressed in standardised units attained threshold of moderate \(ES > 0.6\).](#)

Results

Mesocycle

Total number of games during each of the 6 x 6-week mesocycle ranged from 6-10 (mesocycle 1 = 6; mesocycle 2 = 9; mesocycle 3 = 10; mesocycle 4 = 6; mesocycle 5 = 9, and mesocycle 6 = 9). Mean daily sRPE-TL, total distance, high-speed distance and very high-speed distance across each of the 6 x 6-week mesocycles by playing position are presented in Table 1. A statistically significant change in all variables was observed across the six mesocycles (all $p < 0.001$). Daily sRPE-TL was higher during the early stages of the season with greater values observed in mesocycle 1 than all other mesocycles (95% CI range, 16 to 111) and greater values observed in mesocycle 2 than mesocycles 3 and 4 (95% CI range, 15 to 91 AU). Total distance was higher in mesocycles 1 and 2 than mesocycles 3, 4, and 6 (95% CI range, 179 to 949 AU). Meaningful differences in high-speed distance were only observed in mesocycle 5 compared to mesocycle 4 (95% CI, 66 to 228 m) with greater very high-speed distance observed in mesocycle 2 than mesocycle 4 (95% CI, 21 to 64 m). No meaningful or statistically significant main effects of playing position or interaction between playing position and mesocycle were observed for any variable (all $p > 0.05$).

**** Table 1 near here****

Day Type

Mean daily sRPE-TL, total distance, total high-speed distance and total very high-speed distance across all day types are represented in Figures 4-7. No meaningful or statistically significant main effect of playing position were observed for any

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

variable ($p > 0.05$). There was a statistically significant main effect of day-type for all variables (all $p < 0.001$). Session RPE-TL (MD vs. other days: 95% CI range, 208 to 409 AU; G-1 vs. other days: 95% CI range, -409 to -47 AU), total distance (MD vs. other days: 95% CI range, 4188 to 6069 m; G-1 vs. other days: 95% CI range, -6070 to -430 m), total high-speed distance (MD vs. other days: 95% CI range, 1466 to 1875 AU; G-1 vs. other days: 95% CI range, -1875 to -35 m) and total very high-speed distance (MD vs. other days: 95% CI range, 425 to 542 AU; G-1 vs. other days: 95% CI range, -542 to -20 m) were higher on MD and lower on G-1 compared to all other days. sRPE-TL (~70-90 AU per day) and total distance (~700-800 m per day) progressively reduced over the three days before a match ($p < 0.001$). High-speed distance was greater on G-3 than G-1 (95% CI, 140 to 336 m) and very high-speed distance was greater on G-3 and G-2 vs G-1 (95% CI range, 8 to 62 m; $p < 0.001$; Figure 6 and 7).

There was a statistically significant interaction between day-type and playing position for all variables predominantly reflecting positional differences on MD (all $p < 0.001$; Figures 4-7). During training, sRPE-TL was lower in WM than WD on G-3 (95% CI, -208 to -18 AU). sRPE-TL was higher in A than WD and CM on G-2 (95% CI range, -29 to 129 AU) and higher than all other positions on G-1 (95% CI range, -2 to 156 AU). Attackers covered greater total distance than on CD and WD on G-1 (95% CI range, 102 to 1387 m). Differences in high-speed activity between positions were only observed on MD.

****Figure 4 near here****

1
2
3 ****Figure 5 near here****
4
5
6

7
8 ****Figure 6 near here****
9
10

11
12 ****Figure 7 near here****
13
14
15

16 Discussion

17
18
19 The aim of the current study was to examine the external and internal load incurred
20 by elite soccer players across both the larger and smaller units of the annual
21 competition period. Across the competition period there was limited variation in
22 loading between the mesocycles with similar loads observed between playing
23 positions. In contrast, marked fluctuations in external and internal load were evident
24 within the weekly microcycle phase which was further influenced by playing
25 position. This was generally characterised by a post-match recovery day (low load)
26 followed by an increase in loading (G+2 through to G+3 and G-3) and subsequent
27 taper through G-2, and G-1. The findings of the present study provide novel insights
28 into the training periodisation undertaken by an elite English Premier League team
29 during a championship winning season. Further studies of this type are required to
30 enable a more comprehensive examination and subsequent development of the
31 training methodologies adopted by elite coaches.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 In the present study, total distance and sRPE-TL were ~ 470 m (95 % CI, 228 to 724
52 m), and 40 AU (95 % CI, 19 to 62 AU) higher at the start of the competitive phase
53 (mesocycle 1) versus the end (mesocycle 6). These changes in total distance are
54 lower than those previously observed by Malone and colleagues (2015), where
55
56
57
58
59
60

1
2
3 players covered ~1300 m more total distance in mesocycle 1 than mesocycle 6.
4
5 Mean daily total distance (95 % CI, 472 to 947 m), sRPE-TL (95 % CI, 67 to 111
6
7 AU) and high-speed distance (95 % CI, -19 to 159 m) were also ~700 m, 90 AU and
8
9 70 m higher respectively at the start of the season (mesocycle 1) compared with mid-
10
11 season (mesocycle 4) across all positions in the present study. Greater training loads
12
13 at the beginning of the in-season competitive phase may often reflect the coaches'
14
15 desire to maintain the emphasis on the development of fitness levels following the
16
17 pre-season training period (Malone et al., 2015).
18
19
20
21
22
23

24 The middle phase of the season (mesocycle 4 - mid-December) is associated with the
25
26 lead into the Christmas period, which typically has a highly congested fixture
27
28 schedule in the English Premier League. We presently observed the highest number
29
30 of matches ($n = 7$) and the greatest average number of training session observations
31
32 62 (range: [n], 40-62) during this period. However, the average training session
33
34 duration (48 ± 5 min) was greatly reduced across December compared to all other
35
36 periods of the season which resulted in the lowest sRPE-TL, total distance, high-
37
38 speed distance, and very high-speed distances. These changes were consistent with
39
40 the strategy employed by the head coach which aimed to offset the increased
41
42 frequency of matches by reducing training induced fatigue in order to maintain
43
44 match readiness. Our findings are in-line with Malone and colleagues (2015) who
45
46 also reported reductions in training volume during the mid-season phase, whereby
47
48 sRPE-TL was lower by ~80 AU across this period.
49
50
51
52
53
54
55

56 Training load prescription in soccer is largely influenced by the competition
57
58 frequency, with in-season microcycles of typically 3 to 7 days in duration repeatedly
59
60

1
2
3 occurring around matches (Morgans et al., 2014; Malone et al., 2015; Akenhead et
4 al., 2016). sRPE-TL (~70-90 AU per day) and total distance (~700-800 m per day)
5 progressively reduced over the three days before a match. High-speed distance was
6 also greater on G-3 than G-1 (95% CI, 140-336 m) and very high-speed distance was
7 greater on G-3 and G-2 vs G-1. The higher training loads observed on G-3 reflected
8 training sessions incorporating drills undertaken on larger pitch sizes (i.e. extensive
9 endurance position-specific practices) with a greater number of players (7 v 7 – 11 v
10 11). More intensive endurance drills were undertaken in smaller training areas with a
11 reduced number of players (e.g. 3 v 2, 5 v 4, and 1 v 1 – 3 v 3) as part of training
12 sessions undertaken on G-2. The aim of these training sessions was to elicit
13 intensities deemed suitable to produce the physiological adaptations required for
14 soccer-specific endurance (Little and Williams, 2006) while simultaneously aiding
15 the development of technical and tactical skills similar to situations experienced
16 during the game. All variables were lowest on G-1 as a consequence of the
17 implementation of lower intensity and shorter training sessions the day before a
18 match, consisting mainly of activation and reactive speed training type drills. The
19 decline in daily load from G-3 to G-1 in the current study is in agreement with recent
20 observations in Spanish La Liga reserve team players who showed a marked
21 reduction in total distance (~3000 m) and high-speed distance (~170 m) across the
22 three-day period (Martin-Garcia et al., 2018). In contrast, Malone and colleagues
23 (2015) reported greater high-speed distances on G-1 than G-2 in English Premier
24 League players. The rationale for this approach was not reported by the authors,
25 however, it would seem counterproductive and contrary to ‘tapering’ approaches
26 previously discussed in the literature (Owen et al., 2017). Reducing training load on
27 the day preceding a competitive match may enhance the capability of significantly
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 decreasing physical stressors upon players, whilst leading to reductions in an
4
5 accumulative fatigue response (Owen et al., 2017).
6
7
8
9

10 The present findings demonstrate that a gradual reduction in external and internal
11
12 load across the three-day period leading into a game may constitute an important
13
14 element of training periodisation adopted in the elite game. The 'three-day' pre-
15
16 match tapering strategy facilitates the gradual 'unloading' of players which will
17
18 serve to increase player readiness for the game. It is acknowledged that this type of
19
20 three-day load reduction approach does not concur with the traditional tapering
21
22 strategies reported for individual sports, whereby training load is typically reduced
23
24 over the course of 7 to 28 days pre-competition (Mujika et al., 2004). This may be a
25
26 consequence of several factors. A congested and 'ever changing' fixture schedule
27
28 restricts the amount of time available to fully prepare players, making a 'one-size
29
30 global approach' to periodisation unfeasible within elite soccer. There is also the
31
32 need for constant flexibility to allow for the management of playing times,
33
34 demanding travel schedules, and individual player 'micro-management'.
35
36
37
38
39
40
41

42 Training and match load in the current study showed limited variation between
43
44 playing positions across the season's six mesocycles. This likely reflected the
45
46 inclusion of match data in the analysis which may have masked any potential
47
48 differences in training load per se. Analysis of the loading patterns during the weekly
49
50 microcycle training days in the present study provides a more precise comparison of
51
52 positional loads. For example, sRPE-TL was lower in wide midfielders than wide
53
54 defenders on G-3 while attackers reported higher sRPE-TL on G-2 vs wide and
55
56 central midfielders and higher sRPE-TL compared with all other positions on G-1.
57
58
59
60

1
2
3 Attacking players also covered ~600 m and ~650 m more total distance compared to
4 CD and WD on G-1. In contrast to the present observations, in English Premier
5
6 League players, Malone et al. (2015) reported limited positional differences in the
7
8 days leading into a game. In Spanish reserve team players, Martin-Garcia and
9
10 colleagues (2018) reported the highest total distance in central and offensive
11
12 midfielders during the three-day lead into competition whilst wide defenders covered
13
14 the greatest high-speed running distance during the same period. Collectively, these
15
16 positional differences likely reflect the diversity in training strategies adopted by
17
18 different coaching teams which are often driven by the head coach (Akenhead and
19
20 Nassis, 2016; Weston, 2018).
21
22
23
24
25
26
27
28

29 **Conclusions**

30
31 In summary, our study has systematically quantified the training and match loads
32
33 employed by an English Premier League club during a championship winning
34
35 season. Training load across the mesocycle periods showed limited variation and
36
37 suggests that training schedules employed in elite soccer may be highly repetitive
38
39 likely reflecting the nature of the competition demands. Periodisation of training load
40
41 was evident within the weekly microcycle including the three-day period leading
42
43 into competition. This reflected the coaching teams approach to match recovery and
44
45 preparation across the long competitive period. Further research is needed to expand
46
47 our understanding of the loads encountered by elite players and the different
48
49 periodisation models adopted by coaching teams.
50
51
52
53
54
55
56
57
58
59
60

Practical Implications

The present data centers on a championship winning season and extends the limited literature by providing novel insights into the training loads encountered by elite soccer players. The present findings provide coaches and practitioners with insights into a successful periodisation strategy that was adopted during weekly microcycles in an attempt to facilitate match recovery and preparation. Such strategies are likely to be important in the modern game due to the relatively constant loading incurred across the season as a consequence of the high frequency of matches encountered by elite teams. Methodological challenges inherent in soccer, limit the ability to determine the direct influence of training load on team match physical performance and/or success and therefore our understanding of what may constitute optimal periodisation of training. Future work could therefore focus on the analysis of training load encountered by the same players under different coaches and/or periodisation strategies across extended periods of time or between seasons. By examining both the variation in load as well as factors such as performance testing, player wellness and injury rates, such approaches could represent a move towards a better understanding of how to best prepare elite players.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

Akenhead, R., and Nassis, G. P. (2016). Training load and player monitoring in high-level football: current practice and perceptions. *International Journal of Sports Physiology and Performance*, **11 (5)**, 587-593.

Akenhead, R., Harley, J., and Tweddle, S. (2016). Examining the external training load of an English Premier League football team with special reference to acceleration. *Journal of Strength and Conditioning Research*, **30 (9)**, 2424-2432.

Anderson, L., Orme, P., Di Michele, R., Close, G. L., Milsom, J., Morgans, R., Drust, B., and Morton, J. P. (2016). Quantification of seasonal long physical load in soccer players with different starting status from the English Premier League: implications for maintaining squad physical fitness. *International Journal of Sports Physiology and Performance*, **11 (8)**, 1038-1046.

Aoki, Marcelo S., Ronda, Lorena T., Marcelino, Pablo R., Drago, Gustavo., Carling, Chris., Bradley, Paul S., and Moreira, Alexandre. (2017). Monitoring Training Loads in Professional Basketball Players Engaged in a Periodized Training Program. *Journal of Strength and Conditioning Research*, **31 (2)**, 348–358.

Buchheit, M., Manouvrier, C., Cassirame, J., and Morin, J. B. (2015). Monitoring Locomotor Load in Soccer: Is Metabolic Power, Powerful? *International Journal of Sports Medicine*, **36 (14)**, 1149-1155.

1
2
3 Buchheit, M., Allen, A., Poon, T, K., Modonutti, M., Gregson, W., and Di Salvo, V.
4
5 (2014). Integrating different tracking systems in football: multiple camera semi-
6
7 automatic system, local position measurement and GPS technologies. *Journal of*
8
9 *Sports Sciences*, **32 (20)**, 1844-1857.

10
11
12
13
14 Campos-Vazquez , M. A., Mendez-Villanueva , A., Gonzalez-Jurado , J. A., León-
15
16 Prados , J. A., and Santalla, A. (2015). Relationships Between Rating-of-Perceived-
17
18 Exertion- and Heart- Rate-Derived Internal Training Load in Professional Soccer
19
20 Players: A Comparison of On-Field Integrated Training Sessions. *International*
21
22 *Journal of Sports Physiology and Performance*, **10 (5)**, 587-592.

23
24
25
26
27
28
29 Casamichana, D., and Castellano, J. (2010). Time-motion, heart rate, perceptual and
30
31 motor behaviour demands in small-sides soccer games: effects of pitch size. *Journal*
32
33 *of Sports Sciences*, **28 (14)**, 1615-1623.

34
35
36
37
38 Casamichana, D., Castellano, J., Calleja, J., San Roman, J., and Castagna, C. (2013).
39
40 Relationship between Indicators of Training Load in Soccer Players. *Journal of*
41
42 *Strength and Conditioning Research*, **27 (2)**, 369–374.

43
44
45
46
47 Clemente, F. M., Owen, A., Serra-Olivares, J., Nikolaidis, P. T., van der Linden, C.
48
49 M. I., and Mendes, B. (2019). Characterization of the weekly external load profile of
50
51 professional soccer teams from Portugal and the Netherlands. *Journal of Human*
52
53 *Kinetics*, **66**, 155-164.

1
2
3 Cnaan A, Laird NM, Slasor P. (1997). Using the general linear mixed model to
4 analyse unbalanced repeated measures and longitudinal data. *Stat Med.* **16**, 2349–
5
6 2380.
7
8

9
10
11
12 [Cook, J.A., Julious, S.A., Sones, W., Hampson, L.V., Hewitt, C., Berlin, J.A.,](#)
13 [Ashby, D., Emsley, R., Fergusson, D.A. and Walters, S.J. \(2018\). DELTA 2 guidance](#)
14 [on choosing the target difference and undertaking and reporting the sample size](#)
15 [calculation for a randomised controlled trial. *British Medical Journal*, **363**: k3750–](#)
16 [k3757.](#)
17
18
19
20
21
22
23
24

25
26 Coutts, A. J., Rampinini, E., Marcora, S. M., Castagna, C., and Impellizzeri, F. M.
27 (2009). Heart and blood lactate correlates of perceived exertion during small-sided
28 soccer games. *Journal of Science and Medicine in Sport*, **12 (1)**, 79-84.
29
30
31

32
33
34
35 Ebben, W. P., Carroll, R. M., and Simenz, C. J. (2004). Strength and conditioning
36 practices of National Hockey League strength and conditioning coaches. *Journal of*
37 *Strength and Conditioning Research*, **18 (4)**, 889-897.
38
39
40
41

42
43
44
45 Ebben, W. P., Hintz, M. J., and Simenz, C. J. (2005). Strength and conditioning
46 practices of Major League Baseball strength and conditioning coaches. *Journal of*
47 *Strength and Conditioning Research*, **19 (3)**, 538-546.
48
49
50

51
52
53 Foster, C., Florhaug, J. A., Franklin, J., Gottschall, L., Hrovatin, L. A., Parker, S.,
54 Doleshal, P., and Dodge, C. (2001). A new approach to monitoring exercise training.
55 *Journal of Strength and Conditioning Research*, **15 (1)**, 109-115.
56
57
58
59
60

1
2
3 Gabbett, T. J., and Jenkins, D. G. (2011). Relationship between training load and
4 injury in professional rugby league players. *Journal of Science and Medicine in*
5
6
7
8 Sport, 14, 204-209.
9

10
11
12 Gaudino, P., Iaia, F. M., Alberti, G., Strudwick, A. J., Atkinson, G., and Gregson, W.
13
14 (2013). Monitoring training in elite soccer players: systematic bias between running
15 speed and metabolic power data. *International Journal of Sports Medicine*, **34 (11)**,
16
17
18 963-968.
19

20
21
22
23 Hopkins, W. G., Marshall, S. W., Batterham, A. M., and Hanin, J. (2009).
24
25 Progressive Statistics for Studies in sports Medicine and Exercise Science. *Medicine*
26
27
28 *and Science in Sports and Exercise*, **41 (1)**, 3-12.
29

30
31
32
33 Impellizzeri, F. M., Rampinini, E., Coutts, A. J., Sassi, A. and Marcora, S. M.
34
35 (2004). Use of RPE-Based Training Load in Soccer. *Medicine and Science in Sports*
36
37
38 *and Exercise*, **36 (6)**, 1042-1047.
39

40
41
42 Impellizzeri, F. M., Rampinini, E., Marcora, S. M. (2005). Physiological assessment
43
44
45 of aerobic training in soccer. *Journal of Sports Sciences*, **23**, 583-592.
46

47
48
49 Jennings, D., Cormack, S., Coutts, A. J., Boyd, L. J., and Aughey, R. J. (2010).
50
51
52 Variability of GPS units for measuring distance in team sport movements.
53
54
55 *International Journal of Sports Physiology and Performance*. **5**, 565-569.
56
57
58
59
60

1
2
3 Kelly, D. M., Strudwick, A. J., Atkinson, G., Drust, B., and Gregson, W. (2016). The
4 within-participant correlation between perception of effort and heart rate-based
5 estimations of training load in elite soccer players. *Journal of Sports Sciences*, **34**
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(14), 1328-1332.

[Lenth, R.V. \(2006\). Some practical guidelines for effective sample size. *The American Statistician*, 55 \(3\), 187-193.](#)

Little, T., and Williams, A. G. (2006). Suitability of soccer Training Drills for Endurance Training. *Journal of Strength and Conditioning Research*, **20 (2)**, 316-319.

Malone, J. J., Rocco, D. M., Morgans, R., Burgess, D., Morton, J. P., and Drust, B. (2015). Seasonal Training-Load Quantification in Elite English Premier League Soccer Players, *International Journal of Sports Physiology and Performance*, **10**, 489-497.

Martin-Garcia, A., Gomez Diaz, A., Bradley, P. S., Morera, F., and Casamichana, D. (2018). Quantification of a professional football team's external load using a microcycle structure. *Journal of Strength and Conditioning Research*, (In Press.).

McGahan, J., O'Neill, C., and Burns, C. (2017). Seasonal variation of training, competition load and markers of wellness in an elite Gaelic football team. *Physical Therapy in Sport*, **28**, e24.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Moreira, A., Bilsborough, J. C., Sullivan, C. J., Cianciosi, M., Aoki, M. S., and Coutts, A. J. (2016). Training Periodization of Professional Australian Football Players During an Entire Australian Football League Season. *International Journal of Sports Physiology and Performance*, **10**, 566-571.

Morgans, R., Orme, P., Anderson, L., and Drust, B. (2014). Principles and practices of training for soccer. *Journal of Sport and Health Science*, **3** (4), 251-257.

Mujika, I., Padilla, S., Pyne, D., and Busso, T. (2004). Physiological changes associated with the pre-event taper in athletes. *Sports Medicine*, **34** (13), 891-927.

Owen, A. L., Lago-Penas, C., Miguel-Angel, G., Mendes, B., and Dellal, A. (2017). Analysis of a training mesocycle and positional quantification in elite European soccer players. *International Journal of Sports Science & Coaching*, **12** (5), 665-676.

Perneger, T. V. (1998). Whats wrong with Bonferroni adjustments. *British Medical Journal*, **316**, 1236-1238.

Portas, M. D., Harley, J. A., Barnes, C. A., and Rush, C. J. (2010). The validity and reliability of 1-Hz and 5-Hz global positioning systems for linear, multidirectional, and soccer-specific activities. *International Journal of Sports Physiology and Performance*, **5**, 448-458.

1
2
3 Randers, M. B., Mujika, I., Hewitt, A., Santisteban, J., Bischoff, R., Solano, R.,
4
5 Zubillaga, A., Peltola, E., Krstrup, P., and Mohr, M. (2010). Application of four
6
7 different football match analysis systems: a comparative study. *Journal of Sports*
8
9 *Sciences*. **28**, 171-182.

10
11
12
13
14 Scott, B. R., Lockie, R. G., Knight, T. J., Clark, A. C., and Xanne, A. K. Janse de
15
16 Jonge. (2013). A Comparison of Methods to Quantify the In-Season Training Load
17
18 of Professional Soccer Players. *International Journal of Sports Physiology and*
19
20 *Performance*, **8 (2)**, 195-202.

21
22
23
24
25
26 Stevens, T. G. A., de Ruiter, C. J., Twisk, J. W. R., Savelsbergh, G. J. P., and beek,
27
28 P. J. (2017). Quantification of in-season training load relative to match load in
29
30 professional Dutch Eredivisie football players. *Science and Medicine in Football*, **1**
31
32 **(2)**, 117-125.

33
34
35
36
37 Stoszkowski, J., and Collins, D. (2016). Sources, topics and use of knowledge by
38
39 coaches. *Journal of Sports Sciences*, **34 (9)**, 794-802.

40
41
42
43
44 Varley, M. C., Fairweather, I. H., and Aughey, R. J. (2012). Validity and reliability
45
46 of GPS for measuring instantaneous velocity during acceleration, deceleration, and
47
48 constant motion. *Journal of Sports Sciences*. **30 (2)**, 121-127.

49
50
51
52
53 Viru, A. & Viru, M., (2000). Nature of training effects. In W. E. Garrett, D. T.
54
55 Kirkundall (Eds.) *Exercise and Sport Science* (pp. 67-95). Philadelphia: Lippincott
56
57
58 Williams and Wilkins.
59
60

1
2
3 Waldron, M., Worsfold, P., Twist, C., and Lamb, K. (2011). Concurrent validity and
4 test-retest reliability of a global positioning system (GPS) and timing gate to assess
5 sprint performance variables. *Journal of Sports Sciences*, **29**, 1613-1619.
6
7
8
9

10
11
12 Weston, M. (2018). Training load monitoring in elite English soccer: a comparison
13 of practices and perceptions between coaches and practitioners. *Science and*
14 *Medicine in Football*, 2 (3), 216-224.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only




Figure 1. Diagrammatic representation of the experimental design used in the current study. Each small block represents individual weeks within the annual training cycle, with larger blocks showing the 6-week mesocycle phases of the competitive season.




Figure 2. Mean \pm SD number of training sessions and competitive games by playing position during the 2012-13 season.




Figure 3. Mean \pm SD training session duration by playing position during the 2012-13 season (central defender [CD]; wide defender [WD]; central midfielder [CM]; wide midfielder [WM]; attacker [A]).




Figure 4. Mean \pm SD sRPE-TL for training day's pre- and post-competitive match and match-day between positions. Subscripts denote moderate (M), large (L), and very large (V).

Day Type: **G-3**; L vs. G-2, V vs. G-1, M vs. G+2 and G+3. **G-1**; V vs. G-3, M vs. G-2, L vs. G+2, and G+3. **MD**; V vs. G-3, G-2, G-1, G+2 and G+3. Day Type x Playing Position: **G-3**; **WM**, M vs. WD. **G-2**; **A**, M vs. WM and CM. **G-1**; **A**, M vs. CD, WD, and WM. **MD**; **CD**, L vs. CM, M vs. WM and A. **WD**, L vs. CM, M vs. WM and A. **G+2**; **A**, M vs. CD, CM, and WM.




Figure 5. Mean \pm SD total distance for training day's pre- and post-competitive match and match day between positions. Subscripts denote moderate (M), large (L), and very large (V). Day Type: G-3; M vs. G-2 and G+2. G-1; V vs. G-3, M vs. G-2, L vs. G+2 and G+3. MD; V vs. G-3, G-2, G-1, G+2, and G+3. Day Type x Playing Position: G-1; A, M vs. CD and WD. MD; CD, M vs. CM, WM, and A. WD, M vs. CM, WM, and A. G+2; A, M vs. CD.




Figure 6. Mean \pm SD total high-speed distance for training day's pre- and post-competitive match and match day between positions. Subscripts denote (M), large (L), and very large (V).

Day Type: **G-1;** M vs. G-3, G+2 and G+3. **MD;** V vs. G-3, G-2, G-1, G+2, and G+3. Day

Type x Playing Position: **MD;** WD, M vs. CD and A. **CM,** M vs. CD and A.




Figure 7. Mean \pm SD total very high-speed distance for training day's pre- and post-competitive match and match day between positions. Subscripts denote moderate (M), large (L), and very large (V). Day Type: **G-1**; L vs. G-3, M vs. G-2, G+2, and G+3. **MD**; V vs. G-3, G-2, G-1, G+2, and G+3. Day Type x Playing Position: **MD**; **CD**, L vs. WD and CM, M vs. WM and A.