

CHEMISTRY 
A EUROPEAN JOURNAL

Supporting Information

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2006

Quantification of Ultrasound-Induced Chain Scission in Pd^{II}-Phosphine Coordination Polymers

Jos M.J. Paulusse, Jeroen P.J. Huijbers, and Rint P. Sijbesma*^[a]

[a] Laboratory of Macromolecular and Organic Chemistry, Eindhoven University of Technology

P.O. Box 513, 5600 MB Eindhoven (The Netherlands)

E-mail: R.P.Sijbesma@tue.nl; Fax: +40-2451036; Tel: +40-2472655




Figure S1. Ligand exchange of coordination polymer **3** without sonication in the presence of 1.7 equivalents of stopper complex **4** (quantified with SEC-RI).




Figure S2. SEC-traces measured at regular intervals during sonication of coordination polymer **3** in toluene (10 g/L, 1.45 mM).




Figure S3. SEC-traces measured at regular intervals during sonication of coordination polymer **3** in toluene (10 g/L, 1.45 mM) in the presence of 1.7 equivalents of stopper **4**.




Figure S4. SEC-traces measured at regular intervals during sonication of coordination polymer **3** in toluene (10 g/L, 1.45 mM) in the presence of 10 equivalents of stopper **4**.




Figure S5. SEC-traces measured at regular intervals during sonication of coordination polymer **3** in toluene (10 g/L, 1.45 mM) in the presence of 60 equivalents of stopper **4**.




Figure S6. ^{31}P $\{^1\text{H}\}$ NMR spectrum of coordination polymer **3** in toluene (10 g/L, 1.45 mM) in the presence of 10 equivalents of stopper **4**, before sonication




Figure S7. ^{31}P $\{^1\text{H}\}$ NMR spectrum of coordination polymer **3** in toluene (10 g/L, 1.45 mM) in the presence of 10 equivalents of stopper **4**, after 3 hours of sonication




Figure S8. ^{31}P $\{^1\text{H}\}$ NMR spectra of coordination polymer **8** (top), stopper complex **4** (middle), mixture of complexes **4** and **8** after 30 minutes (bottom).

Ligand exchange with bridged complex 8: Bridged palladium(II) complex **8** was mixed with 0.5 equivalents of stopper **4**. After 30 minutes new signals appeared, approximately 15% in total. Signals corresponding to the heterocomplex (10% of total) were identified by their characteristic coupling constant of $J_{\text{P-P}} = 527$ Hz. Peaks A, B and C could not be identified, but most probably correspond to a hetero-bridged complex (A and C) and the diphenylphosphine homo-bridged complex (B).