

 Open access • Posted Content • DOI:10.1101/078428

## Quantifying the evolutionary potential and constraints of a drug-targeted viral protein — [Source link](#)

[Lei Dai](#), [Yushen Du](#), [Hangfei Qi](#), [Nicholas C. Wu](#) ...+2 more authors

**Institutions:** [University of California, Los Angeles](#)

**Published on:** 30 Sep 2016 - [bioRxiv](#) (Cold Spring Harbor Laboratory)

**Topics:** [Robustness \(evolution\)](#), [Mutation rate](#), [Fitness landscape](#), [Selection coefficient](#) and [Viral protein](#)

Related papers:

- [Adaptive potential of a drug-targeted viral protein as a function of positive selection](#)
- [Adaptive potential of a drug-targeted viral protein as a function of environmental stress](#)
- [Assessing in vivo mutation frequencies and creating a high-resolution genome-wide map of fitness costs of Hepatitis C virus](#)
- [Evolutionary dynamics of viral escape under antibodies stress: A biophysical model.](#)
- [Costs and benefits of mutational robustness in RNA viruses.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/quantifying-the-evolutionary-potential-and-constraints-of-a-3ualb04rso>

# 1 **Adaptive potential of a drug-targeted viral protein as a function of environmental stress**

2  
3 Lei Dai<sup>1, 2\*</sup>, Yushen Du<sup>1\*</sup>, Hangfei Qi<sup>1</sup>, Christian D. Huber<sup>2</sup>, Nicholas C. Wu<sup>3</sup>, Ergang Wang<sup>1</sup>, James O.  
4 Lloyd-Smith<sup>2</sup>, Ren Sun<sup>1</sup>

5  
6 <sup>1</sup>Department of Molecular and Medical Pharmacology

7 <sup>2</sup>Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los Angeles, United  
8 States

9 <sup>3</sup>Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla,  
10 United States

11  
12 \*These authors contributed equally to this work.

13  
14 Corresponding author:

15 Lei Dai, Ph.D.

16 Email: [leidai@ucla.edu](mailto:leidai@ucla.edu)

17 Ren Sun, Ph.D.

18 Email: [rsun@mednet.ucla.edu](mailto:rsun@mednet.ucla.edu)

## 19 **Abstract**

20 RNA viruses are notorious for their ability to evolve rapidly under selection in novel environments. It is known  
21 that the high mutation rate of RNA viruses can generate huge genetic diversity to facilitate viral adaptation.  
22 However, less attention has been paid to the underlying fitness landscape that represents the selection  
23 forces on viral genomes. Here we systematically quantified the distribution of fitness effects (DFE) of about  
24 1,600 single amino acid substitutions in the drug-targeted region of NS5A protein of Hepatitis C Virus (HCV).  
25 We found that the majority of non-synonymous substitutions incur large fitness costs, suggesting that NS5A  
26 protein is highly optimized in natural conditions. We characterized the adaptive potential of HCV by  
27 subjecting the mutant viruses to selection by the antiviral drug Daclatasvir. Both the selection coefficient and  
28 the number of beneficial mutations are found to increase with the level of environmental stress, which is  
29 modulated by the concentration of Daclatasvir. The changes in the spectrum of beneficial mutations in NS5A  
30 protein can be explained by a pharmacodynamics model describing viral fitness as a function of drug  
31 concentration. We test theoretical predictions regarding the distribution of beneficial fitness effects of mutations.  
32 We also interpret the data in the context of Fisher's Geometric Model and find an increased distance to optimum  
33 as a function of environmental stress. Finally, we show that replication fitness of viruses is correlated with the  
34 pattern of sequence conservation in nature and viral evolution is constrained by the need to maintain protein  
35 stability.

## 37 Introduction

38 In our evolutionary battles with microbial pathogens, RNA viruses are among the most formidable foes.  
39 HIV-1 and Hepatitis C Virus acquire drug resistance in patients under antiviral therapy. Influenza and Ebola  
40 virus cross the species barrier to infect human hosts. Understanding the evolution of RNA viruses is  
41 therefore of paramount importance for developing antivirals and vaccines and assessing the risk of future  
42 emergence events (Goldberg *et al.* 2012; Domingo *et al.* 2012; Metcalf *et al.* 2015). Comprehensive  
43 characterization of viral fitness landscapes, and the principles underpinning them, will provide us with a map  
44 of evolutionary pathways accessible to viruses and guide our design of effective strategies to limit antiviral  
45 resistance, immune escape and cross-species transmission (Turner and Elena 2000; Ke *et al.* 2015; Barton  
46 *et al.* 2016).

47 Although the concept of fitness landscapes has been around for a long time (Wright 1932), we still know  
48 little about their properties in real biological systems. Previous empirical studies of fitness landscapes have  
49 been constrained by very limited sampling of sequence space. In a typical study, mutants are generated by  
50 site-directed mutagenesis and assayed for growth rate individually. We and others have recently developed  
51 a high-throughput technique, often referred to as “deep mutational scanning” or “quantitative high-resolution  
52 genetics”, to profile the fitness effect of mutations by integrating deep sequencing with selection experiments  
53 in vitro or in vivo (Hietpas *et al.* 2011; Wu *et al.* 2013; Thyagarajan and Bloom 2014; Qi *et al.* 2014; Fowler  
54 and Fields 2014). This novel application of next generation sequencing has raised an exciting prospect of  
55 large-scale fitness measurements (Olson *et al.* 2014; Puchta *et al.* 2015; Li *et al.* 2016; Wu *et al.* 2016) and a  
56 revolution in our understanding of molecular evolution (He and Liu 2016).

57 The distribution of fitness effects (DFE) of mutations is a fundamental entity in genetics and reveals the  
58 local structure of a fitness landscape (Burch and Chao 2000; Eyre-Walker and Keightley 2007; Hietpas *et al.*  
59 2011; Desai 2013; Jacquier *et al.* 2013; Bataillon and Bailey 2014; Chevereau *et al.* 2015; Bank *et al.* 2015).  
60 Deleterious mutations are usually abundant and impose severe constraints on the accessibility of fitness  
61 landscapes. In contrast, beneficial mutations are rare and provide the raw materials of adaptation.  
62 Quantifying the DFE of viruses is crucial for understanding how these pathogens evolve to acquire drug  
63 resistance and surmount other evolutionary challenges.

64 Most empirical studies of the DFE have been performed in a single, static environment (Eyre-Walker  
65 and Keightley 2007; Bataillon and Bailey 2014). A central challenge is to characterize the DFE, and its  
66 determinants, in fluctuating or heterogeneous environments where evolution typically occurs (e.g. fluctuating

67 drug concentrations or a gradient across space). Previous studies on yeast have investigated the change in  
68 DFE across different levels of temperature and salinity (Hietpas *et al.* 2013; Bank *et al.* 2014). For bacteria,  
69 the fitness effects of mutations at different drug concentrations have been studied (Firnberg *et al.* 2014).  
70 One recent study has demonstrated that drug concentration modulates the shape of the DFE and  
71 determines the evolvability under new environments (Stiffler *et al.* 2015). In another study, the implications of  
72 differing drug concentrations on the adaptive landscape have been examined in the context of resistance  
73 evolution (Ogbunugafor *et al.* 2016). For viruses, the fitness effects of mutations have been measured  
74 across different hosts (Lalić *et al.* 2011; Vale *et al.* 2012). The shape of DFE of viruses has been inferred  
75 from experimentally passaged populations (Foll *et al.* 2014) and from patient data (Renzette *et al.* 2017).

76 In this study, we profile the DFE of ~1,600 single amino acid substitutions in a drug-targeted viral protein  
77 by combining selection experiment of a mutant library and deep sequencing. We examine the changes in  
78 DFE under varying levels of environmental stress by tuning the concentration of an antiviral drug. We test  
79 theoretical predictions regarding the distribution of beneficial fitness effects of mutations (Orr 2003). We also  
80 interpret the data in the context of Fisher's Geometric Model (Martin and Lenormand 2006b) and find an  
81 increased distance to optimum as a function of environmental stress. Finally, we show that replication fitness  
82 of viruses is correlated with the pattern of sequence conservation in nature and viral evolution is constrained  
83 by the need to maintain protein stability.

## 84 85 **Results**

### 86 **Profiling the fitness landscape of the drug-interacting domain of HCV NS5A protein**

87 The system used in our study is Hepatitis C Virus (HCV), a positive sense single-stranded RNA virus with a  
88 genome of ~9.6 kb. HCV has been studied extensively in the past two decades in patients and in laboratory  
89 and provides an excellent model system to study viral evolution. We applied high-throughput assays to map  
90 the fitness effects of all single amino acid substitutions in domain IA (amino acid 18-103) of HCV NS5A  
91 protein (Methods). This domain is the target of several directly-acting antiviral drugs, including the potent  
92 HCV NS5A inhibitor Daclatasvir (DCV) (Gao *et al.* 2010).

93 To study the DFE of mutations of HCV NS5A protein, we conducted new selection experiments using a  
94 previously constructed saturation mutagenesis library of mutant viruses (Qi *et al.* 2014). Briefly, each codon  
95 in the mutated region was randomized to cover all possible single amino acid substitutions. We observed  
96 2520 non-synonymous mutations in the plasmid library, as well as 105 synonymous mutations. After  
97 transfection to reconstitute mutant viruses, we performed selection in an HCV cell culture system

(Lindenbach *et al.* 2005; Wakita *et al.* 2005). The relative fitness of a mutant virus to the wild-type virus was calculated based on the changes in frequency of the mutant virus and the wild-type virus after one round of selection in cell culture (Supplementary Figure 1). In our selection experiment, we grew 5 small sub-libraries (~500 mutants each) separately to reduce the noise in fitness measurements (Methods). The fitness data reported in this study is highly correlated to an independent experiment using the same plasmid library (Supplementary Figure 2) (Qi *et al.* 2014).

Our experiment provides a comprehensive profiling of the fitness effect of single amino acid substitutions (1565 out of 1634 possible substitutions, after filtering out low frequency mutants in the plasmid library). We grouped together non-synonymous mutations leading to the same amino acid substitution. As expected, the fitness effects of synonymous mutations were nearly neutral, while most non-synonymous mutations were deleterious (Figure 1). We found that the majority of single amino acid mutations had fitness costs and more than half of them were found to be significantly deleterious, or “lethal” (Methods). The fraction of lethal mutations (not shown explicitly in Figure 1) is 57.0% (932/1634) for single amino acid substitutions, 1.0% (1/105) for synonymous mutations and 90.6% (77/85) for nonsense mutations. The low tolerance of non-synonymous mutations in HCV NS5A, which is an essential protein for viral replication, is consistent with previous small-scale mutagenesis studies of RNA viruses (Sanjuan *et al.* 2004). Our data support the view that RNA viruses are very sensitive to the effect of deleterious mutations, possibly due to the compactness of their genomes (Elena *et al.* 2006; Rihn *et al.* 2013).

Using the distribution of fitness effects of synonymous mutations as a benchmark for neutrality, we identified that only 2.3% (37/1634) of single amino acid mutations are beneficial (Methods). The estimated fraction of beneficial mutations is consistent with previous small-scale mutagenesis studies in viruses including bacteriophages, vesicular stomatitis virus, etc. (Sanjuan *et al.* 2004; Burch *et al.* 2007; Silander *et al.* 2007; Eyre-Walker and Keightley 2007). Our results indicate that HCV NS5A protein is under strong purifying selection, suggesting that viral proteins are highly optimized in their natural conditions.

### **Adaptive potential as a function of environmental stress**

Beneficial mutations are the raw materials of protein adaptation (Eyre-Walker and Keightley 2007). In this study, we aimed to study the role of environmental stress in modulating the adaptive potential of drug-targeted viral proteins. In an independent study (Qi *et al.* 2014), the mutant library of HCV NS5A protein was selected under a single drug concentration ([DCV]=20 pM) to profile the effects of mutations on drug resistance. In this study, we selected the mutant library at 10, 40 and 100 pM of DCV. The drug concentrations were chosen based on in vitro IC<sub>50</sub> of wild type HCV virus (~20 pM) to represent different

129 levels of environmental stress (mild, intermediate and strong).

130 By tuning the concentration of DCV, we observed a change in the DFE (Supplementary Table 1&2),  
131 particularly of beneficial mutations (Figure 2A). At higher drug concentrations, we observed an increase in  
132 the median selection coefficient (Figure 2B) as well as the total number of beneficial mutations (Figure 2C,  
133 Supplementary Table 3). We further tested whether the shape of this distribution changed under drug  
134 selection. Previous empirical studies supported the hypothesis that the DFE of beneficial mutations is  
135 exponential (Orr 1998, 2003, 2006; Imhof and Schlötterer 2001; Sanjuan *et al.* 2004; Rokyta *et al.* 2005;  
136 Cowperthwaite *et al.* 2005; Kassen and Bataillon 2006; Burch *et al.* 2007; Carrasco *et al.* 2007; MacLean  
137 and Buckling 2009; Peris *et al.* 2010; Bataillon *et al.* 2011). Following a maximum likelihood approach, we fit  
138 the DFE of beneficial mutations to the Generalized Pareto Distribution (Supplementary Figure 3, Methods).  
139 The fitted distribution is described by two parameters: a scale parameter ( $\tau$ ), and a shape parameter ( $\kappa$ ) that  
140 determines the behavior of the distribution's tail. Using a likelihood-ratio test (Beisel *et al.* 2007), we found  
141 that our data are consistent with the null hypothesis that the DFE of beneficial mutations is exponential ( $\kappa = 0$ )  
142 (Supplementary Table 4).

143 Furthermore, we used a maximum-likelihood approach to fit a displaced-gamma distribution to the DFE  
144 to estimate the distance to the phenotypic optimum in Fisher's Geometric Model (FGM) (Martin and  
145 Lenormand 2006b; Bank *et al.* 2014) (Supplementary Figure 4). The displaced-gamma distribution has the  
146 shape of a negative gamma distribution, shifted by a parameter  $s_0$  that indicates the distance of the initial  
147 genotype (i.e. wild-type) to the optimum (Methods). Estimated distances to the optimum under different  
148 conditions are summarized in Supplementary Table 5. In accordance with theoretical expectations, we found  
149 that the distance to the optimum increased as the level of environmental stress increased (i.e. increasing  
150 drug concentration).

### 151 **The effects of mutations on drug resistance and replication fitness**

152 Our results show that the adaptive potential of proteins is modulated by the strength of environmental stress.  
153 The changing spectra of beneficial mutations upon drug treatment can be explained by a  
154 pharmacodynamics model describing viral fitness as a function of drug concentration (i.e. phenotype-fitness  
155 mapping) (Figure 3A).

$$156 \quad f = f_0 \frac{IC_{50}}{IC_{50} + [drug]}$$

157 where  $f_0$  is the fitness without drug selection and  $IC_{50}$  is the half inhibitory concentration. The absolute  
158 fitness  $f$  decreases with drug concentration [*drug*]. In this paper, we define a drug-resistant mutant as any  
159 viral variant that is less inhibited than the wild type for some drug concentration, i.e. higher  $IC_{50}$  than  
160 wild-type (Rosenbloom *et al.* 2012).

161 Mutations that reduce a protein's binding affinity to drug molecules (i.e. less inhibited by the drug) may  
162 come with a fitness cost (i.e. smaller  $f_0$  than wild-type). Thus, a drug-resistant mutant that is deleterious in  
163 the absence of drug may become beneficial under drug selection, leading to an increase in the number of  
164 beneficial mutations. Moreover, the relative fitness of the drug-resistant mutant is expected to increase with  
165 stronger selection pressure (Figure 3A, dashed line). The dose response curves were previously measured  
166 for a set of mutants constructed by site-directed mutagenesis (Supplementary Figure 5) (Qi *et al.* 2014).  
167 Indeed, we found that the relative fitness of validated drug-resistant mutants increased at higher drug  
168 concentration (Figure 3B); in contrast, drug-sensitive mutants became less fit under drug selection.

169 Furthermore, we showed that the effects of mutations on drug resistance can be estimated from the  
170 fitness data and the results were generally consistent with estimates based on the dose response curves  
171 (Supplementary Figure 6, Methods). Among all the non-lethal single amino acid substitutions profiled in our  
172 HCV NS5A protein library, we found that roughly half of the mutations increased resistance to DCV (i.e.  
173 improved new function) at the expense of replication fitness without drug (Figure 3C, Spearman's  $\rho = -0.13$ ,  
174  $p = 8.3 \times 10^{-4}$ ). This group of resistance mutations (lower right section in Figure 3C) can become beneficial  
175 when the positive selection imposed by the antiviral drug is strong, leading to an increase in the supply of  
176 beneficial mutations at higher drug concentrations. We found no association between drug resistance and  
177 fitness cost (Fisher's exact test,  $p = 0.26$ ), suggesting that there is no or very weak tradeoff in adaptation of  
178 NS5A protein under the two different environments (i.e. with and without DCV selection).

### 179 **Deleterious mutations as evolutionary constraints**

180 While beneficial mutations open up adaptive pathways to genotypes with higher fitness, mutations that  
181 severely reduce replication fitness impose constraints on the evolution of viruses and are less likely to  
182 contribute to adaptation through gain of function. We analyzed sequence diversity of HCV sequences  
183 identified in patients from the HCV sequence database of Los Alamos National Lab (Methods). As expected,  
184 we found that amino acid sites with high fitness costs are often highly conserved (Figure 4A). The sequence  
185 diversity at each site was highly correlated to the replication fitness (the median fitness of observed mutants



at each site) measured in our study (Spearman's  $\rho=0.82$ ,  $p=1.8\times 10^{-21}$ ).

To understand the biophysical basis of mutational effects (Liberles *et al.* 2012), we took advantage of the available structural information (Supplementary Figure 7A). The crystal structure of NS5A domain I is available excluding the amphipathic helix at N-terminus (Tellinghuisen *et al.* 2005; Love *et al.* 2009). We found that the fitness effects of deleterious mutations at buried sites (i.e. with lower solvent accessibility) were more pronounced than those at surface exposed sites (Figure 4B, Spearman's  $\rho=0.51$ ,  $p=5.1\times 10^{-6}$ ; Supplementary Figure 8A) (Ramsey *et al.* 2011). Moreover, we performed simulations of protein stability for individual mutants using PyRosetta (Methods) (Das and Baker 2008; Chaudhury *et al.* 2010). A mutation with  $\Delta\Delta G>0$ , i.e. shifting the free energy difference to favor the unfolded state, is expected to destabilize the protein. We found that mutations that decreased protein stability led to reduced viral fitness (Figure 4C, Spearman's  $\rho= -0.57$ ,  $p=1.5\times 10^{-7}$ ). For example, mutations at a stretch of highly conserved residues (F88-N91) that run through the core of NS5A protein tended to destabilize the protein and significantly reduced the viral fitness. Mutations that increase  $\Delta\Delta G$  beyond a threshold ( $\sim 5$  Rosetta Energy Unit) were mostly lethal (Supplementary Figure 8B). This is consistent with the threshold robustness model, which predicts that proteins become unfolded after using up the stability margin (Bloom *et al.* 2005; Wylie and Shakhnovich 2011; Olson *et al.* 2014). Also, we note that mutations can be deleterious because they impair protein function rather than destabilize the protein, so the correlation between protein stability and fitness is not expected to be perfect. The level of correlation between  $\Delta\Delta G$  and fitness that we observed is similar to previous studies in other proteins (Firnberg *et al.* 2014; Wu *et al.* 2015).

## Discussion

Site-directed mutagenesis and experimental evolution are traditional approaches to examine the DFE (Domingo-Calap *et al.* 2009; Sanjuán 2010; Levy *et al.* 2015; Visher *et al.* 2016). Both methods provide pivotal insights into the shape of the DFE, yet with limitations. The site-directed mutagenesis approach requires fitness assays for each individual mutant and can only provide a sparse sampling of mutations. In experimental evolution, the sampling of sequence space via de novo mutations is biased towards large-effect beneficial mutations, as they are more likely to fix in the population. In contrast, the deep mutational scanning approach (Fowler and Fields 2014), which utilizes high-throughput sequencing to simultaneously assay the fitness or phenotype of a library of mutants, allows for unbiased and large-scale sampling of fitness landscapes and thus is ideal for studying the characteristics of empirical DFE. The downside of this high-throughput approach is that the fitness measurements can be noisy, especially for

217 large mutant libraries (Matuszewski *et al.* 2016). In our experiment, we divided the mutant library into smaller  
218 sub-libraries (~500 mutants) in selection experiments. We compared the data to an independent experiment  
219 and found that the fitness estimates were largely reproducible (Supplementary Figure 2). We also showed  
220 that the observed change in the DFE under different conditions was consistent with validation experiments  
221 (Figure 3). Since this study is focused on the properties of the entire distribution of mutations rather than the  
222 effects of specific mutations, our findings on the general patterns of DFE are robust to the errors in fitness  
223 estimates. Our study quantified the fitness effects of single amino acid substitutions in the drug-targeted  
224 region of an essential viral protein. In general, the empirical DFE of HCV NS5A was consistent with previous  
225 findings that viral proteins were highly optimized in the natural condition and very sensitive to the effects of  
226 deleterious mutations.

227 One crucial but often overlooked point is that DFE will vary as a function of the environment (Martin and  
228 Lenormand 2006a; Lalić *et al.* 2011; Stiffler *et al.* 2015). In the study by Stiffler *et al.* 2015, the level of  
229 environmental stress is controlled by ampicillin concentration. Because TEM-1's function is to degrade  
230 ampicillin, deleterious mutations that impair the enzyme function ("loss-of-function") would become more  
231 deleterious at higher dose of ampicillin. In our system, we do not expect the dose of Daclatasvir to alter the  
232 strength of purifying selection on maintaining HCV NS5A protein's function in viral replication. Indeed, we do  
233 not find much difference on the deleterious side of DFE across different environments. Instead, we have  
234 observed significant changes on the beneficial side of DFE as a function of the drug dose. Because HCV  
235 NS5A protein is not well adapted in the novel environment of Daclatasvir selection, the effect of drug  
236 resistance mutations ("gain-of-function") becomes more beneficial at higher drug dose. Moreover, due  
237 to the pleiotropic effect of mutations on drug resistance and replication fitness (Figure 3), there is an  
238 increasing supply of beneficial mutations at higher drug dose.

239 Although different systems have distinct protein-drug interactions that lead to different resistance  
240 profiles (Robinson *et al.* 2011), the results in our study provide a general framework to study DFE of  
241 drug-targeted proteins. Future studies along this line will further our understanding of how proteins evolve  
242 new functions under the constraint of maintaining their original function (Soskine and Tawfik 2010), as  
243 exemplified in the evolution of resistance to directly-acting antiviral drugs (Rosenbloom *et al.* 2012).  
244 Quantifying the characteristics of DFE of drug-targeted proteins under different environments (e.g. varying  
245 levels of environmental stress, or conflicting selection pressures), would allow us to assess repeatability in  
246 the outcomes of viral evolution (de Visser and Krug 2014) and guide the design of therapies to minimize  
247 drug resistance (Ogbunugafor *et al.* 2016).

248

## 249 **Conclusions**

250 Many viruses adapt rapidly to novel selection pressures, such as antiviral drugs. Understanding how  
251 pathogens evolve under drug selection is critical for the success of antiviral therapy against human  
252 pathogens. By combining deep sequencing with selection experiments in cell culture, we have quantified the  
253 distribution of fitness effects of mutations in the drug-targeted domain of Hepatitis C Virus NS5A protein. Our  
254 results indicate that the majority of single amino acid substitutions in NS5A protein incur large fitness costs.  
255 By subjecting the mutant viruses to selection under an antiviral drug, we find that the adaptive potential of  
256 viral proteins in a novel environment is modulated by the level of environmental stress. We test theoretical  
257 predictions regarding the distribution of fitness effects of mutations. Finally, we show that viral evolution is  
258 constrained by the need to maintain protein stability.

259

## 260 **Materials and Methods**

### 261 **Mutagenesis**

262 The mutant library of HCV NS5A protein domain IA (86 amino acids) was constructed using saturation  
263 mutagenesis as previously described (Qi *et al.* 2014). In brief, the entire region was divided into five  
264 sub-libraries each containing 17-18 amino acids (~500 mutants in each sub-library). NNK (N: A/T/C/G, K:  
265 T/G) was used to replace each amino acid. The oligos, each of which contains one random codon, were  
266 synthesized by IDT. The mutated region was ligated to the flanking constant regions, subcloned into the  
267 pFNX-HCV plasmid and then transformed into bacteria. The pFNX-HCV plasmid carrying the viral genome  
268 was synthesized in Dr. Ren Sun's lab based on the chimeric sequence of genotype 2a HCV strains J6/JFH1.

### 269 **Cell culture**

270 The human hepatoma cell line (Huh-7.5.1) was provided by Dr. Francis Chisari from the Scripps Research  
271 Institute, La Jolla. The cells were cultured in T-75 tissue culture flasks (Genesee Scientific) at 37 °C with 5%  
272 CO<sub>2</sub>. The complete growth medium contained Dulbecco's Modified Eagle's Medium (Corning Cellgro), 10%  
273 heat-inactivated Fetal Bovine Serum (Omega Scientific), 10 mM HEPES (Life Technologies), 1x MEM  
274 Non-Essential Amino Acids Solution (Life Technologies) and 1x Penicillin-Streptomycin-Glutamine (Life  
275 Technologies).

### 276 **Selection of mutant viruses**

277 Plasmid mutant library was transcribed *in vitro* using T7 RiboMAX Express Large Scale RNA Production  
278 System (Promega) and purified by PureLink RNA Mini Kit (Life Technologies). 10 µg of *in vitro* transcribed

279 RNA was used to transfect 4 million Huh-7.5.1 cells via electroporation by Bio-Rad Gene Pulser (246 V, 950  
280  $\mu$ F). The supernatant was collected 6 days post transfection and virus titer was determined by  
281 immunofluorescence assay. The viruses collected after transfection were used to infect ~2 million Huh-7.5.1  
282 cells with an MOI at around 0.1-0.2. The five sub-libraries were passaged for selection separately. For the  
283 three different levels of selection pressure, the growth media was supplemented with 10 pM, 40 pM and 100  
284 pM HCV NS5A inhibitor Daclatasvir (BMS-790052), respectively. The supernatant was collected at 6 days  
285 post infection.

### 286 **Preparation of Illumina sequencing samples**

287 For each sample, viral RNA was extracted from 700  $\mu$ l supernatant collected after transfection and after  
288 selection using QIAamp Viral RNA Mini Kit (Qiagen). Extracted RNA was reverse transcribed into cDNA by  
289 SuperScript III Reverse Transcriptase Kit (Life Technologies). The targeted region in NS5A (51-54 nt) was  
290 PCR amplified using KOD Hot Start DNA polymerase (Novagen). The Eppendorf thermocycler was set as  
291 following: 2 min at 95 °C; 25 to 35 three-step cycles of 20 s at 95 °C, 15 s at 52-56 °C (sub-library #1, 52 °C;  
292 #2, 52 °C; #3, 52 °C; #4, 56 °C; #5, 54 °C) and 25s at 68 °C; 1 min at 68 °C. The number of PCR cycles are  
293 chosen based on the copy number of cDNA templates as determined by qPCR (Bio-Rad). The PCR primers  
294 are listed in Supplementary Table 6. The PCR products were purified using PureLink PCR Purification Kit  
295 (Life Technologies) and prepared for Illumina HiSeq 2000 sequencing (paired-end 100 bp) following  
296 5'-phosphorylation using T4 Polynucleotide Kinase (New England BioLabs), 3' dA-tailing using dA-tailing  
297 module (New England BioLabs), and TA ligation of the adapter using T4 DNA ligase (Life Technologies).  
298 Each sample was tagged with a unique 3-bp customized barcodes, which were part of the adapter sequence  
299 and were sequenced as the first three nucleotides in both the forward and reverse reads (Wu *et al.* 2015)  
300 (Supplementary Table 7).

### 301 **Analysis of Illumina sequencing data**

302 The sequencing data were parsed by SeqIO function of BioPython. The reads from different samples were  
303 de-multiplexed by the barcodes and mapped to the entire mutated region in NS5A by allowing at maximum 5  
304 mismatches with the reference genome (Supplementary Data 3) (Qi *et al.* 2014). Since both forward and  
305 reverse reads cover the whole amplicon, we used paired reads to correct for sequencing errors. A mutation  
306 was called only if it was observed in both reads and the quality score at the corresponding position was at  
307 least 30. Sequencing reads containing mutations not supposed to appear in our single-codon mutant library  
308 were excluded from downstream analysis. The sequencing depth for each sub-library is at least  $\sim 10^5$  and  
309 two orders of magnitude higher than the library complexity.

## 310 Calculation of relative fitness

311 For each condition of selection experiments (i.e. different concentration of Daclatasvir [DCV]), the relative  
312 fitness (RF) of a mutant virus to the wild-type virus is calculated by the relative changes in frequency after  
313 selection,

$$314 \quad RF_{mut}([DCV]) = \left( \frac{f_{mut}^{T=2}}{f_{mut}^{T=1}} \right) / \left( \frac{f_{WT}^{T=2}}{f_{WT}^{T=1}} \right)$$

315 where  $f_{mut}^{T=round}$  and  $f_{WT}^{T=round}$  is the frequency of the mutant virus and the wild-type virus at round 1 (after  
316 transfection) or round 2 (after infection). The fitness of wild-type virus is normalized to 1. The fitness values  
317 estimated from one round (round 1 to round 2) have been shown to be highly consistent to estimated based  
318 round 0 to round 1 (Supplementary Figure 2), and estimates from multiple rounds of selection (Qi *et al.*  
319 2014). A mutant was labeled as “missing” if the mutant’s frequency in the plasmid library was less than  
320 0.0005 (RF=NaN, see Supplementary Data 1 and 2). A mutant was labeled as “lethal” if the mutant’s  
321 frequency after transfection was less than 0.0005, or its frequency after infection was 0 (RF=0) (Qi *et al.*  
322 2014).

323 The selection coefficient is defined in the context of discrete generations (Chevin 2010)

$$324 \quad s_{mut} = \log(RF_{mut})$$

325 The threshold for beneficial mutations is chosen as  $2\sigma_{silent}$ , where  $\sigma_{silent}$  is the standard deviation of the  
326 selection coefficients of synonymous mutations (Figure 1). The fitness effects of non-synonymous mutations  
327 leading to the same amino acid substitution were averaged to estimate the fitness effect of the given single  
328 amino acid substitution.

## 329 Fitting the distribution of fitness effects of beneficial mutations

330 The distribution of selection coefficients of beneficial mutations were fitted to a Generalized Pareto  
331 Distribution following a maximum likelihood approach (Beisel *et al.* 2007),

$$332 \quad F(x|\kappa, \tau) = \begin{cases} 1 - \left(1 + \frac{\kappa}{\tau}x\right)^{-1/\kappa}, x \geq 0, \text{ if } \kappa > 0 & \text{(Frechet)} \\ 1 - \left(1 + \frac{\kappa}{\tau}x\right)^{-1/\kappa}, 0 \leq x < -\frac{\tau}{\kappa}, \text{ if } \kappa < 0 & \text{(Weibull)} \\ 1 - e^{-x/\tau}, x \geq 0, \text{ if } \kappa = 0 & \text{(Gumbel)} \end{cases}$$

333 Only mutations with selection coefficients higher than the beneficial threshold  $2\sigma_{\text{silent}}$  were included in the  
334 distribution of beneficial mutations. The selection coefficients were normalized to the beneficial threshold.  
335 The shape parameter  $\kappa$  determines the tail behavior of the distribution, which can be divided into three  
336 domains of attraction: Gumbel domain (exponential tail,  $\kappa = 0$ ), Weibull domain (truncated tail,  $\kappa < 0$ ) and  
337 Fréchet domain (heavy tail,  $\kappa > 0$ ). For each selection condition, a likelihood ratio test is performed to  
338 evaluate whether the null hypothesis  $\kappa = 0$  (exponential distribution) can be rejected.

### 339 **Fitting the distribution of fitness effects to Fisher's Geometrical model**

340 Fisher's Geometrical Model predicts that the distribution of fitness effects of mutations is distributed  
341 according to a negative displaced gamma distribution (Martin and Lenormand 2006a, Bank et al. 2014). This  
342 distribution has a shape parameter ( $\alpha$ ), a scale parameter ( $\beta$ ), and a displacement parameter ( $s_0$ ). We  
343 assume that selection coefficients are measured with a normally distributed measurement error with  
344 standard deviation  $\sigma_{\text{silent}}$ . Thus, the observed distribution of selection coefficients is modeled as the sum of a  
345 gamma and normally distributed random variable. We use the NormalGamma package in R to numerically  
346 compute the normal-gamma density function (Plancade *et al.* 2012). Maximum likelihood estimates of the  
347 parameters of the negative displaced gamma distribution are obtained with L-BFGS-B optimization  
348 implemented in the R function `optim`.

### 349 **Inferring drug resistance from fitness data**

350 We can quantify the drug resistance of each mutant in the library by computing its fold change in relative  
351 fitness,

$$352 \quad W([DCV]) = \frac{RF_{\text{mut}}([DCV])}{RF_{\text{mut}}}$$

353 Here  $RF_{\text{mut}}$  is the relative fitness of a mutant under the natural condition (i.e. no drug).  $W$  is the fold change  
354 in relative fitness and represents the level of drug resistance relative to the wild type.  $W > 1$  indicates drug  
355 resistance, and  $W < 1$  indicates drug sensitivity.

356 This empirical measure of drug resistance can be directly linked to a simple pharmacodynamics model  
357 (Rosenbloom *et al.* 2012), where the viral replicative fitness is modeled as a function of drug dose,

$$358 \quad W_{\text{predict}}([DCV]) = \left( \frac{IC_{\text{mut}}}{[DCV] + IC_{\text{mut}}} \right) \Bigg/ \left( \frac{IC_{\text{wt}}}{[DCV] + IC_{\text{wt}}} \right)$$

359 Here  $IC$  denotes the half-inhibitory concentration. The Hill coefficient describing the sigmoidal shape of the  
360 dose response curve is fixed to 1, as used in fitting the dose response curves of wild-type virus and validated



361 mutant viruses (Supplementary Figure 5). The drug resistance score  $W$  inferred from fitness data is  
362 consistent with the drug resistance score  $W_{predict}$  predicted from dose response curves of validated mutants  
363 (Supplementary Figure 6).

### 364 **Calculation of relative solvent accessibility**

365 DSSP (<http://www.cmbi.ru.nl/dssp.html>) was used to compute the Solvent Accessible Surface Area (SASA)  
366 (Kabsch and Sander 1983) from the HCV NS5A protein structure (PDB: 3FQM) (Love *et al.* 2009). SASA  
367 was then normalized to Relative Solvent Accessibility (RSA) using the empirical scale reported in (Tien *et al.*  
368 2013).

### 369 **Predictions of protein stability**

370  $\Delta\Delta G$  (in Rosetta Energy Unit) of HCV NS5A mutants was predicted by PyRosetta (version:  
371 “monolith.ubuntu.release-104”) as the difference in scores between the monomer structure of mutants  
372 (single amino acid mutations from site 32 to 103) and the reference (PDB: 3FQM). The score is designed to  
373 capture the change in thermodynamic stability caused by the mutation ( $\Delta\Delta G$ ) (Das and Baker 2008). The  
374 reference sequence of NS5A in the PDB file (PDB: 3FQM) is different from the WT sequence in our  
375 experiment by 20 amino acid substitutions. Thus instead of directly comparing  $\Delta\Delta G$  to fitness effects of  
376 individual mutations, we used the median  $\Delta\Delta G$  caused by amino acid substitutions at each site.

377 The PDB file of NS5A dimer was cleaned and trimmed to a monomer (chain A). Next, all side chains were  
378 repacked (sampling from the 2010 Dunbrack rotamer library (Shapovalov and Dunbrack 2011)) and  
379 minimized for the reference structure using the talaris2014 scoring function. After an amino acid mutation  
380 was introduced, the mutated residue was repacked, followed by quasi-Newton minimization of the backbone  
381 and all side chains (algorithm: “lbfgs\_armijo\_nonmonotone”). This procedure was performed 50 times, and  
382 the predicted  $\Delta G$  of a mutant structure is the average of the three lowest scoring structures.

383 We note that predictions based on NS5A monomer structure were only meant to provide a crude profile of  
384 how mutations at each site may impact protein stability. Potential structural constraints at the dimer interface  
385 have been ignored, which is further complicated by the observations of two different NS5A dimer structures  
386 (Tellinghuisen *et al.* 2005; Love *et al.* 2009).

### 387 **Diversity of HCV sequences identified in patients**

388 Aligned nucleotide sequences of HCV NS5A protein were downloaded from Los Alamos National Lab  
389 database (Kuiken *et al.* 2005) (all HCV genotypes, ~2600 sequences total) and clipped to the region of

390 interest (amino acid 18-103 of NS5A). Sequences that caused gaps in the alignment of H77 reference  
391 genome were manually removed. After translation to amino acid sequences, sequences with ambiguous  
392 amino acids were removed (~2300 amino acid sequences after filtering). The sequence diversity at each  
393 amino acid site was quantified by Shannon entropy.

#### 394 **Data and reagent availability**

395 All research materials are available upon request. Raw sequencing data have been submitted to the NIH  
396 Short Read Archive (SRA) under accession number: BioProject PRJNA395730. All scripts have been  
397 deposited to <https://github.com/leidai-evolution/DFE-HCV>.

#### 398 **Ethics Statement**

399 The use of human cell lines and infectious agents in this paper is approved by Institutional Biosafety  
400 Committee at University of California, Los Angeles (IBC #40.10.2-f).

#### 402 **Acknowledgements**

403 We thank Daniel Weinreich and two anonymous reviewers for constructive comments on the manuscript.  
404 L.D. was supported by HHMI Postdoctoral Fellowship from Jane Coffin Childs Memorial Fund for Medical  
405 Research. N.C.W. was supported by Croucher Foundation Fellowship. R.S. was supported by NSFC  
406 81172314, NIH DE023591 and NIH CA177322.

#### 408 **Author contributions**

409 L.D., Y.D., H.Q. and R.S. designed the experiments. L.D., H.Q. and Y.D. performed the experiments. L.D.  
410 and Y.D. analyzed the experimental data. L.D., E.W. and Y.D. performed the bioinformatics analyses. C.D.H.  
411 and L.D. performed the analysis on FGM. L.D. wrote the first draft of the manuscript, with revisions from Y.D.,  
412 H.Q., N.C.W., J.O.L-S., and R.S.. All authors discussed the results and commented on the manuscript.



## References

- 413  
414 Bank C., Hietpas R. T., Wong A., Bolon D. N., Jensen J. D., 2014 A Bayesian MCMC Approach to Assess  
415 the Complete Distribution of Fitness Effects of New Mutations: Uncovering the Potential for Adaptive  
416 Walks in Challenging Environments. *Genetics* 196.
- 417 Bank C., Hietpas R. T., Jensen J. D., Bolon D. N. A., 2015 A systematic survey of an intragenic epistatic  
418 landscape. *Mol. Biol. Evol.* 32: 229–38.
- 419 Barton J. P., Goonetilleke N., Butler T. C., Walker B. D., McMichael A. J., *et al.*, 2016 Relative rate and  
420 location of intra-host HIV evolution to evade cellular immunity are predictable. *Nat. Commun.* 7: 11660.
- 421 Bataillon T., Zhang T., Kassen R., 2011 Cost of adaptation and fitness effects of beneficial mutations in  
422 *Pseudomonas fluorescens*. *Genetics* 189: 939–949.
- 423 Bataillon T., Bailey S., 2014 Effects of new mutations on fitness: insights from models and data. *Ann. N. Y.*  
424 *Acad. Sci.* 1320: 76–92.
- 425 Beisel C. J., Rokyta D. R., Wichman H. A., Joyce P., 2007 Testing the extreme value domain of attraction for  
426 distributions of beneficial fitness effects. *Genetics* 176: 2441–9.
- 427 Bloom J. D., Silberg J. J., Wilke C. O., Drummond D. A., Adami C., *et al.*, 2005 Thermodynamic prediction of  
428 protein neutrality. *Proc. Natl. Acad. Sci. U. S. A.* 102: 606–11.
- 429 Burch C. L., Chao L., 2000 Evolvability of an RNA virus is determined by its mutational neighbourhood.  
430 *Nature* 406: 625–8.
- 431 Burch C., Guyader S., Samarov D., Shen H., 2007 Experimental estimate of the abundance and effects of  
432 nearly neutral mutations in the RNA virus  $\phi 6$ . *Genetics* 476: 467–476.
- 433 Carrasco P., Iglesia F. de la, Elena S., 2007 Distribution of fitness and virulence effects caused by  
434 single-nucleotide substitutions in Tobacco etch virus. *J. Virol.* 81: 12979–12984.
- 435 Chaudhury S., Lyskov S., Gray J. J., 2010 PyRosetta: a script-based interface for implementing molecular  
436 modeling algorithms using Rosetta. *Bioinformatics* 26: 689–91.
- 437 Chevereau G., Dravecká M., Batur T., Guvenek A., Ayhan D. H., *et al.*, 2015 Quantifying the Determinants of  
438 Evolutionary Dynamics Leading to Drug Resistance. *PLoS Biol.* 13: e1002299.
- 439 Chevin L.-M., 2010 On measuring selection in experimental evolution. *Biol. Lett.*
- 440 Cowperthwaite M. C., Bull J. J., Meyers L. A., 2005 Distributions of beneficial fitness effects in RNA.  
441 *Genetics* 170: 1449–57.
- 442 Das R., Baker D., 2008 Macromolecular Modeling with Rosetta. *Annu. Rev. Biochem.* 77: 363–382.
- 443 Desai M. M., 2013 Statistical questions in experimental evolution. *J. Stat. Mech. Theory Exp.* 2013: P01003.


- 444 Domingo E., Sheldon J., Perales C., 2012 Viral quasispecies evolution. *Microbiol. Mol. Biol. Rev.* 76: 159–  
445 216.
- 446 Domingo-Calap P., Cuevas J. M., Sanjuán R., 2009 The Fitness Effects of Random Mutations in  
447 Single-Stranded DNA and RNA Bacteriophages (DJ Begun, Ed.). *PLoS Genet.* 5: e1000742.
- 448 Elena S. F., Carrasco P., Daròs J.-A., Sanjuán R., 2006 Mechanisms of genetic robustness in RNA viruses.  
449 *EMBO Rep.* 7: 168–73.
- 450 Eyre-Walker A., Keightley P. D., 2007 The distribution of fitness effects of new mutations. *Nat. Rev. Genet.* 8:  
451 610–8.
- 452 Firnberg E., Labonte J. W., Gray J. J., Ostermeier M., 2014 A comprehensive, high-resolution map of a  
453 gene's fitness landscape. *Mol. Biol. Evol.* 31: 1581–92.
- 454 Foll M., Poh Y.-P., Renzette N., Ferrer-Admetlla A., Bank C., *et al.*, 2014 Influenza virus drug resistance: a  
455 time-sampled population genetics perspective. *PLoS Genet.* 10: e1004185.
- 456 Fowler D. M., Fields S., 2014 Deep mutational scanning: a new style of protein science. *Nat. Methods* 11:  
457 801–807.
- 458 Gao M., Nettles R. E., Belema M., Snyder L. B., Nguyen V. N., *et al.*, 2010 Chemical genetics strategy  
459 identifies an HCV NS5A inhibitor with a potent clinical effect. *Nature* 465: 96–100.
- 460 Goldberg D. E., Siliciano R. F., Jacobs W. R., 2012 Outwitting evolution: fighting drug-resistant TB, malaria,  
461 and HIV. *Cell* 148: 1271–83.
- 462 He X., Liu L., 2016 Toward a prospective molecular evolution. *Science* 352: 769–70.
- 463 Hietpas R. T., Jensen J. D., Bolon D. N. A., 2011 Experimental illumination of a fitness landscape. *Proc. Natl.*  
464 *Acad. Sci. U. S. A.* 108: 7896–901.
- 465 Hietpas R. T., Bank C., Jensen J. D., Bolon D. N. A., 2013 SHIFTING FITNESS LANDSCAPES IN  
466 RESPONSE TO ALTERED ENVIRONMENTS. *Evolution (N. Y.)*. 67: 3512–3522.
- 467 Imhof M., Schlötterer C., 2001 Fitness effects of advantageous mutations in evolving *Escherichia coli*  
468 populations. *Proc. Natl. Acad. Sci. U. S. A.* 98: 1113–1117.
- 469 Jacquier H., Birgy A., Nagard H. Le, Mechulam Y., Schmitt E., *et al.*, 2013 Capturing the mutational  
470 landscape of the beta-lactamase TEM-1. *Proc. Natl. Acad. Sci. U. S. A.* 110: 13067–72.
- 471 Kabsch W., Sander C., 1983 Dictionary of protein secondary structure: Pattern recognition of  
472 hydrogen-bonded and geometrical features. *Biopolymers* 22: 2577–2637.
- 473 Kassen R., Bataillon T., 2006 Distribution of fitness effects among beneficial mutations before selection in  
474 experimental populations of bacteria. *Nat. Genet.* 38: 484–8.

- 475 Ke R., Loverdo C., Qi H., Sun R., Lloyd-Smith J. O., 2015 Rational Design and Adaptive Management of  
476 Combination Therapies for Hepatitis C Virus Infection. *PLoS Comput. Biol.* 11: e1004040.
- 477 Kuiken C., Yusim K., Boykin L., Richardson R., 2005 The Los Alamos hepatitis C sequence database.  
478 *Bioinformatics* 21: 379–384.
- 479 Lalić J., Cuevas J. M., Elena S. F., 2011 Effect of host species on the distribution of mutational fitness effects  
480 for an RNA virus. *PLoS Genet.* 7: e1002378.
- 481 Levy S. F., Blundell J. R., Venkataram S., Petrov D. A., Fisher D. S., *et al.*, 2015 Quantitative evolutionary  
482 dynamics using high-resolution lineage tracking. *Nature* 519: 181–6.
- 483 Li C., Qian W., Maclean C. J., Zhang J., 2016 The fitness landscape of a tRNA gene. *Science* 352: 837–840.
- 484 Liberles D. A., Teichmann S. A., Bahar I., Bastolla U., Bloom J., *et al.*, 2012 The interface of protein structure,  
485 protein biophysics, and molecular evolution. *Protein Sci.* 21: 769–785.
- 486 Lindenbach B. D., Evans M. J., Syder A. J., Wölk B., Tellinghuisen T. L., *et al.*, 2005 Complete Replication of  
487 Hepatitis C Virus in Cell Culture. *Science* 309: 623–626.
- 488 Love R. A., Brodsky O., Hickey M. J., Wells P. A., Cronin C. N., 2009 Crystal structure of a novel dimeric  
489 form of NS5A domain I protein from hepatitis C virus. *J. Virol.* 83: 4395–403.
- 490 MacLean R. C., Buckling A., 2009 The distribution of fitness effects of beneficial mutations in *Pseudomonas*  
491 *aeruginosa*. *PLoS Genet.* 5: e1000406.
- 492 Martin G., Lenormand T., 2006a The fitness effect of mutations across environments: a survey in light of  
493 fitness landscape models. *Evolution* 60: 2413–2427.
- 494 Martin G., Lenormand T., 2006b A general multivariate extension of Fisher’s geometrical model and the  
495 distribution of mutation fitness effects across species. *Evolution* 60: 893–907.
- 496 Matuszewski S., Hildebrandt M. E., Ghenu A.-H., Jensen J. D., Bank C., 2016 A Statistical Guide to the  
497 Design of Deep Mutational Scanning Experiments. *Genetics*.
- 498 Metcalf C. J. E., Birger R. B., Funk S., Kouyos R. D., Lloyd-Smith J. O., *et al.*, 2015 Five challenges in  
499 evolution and infectious diseases. *Epidemics* 10: 40–44.
- 500 Ogbunugafor C. B., Wylie C. S., Diakite I., Weinreich D. M., Hartl D. L., 2016 Adaptive Landscape by  
501 Environment Interactions Dictate Evolutionary Dynamics in Models of Drug Resistance. *PLoS Comput.*  
502 *Biol.* 12: e1004710.
- 503 Olson C. A., Wu N. C., Sun R., 2014 A comprehensive biophysical description of pairwise epistasis  
504 throughout an entire protein domain. *Curr. Biol.* 24: 2643–51.
- 505 Orr H. A., 1998 The population genetics of adaptation: the distribution of factors fixed during adaptive

- 506 evolution. *Evolution* (N. Y). 52: 935–949.
- 507 Orr H., 2003 The distribution of fitness effects among beneficial mutations. *Genetics* 1526: 1519–1526.
- 508 Orr H., 2006 The distribution of fitness effects among beneficial mutations in Fisher’s geometric model of  
509 adaptation. *J. Theor. Biol.* 238: 279–285.
- 510 Peris J. B., Davis P., Cuevas J. M., Nebot M. R., Sanjuán R., 2010 Distribution of fitness effects caused by  
511 single-nucleotide substitutions in bacteriophage f1. *Genetics* 185: 603–9.
- 512 Plancade S., Rozenholc Y., Lund E., 2012 Generalization of the normal-exponential model: exploration of a  
513 more accurate parametrisation for the signal distribution on Illumina BeadArrays. *BMC Bioinformatics*  
514 13: 329.
- 515 Puchta O., Cseke B., Czaja H., Tollervey D., Sanguinetti G., *et al.*, 2015 Network of epistatic interactions  
516 within a yeast snoRNA. *Science* 352: 840–844.
- 517 Qi H., Olson C. A., Wu N. C., Ke R., Loverdo C., *et al.*, 2014 A quantitative high-resolution genetic profile  
518 rapidly identifies sequence determinants of hepatitis C viral fitness and drug sensitivity. *PLoS Pathog.*  
519 10: e1004064.
- 520 Ramsey D. C., Scherrer M. P., Zhou T., Wilke C. O., 2011 The Relationship Between Relative Solvent  
521 Accessibility and Evolutionary Rate in Protein Evolution. *Genetics* 188: 479–488.
- 522 Renzette N., Pfeifer S. P., Matuszewski S., Kowalik T. F., Jensen J. D., 2017 On the Analysis of Intra-host and  
523 Inter-host Viral Populations: Human Cytomegalovirus as a Case Study of Pitfalls and Expectations. *J.*  
524 *Virology* 91: e01976-16.
- 525 Rihn S. J., Wilson S. J., Loman N. J., Alim M., Bakker S. E., *et al.*, 2013 Extreme genetic fragility of the HIV-1  
526 capsid. *PLoS Pathog.* 9: e1003461.
- 527 Robinson M., Tian Y., Delaney W. E., Greenstein A. E., 2011 Preexisting drug-resistance mutations reveal  
528 unique barriers to resistance for distinct antivirals. *Proc. Natl. Acad. Sci. U. S. A.* 108: 10290–5.
- 529 Rokyta D., Joyce P., Caudle S., Wichman H., 2005 An empirical test of the mutational landscape model of  
530 adaptation using a single-stranded DNA virus. *Nat. Genet.* 37: 441–444.
- 531 Rosenbloom D. I. S., Hill A. L., Rabi S. A., Siliciano R. F., Nowak M. A., 2012 Antiretroviral dynamics  
532 determines HIV evolution and predicts therapy outcome. *Nat. Med.* 18: 1378–85.
- 533 Sanjuan R., Moya A., Elena S. F., 2004 The distribution of fitness effects caused by single-nucleotide  
534 substitutions in an RNA virus. *Proc. Natl. Acad. Sci.* 101: 8396–8401.
- 535 Sanjuán R., 2010 Mutational fitness effects in RNA and single-stranded DNA viruses: common patterns  
536 revealed by site-directed mutagenesis studies. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 365: 1975–82.

- 537 Shapovalov M. V., Dunbrack R. L., 2011 A smoothed backbone-dependent rotamer library for proteins  
538 derived from adaptive kernel density estimates and regressions. *Structure* 19: 844–58.
- 539 Silander O. K., Tenailon O., Chao L., 2007 Understanding the Evolutionary Fate of Finite Populations: The  
540 Dynamics of Mutational Effects. *PLoS Biol.* 5: e94.
- 541 Soskine M., Tawfik D. S., 2010 Mutational effects and the evolution of new protein functions. *Nat. Rev. Genet.*  
542 11: 572–82.
- 543 Stiffler M. A., Hekstra D. R., Ranganathan R., 2015 Evolvability as a Function of Purifying Selection in  
544 TEM-1  $\beta$ -Lactamase. *Cell* 160: 882–892.
- 545 Tellinghuisen T. L., Marcotrigiano J., Rice C. M., 2005 Structure of the zinc-binding domain of an essential  
546 component of the hepatitis C virus replicase. *Nature* 435: 374–9.
- 547 Thyagarajan B., Bloom J. D., 2014 The inherent mutational tolerance and antigenic evolvability of influenza  
548 hemagglutinin. *Elife* 3: e03300.
- 549 Tien M. Z., Meyer A. G., Sydykova D. K., Spielman S. J., Wilke C. O., 2013 Maximum allowed solvent  
550 accessibilities of residues in proteins. *PLoS One* 8: e80635.
- 551 Turner P. E., Elena S. F., 2000 Cost of Host Radiation in an RNA Virus. *Genetics* 156: 1465–1470.
- 552 Vale P. F., Choisy M., Froissart R., Sanjuán R., Gandon S., 2012 THE DISTRIBUTION OF MUTATIONAL  
553 FITNESS EFFECTS OF PHAGE  $\phi$ X174 ON DIFFERENT HOSTS. *Evolution (N. Y.)*. 66: 3495–3507.
- 554 Visher E., Whitefield S. E., McCrone J. T., Fitzsimmons W., Luring A. S., 2016 The Mutational Robustness  
555 of Influenza A Virus. *PLOS Pathog.* 12: e1005856.
- 556 Visser J. A. G. M. de, Krug J., 2014 Empirical fitness landscapes and the predictability of evolution. *Nat. Rev.*  
557 *Genet.* 15: 480–90.
- 558 Wakita T., Pietschmann T., Kato T., Date T., Miyamoto M., *et al.*, 2005 Production of infectious hepatitis C  
559 virus in tissue culture from a cloned viral genome. *Nat. Med.* 11: 791–796.
- 560 Wright S., 1932 The Roles of Mutation, Inbreeding, Crossbreeding and Selection in Evolution. 1: 356–366.
- 561 Wu N. C., Young A. P., Dandekar S., Wijersuriya H., Al-Mawsawi L. Q., *et al.*, 2013 Systematic identification  
562 of H274Y compensatory mutations in influenza A virus neuraminidase by high-throughput screening. *J.*  
563 *Viro.* 87: 1193–9.
- 564 Wu N. C., Olson C. A., Du Y., Le S., Tran K., *et al.*, 2015 Functional Constraint Profiling of a Viral Protein  
565 Reveals Discordance of Evolutionary Conservation and Functionality. *PLOS Genet.* 11: e1005310.
- 566 Wu N. C., Dai L., Olson C. A., Lloyd-Smith J. O., Sun R., 2016 Adaptation in protein fitness landscapes is  
567 facilitated by indirect paths. *Elife* 5: e16965.

568 Wylie C. S., Shakhnovich E. I., 2011 A biophysical protein folding model accounts for most mutational fitness  
569 effects in viruses. Proc. Natl. Acad. Sci. 108: 9916–9921.  
570




571

572

573

574


**Figure 1. Distribution of fitness effects (DFE) of single amino acid substitutions in domain IA of HCV NS5A protein without drug selection.** DFE of single amino acid substitutions (A) and synonymous substitutions (B). Lethal mutations are not shown in the histogram.



575

576 **Figure 2. The spectrum of beneficial mutations changes under increasing environmental stress**  
577 **imposed by the antiviral drug Daclatasvir.** (A) DFE of single amino acid substitutions in domain IA of HCV  
578 NS5A protein under increasing environmental stress by Daclatasvir. The black line indicates the threshold  
579 used for classifying beneficial mutations (Methods). (B) The cumulative distribution function of the fitness  
580 effect of beneficial mutations. (C) The number of beneficial mutations as a function of environmental stress  
581 imposed by Daclatasvir.





582

583

**Figure 3. The adaptive potential under drug selection is determined by the effects of mutations on replication fitness and drug resistance.** (A) Hypothetical dose response curves of the wild-type virus and

584

a drug-resistant mutant virus. The absolute fitness  $f$  decreases with drug concentration  $[drug]$

585

586  $f = f_0 \frac{IC_{50}}{IC_{50} + [drug]}$ , where  $f_0$  is the fitness without drug selection and  $IC_{50}$  is the half inhibitory

587

concentration. Compared to the wild-type virus, the hypothetical drug-resistant mutant carries a fitness cost (smaller  $f_0$ ) but is less sensitive to drug inhibition (larger  $IC_{50}$ ). Relative fitness of the drug-resistant mutant

588


is expected to increase with drug concentration. (B) Relative fitness of validated drug-resistant and drug-sensitive mutants (Supplementary Figure 5) as a function of [DCV]. (C) The effects of mutations on

589

replication fitness (i.e. fitness without drug) and drug resistance score  $W$  at [DCV]=40 pM (Methods).

590

591



592

593

594

595

596

597

598

599

600

**Figure 4. Mutations with deleterious fitness effects reveal constraints of protein evolution.** (A) The pattern of sequence conservation observed in patient sequences is highly correlated to the replication fitness measured in cell culture. (B) Mutations at amino acid sites with lower solvent accessibility tend to incur larger fitness costs. (C) Mutations at amino acid sites with larger effects on destabilizing protein stability ( $\Delta\Delta G > 0$ ) tend to reduce the viral replication fitness. Changes in folding free energy  $\Delta\Delta G$  (Rosetta Energy Unit) of NS5A monomer were predicted by PyRosetta. The median  $\Delta\Delta G$  at each amino acid site is shown. In (A-C), the median fitness of observed mutants at each amino acid site is shown. In (B) and (C), red lines represent the fits by linear regression and are only used to guide the eye.