

Quantization via mirror symmetry^{*}

Sergei Gukov

Received: 5 November 2010 / Revised: 1 November 2011 / Accepted: 8 November 2011

Published online: 25 December 2011

© The Mathematical Society of Japan and Springer 2011

Communicated by: Toshiyuki Kobayashi

Abstract. When combined with mirror symmetry, the A -model approach to quantization leads to a fairly simple and tractable problem. The most interesting part of the problem then becomes finding the mirror of the coisotropic brane. We illustrate how it can be addressed in a number of interesting examples related to representation theory and gauge theory, in which mirror geometry is naturally associated with the Langlands dual group. Hyperholomorphic sheaves and (B, B, B) branes play an important role in the B -model approach to quantization.

Keywords and phrases: mirror symmetry, derived category, branes, quantization, symplectic geometry

Mathematics Subject Classification (2010): 18F20, 51P05, 81S10

^{*} This article is based on the 8th Takagi Lectures that the author delivered at Research Institute for Mathematical Sciences, Kyoto University on November 23, 2010.

S. GUKOV

California Institute of Technology, Pasadena, CA 91125, USA and
Max-Planck-Institut für Mathematik, Vivatsgasse 7, D-53111 Bonn, Germany
(e-mail: gukov@theory.caltech.edu)