

# Quantum Belief Propagation

M. B. Hastings, T-13 LANL

# Classical Belief Propagation: the transfer matrix

- Solve system of  $N$  sites, compute probability distribution of last site
- Add one more site and repeat



Discrete set of states on each site:  $\sigma_i$

Nearest neighbor Hamiltonian:  $H = \sum_i h_{i,i+1}$

Hamiltonian on first  $N$  sites:  $h^{(N)} \equiv \sum_{i=1}^{N-1} h_{i,i+1}$

**Partition function for chain of N sites:**

$$Z^{(N)} = \sum_{\{\sigma_1, \dots, \sigma_N\}} \exp(-\beta h^{(N)})$$

**Probability distribution:**

$$P^{(N)}(\sigma_1, \dots, \sigma_N) = \frac{1}{Z^{(N)}} \exp(-\beta h^{(N)})$$

**Probability of last site:**


$$P^{(N)}(\sigma_N) = \sum_{\{\sigma_1, \dots, \sigma_{N-1}\}} P^{(N)}(\sigma_1, \dots, \sigma_N)$$

**Recursion relation:**

$$P^{(N+1)}(\sigma_{N+1}) \propto \sum_{\sigma_N} P^{(N)}(\sigma_N) \exp(-\beta h_{N,N+1})$$

# Quantum belief propagation:

- Analogue of probability on last site is reduced density matrix
- Need window of several sites. Window size is  $l_0 - 1$




$$\rho^{(N)} = \exp(-\beta h^{(N)})$$

$$\rho_{red}^{(N)} = \text{Tr}_{1, \dots, N+1-l_0}(\rho^{(N)})$$

# Quantum belief propagation:

Construct  $O$  such that:  $\rho^{(N+1)} \approx O_{N+1} \rho^{(N)} O_{N+1}^\dagger$



$$\rho^{(N+1)} = O_{N+1} O_N O_{N-1} \dots O_{N-1}^\dagger O_N^\dagger O_{N+1}^\dagger$$

$$\rho_{red}^{(N+1)} = \text{Tr}_{N+2-l_0} \left( O_{N+1} (\rho_{red}^{(N)} \otimes 1_{N+1}) O_{N+1}^\dagger \right)$$

# Algorithm:

- Initialize reduced density matrix on first  $l_0 - 1$  sites
- Iterate completely positive map until convergence
- Compute partition function from normalization
- Observables: insert operator before tracing on first site

Completely positive map:  $\rho_{red} \rightarrow \text{Tr}_1 \left( O_{N+1} (\rho_{red}^{(N)} \otimes 1_{N+1}) O_{N+1}^\dagger \right)$

Observables:  $\rho_{red} \rightarrow \text{Tr}_1 \left( S_1^z O_{N+1} (\rho_{red}^{(N)} \otimes 1_{N+1}) O_{N+1}^\dagger \right)$

# Performance:

- Computational effort : diagonalize operators of dimension  $2^{l_0}$  to compute  $\mathcal{O}$ .
- Window size needed scales as  $l_0 \sim v_{LR}\beta$
- No Trotter error, very accurate at high temperature
- Handle disorder by precomputing operators.


Heisenberg chain:  $\chi_{exact}(T_{max}) = 0.146926279\dots$

$\chi_{QBP}(T_{max}) = 0.146927031\dots$

for 10-by-10 matrices diagonalized (<.1 second CPU time)

# Spin-1/2 Heisenberg Chain

$$H = \sum \vec{S}_i \cdot \vec{S}_{i+1}$$



thanks A. C.  
Klumper for data

FIG. 1: Specific heat against temperature for  $l_0 = 3$  (dashed line), 5 (dotted line), 7 (solid line). Curves that go negative are from  $-3\beta^2\partial_\beta\langle S_i^z S_{i+1}^z \rangle$ , while those that diverge positively are from  $\beta^2\partial_\beta^2 \log(Z)$ . Inset:  $l_0 = 9$  and Bethe ansatz.



# Spin-1/2 disordered chain:

$$H = \sum_i J_i \vec{S}_i \cdot \vec{S}_{i+1} + \sum_i K_i \vec{S}_i \cdot \vec{S}_{i+2}$$

1




FIG. 1: Top: specific heat for the pure system. Middle: uniform susceptibility for the pure system ( $l_0 = 5, 7, 9$  are black, red, green respectively) and the disordered system ( $l_0 = 5, 7, 9$  are blue, yellow, brown respectively). Bottom: dimer susceptibility for the pure system ( $l_0 = 5, 7, 9$  are black, red, green respectively) and the disordered system ( $l_0 = 5, 7, 9$  are blue, yellow, brown respectively).

# QBP Equations:

Exact result for small change in H:

$$\partial_s \exp[-\beta(H + sA)] = \eta_s \exp(-\beta H_s) + \exp(-\beta H_s) \eta_s^\dagger$$

$$\eta_s = - \int d\omega \left( \frac{\beta}{2} + \beta F(\omega) \right) A^{\omega, s}$$

classical term

quantum term

Perturbation at  
frequency omega

Integrate to add site:

$$O = \exp\left(\int_0^1 ds \eta_s\right)$$

# Conclusions

- Accurate at high temperature
- Works well on loopless models
- Can we extend to loopy models?
- How do the different QBPs relate? (Poulin, Leifer)
- Works well for disorder by precomputing  $\mathcal{O}$
- A new kind of transfer matrix