

QUANTUM MULTILINEAR ALGEBRA

MITSUYASU HASHIMOTO AND TAKAHIRO HAYASHI

(Received August 12, 1991, revised May 7, 1992)

Abstract. We construct a quantized version of the theory of multilinear algebra, based on Jimbo's solution of Yang-Baxter equation of type $A_{N-1}^{(1)}$. Using this, we discuss the polynomial representations of quantum general linear groups.

CONTENTS

Introduction	471
1. Yang-Baxter operators	473
2. Algebras associated with Yang-Baxter pairs	476
3. Quantum matrices	477
4. Symmetric algebras, exterior algebras and the fusion procedure	480
5. Divided power algebra	486
6. Schur modules and Weyl modules	493
7. Universality of Weyl modules	505
8. Polynomial representation of GL_q over a field	510
9. Quantum determinants and straightening formulas	515
References	519

Introduction. Quantum groups are mathematical objects which arose from the study of the quantum inverse scattering method, especially the Yang-Baxter equation. They are very remarkable Hopf algebras and can be considered as q -analogues of Kac-Moody enveloping algebras or of coordinate rings of Lie groups. Not only have they added new aspects to representation theory, but also they have brought to *non-commutative geometry* a remarkable progress, i.e. the discovery of many new examples such as quantum linear algebraic groups, quantum spheres and so on.

In this article, we study quantum analogues of some linear-algebraic objects such as matrices, symmetric and alternating tensors, and determinants. We construct these from Jimbo's solution of Yang-Baxter (YB) equation of type $A_{N-1}^{(1)}$ and investigate their structure via the notion which we call *Yang-Baxter bialgebras*. As applications, we give realizations and free bases of Weyl modules $K_\lambda V$ and their dual modules (Schur modules) of quantum general linear groups $GL_q(N)$, and give a criterion for the irreducibility of $K_\lambda V$. We also give an analogue of the straightening formula for quantum matrix bialgebras. We would like to emphasize that these objects are defined over any commutative ring R and any unit element $q \in R^\times$ and are compatible with extensions

of the base ring R . Hence, we can get the representation theory of quantum general group ‘defined over $\mathbb{Z}[Q, Q^{-1}]$ ’, where Q denotes an indeterminate.

In Section 1, we introduce operations on YB operators (i.e. solutions of the YB equation) called the product \times , dual \vee , and fusion procedure. In Section 2, we associate with a YB operator two algebras which we call the symmetric and the exterior algebras. In Section 3, we apply these tools to the construction of bialgebras SE^\vee which are called quantum matrix bialgebras (cf. [12]).

In Section 4, we construct two YB operators φ and ψ on the symmetric and exterior algebras of Jimbo’s YB operators of type $A_{N-1}^{(1)}$ using the fusion procedure. With the YB operator φ , we introduce unusual algebra structures into tensor products of these algebras, and prove that these algebras have some structures which we call *YB-bialgebras*. Though a YB-bialgebra has structures of an algebra and a coalgebra, it is not necessarily a bialgebra in the usual sense. The ‘commutativity’ and the ‘cocommutativity’ of these YB-bialgebras are described in terms of the YB-operator ψ .

In Section 5, we discuss an analogue of the theory of graded multilinear bialgebras. Recall that divided power algebras have been important in the study of characteristic-free representation theory (cf. [2], [3], [4]). Here we introduce *divided power algebras* of the YB pairs treated in Section 4. They are defined to be graded duals of symmetric algebras defined in Section 4. This concept enables us to define Weyl modules and Schur algebras in a natural way without assuming that q is not a root of unity.

From Sections 6 to 9, our interest is concentrated on the study of the representation theory of quantum deformations of general linear groups.

In Section 6, we define (deformed) Weyl modules $K_\lambda V$ and Schur modules $L_\lambda V$ associated to a partition λ , using the YB-bialgebra structure of the symmetric and exterior algebras defined in Section 4. We prove that $L_\lambda V$ and $K_\lambda V$ are finite free R -modules and that they are ‘defined over $\mathbb{Z}[Q, Q^{-1}]$ ’ in the sense that they are compatible with base extensions. This property is an analogue of the so-called *universal freeness*. For the original version of this result, we refer the reader to [4]. Though our definition of $L_\lambda V$ is different from $L_q^\lambda(B)$ in Taft-Towber [37], they will turn out to be equivalent (cf. Proposition 9.7). Though our construction and argument in this section are nothing but the deformed versions of those in [4, Chapter II], we include some details, since they do not seem to be so obvious.

In Section 7, we introduce the deformed versions of the Schur algebra, weights, and the contravariant dual functor in our language. We will show that (deformed) Weyl modules are universal highest weight modules in the sense of Theorem 7.12.

In Section 8, we work over a base field K , and discuss the irreducibility and complete reducibility of SE^\vee -comodules. The notion of formal characters is introduced in this section. Theorem 8.9 is a q -analogue of the hook length criterion for the irreducibility of Weyl modules. In the proof, Gyoja’s q -analogues of Young symmetrizers [14] play important roles.

In Section 9, we prove a q -analogue of the straightening formula using quantum de-

terminants. Namely, we prove that the degree k component $S_k E^\vee$ of SE^\vee admits a filtration of SE^\vee -subbicomodules whose associated graded object is $\bigoplus_{|\lambda|=k} (K_\lambda V)^* \otimes L_{\tilde{\lambda}} V$, where $\tilde{\lambda}$ is the transpose of λ . This formula was originated by Doubilet-Rota-Stein [11] in the case $q=1$. Our approach is a q -analogue of the treatment in [4].

Lastly, we remark on the relation between our construction and the quantum enveloping algebra $U_q\mathfrak{gl}(N)$, or rather the “quantum hyperalgebra” such as that defined in Lusztig [23]. Since these Hopf algebras satisfy the conditions in Theorem 3.3, there are pairings of bialgebras between the quantum matrix bialgebra over Jimbo’s YB pair of type $A_{N-1}^{(1)}$ and the quantum hyperalgebra. Hence, thanks to the general theory of bialgebras, the quantum hyperalgebra acts on our Schur and Weyl modules. Moreover, the criterion for the irreducibility (Theorem 8.9) is also valid when they are viewed as representations over the quantum hyperalgebra.

After the submission of the first version of this work, the authors received preprints by Dipper-James [10], Noumi-Yamada-Mimachi [29], Parshall-Wang [30], which have some overlap with our paper.

Our special thanks are due to Professors Mitsuo Hoshino, Yukihiro Kanie, Hideyuki Matsumura and Akihiro Tsuchiya for valuable advice.

1. Yang-Baxter operators. Let V be a free R -module. A *Yang-Baxter* (or YB) operator on V is an automorphism $\beta_V \in \text{End}_R(V \otimes V)$ such that

$$(1.1) \quad \begin{aligned} (\beta_V)_1 \circ (\beta_V)_2 \circ (\beta_V)_1 &= (\beta_V)_2 \circ (\beta_V)_1 \circ (\beta_V)_2, \\ (\beta_V)_1 &:= \beta_V \otimes \text{id}_V, \quad (\beta_V)_2 := \text{id}_V \otimes \beta_V. \end{aligned}$$

A *Yang-Baxter pair* $V = (V, \beta_V)$ is a free R -module V equipped with a Yang-Baxter operator β_V on V .

Here we give some examples of Yang-Baxter pairs.

EXAMPLE (1) (trivial twisting). Let V be a free R -module. Then the map $\tau_V := \tau_{V,V}$ defined by $\tau_V(u \otimes u') = u' \otimes u$ is a YB operator on V . We call τ_V the *trivial twisting* on V .

EXAMPLE (2) (Jimbo’s operators of type $A_{N-1}^{(1)}$). Let V be a free R -module with a basis $\{u_1, u_2, \dots, u_N\}$ and let q be an invertible element of R . Then Jimbo’s YB operators of type $A_{N-1}^{(1)}$ is the map defined by the following formula:

$$(1.2) \quad \beta_V(u_i \otimes u_j) = \begin{cases} u_i \otimes u_i & (i=j), \\ qu_j \otimes u_i & (i < j), \\ (1-q^2)u_i \otimes u_j + qu_j \otimes u_i & (i > j). \end{cases}$$

We call $\{u_i\}$ the *standard basis* of (V, β_V) . This operator also satisfies *Iwahori’s quadratic equation*

$$(1.3) \quad (\text{id}_{V \otimes V} - \beta_V) \circ (\text{id}_{V \otimes V} + q^{-2} \beta_V) = 0.$$

As for Jimbo's other operators, we refer the reader to [16] and [31].

EXAMPLE (3) (A 'super symmetric extension' of Jimbo's type $A_{N-1}^{(1)}$ operator). Let V be a free module with a basis $\{u_1^0, u_2^0, \dots, u_M^0, u_1^1, \dots, u_N^1\}$ and q an invertible element of R such that $q^2 + 1$ is also invertible. Then the following formulas define a YB operator β_V on V :

$$(1.4) \quad \begin{aligned} \beta_V(u_i^0 \otimes u_i^0) &= u_i^0 \otimes u_i^0, & \beta_V(u_i^1 \otimes u_i^1) &= -q^2 u_i^1 \otimes u_i^1, \\ \beta_V(u_i^0 \otimes u_i^1) &= qu_i^1 \otimes u_i^0, & \beta_V(u_i^1 \otimes u_i^0) &= (1-q^2)u_i^1 \otimes u_i^0 + qu_i^0 \otimes u_i^1, \\ \beta_V(u_i^a \otimes u_j^b) &= \begin{cases} (-1)^{ab}qu_j^b \otimes u_i^a & (i < j) \\ (1-q^2)u_i^a \otimes u_j^b + (-1)^{ab}qu_j^b \otimes u_i^a & (i > j) \end{cases} \end{aligned}$$

The restriction of this operator to $(\bigoplus_i Ru_i^0)^{\otimes 2}$ coincides with the operator in Example (2) under the identification $u_i^0 = u_i$. This operator also satisfies the relation (1.3).

EXAMPLE (4) (YB operators associated with distributive lattices). Let (L, \wedge, \vee) be a distributive lattice, i.e. a set together with two maps $\wedge, \vee: L \times L \rightarrow L$ satisfying the following laws:

$$(1.5) \quad \begin{aligned} a \wedge b &= b \wedge a, & a \vee b &= b \vee a, & a \wedge (b \wedge c) &= (a \wedge b) \wedge c, & a \vee (b \vee c) &= (a \vee b) \vee c, \\ a \vee (a \wedge b) &= (a \vee b) \wedge a = a, & a \wedge (b \vee c) &= (a \wedge b) \vee (a \wedge c) & (a, b, c \in L). \end{aligned}$$

Let V be the free R -module on L and define $\beta_V \in \text{End}_R(V \otimes V)$ by $\beta_V(a \otimes b) = a \wedge b \otimes a \vee b$ ($a, b \in L$). Then β_V satisfies the condition for a YB operator except the existence of β_V^{-1} .

Let \mathfrak{B}_k be the E. Artin's braid group. Namely, \mathfrak{B}_k is a group generated by elements $\{b_i \mid 1 \leq i < k\}$ with the following fundamental relations:

$$(1.6) \quad b_i b_j = b_j b_i \quad (\mid i-j \mid \geq 2), \quad b_i b_{i+1} b_i = b_{i+1} b_i b_{i+1} \quad (1 \leq i \leq k-2).$$

For each YB pair (V, β_V) , there is a representation of \mathfrak{B}_k on the k -fold tensor product $T_k V = V^{\otimes k}$ defined by

$$(1.7) \quad b_i \mapsto (\beta_V)_i := (\text{id}_V)^{\otimes i-1} \otimes \beta_V \otimes (\text{id}_V)^{\otimes k-i-1}.$$

There is an important observation which is due to Iwahori in the Coxeter group case. Let \mathfrak{S}_k be the symmetric group of degree k . The length $\ell(\sigma)$ of $\sigma \in \mathfrak{S}_k$ is the number of the pairs (i, j) such that $1 \leq i < j \leq k$ and $\sigma i > \sigma j$. It is well-known that \mathfrak{S}_k is generated by the transpositions $\sigma_i = (i, i+1)$ ($1 \leq i < k$) and that $\ell(\sigma)$ coincides with the minimal number of factors needed to express σ as a product of the σ_i 's. An expression $\sigma = s_1 s_2 \cdots s_i$ ($s_1, \dots, s_i \in \mathfrak{S}_k$) of $\sigma \in \mathfrak{S}_k$ is said to be reduced if $\ell(\sigma) = \ell(s_1) + \cdots + \ell(s_i)$. Then Iwahori's theorem says that there is a well-defined map $\mathfrak{S}_k \rightarrow \mathfrak{B}_k$ which sends $\sigma \in \mathfrak{S}_k$ in reduced expression $\sigma = \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_{\ell(\sigma)}}$ ($1 \leq i_1, \dots, i_{\ell(\sigma)} \leq k$) to the element $b_{i_1} b_{i_2} \cdots b_{i_{\ell(\sigma)}}$ of the braid group. Combining this map with the homomorphism (1.7), we get a map from \mathfrak{S}_k into $\text{End}_R(T_k V)$. We denote the image of $\sigma \in \mathfrak{S}_k$ by $\beta_V(\sigma)$. Note that if $\sigma = s_1 \cdots s_i$ ($s_1, \dots,$

$s_i \in \mathfrak{S}_k$) is reduced, then $\beta_V(\sigma) = \beta_V(s_1) \cdots \beta_V(s_i)$. Let χ_{ij} ($i, j \geq 1$) be an element of \mathfrak{S}_{i+j} defined by

$$(1.8) \quad \chi_{ij} = \begin{pmatrix} 1 & 2 & 3 & \cdots & i & i+1 & i+2 & \cdots & i+j \\ j+1 & j+2 & j+3 & \cdots & j+i & 1 & 2 & \cdots & j \end{pmatrix}.$$

Then, $\beta_V(\chi_{ij})$ defines a linear isomorphism from $T_i V \otimes T_j V$ onto $T_j V \otimes T_i V$. By Iwahori's theorem we get:

PROPOSITION 1.1 (fusion procedure). *The R -endomorphism $T\beta_V := \bigoplus_{i,j \geq 0} \beta_V(\chi_{ij})$ (resp. $T_i \beta_V := \beta_V(\chi_{ii})$) defines a YB operator on the tensor algebra $TV = \bigoplus_{k \geq 0} T_k V$ (resp. $T_i V$).*

We denote the YB pair $(TV, T\beta_V)$ (resp. $(T_i V, T_i \beta_V)$) by $T(V, \beta_V)$ (resp. $T_i(V, \beta_V)$).

Now we will define a category \mathcal{YB}_R as follows:

(1) The objects of \mathcal{YB}_R consist of all YB pairs on free R -modules of finite rank.

(2) For Yang-Baxter pairs $V = (V, \beta_V)$ and $W = (W, \beta_W)$, the set of morphisms $\mathcal{YB}_R(V, W)$ consists of all R -module maps $f: V \rightarrow W$ satisfying the following commutative diagram:

$$(1.9) \quad \begin{array}{ccc} V \otimes V & \xrightarrow{f \otimes f} & W \otimes W \\ \downarrow \beta_V & & \downarrow \beta_W \\ V \otimes V & \xrightarrow{f \otimes f} & W \otimes W. \end{array}$$

It is easy to see that T_i is a functor from \mathcal{YB}_R to itself for $i \geq 0$.

Now we will give more operations on \mathcal{YB}_R . Let $V = (V, \beta_V)$ and $W = (W, \beta_W)$ be Yang-Baxter pairs.

DEFINITION 1.2. The product $\beta_V \times \beta_W$ of β_V and β_W is a YB operator on $V \otimes W$ defined by the following commutative diagram:

$$(1.10) \quad \begin{array}{ccc} V^{\otimes 2} \otimes W^{\otimes 2} & \xrightarrow{1 \otimes \tau \otimes 1} & V \otimes W \otimes V \otimes W \\ \downarrow \beta_V \otimes \beta_W & & \downarrow \beta_V \times \beta_W \\ V^{\otimes 2} \otimes W^{\otimes 2} & \xrightarrow{1 \otimes \tau \otimes 1} & V \otimes W \otimes V \otimes W. \end{array}$$

We denote the corresponding YB-pair by $V \times W$.

PROPOSITION 1.3. *The product defines a bifunctor $\mathcal{YB}_R \times \mathcal{YB}_R \rightarrow \mathcal{YB}_R$ satisfying the following conditions:*

(1) *associativity constraint*

$$(U \times V) \times W = U \times (V \times W)$$

(2) *existence of the identity object*

$$(R, \text{id}_{R \otimes R}) \times V = V \times (R, \text{id}_{R \otimes R}) = V$$

(3) *commutativity constraint*

$$\tau_{V,W} : V \times W \simeq W \times V.$$

The proof is straightforward.

DEFINITION 1.4. Let (V, β_V) be an object of \mathcal{YB}_R . We define a YB pair $'V = (V^*, \beta_V)$ (resp. V^{-1} , resp. $V^\vee = (V^*, \beta_V^\vee)$) by

$$(1.11) \quad \text{ev}'(\beta_V(v \otimes v') \otimes (u \otimes u')) = \text{ev}((v \otimes v') \otimes \beta_V(u \otimes u')) \quad (v, v' \in V^*, w, w' \in V).$$

(resp. $V^{-1} = (V, \beta_V^{-1})$, resp. $\beta_V^\vee = (\beta_V^{-1})$) and call it the *transpose* (resp. *inverse*, resp. *dual*) of V .

It is important that the evaluation map $\text{ev} : V^* \otimes V \rightarrow R$ gives a morphism of \mathcal{YB}_R from $V^\vee \times V$ onto the unit YB pair $(R, \text{id}_{R \otimes R})$. The following theorem shows that \mathcal{YB}_R is a rigid tensor category (see [9]):

THEOREM 1.5. *We have the following isomorphisms of functors.*

- (1) $(V^\vee)^\vee = V$,
- (2) $(V \times W)^\vee = V^\vee \times W^\vee$,
- (3) $\mathcal{YB}_R(U \times V, W) \simeq \mathcal{YB}_R(V, U^\vee \times W) \quad (U, V, W \in \mathcal{YB}_R)$.

PROOF. Parts (1) and (2) are obvious. To see Part (3), we note that under the isomorphism $\text{Hom}_R(U \otimes V, W) \simeq \text{Hom}_R(V, U^* \otimes W)$, a map $f : U \otimes V \rightarrow W$ corresponds to the composite

$$(1.12) \quad f' : V \xrightarrow{\text{ev}^* \otimes \text{id}_V} U^* \otimes U \otimes V \xrightarrow{\text{id}_{U^*} \otimes f} U^* \otimes W$$

and the map $f' : V \rightarrow U^* \otimes W$ corresponds to the composite

$$(1.13) \quad f : U \otimes V \xrightarrow{\text{id}_U \otimes f'} U \otimes U^* \otimes W \xrightarrow{\text{ev} \otimes \text{id}_W} W.$$

Since $\text{ev}^* \otimes \text{id}_V \in \mathcal{YB}_R(V, U^\vee \times U \times V)$ and $\text{ev} \otimes \text{id}_W \in \mathcal{YB}_R(U^\vee \times U \times W, W)$, the map f is a morphism of \mathcal{YB}_R if and only if so is f' . Therefore (1.12) and (1.13) give the desired isomorphism. \square

2. Symmetric and exterior algebras associated with Yang-Baxter pairs. In this section, we introduce two classes of algebras associated with YB pairs.

DEFINITION 2.1. Let $V = (V, \beta_V)$ be a YB pair. Define quotient graded algebras $SV = \bigoplus_{i \geq 0} S_i V$ and $\bigwedge V = \bigoplus_{i \geq 0} \bigwedge_i V$ of the tensor algebra $TV = \bigoplus_{i \geq 0} V^{\otimes i}$ as follows:

$$(2.1) \quad \begin{aligned} SV &:= TV/(\text{Im}(\text{id}_{V \otimes V} - \beta_V)), \\ \bigwedge V &:= TV/(\text{Ker}(\text{id}_{V \otimes V} - \beta_V)). \end{aligned}$$

We call SV and $\bigwedge V$ the *symmetric* and the *exterior algebra* of V , respectively.

To distinguish elements of $\bigwedge V$ with those of SV , we denote the product of $a, b \in \bigwedge V$ by $a \wedge b$.

If β_V is the trivial twisting τ_V , then SV (resp. $\bigwedge V$) coincides with the usual symmetric (resp. exterior) algebra of the free R -module V . For Jimbo's YB pair of type $A_{N-1}^{(1)}$, the fundamental relations of these algebras are given by

$$(2.2) \quad \begin{aligned} SV &= \langle u_1, \dots, u_N \mid u_i u_j = q u_j u_i \quad (i < j) \rangle, \\ \bigwedge V &= \langle u_1, \dots, u_N \mid u_i \wedge u_i = 0, \quad q u_i \wedge u_j + u_j \wedge u_i = 0 \quad (i < j) \rangle. \end{aligned}$$

Hence, in this case, we have:

$$(2.3) \quad \text{If } i_1, \dots, i_k \in [1, N] \text{ has a repetition, then } u_{i_1} \wedge \cdots \wedge u_{i_k} = 0 \text{ in } \bigwedge V.$$

$$(2.4) \quad \text{If } 1 \leq i_1 < \cdots < i_k \leq N \text{ and } \sigma \in \mathfrak{S}_k, \text{ then } u_{i_1} \wedge \cdots \wedge u_{i_k} = (-q)^{-\ell(\sigma)} u_{i_{\sigma 1}} \wedge \cdots \wedge u_{i_{\sigma k}}$$

for $k \in \mathbb{N}$.

The R -modules $S_r V$ and $\bigwedge_r V$ are free with respective bases

$$(2.5) \quad \{u_{j_1} u_{j_2} \cdots u_{j_r} \mid 1 \leq j_1 \leq j_2 \leq \cdots \leq j_r \leq N\},$$

$$(2.6) \quad \{u_{j_1} \wedge u_{j_2} \wedge \cdots \wedge u_{j_r} \mid 1 \leq j_1 < j_2 < \cdots < j_r \leq N\}.$$

For the YB pair associated with a distributive lattice L , the symmetric algebra is the commutative algebra generated by elements of L with the fundamental relations $a \cdot b = (a \wedge b) \cdot (a \vee b)$ ($a, b \in L$). This algebra was introduced by Hibi [16] to study lattices by means of commutative ring theory.

3. Quantum matrices. Let V be a finite free R -module. Then, the dual coalgebra of $\text{End}(V)$ is identified with $V^* \otimes V$ as an R -module. With this identification, $V^* \otimes V$ is an R -coalgebra with the coproduct

$$(3.1) \quad \bar{\delta}: V^* \otimes V \simeq V^* \otimes R \otimes V \xrightarrow{1 \otimes \text{ev}^* \otimes 1} V^* \otimes V \otimes V^* \otimes V$$

and the counit $\bar{e} = \text{ev}: V^* \otimes V \rightarrow R$. More explicitly, these maps are expressed as

$$(3.2) \quad \bar{\delta}(x_{ij}) = \sum_{k=1}^N x_{ik} \otimes x_{kj}, \quad \bar{e}(x_{ij}) = \delta_{ij},$$

where $\{u_1, \dots, u_N\}$ is a free basis of V , $\{v_1, \dots, v_N\}$ is its dual basis and $x_{ij} = v_i \otimes u_j$, and δ_{ij} is Kronecker's delta.

The R -module V is a $V^* \otimes V$ -comodule with the coaction

$$(3.3) \quad \bar{\omega}: V \simeq R \otimes V \xrightarrow{\text{ev}^* \otimes 1} V \otimes V^* \otimes V; \quad u_j \longmapsto \sum_{i=1}^N u_i \otimes x_{ij}.$$

The maps δ and \bar{e} are extended to the (unique) algebra maps $\delta: T(V^* \otimes V) \rightarrow T(V^* \otimes V) \otimes T(V^* \otimes V)$ and $\bar{e}: T(V^* \otimes V) \rightarrow R$, respectively, and $T(V^* \otimes V)$ is a bialgebra. Hence, $V^{\otimes k}$ is a $T(V^* \otimes V)$ -comodule with the coaction $\bar{\omega}$ given by

$$(3.4) \quad \bar{\omega}(u_{j_1} \otimes \cdots \otimes u_{j_k}) = \sum_{i_1, \dots, i_k} (u_{i_1} \otimes \cdots \otimes u_{i_k}) \otimes (x_{i_1 j_1} \otimes \cdots \otimes x_{i_k j_k}).$$

Let A be an R -coalgebra, and V an A -comodule with coaction $\omega_V: V \rightarrow V \otimes A$. We define a coalgebra homomorphism $\text{cf}_V: V^* \otimes V \rightarrow A$ by $\text{cf}_V = (\text{ev} \otimes \text{id}_A) \circ (\text{id}_{V^*} \otimes \omega_V)$ and call it the *coefficient map* of V . The coaction ω_V is determined by cf_V by the formula $\omega_V = (\text{id}_V \otimes \text{cf}_V) \circ \bar{\omega}$.

Now we consider a YB operator β_V on V . The YB pair $(E, \beta_E) := (V, \beta_V) \times (V, \beta_V)^\vee$ is called the *internal End* of the YB pair (V, β_V) . The R -module E (resp. E^\vee) is nothing but $V \otimes V^*$ (resp. $V^* \otimes V$). By Theorem 1.5, E^\vee is identified with the internal End $V^\vee \times (V^\vee)^\vee$ of V^\vee . Since $\text{ev}^* \in \mathcal{YB}_R(R, V^\vee \times V)$ and $\text{ev} \in \mathcal{YB}_R(V \times V^\vee, R)$, the maps δ , \bar{e} and $\bar{\omega}$ are contained in $\mathcal{YB}_R(E^\vee, E^\vee \times E^\vee)$, $\mathcal{YB}_R(E^\vee, R)$ and $\mathcal{YB}_R(V, V \times E^\vee)$, respectively. It is easy to see that β_{E^\vee} is a coalgebra map. Hence, by [34, Proposition 1.4.8], $(\text{Im}(\text{id}_{E^\vee} \otimes E^\vee - \beta_{E^\vee}))$ is a bi-ideal of TE^\vee . Thus the symmetric algebra SE^\vee becomes a quotient bialgebra of the bialgebra $(TE^\vee, \delta, \bar{e})$. We call this the *quantum matrix bialgebra* for the YB pair (V, β_V) and denote its coproduct and counit by δ_S and e_S , respectively. Note that the exterior algebra $\bigwedge E^\vee$ also becomes a quotient bialgebra of $(TE^\vee, \delta, \bar{e})$.

Let us write down the fundamental relations of SE^\vee and $\bigwedge E^\vee$ for Jimbo's YB pair of type $A_{N-1}^{(1)}$. For Jimbo's YB pair of type $A_{N-1}^{(1)}$, the internal End YB operator β_{E^\vee} is given by

$$(3.5) \quad \begin{aligned} \beta_E(x_{ik} \otimes x_{ik}) &= x_{ik} \otimes x_{ik}, \quad \beta_E(x_{il} \otimes x_{ik}) = (1-q^2)x_{il} \otimes x_{ik} + qx_{ik} \otimes x_{il}, \\ \beta_E(x_{ik} \otimes x_{il}) &= qx_{il} \otimes x_{ik}, \quad \beta_E(x_{jk} \otimes x_{ik}) = q^{-1}x_{ik} \otimes x_{jk}, \\ \beta_E(x_{ik} \otimes x_{jk}) &= (1-q^{-2})x_{ik} \otimes x_{jk} + q^{-1}x_{jk} \otimes x_{ik}, \\ \beta_E(x_{jl} \otimes x_{ik}) &= (q^{-1}-q)x_{il} \otimes x_{jk} + x_{ik} \otimes x_{jl}, \quad \beta_E(x_{jk} \otimes x_{il}) = x_{il} \otimes x_{jk}, \\ \beta_E(x_{il} \otimes x_{jk}) &= -(q-q^{-1})^2 x_{il} \otimes x_{jk} + x_{jk} \otimes x_{il} + (q-q^{-1})(x_{jl} \otimes x_{ik} - x_{ik} \otimes x_{jl}), \\ \beta_E(x_{ik} \otimes x_{jl}) &= -(q-q^{-1})x_{il} \otimes x_{jk} + x_{jl} \otimes x_{ik}, \quad (1 \leq i < j \leq N, 1 \leq k < l \leq N). \end{aligned}$$

Hence, the fundamental relations of SE^\vee are

$$(3.6) \quad \begin{aligned} x_{ik}x_{il} &= qx_{il}x_{ik}, \quad x_{ik}x_{jk} = qx_{jk}x_{ik}, \quad x_{il}x_{jk} = x_{jk}x_{il}, \\ x_{jl}x_{ik} - x_{ik}x_{jl} + (q-q^{-1})x_{il}x_{jk} &= 0, \quad (1 \leq i < j \leq N, 1 \leq k < l \leq N), \end{aligned}$$

and those of $\bigwedge E^\vee$ are

$$(3.7) \quad \begin{aligned} x_{ik} \wedge x_{ik} &= 0, \quad x_{ik} \wedge x_{il} + qx_{il} \wedge x_{ik} = 0, \quad x_{ik} \wedge x_{jk} + qx_{jk} \wedge x_{ik} = 0, \\ x_{il} \wedge x_{jk} + x_{jk} \wedge x_{il} - (q - q^{-1})x_{ik} \wedge x_{jl} &= 0, \quad x_{ik} \wedge x_{jl} + x_{jl} \wedge x_{ik} = 0, \\ (1 \leq i < j \leq N, \quad 1 \leq k < l \leq N). \end{aligned}$$

Using [5], one can easily see that $S_r E^\vee$ and $\bigwedge_r E^\vee$ have the following R -free bases, respectively:

$$(3.8) \quad \begin{aligned} &\{x_{i_1 j_1} x_{i_2 j_2} \cdots x_{i_r j_r} \mid (i_1, j_1) \leq (i_2, j_2) \leq \cdots \leq (i_r, j_r)\}, \\ &\{x_{i_1 j_1} \wedge x_{i_2 j_2} \wedge \cdots \wedge x_{i_r j_r} \mid (i_1, j_1) < (i_2, j_2) < \cdots < (i_r, j_r)\}. \end{aligned}$$

Here \leq denotes the lexicographic ordering of $\{1, \dots, N\}^2$.

In the remainder of this section, we will consider representations of the quantum matrix bialgebra SE^\vee for an arbitrary YB pair (V, β_V) . Since V is a comodule for the bialgebra SE^\vee with the structure map $\omega_S(u_j) = \sum_i u_i \otimes x_{ij}$, the k -fold tensor power $T_k V$ of V is also an SE^\vee -comodule with the structure map

$$(3.9) \quad \omega_S(u_{j_1} \otimes \cdots \otimes u_{j_k}) = \sum_{i_1, \dots, i_k} (u_{i_1} \otimes \cdots \otimes u_{i_k}) \otimes x_{i_1 j_1} \cdots x_{i_k j_k}.$$

PROPOSITION 3.1. *The coaction of SE^\vee on $T_k V$ commutes with the action of the braid group \mathfrak{B}_k . In other words, the following diagram is commutative for $1 \leq i < k$:*

$$(3.10) \quad \begin{array}{ccc} T_k V & \xrightarrow{\omega_S} & T_k V \otimes SE^\vee \\ \downarrow (\beta_V)_i & & \downarrow (\beta_V)_i \otimes \text{id} \\ T_k V & \xrightarrow{\omega_S} & T_k V \otimes SE^\vee. \end{array}$$

PROOF. Clearly, it is enough to consider the case $k=2$. Let $\bar{\omega}$ be the coaction of (TE^\vee, δ) on TV . Since $\bar{\omega} \in \mathcal{YB}_R(V, V \times E^\vee)$, we have

$$(3.11) \quad (\beta_V \otimes \beta_{E^\vee}) \circ \bar{\omega}|_{V \otimes V} = \bar{\omega}|_{V \otimes V} \circ \beta_V.$$

Let p be the projection from $T_2 E^\vee$ onto $S_2 E^\vee$. Since $\omega_S|_{V \otimes V} = (\text{id}_{V \otimes V} \otimes p) \circ \bar{\omega}|_{V \otimes V}$, we get

$$(3.12) \quad \begin{aligned} \omega_S|_{V \otimes V} \circ \beta_V &= p \otimes \text{id}_{V \otimes V} \circ \beta_E \otimes \beta_V \circ \bar{\omega}|_{V \otimes V} \\ &= p \otimes \beta_V \circ \bar{\omega}|_{V \otimes V} = \text{id}_{SE^\vee} \otimes \beta_V \circ \omega_S|_{V \otimes V}. \end{aligned}$$

□

THEOREM 3.2 (Universal mapping property of SE^\vee). *Suppose there is a bialgebra (A, δ_A) and coaction $\omega_A: V \rightarrow V \otimes A$ such that the coaction of A on $T_k V$ commutes with the action of the braid group \mathfrak{B}_k . Then there exists a unique bialgebra homomorphism $f: SE^\vee \rightarrow A$ such that the following diagram is commutative.*

$$(3.13) \quad \begin{array}{ccc} T_k V & \xrightarrow{\omega_s} & T_k V \otimes SE^\vee \\ \downarrow = & & \downarrow \text{id} \otimes f \\ T_k V & \xrightarrow{\omega_A} & T_k V \otimes A . \end{array}$$

PROOF. The coalgebra map $\text{cf}_V: E^\vee = V^* \otimes V \rightarrow A$ uniquely extends to a bialgebra map $\tilde{f}: TE^\vee \rightarrow A$. By (3.10), it is easy to see that $\tilde{f}(\text{Im}(\text{id} - \beta_{E^\vee})) = 0$ and that the theorem follows. \square

Similarly, we get:

THEOREM 3.3. Suppose there is a bialgebra (U, δ_U) and an action $\theta_U: U \otimes V \rightarrow V$ such that the action of U on $T_k V$ commutes with the action of the braid group. Then there exists a unique bilinear pairing $\langle , \rangle: U \otimes SE^\vee \rightarrow R$ satisfying

- (1) $\langle X, ab \rangle = (\langle , a \rangle \otimes \langle , b \rangle)(\delta_U X)$,
- (2) $\langle XY, a \rangle = (\langle X, \rangle \otimes \langle Y, \rangle)(\delta_S a)$,
- (3) $\theta_U(X \otimes w) = (\text{id}_{T(V)} \otimes \langle X, \rangle)(\omega_S(w)) \quad (X, Y \in U, a, b \in SE^\vee, w \in TV)$.

In particular, the bialgebra U acts on SE^\vee by $Xa = (\text{id}_{S(E)} \otimes \langle X, \rangle) \circ \delta_S(a)$ ($X \in U, a \in SE$).

4. Symmetric algebras, exterior algebras and the fusion procedure.

DEFINITION 4.1. Let $A = (A, m_A, \eta_A)$ (resp. $C = (C, \Delta_C, \varepsilon_C)$) be an R -algebra with product m_A and unit η_A (resp. an R -coalgebra with coproduct Δ_C and counit ε_C). Let φ_A (resp. φ_C) be a YB operator on A (resp. C). We call (A, φ_A) (resp. (C, φ_C)) a YB algebra (resp. YB coalgebra) if the following diagrams (4.1) (resp. (4.2)) are commutative:

$$(4.1) \quad \begin{array}{ccccc} A \otimes A \otimes A & \xrightarrow{\varphi_A(\chi_{21})} & A \otimes A \otimes A & \xrightarrow{\varphi_A(\chi_{12})} & A \otimes A \otimes A \\ \downarrow m_A \otimes \text{id}_A & & \downarrow \text{id}_A \otimes m_A & & \downarrow m_A \otimes \text{id}_A \\ A \otimes A & \xrightarrow{\varphi_A} & A \otimes A & \xrightarrow{\varphi_A} & A \otimes A \\ \uparrow \eta_A \otimes \text{id}_A & & \uparrow \text{id}_A \otimes \eta_A & & \uparrow \eta_A \otimes \text{id}_A \\ R \otimes A & \xrightarrow{\cong} & A \otimes R & \xrightarrow{\cong} & R \otimes A , \end{array}$$

$$(4.2) \quad \begin{array}{ccccc} C \otimes C \otimes C & \xrightarrow{\varphi_C(\chi_{21})} & C \otimes C \otimes C & \xrightarrow{\varphi_C(\chi_{12})} & C \otimes C \otimes C \\ \uparrow \Delta_C \otimes \text{id}_C & & \uparrow \text{id}_C \otimes \Delta_C & & \uparrow \Delta_C \otimes \text{id}_C \\ C \otimes C & \xrightarrow{\varphi_C} & C \otimes C & \xrightarrow{\varphi_C} & C \otimes C \\ \downarrow \varepsilon_C \otimes \text{id}_C & & \downarrow \text{id}_C \otimes \varepsilon_C & & \downarrow \varepsilon_C \otimes \text{id}_C \\ R \otimes C & \xrightarrow{\cong} & C \otimes R & \xrightarrow{\cong} & R \otimes C . \end{array}$$

PROPOSITION 4.2. (1) For a YB algebra (A, φ_A) , the YB pair $T_i(A, \varphi_A)$ (see Proposition 1.1) becomes a YB algebra with product $m_{T_i A} := m_A^{\otimes i} \circ \varphi_A(w_i)$ and unit $(\eta_A)^{\otimes i} : R \simeq R^{\otimes i} \rightarrow T_i A$, where w_i is the element of \mathfrak{S}_{2i} defined by

$$(4.3) \quad w_i = \begin{pmatrix} 1 & 2 & 3 & \cdots & i & i+1 & i+2 & \cdots & 2i \\ 1 & 3 & 5 & \cdots & 2i-1 & 2 & 4 & \cdots & 2i \end{pmatrix}.$$

(2) For a YB coalgebra (C, φ_C) , the YB pair $T_i(C, \varphi_C)$ becomes a YB coalgebra with coproduct $\Delta_{T_i C} := \varphi_C(w_i^{-1}) \circ \Delta_C^{\otimes i}$ and counit $(\varepsilon_C)^{\otimes i} : T_i C \rightarrow R^{\otimes i} \simeq R$.

PROOF. By (4.1), we have

$$(4.4) \quad \varphi_A(\chi_{ij}) \circ (m_{T_i A} \otimes \text{id}_A^{\otimes j}) = (\text{id}^{\otimes j} \otimes m_A^{\otimes i}) \circ \varphi_A(\chi_{2i,j}) \circ (\varphi_A(w_i) \otimes \text{id}_A^{\otimes j}).$$

Since both sides of the equality $\chi_{2i,j}(w_i \times 1_{\mathfrak{S}_j}) = (1_{\mathfrak{S}_j} \times w_i)\chi_{2i,j}$ are reduced, the right hand side of (4.4) becomes

$$(\text{id}^{\otimes j} \otimes m_A^{\otimes i}) \circ (\text{id}_A^{\otimes j} \otimes \varphi_A(w_i)) \circ \varphi_A(\chi_{2i,j}),$$

hence we have

$$(4.5) \quad \varphi_A(\chi_{ij}) \circ (m_{T_i A} \otimes \text{id}_A^{\otimes j}) = (\text{id}^{\otimes j} \otimes m_{T_i A}) \circ \varphi_A(\chi_{2i,j}).$$

Similarly

$$(4.6) \quad \varphi_A(\chi_{ji}) \circ (\text{id}_A^{\otimes j} \otimes m_{T_i A}) = (m_{T_i A} \otimes \text{id}_A^{\otimes j}) \circ \varphi_A(\chi_{j,2i}).$$

Taking $j=i$, we find the commutativity of the upper two diagrams of (4.1) for $T_i A$. Now we will prove the associativity of $m_{T_i A}$. Since $(1 \times 1 \times w_i)(1 \times \chi_{i1} \times 1)$ is a reduced expression for w_{i+1} , we have

$$(4.7) \quad m_{T_{i+1} A} = (m_A \otimes m_{T_i A}) \circ (\text{id}_A \otimes \varphi_A(\chi_{i1}) \otimes \text{id}_{T_i A}).$$

Hence by (4.5), (4.6), we get

$$(4.8) \quad \begin{aligned} m_{T_{i+1} A} \circ (m_{T_{i+1} A} \otimes \text{id}_{T_{i+1} A}) &= ((m_A \circ (m_A \otimes \text{id}_A)) \otimes (m_{T_i A} \circ (m_{T_i A} \otimes \text{id}_{T_i A}))) \circ \varphi_A(\sigma), \\ m_{T_{i+1} A} \circ (\text{id}_{T_{i+1} A} \otimes m_{T_{i+1} A}) &= ((m_A \circ (\text{id}_A \otimes m_A)) \otimes (m_{T_i A} \circ (\text{id}_{T_i A} \otimes m_{T_i A}))) \circ \varphi_A(\sigma), \end{aligned}$$

where $\sigma = (1_{\mathfrak{S}_2} \times \chi_{2i,1} \times 1_{\mathfrak{S}_i})(1_{\mathfrak{S}_1} \times \chi_{i1} \times 1_{\mathfrak{S}_{2i+1}})$. Hence the associativity of $m_{T_{i+1} A}$ follows from that of m_A and $m_{T_i A}$. The rest of the proof is similar and easy. \square

EXAMPLE. For an arbitrary algebra A , the YB pair (A, τ_A) becomes a YB algebra (cf. §1 Example (1)). The product of $T_i(A, \tau_A)$ is given by $m_{T_i A}((a_1 \otimes \cdots \otimes a_i) \otimes (b_1 \otimes \cdots \otimes b_i)) = a_1 b_1 \otimes \cdots \otimes a_i b_i$.

DEFINITION 4.3. A sextuple $A = (A, \Delta_A, \varepsilon_A, m_A, \eta_A, \varphi_A)$ is called a *graded YB-bialgebra* over R , if:

(1) $(A, m_A, \eta_A, \varphi_A)$ is an YB algebra over R and $(A, \Delta_A, \varepsilon_A, \varphi_A)$ is an YB coalgebra over R .

- (2) The R -module A is a direct sum of submodules A_i ($i \geq 0$) such that $m_A(A_i \otimes A_j) \subset A_{i+j}$, $\Delta_A(A_k) \subset \sum_{i+j=k} A_i \otimes A_j$ and $\varphi_A(A_i \otimes A_j) \subset A_j \otimes A_i$.
 (3) The unit η_A and the counit ε_A give an isomorphism $A_0 \simeq R$.
 (4) Both $\Delta_A: A \rightarrow T_2(A, m_A, \eta_A, \varphi_A)$ and ε_A are algebra maps.

Note that in general, A is *not* a bialgebra in the usual sense. But the following lemmas still hold.

LEMMA 4.4. *Under the conditions (1), (2), (3), the condition (4) in Definition 4.3 is equivalent to*

- (4') *Both η_A and m_A are coalgebra maps, where the coalgebra structure of $A \otimes A$ is as in Proposition 4.2 (2).*

LEMMA 4.5. *For a graded YB bialgebra (A, φ_A) , the tensor product $T_i(A, \varphi_A)$ ($i \geq 0$) becomes a graded YB bialgebra with a product $m_{T_i A}$ and a coproduct $\Delta_{T_i A}$ (see Proposition 4.2).*

Let (A, φ_A) and (B, φ_B) be YB algebras (resp. coalgebras, resp. bialgebras), and $f: A \rightarrow B$ a homomorphism of algebras (resp. coalgebras, resp. both algebras and coalgebras). We say that f is a homomorphism of YB algebras (resp. coalgebras, resp. bialgebras) when $(f \otimes f) \circ \varphi_A = \varphi_B \circ (f \otimes f)$ (cf. (1.10)). For a homomorphism $f: (A, \varphi_A) \rightarrow (B, \varphi_B)$ of YB algebras (resp. coalgebras, resp. bialgebras), $T_i f: (T_i A, T_i \varphi_A) \rightarrow (T_i B, T_i \varphi_B)$ is again a homomorphism of YB algebras (resp. coalgebras, resp. bialgebras) for $i \geq 0$.

Let β_V and γ_V be YB operators on a free R -module V . We call (V, β_V, γ_V) a YB triple if the following two conditions are satisfied.

$$(4.9) \quad (\text{id}_{V \otimes V} - \beta_V)(\text{id}_{V \otimes V} - \gamma_V) = (\text{id}_{V \otimes V} - \beta_V)(\text{id}_{V \otimes V} - \gamma_V) = 0,$$

$$(4.10) \quad \begin{aligned} \beta_1 \circ \beta_2 \circ \gamma_1 &= \gamma_2 \circ \beta_1 \circ \beta_2, & \gamma_1 \circ \gamma_2 \circ \beta_1 &= \beta_2 \circ \gamma_1 \circ \gamma_2, \\ \beta_2 \circ \beta_1 \circ \gamma_2 &= \gamma_1 \circ \beta_2 \circ \beta_1, & \gamma_2 \circ \gamma_1 \circ \beta_2 &= \beta_1 \circ \gamma_2 \circ \gamma_1. \end{aligned}$$

For a YB triple $V = (V, \beta_V, \gamma_V)$, we set $SV = S(V, \beta_V)$ and $\bigwedge V = \bigwedge(V, \beta_V)$. There exist two important examples of YB triples.

EXAMPLE (1). Let (V, β_V) be Jimbo's YB pair of type $A_{N-1}^{(1)}$ for a fixed ring R and its invertible element q . By (1.3), the YB operator $\gamma_V := -q^{-2}\beta_V$ satisfies the equation (4.9).

EXAMPLE (2). Let (V, β_V) be as in Example (1) and let (E, β_E) be its internal End YB pair $V \times V^\vee$. Set $\gamma_E = \beta_V \times {}^t \gamma_V$. It is easy to verify that (E, β_E, γ_E) is a YB triple.

Let φ_A and ψ_A be YB operators on an algebra A . We call (A, φ_A, ψ_A) a YB algebra triple if (1) both (A, φ_A) and (A, ψ_A) are YB algebras and (2) $\beta = \varphi_A$ and $\gamma = \psi_A$ satisfy (4.10).

PROPOSITION 4.6. (1) Let (V, β_V) be a YB pair. Then the tensor algebra $TV = \bigoplus_{i \geq 0} V^{\otimes i}$ becomes a graded YB bialgebra with a YB operator $T\beta_V$ and a coproduct $\Delta_{TV}: TV \rightarrow T_2(TV, T\beta_V)$ defined by $\Delta(u) = u \otimes 1 + 1 \otimes u$ ($u \in V$). (2) If (V, β_V, γ_V) is a YB triple, then $(TV, T(\pm \beta_V), T(\pm \gamma_V))$ is a YB algebra triple.

PROOF. By Proposition 1.1, $(TV, T\beta_V)$ is a YB algebra. By the universal mapping property of the tensor algebra, there exists a unique algebra map $\Delta_{TV}: TV \rightarrow T_2(V, T\beta_V)$ such that $\Delta(u) = u \otimes 1 + 1 \otimes u$ for $u \in V$. The commutativity of (4.2) follows from Proposition 4.8 below. Next, we show the coassociativity of Δ_{TV} . Because the restrictions of $\Delta_{TV} \otimes \text{id} \circ \Delta_{TV}$ and $\text{id} \otimes \Delta_{TV} \circ \Delta_{TV}$ on V coincide, it suffices to show that these are algebra maps. Since Δ_{TV} is an algebra map, this follows easily from the commutativity of (4.2). Part (2) follows from the following lemma. \square

LEMMA 4.7. For a YB triple (V, β_V, γ_V) , we have

$$(4.11) \quad \begin{aligned} \beta(\chi_{ij}) \circ \beta_k &= \beta_{k+j} \circ \beta(\chi_{ij}), & \gamma(\chi_{ij}) \circ \beta_k &= \beta_{k+j} \circ \gamma(\chi_{ij}) \quad (1 \leq k < i), \\ \beta(\chi_{ij}) \circ \beta_{k+i} &= \beta_k \circ \beta(\chi_{ij}), & \gamma(\chi_{ij}) \circ \beta_{k+i} &= \beta_k \circ \gamma(\chi_{ij}) \quad (1 \leq k < j). \end{aligned}$$

PROOF. Since

$$(4.12) \quad \chi_{ij} = (1_{\mathfrak{S}_{j-1}} \times \chi_{i1}) \cdots (1_{\mathfrak{S}_1} \times \chi_{i1} \times 1_{\mathfrak{S}_{j-2}})(\chi_{i1} \times 1_{\mathfrak{S}_{j-1}})$$

is a reduced expression, we have

$$(4.13) \quad \gamma(\chi_{ij}) = \gamma(1_{\mathfrak{S}_{j-1}} \times \chi_{i1}) \cdots \gamma(1_{\mathfrak{S}_1} \times \chi_{i1} \times 1_{\mathfrak{S}_{j-2}})\gamma(\chi_{i1} \times 1_{\mathfrak{S}_{j-1}}).$$

Hence the second identity follows from

$$(4.14) \quad \gamma(1_{\mathfrak{S}_m} \times \chi_{i1} \times 1_{\mathfrak{S}_{j-m-1}})\beta_{k+m} = \beta_{k+m+1}\gamma(1_{\mathfrak{S}_m} \times \chi_{i1} \times 1_{\mathfrak{S}_{j-m-1}}).$$

The other identities are obtained similarly. \square

To give an explicit formula for Δ_{TV} , we need some notation. Let $k \geq 1$ and let $\alpha = (\alpha_1, \dots, \alpha_s)$ be a sequence of non-negative integers with $\alpha_1 + \dots + \alpha_s = k$. Let $C = \bigoplus_{i \geq 0} C_i$ be a graded coalgebra. We denote by Δ_C^α the composite map $C \rightarrow C^{\otimes s} \rightarrow C_\alpha$ of the s -th iteration of Δ_C and the projection, where $C_\alpha = C_{\alpha_1} \otimes \dots \otimes C_{\alpha_s}$. If α and C in consideration are clear by context, then we will drop them, and simply denote it by Δ . Let \mathfrak{S}_α be the Young subgroup $\mathfrak{S}_{\alpha_1} \times \dots \times \mathfrak{S}_{\alpha_s}$ of \mathfrak{S}_k . We denote by \mathfrak{S}^α the following complete set of representatives of $\mathfrak{S}_k/\mathfrak{S}_\alpha$ in \mathfrak{S}_k .

$$(4.15) \quad \mathfrak{S}^\alpha := \{\sigma \in \mathfrak{S}_k \mid \sigma 1 < \dots < \sigma \alpha_1, \sigma(\alpha_1 + 1) < \dots < \sigma(\alpha_1 + \alpha_2), \dots, \sigma(\alpha_1 + \dots + \alpha_{s-1} + 1) < \dots < \sigma(\alpha_1 + \dots + \alpha_s)\}.$$

Any $\sigma \in \mathfrak{S}_k$ is uniquely expressed as $\sigma = \sigma' \sigma''$ with $\sigma' \in \mathfrak{S}^\alpha$ and $\sigma'' \in \mathfrak{S}_\alpha$, and this expression is reduced.

PROPOSITION 4.8. For a YB pair (V, β_V) , the coproduct of the YB bialgebra $T(V, \beta_V)$ satisfies the following formula.

$$(4.16) \quad \Delta_{TV}^{\alpha} = \sum_{\sigma \in \mathfrak{S}^{\alpha}} \beta_V(\sigma^{-1}).$$

PROOF. We prove this formula for $\alpha=(i,j)$ by induction on $k=i+j$. Let w and u be elements of $T_{i+j}V$ and V , respectively. Since Δ_{TV} is an algebra map, we have

$$(4.17) \quad \Delta^{(i,j+1)}(w \otimes u) = m_{T_2(TV)}(\Delta^{(i-1,j+1)}(w) \otimes (u \otimes 1)) + m_{T_2(TV)}(\Delta^{(i,j)}(w) \otimes (1 \otimes u)).$$

By the induction hypothesis, the right hand side of (4.17) is

$$\begin{aligned} & (\text{id}_V^{\otimes i-1} \otimes \beta_V(\chi_{j+1,1})) \circ \left(\sum_{\sigma \in \mathfrak{S}^{(i-1,j+1)}} \beta_V(\sigma^{-1}) \otimes \text{id}_V \right) (w \otimes u) \\ & + \sum_{\sigma \in \mathfrak{S}^{(i,j)}} \beta_V(\sigma^{-1}) \otimes \text{id}_V (w \otimes u). \end{aligned}$$

Since

$$\mathfrak{S}^{(i,j+1)} = (\mathfrak{S}^{(i-1,j+1)}(1_{\mathfrak{S}_{i-1}} \times \chi_{1,j+1})) \coprod \mathfrak{S}^{(i,j)},$$

we obtain

$$\Delta^{(i,j+1)} = \sum_{\sigma \in \mathfrak{S}^{(i,j+1)}} \beta_V(\sigma^{-1}).$$

□

Now we fix a YB triple $V=(V, \beta_V, \gamma_V)$ such that $S_i V$ and $\bigwedge_i V$ are free R -modules for each $i \geq 0$. The next lemma follows easily from Lemma 4.7.

LEMMA 4.9. (1) *There exist YB operators φ_{SV} and ψ_{SV} on SV (resp. $\varphi_{\bigwedge V}$ and $\psi_{\bigwedge V}$ on $\bigwedge V$) which satisfy*

$$(4.18) \quad \begin{aligned} \varphi_{SV} \circ (p \otimes p) &= (p \otimes p) \circ T(-\gamma_V), & \psi_{SV} \circ (p \otimes p) &= (p \otimes p) \circ T\beta_V, \\ \varphi_{\bigwedge V} \circ (p \otimes p) &= (p \otimes p) \circ T(-\beta_V), & \psi_{\bigwedge V} \circ (p \otimes p) &= (p \otimes p) \circ T\gamma_V, \end{aligned}$$

where p denotes the projection from TV onto SV (resp. $\bigwedge V$).

(2) *Both $(SV, \varphi_{SV}, \psi_{SV})$ and $(\bigwedge V, \varphi_{\bigwedge V}, \psi_{\bigwedge V})$ are YB algebra triples.*

Let A be either SV or $\bigwedge V$. Then we have the following:

THEOREM 4.10. (1) *There exists a unique algebra map $\Delta_A: A \rightarrow A \otimes A$ such that $\Delta_A(u) = u \otimes 1 + 1 \otimes u$ for $u \in A_1 \simeq V$, where the algebra structure of $A \otimes A$ is given by Proposition 4.2 in terms of the YB operator φ_A .*

(2) *Define a map $\varepsilon_A: A \rightarrow R$ by $\varepsilon_A(\sum_i a_i) = a_0$ ($a_i \in A_i$). Then $(A, m_A, \eta_A, \Delta_A, \varepsilon_A, \varphi_A)$ becomes a graded YB bialgebra.*

PROOF. We prove this theorem only in the case $A=SV$. By Proposition 4.6 (1), it suffices to show that the ideal $(\text{Im}(\text{id} - \beta_V))$ is a coideal of TV . By (4.11),

$(\text{Im}(\text{id} - \beta_V)) \otimes TV + TV \otimes (\text{Im}(\text{id} - \beta_V))$ is a two-sided ideal of $T_2(TV, T(-\gamma_V))$. On the other hand, by (4.9) and (4.16),

$$\begin{aligned}\Delta_{TV}((\text{id} - \beta_V)(w)) &= (\text{id} - \beta_V)(w) \otimes 1 + 1 \otimes (\text{id} - \beta_V)(w) \\ &\in (\text{Im}(\text{id} - \beta_V)) \otimes TV + TV \otimes (\text{Im}(\text{id} - \beta_V))\end{aligned}$$

for $w \in T_2 V$. \square

PROPOSITION 4.11. *Let A be as above. Then, (1) $m_A \circ \psi_A = m_A$. (2) If $\Delta: A_i \rightarrow A_1^{\otimes i}$ is injective for any $i \geq 0$, then, $\psi_A \circ \Delta_A = \Delta_A$.*

PROOF. We show this proposition only for $A = SV$. Using (4.2), we obtain

$$\begin{aligned}(4.19) \quad \Delta^{(1^i)} \otimes \Delta^{(1^j)} \circ \psi_{SV} \circ \Delta^{(i,j)} \circ p_{i+j} &= \beta(\chi_{ij}) \circ \Delta^{(1^i)} \otimes \Delta^{(1^j)} \circ \Delta^{(i,j)} \circ p_{i+j} \\ &= \beta(\chi_{ij}) \circ \Delta^{(1^{i+j})} \circ p_{i+j},\end{aligned}$$

where $(1^i) = (1, \dots, 1) \in \mathbb{Z}^i$. By (4.16), the right hand side of (4.19) is $\beta(\chi_{ij}) \circ \sum_{\sigma \in \mathfrak{S}_k} (-\gamma_V)(\sigma)$. Since

$$(\beta_V)_r \sum_{\sigma \in \mathfrak{S}_k} (-\gamma_V)(\sigma) = (\beta_V)_r \sum_{\ell(\sigma) < \ell((r, r+1)\sigma)} (\text{id} - (\gamma_V)_r)(-\gamma_V)(\sigma) = \sum_{\sigma \in \mathfrak{S}_k} (-\gamma_V)(\sigma)$$

for $1 \leq r \leq i+j$, we get

$$\Delta^{(1^i)} \otimes \Delta^{(1^j)} \circ \psi_{SV} \circ \Delta^{(i,j)} \circ p_{i+j} = \Delta^{(1^i)} \otimes \Delta^{(1^j)} \circ \Delta^{(i,j)} \circ p_{i+j}.$$

This proves Part (2). Part (1) follows from

$$m_A \circ \psi_A \circ p_i \otimes p_j = m_A \circ p_j \otimes p_i \circ \beta(\chi_{ij}) = p_{i+j} \circ \beta(\chi_{ij}) = p_{i+j} = m_A \circ p_i \otimes p_j.$$

\square

Now we shall look at the representation-theoretical aspects of the YB algebras we discussed so far.

DEFINITION 4.12. Let H be an R -bialgebra, and $A = (A, \varphi_A)$ be a YB algebra with a H -comodule structure. We say that A is H -equivariant YB algebra when m_A , η_A and φ_A are homomorphisms of H -comodules. We define H -equivariance of YB algebra triples, YB coalgebras, and YB bialgebras similarly. All of the structure morphisms (including YB operators) are required to be homomorphisms of H -comodules.

If (A, φ_A) is a H -equivariant YB algebra (resp. coalgebra, resp. bialgebra), then $T_i(A, \varphi_A)$ is also H -equivariant for any $i \geq 0$.

Let (V, β, γ) be a YB triple such that SV and $\bigwedge V$ are free, that V is a H -comodule, and that β is a homomorphism of H -comodules. Then it is easy to see that $(TV, T(\pm \beta))$ is a H -equivariant YB algebra. The map $V \rightarrow V \otimes R \oplus R \otimes V$; $u \mapsto u \otimes 1 + 1 \otimes u$ is clearly a H -homomorphism. Since $T_2(TV, T(\pm \beta))$ is a H -equivariant YB algebra, the coproduct Δ_{TV} of $(TV, T(\pm \beta))$ is also a H -homomorphism. Hence, $(TV, T(\pm \beta))$ is a H -equivariant

YB bialgebra. Since the ideals $(\text{Im}(1-\beta))$ and $(\text{Ker}(1-\beta))$ are subcomodules of TV , we see that SV and $\bigwedge V$ are quotient comodules of TV . The YB bialgebra structure of $(\bigwedge V, \varphi_{\bigwedge V})$ is induced by the structure of $(TV, T(-\beta))$. Hence, $\bigwedge V$ is also a H -equivariant YB bialgebra. Though m_{SV} , η_{SV} , ε_{SV} , and ψ_{SV} are H -homomorphisms, Δ_{SV} , φ_{SV} and $\psi_{\bigwedge V}$ may not be H -homomorphisms.

The most important example of H is the quantum matric bialgebra SE^\vee (see Proposition 3.1 and Theorem 3.2). If γ is also a H -homomorphism, then $(TV, T(\pm\gamma))$ is also a H -equivariant YB bialgebra. It is easy to see that $(\bigwedge V, \varphi_{\bigwedge V}, \psi_{\bigwedge V})$ and $(SV, \varphi_{SV}, \psi_{SV})$ are H -equivariant both as YB algebra triples and as YB bialgebras in this case. Jimbo's YB triple of type $A_{N-1}^{(1)}$ and its matric bialgebra satisfy this condition, since $\gamma_V = -q^{-2}\beta_V$.

5. Divided power algebra. For a graded R -module $M = \bigoplus_{k \geq 0} M_k$, the graded dual M^\dagger of M is $\bigoplus_{k \geq 0} M^*$ by definition. The graded dual f^\dagger of a homogeneous map $f: M \rightarrow N = \bigoplus_k N_k$ is defined to be $\sum_k f_k^*$, where f_k is the degree k component of f .

Let $A = (A, \Delta_A, \varepsilon_A, m_A, \eta_A, \varphi_A)$ be a graded YB-bialgebra with each homogeneous component A_i being free of finite rank for each $i \geq 0$. We introduce into A^\dagger a structure of a graded YB-bialgebra. There is a canonical isomorphism $A^\dagger \otimes A^\dagger \simeq (A \otimes A)^\dagger$ via the pairing

$$A_i^* \otimes A_j^* \otimes A_i \otimes A_j \xrightarrow{1 \otimes \tau \otimes 1} A_i^* \otimes A_i \otimes A_j^* \otimes A_j \xrightarrow{\text{ev} \otimes \text{ev}} R \otimes R = R,$$

where τ is the trivial twisting. The structure morphisms Δ_{A^\dagger} , ε_{A^\dagger} , m_{A^\dagger} , η_{A^\dagger} and φ_{A^\dagger} are defined to be m_A^\dagger , η_A^\dagger , Δ_A^\dagger , ε_A^\dagger and φ_A^\dagger , respectively. For example,

$$A_i^* \otimes A_j^* \otimes A_j \otimes A_i \xrightarrow{\varphi_{A^\dagger} \otimes 1} A_j^* \otimes A_i^* \otimes A_j \otimes A_i \xrightarrow{\text{ev}} R$$

is equal to

$$A_i^* \otimes A_j^* \otimes A_j \otimes A_i \xrightarrow{1 \otimes \varphi_A} A_i^* \otimes A_j^* \otimes A_i \otimes A_j \xrightarrow{\text{ev}} R.$$

Thus, for $k \geq 0$ and $\sigma \in \mathfrak{S}_k$, the graded transpose $(\varphi_A(\sigma))^\dagger$ of $\varphi_A(\sigma)$ is equal to $\varphi_{A^\dagger}(\sigma^{-1})$. With these definitions, A^\dagger is a graded YB-bialgebra. In fact, it is well-known that $(A^\dagger, m_{A^\dagger}, \eta_{A^\dagger})$ is an R -algebra, and that $(A^\dagger, \Delta_{A^\dagger}, \varepsilon_{A^\dagger})$ is an R -coalgebra (see e.g., [28]). Taking the dual of the diagram (4.1), we easily see that A^\dagger is a YB coalgebra. Similarly, taking the dual of the diagram (4.2) for A , we see that A^\dagger is a YB algebra. It is clear that the conditions (2) and (3) in Definition 4.3 for A imply the conditions (2) and (3) for A^\dagger , respectively. By the condition (4) in Definition 4.3 for A , we have the condition (4') in Lemma 4.4 for A^\dagger .

Let $V = (V, \beta, \gamma)$ be a YB triple. We assume that V is a free R -module of finite rank and both SV and $\bigwedge V$ are free. As in Section 4, the exterior algebra $\bigwedge V =$

$T(V, -\beta)/(\text{Ker}(1-\beta))$ and the symmetric algebra $SV = T(V, -\gamma)/(\text{Im}(1-\beta))$ has a structure of YB bialgebras. By (4.16), we have:

$$(5.1) \quad \Delta_{\bigwedge V}^{(1, \dots, 1)}(w_1 \wedge \cdots \wedge w_k) = \sum_{\sigma \in \mathfrak{S}_k} (-\beta)(\sigma^{-1})(w_1 \otimes \cdots \otimes w_k)$$

$$(5.2) \quad \Delta_{SV}^{(1, \dots, 1)}(w_1 \cdots w_k) = \sum_{\sigma \in \mathfrak{S}_k} (-\gamma)(\sigma^{-1})(w_1 \otimes \cdots \otimes w_k)$$

for any $k \geq 0$ and $w_1, \dots, w_k \in V$.

We denote the YB triple $(V^*, {}^t\beta, {}^t\gamma)$ by tV , where ${}^t\beta$ (resp. ${}^t\gamma$): $V^* \otimes V^* \rightarrow V^* \otimes V^*$ is the transpose of β (see Definition 1.4). We assume that SV and $\bigwedge {}^tV$ are free. Let $A = \bigoplus_{k \geq 0} A_k$ be either $\bigwedge V$ or SV . Note that the i -fold product $m_A: V^{\otimes i} \subseteq A^{\otimes i} \rightarrow A_i$ is nothing but the projection map $p: T_i V = V^{\otimes i} \rightarrow A_i$. We denote by $A' = \bigoplus_{k \geq 0} A'_k$ the YB bialgebra $\bigwedge {}^tV$ or SV , according as $A = \bigwedge V$ or $A = SV$. For a non-negative integer i , we denote the pairing $T_i V \otimes T_i V^* \rightarrow R$; $w_1 \otimes \cdots \otimes w_i \otimes \varphi_1 \otimes \cdots \otimes \varphi_i \mapsto \varphi_1(w_1) \cdots \varphi_i(w_i)$ by $\text{Ev}^{(i)}$. We define a pairing $\text{ev}^{(i)}: A_i \otimes A'_i \rightarrow R$ by the commutativity of the diagram

$$\begin{array}{ccc} T_i V \otimes A'_i & \xrightarrow{1 \otimes \Delta_{A'}} & T_i V \otimes T_i {}^t V \\ \downarrow p \otimes 1 & \swarrow & \downarrow \text{Ev}^{(i)} \\ A_i \otimes A'_i & \xrightarrow{\text{ev}^{(i)}} & R, \end{array}$$

where $p: T_i V \rightarrow A_i$ is the natural projection. This pairing $\text{ev}^{(i)}$ is well-defined, since $\text{Ev}^{(i)}(\text{Ker}(1-\beta_l) \otimes \text{Im}(1-{}^t\beta_l)) = 0$ for $1 \leq l \leq i-1$ (the case $A = \bigwedge V$) and $\text{Ev}^{(i)}(\text{Im}(1-\beta_l) \otimes \text{Im}(1-{}^t\gamma_l)) = 0$ for $1 \leq l \leq i-1$ (the case $A = SV$).

LEMMA 5.1. *In the notation above, the diagram*

$$\begin{array}{ccc} A_i \otimes T_i {}^t V & \xrightarrow{\Delta_A \otimes 1} & T_i V \otimes T_i {}^t V \\ \downarrow 1 \otimes p' & \swarrow & \downarrow \text{Ev}^{(i)} \\ A_i \otimes A'_i & \xrightarrow{\text{ev}^{(i)}} & R \end{array}$$

is commutative, where $p': T_i {}^t V \rightarrow A'_i$ is the natural projection.

PROOF. First, we consider the case $A = \bigwedge V$. It suffices to show that $\text{ev}^{(i)} \circ (p \otimes p') = \text{Ev}^{(i)} \circ ((\Delta_A^{(1, \dots, 1)} \circ p) \otimes 1)$, since $p: T_i V \rightarrow A_i$ is surjective. By (5.1), we have

$$\begin{aligned} \text{Ev}^{(i)} \circ ((\Delta_A^{(1, \dots, 1)} \circ p) \otimes 1) &= \text{Ev}^{(i)} \circ \left(\left(\sum_{\sigma \in \mathfrak{S}_i} (-\beta)(\sigma^{-1}) \right) \otimes 1 \right) \\ &= \text{Ev}^{(i)} \circ \left(1 \otimes \left(\sum_{\sigma \in \mathfrak{S}_i} (-{}^t\beta)(\sigma) \right) \right) = \text{Ev}^{(i)} \circ (1 \otimes (\Delta_{A'}^{(1, \dots, 1)} \circ p')). \end{aligned}$$

Since $\text{Ev}^{(i)} \circ (1 \otimes \Delta_{A'}^{(1, \dots, 1)}) = \text{ev}^{(i)} \circ (p \otimes 1)$ by the definition of $\text{ev}^{(i)}$, we have completed

the proof in this case. The case $A = SV$ is proved similarly by means of (5.2). \square

We denote by Φ_i^A the natural map from A_i to $(A'_i)^*$ induced by the pairing $\text{ev}^{(i)}: A_i \otimes A'_i \rightarrow R$, and denote by Φ^A the sum of Φ_i^A 's.

PROPOSITION 5.2. *In the notation above, $\Phi^A: A \rightarrow (A')^\dagger$ is a homomorphism of YB bialgebra. The map Φ^A is a surjection if and only if $\Delta_{A'}^{(1^k)}$ is a split injection for any $k \geq 0$, where (1^k) is the sequence $(1, \dots, 1)$ (k times 1).*

PROOF. By the definition of $\text{ev}^{(i)}$, $\Phi = \Phi^A$ is a homomorphism of algebras. By Lemma 5.1, Φ is a homomorphism of coalgebras. We set $\alpha = -\beta$ when $A = \bigwedge V$, and $\alpha = -\gamma$ when $A = SV$. We show that $\varphi_{(A')^\dagger} \circ (\Phi \otimes \Phi) = (\Phi \otimes \Phi) \circ \varphi_A$. Since $p \otimes p: T_i V \otimes T_j V \rightarrow A_i \otimes A_j$ is surjective for any $i, j \geq 0$ and

$$\Delta_{(A')^\dagger}^{(1^j)} \otimes \Delta_{(A')^\dagger}^{(1^i)} = (p')^* \otimes (p')^*: (A'_j)^* \otimes (A'_i)^* \rightarrow (T_j V)^* \otimes (T_i V)^* \simeq T_j V \otimes T_i V$$

is injective for any $i, j \geq 0$, it suffices to show

$$(5.3) \quad ((p')^* \otimes (p')^*) \circ \varphi_{(A')^\dagger} \circ (\Phi_i \otimes \Phi_j) \circ (p \otimes p) = ((p')^* \otimes (p')^*) \circ (\Phi_j \otimes \Phi_i) \circ \varphi_A \circ (p \otimes p)$$

on $T_i V \otimes T_j V$ for any $i, j \geq 0$. Since $p: T_i V \rightarrow A_i$ and $p: T_j V \rightarrow A_j$ are nothing but the iterated multiplication, $(\Phi_i \otimes \Phi_j) \circ (p \otimes p)$ agrees with the composite map

$$T_i V \otimes T_j V \simeq (T_i V)^* \otimes (T_j V)^* \xrightarrow{\Delta_{A'}^{*(1^j)} \otimes \Delta_{A'}^{*(1^i)}} (A'_i)^* \otimes (A'_j)^*,$$

thanks to the fact that Φ is a homomorphism of algebras. We identify $(T_j V)^* \otimes (T_i V)^*$ with $(T_j V \otimes T_i V)^*$, and $T_i V \otimes T_j V$ with $(T_i V \otimes T_j V)^*$ in an obvious manner. Then, $(A'_i)^* \otimes (A'_j)^*$, $(\Delta_{A'}^{(1^i)})^* \otimes (\Delta_{A'}^{(1^j)})^*$, $(p')^* \otimes (p')^*$, and the restriction of $\varphi_{(A')^\dagger}$ to $(A'_i)^* \otimes (A'_j)^*$ are identified with $(A'_i \otimes A'_j)^*$, $(\Delta_{A'}^{(1^i)} \otimes \Delta_{A'}^{(1^j)})^*$, $(p' \otimes p')^*$, and $((\varphi_A)_{|A_i \otimes A_j})^*$, respectively. So the left hand side of (5.3) is identified with the dual of the composite map $(\Delta_{A'}^{(1^i)} \otimes \Delta_{A'}^{(1^j)}) \circ \varphi_{A'} \circ (p' \otimes p')$. By (4.18), it holds that $\varphi_{A'} \circ (p' \otimes p') = (p' \otimes p') \circ T(\alpha)$. Since $(p' \otimes p') \circ (\Delta_{A'}^{(1^i)} \otimes \Delta_{A'}^{(1^j)})$ is the action of $\sum_{\sigma \in \mathfrak{S}_{(i,j)}} (\alpha)(\sigma)$ by (5.1) and (5.2), the left hand side of (5.3) agrees with the map $((\sum_{\sigma \in \mathfrak{S}_{(i,j)}} (\alpha)(\sigma)) \circ (\alpha)(\chi_{ji}))^*$. On the other hand, the right hand side of (5.3) is $((\sum_{\sigma \in \mathfrak{S}_{(i,j)}} \alpha(\sigma)) \circ \alpha(\chi_{ij}))^*$ by a similar argument. For any element $\sigma = \sigma_1 \times \sigma_2 \in \mathfrak{S}_{ij}$, it holds that $((\sigma_1 \times \sigma_2) \chi_{ji})^{-1} = (\sigma_2^{-1} \times \sigma_1^{-1}) \chi_{ij}$, and these expressions are reduced. Hence, we have

$$\left(\left(\sum_{\sigma \in \mathfrak{S}_{(i,j)}} (\alpha)(\sigma) \right) \circ (\alpha)(\chi_{ij}) \right)^* = \left(\sum_{\sigma \in \mathfrak{S}_{(i,j)}} \alpha((\sigma \cdot \chi_{ij})^{-1})^* \right)^*.$$

So (5.3) is proved, and the proof of the first assertion is complete.

Since Φ is a homomorphism of algebras, we have $(\Delta_{A'}^{(1^k)})^* = \Phi_k \circ p$ for any $k \geq 0$, where $p: T_k V \rightarrow A_k$ is the projection, or the k -fold product. Hence, $(\Delta_{A'}^{(1^k)})^*$ is split surjective if and only if Φ_k is surjective, since p is surjective and $(A'_k)^*$ is free. So the second assertion is now clear. \square

LEMMA 5.3. *Let (V, β, γ) be Jimbo's YB triple of type $A_{N-1}^{(1)}$ (see Example (1) in Section 4) with the standard basis u_1, \dots, u_N . Then for any $k \geq 0$ and any sequence (i_1, \dots, i_k) with $1 \leq i_1 < \dots < i_k \leq N$, we have*

$$\Delta_{\bigwedge V}^{(1, \dots, 1)}(u_{i_1} \wedge \dots \wedge u_{i_k}) = \sum_{\sigma \in \mathfrak{S}_k} (-q)^{\ell(\sigma)} u_{i_{\sigma 1}} \otimes \dots \otimes u_{i_{\sigma k}}.$$

In particular, $\Delta_{\bigwedge V}: \bigwedge V \rightarrow T_k V$ is a split injection for any $k \geq 0$.

PROOF. To prove the first assertion, it suffices to show that

$$(5.4) \quad \beta(\sigma^{-1})(u_{i_1} \otimes \dots \otimes u_{i_k}) = q^{\ell(\sigma)} u_{i_{\sigma 1}} \otimes \dots \otimes u_{i_{\sigma k}}$$

for $\sigma \in \mathfrak{S}_k$, which is proved easily by an induction on $\ell(\sigma)$. If we define $\pi_k: T_k V \rightarrow \bigwedge_k V$ by

$$\pi_k(u_{i_1} \otimes \dots \otimes u_{i_k}) = \begin{cases} u_{i_1} \wedge \dots \wedge u_{i_k} & (i_1 < \dots < i_k) \\ 0 & (\text{otherwise}) \end{cases}$$

for $1 \leq i_1, \dots, i_k \leq N$, then $\pi_k \circ \Delta_{\bigwedge V}^{(1, \dots, 1)} = \text{id}_{\bigwedge_k V}$. Hence, the second assertion is now clear. \square

Let a be a non-negative integer and $t \in R$. We define $[a]_t = 1 + t + \dots + t^{a-1}$ if $a > 0$, and $[0]_t = 0$. We also define $[a]_t^! = \prod_{i=1}^a [i]_t$ if $a > 0$, and $[0]_t^! = 1$.

LEMMA 5.4. *Let (V, β, γ) and k be as in Lemma 5.3. For any sequence $\alpha = (\alpha_1, \dots, \alpha_N)$ of non-negative integers with $\sum_i \alpha_i = k$, we have*

$$\Delta_{SV}^{(1, \dots, 1)}(u_1^{\otimes \alpha_1} \cdots u_N^{\otimes \alpha_N}) = \prod_{i=1}^N [\alpha_i]_{q^{-2}}^! \cdot \sum_{\sigma \in \mathfrak{S}_{\alpha}} q^{-\ell(\sigma)} u_{v_{\sigma^{-1} 1}} \otimes \dots \otimes u_{v_{\sigma^{-1} k}},$$

where the number v_i is the i -th element in the sequence $(1^{\alpha_1}, \dots, N^{\alpha_N})$. In particular, $\Delta_{SV}: S_k V \rightarrow T_k V$ is a split injection if and only if $[a]_{q^{-2}}^!$ is invertible in R for any $1 \leq a \leq k$.

PROOF. It suffices to show that $\sum_{\sigma \in \mathfrak{S}_k} (-\gamma)(\sigma)(u_1^{\otimes \alpha_1} \otimes \dots \otimes u_N^{\otimes \alpha_N})$ equals the right hand side of the formula. Since $\mathfrak{S}_k = \mathfrak{S}^{\alpha} \cdot \mathfrak{S}_{\alpha}$, and since the expression $\sigma = \sigma' \sigma''$ ($\sigma' \in \mathfrak{S}^{\alpha}$, $\sigma'' \in \mathfrak{S}_{\alpha}$) is reduced for any $\sigma \in \mathfrak{S}_k$, we have

$$\sum_{\sigma \in \mathfrak{S}_k} (-\gamma)(\sigma) = \left(\sum_{\sigma' \in \mathfrak{S}^{\alpha}} (-\gamma)(\sigma') \right) \cdot \left(\sum_{\sigma'' \in \mathfrak{S}_{\alpha}} (-\gamma)(\sigma'') \right).$$

For any $a \geq 0$ and $t \in R$, it holds $\sum_{\sigma \in \mathfrak{S}_a} t^{\ell(\sigma)} = [a]_t^!$. Hence, it is easy to see that

$$\sum_{\sigma'' \in \mathfrak{S}_{\alpha}} (-\gamma)(\sigma'')(u_1^{\otimes \alpha_1} \otimes \dots \otimes u_N^{\otimes \alpha_N}) = \prod_{i=1}^N [\alpha_i]_{q^{-2}}^! \cdot u_1^{\otimes \alpha_1} \otimes \dots \otimes u_N^{\otimes \alpha_N}.$$

So it suffices to show that

$$(-\gamma)(\sigma') u_1^{\otimes \alpha_1} \otimes \dots \otimes u_N^{\otimes \alpha_N} = (-q)^{-\ell(\sigma')} u_{v_{\sigma'^{-1} 1}} \otimes \dots \otimes u_{v_{\sigma'^{-1} N}},$$

which is proved easily by induction on $\ell(\sigma')$.

Now assume that $[a]_{q^{-2}}$ is invertible for any a such that $1 \leq a \leq k$. We define $\pi_k: T_k V \rightarrow S_k V$ by

$$\pi_k(u_{i_1} \otimes \cdots \otimes u_{i_k}) = \begin{cases} \left(\prod_{i=1}^N [a_i]_{q^{-2}}! \right)^{-1} \cdot u_{i_1} \cdots u_{i_k} & (i_1 \leq \cdots \leq i_k) \\ 0 & (\text{otherwise}), \end{cases}$$

for $1 \leq i_1, \dots, i_k \leq N$, where $a_i = \#\{j \mid i_j = i\}$ for $1 \leq i \leq N$. It is clear that $\pi_k \circ \Delta_{SV}^{(1, \dots, 1)} = \text{id}$ by the first assertion, and $\Delta_{SV}^{(1, \dots, 1)}$ is a split injection. The converse is obvious from the first assertion. \square

DEFINITION 5.5. The graded dual $(S'V)^\dagger$ of $S'V$ is denoted by DV and is called the *divided power algebra* of V . The degree i component of DV is denoted by $D_i V$ and is called the *i -th divided power* of V .

By definition, $D_0 V = R$, and $D_1 V = V$. Moreover, the restriction of φ_{DV} to $D_1 V \otimes D_1 V = V \otimes V$ is equal to β_V , since ${}^t(\beta_V) = \beta_V$. By Proposition 5.2 and Lemma 5.3, we have $\bigwedge V \simeq (\bigwedge V)^\dagger$ for Jimbo's YB triple of type $A_{N-1}^{(1)}$. By Proposition 5.2 and Lemma 5.4, we have $SV \simeq DV$ for Jimbo's YB triple when $[n]_{q^2}$ is a unit for any $n \geq 1$, but not in general.

PROPOSITION 5.6. Let $V = (V, \beta, \gamma)$ be a YB triple with finitely generated V and free $S'V$. Then the iterated coproduct $\Delta_{DV}^{(1^k)}: D_k V \rightarrow {}^t(T_k {}^t V) = T_k V$ is a split injection. The image of this map is $(T_k V)^{\mathfrak{B}_k}$.

PROOF. The first assertion is clear, since the coproduct of DV is the dual of the multiplication of $S'V$. By the definition of the symmetric algebra, the sequence

$$\bigoplus_{1 \leq i \leq k-1} T_k {}^t V \xrightarrow{\sum_i (1 - \beta_i)} T_k {}^t V \xrightarrow{p} S_k {}^t V \longrightarrow 0$$

is exact, where p is the projection. Taking the dual, we see that the sequence

$$\bigoplus_{1 \leq i \leq k-1} T_k V \xleftarrow{\sum_i (1 - \beta_i)} T_k V \xleftarrow{\Delta_{DV}^{(1^k)}} D_k V \longleftarrow 0$$

is also exact. This shows that $D_k V$ is the intersection of $\text{Ker}(1 - \beta_i)$ for $1 \leq i \leq k-1$. Since \mathfrak{B}_k is generated by the b_i 's, the second assertion is now clear. \square

EXAMPLE 5.7. Let $V = (V, \beta, \gamma)$ be Jimbo's YB triple of type $A_{N-1}^{(1)}$. If $[2]_{q^2} = 0$, then the R -submodule $\Pi_2 V$ of $S_2 V$ generated by u_1^2, \dots, u_N^2 is also an SE -comodule, and we have an exact sequence

$$0 \rightarrow \Pi_2 V \rightarrow S_2 V \rightarrow \bigwedge_2 V \rightarrow 0.$$

However $\bigwedge_2 V$ is not a quotient of $D_2 V$ (see Section 7). Hence, we have $S_2 V \neq D_2 V$

as SE -comodules. The coproduct $S_2V \rightarrow V \otimes V$ vanishes on Π_2V , so the coalgebra structure of SV is different from that of DV .

Let (W, β_W, γ_W) be another YB triple such that SW , $\bigwedge W$ and S^tW are free, and W is finite. Consider a map $f \in \mathcal{YB}_R((V, \beta_V), (W, \beta_W))$. Then, $Tf: (TV, T(\pm\beta_V)) \rightarrow (TW, T(\pm\beta_W))$ is a homomorphism of YB algebras. Since $T_2Tf: T_2(TV, T(\pm\beta_V)) \rightarrow T_2(TW, T(\pm\beta_W))$ is a homomorphism of algebras, it follows that Tf is also a homomorphisms of coalgebras, hence is a homomorphism of YB bialgebras. It is easy to see that Tf induces $\bigwedge f: (\bigwedge V, \phi_{\bigwedge V}) \rightarrow (\bigwedge W, \phi_{\bigwedge W})$, $Sf: (SV, \psi_{SV}) \rightarrow (SW, \psi_{SW})$ and $Df: (DV, \psi_{DV}) \rightarrow (DW, \psi_{DW})$. The maps $\bigwedge f$, Sf and Df are homomorphisms of YB bialgebras, algebras and coalgebras, respectively. These maps preserve the grading, so we obtain $\bigwedge_i f$, $S_i f$ and $D_i f$ for $i \geq 0$. If f is also an element of $\mathcal{YB}_R((V, \gamma_V), (W, \gamma_W))$, then Sf and Df are homomorphisms of YB bialgebras.

Let M be a finite free R -module. Then $M^* \otimes M$ and $M \otimes M^* = (M^*)^* \otimes M^*$ are coalgebras (see Section 3). It is easy to see that the trivial twisting $\tau_{M, M^*}: M \otimes M^* \rightarrow M^* \otimes M$ is an anti-homomorphism of coalgebras. Consider an R -coalgebra A and a coaction $\omega_M: M \rightarrow M \otimes A$. Then, the composite map $c_{M \otimes A} \circ \tau_{M, M^*}: M \otimes M^* \rightarrow A$ is an anti-homomorphism of coalgebras, and yields a left comodule structure of M^* . The left A -comodule M^* is called the R -dual of the A -comodule M . For another right A -comodule N with finite R -module structure, the identification $(M \otimes N)^* \simeq M^* \otimes N^*$ is an isomorphism of left A -comodules. Similarly, the R -duals of left A -comodules are defined, and we have $M^{***} \simeq M$.

LEMMA 5.8. *Let $(V, \beta_V) \in \text{ob } \mathcal{YB}_R$. Then $S\tau_{V, V^*}: S(V \times V^\vee) \rightarrow S(V^\vee \times V)$ is an isomorphism of algebras, and is an anti-isomorphism of coalgebras.*

PROOF. Since $\tau_{V, V^*} \in \mathcal{YB}_R(V \times V^\vee, V^\vee \times V)$ is an isomorphism, $S\tau_{V, V^*}$ is an isomorphism of algebras. Since $\tau_{V, V^*}: V \otimes V^* \rightarrow V^* \otimes V$ is an anti-homomorphism of coalgebras, and $S(V \times V^\vee)$ is generated by elements of degree one, $S\tau_{V, V^*}$ is also an anti-homomorphism of coalgebras. \square

Note that $E = V \times V^\vee$ is the internal End of V . Hence, TV^* is a right SE -comodule, and $\beta^\vee, \beta' = (\beta^\vee)^{-1}$ are homomorphisms of SE -comodules. So $T'V$, $\bigwedge' V$ and $S'V$ are SE -equivariant graded YB bialgebras. With the anti-coalgebra algebra homomorphism $S\tau_{V, V^*}$ these algebras are viewed as left SE^\vee -equivariant graded YB bialgebras. Taking the dual comodules of each degree, the graded duals $(T'V)^\dagger$, $(\bigwedge' V)^\dagger$ and $DV = (S'V)^\dagger$ are (right) SE -equivariant graded YB bialgebras. It is easy to see that the natural identification $(T'V)^\dagger \simeq TV$ of graded YB bialgebras is also an SE^\vee -comodule isomorphism. Hence, $D_k V$ is a subcomodule of $V^{\otimes k}$ via the injection $\Delta: D_k V \rightarrow V^{\otimes k}$. Since $\Phi^{\bigwedge V}: \bigwedge V \rightarrow (\bigwedge' V)^\dagger$ and $\Phi^{SV}: SV \rightarrow DV$ are induced by $TV \simeq (T'V)^\dagger$ (see Proposition 5.2), these maps are SE^\vee -equivariant.

REMARK 5.9. Since β^{-1} is an SE^\vee -comodule map, TV^{-1} , $\bigwedge V^{-1}$ and SV^{-1} have

SE -comodule structures. Clearly, the identification $TV \simeq TV^{-1}$ is an isomorphism of both algebras and SE^\vee -comodules. Let us compare $\bigwedge V$ and $\bigwedge V^{-1}$. They are the quotient algebras of $TV = TV^{-1}$ defined by the same ideal $(\text{Ker}(1 - \beta)) = (\text{Ker}(1 - \beta^{-1}))$. Hence, $(\bigwedge V, \varphi_{\bigwedge V})$ and $(\bigwedge V^{-1}, \varphi_{\bigwedge V^{-1}}^-)$ are isomorphic as SE^\vee -equivariant YB algebras. For each $k \geq 1$, the image of the coproduct maps $\text{Im } \Delta_{\bigwedge V}^{(1^k)} = \text{Im}(\sum_{\sigma \in S_k} (-\beta)(\sigma))$ and $\text{Im } \Delta_{\bigwedge V^{-1}}^{(1^k)} = \text{Im}(\sum_{\sigma \in S_k} (-\beta^{-1})(\sigma))$ agree. It follows that $(\bigwedge V, \varphi_{\bigwedge V})$ and $(\bigwedge V^{-1}, \varphi_{\bigwedge V^{-1}}^-)$ are also isomorphic as SE^\vee -equivariant YB coalgebras. But they are *not* isomorphic as YB bialgebras in general. To illustrate this, consider Jimbo's YB triple (V, β, γ) of type $A_{N-1}^{(1)}$, and compare the maps

$$\bigwedge_2 V \xrightarrow{\Delta} V \otimes V \xrightarrow{m} \bigwedge_2 V \quad \text{and} \quad \bigwedge_2 V^{-1} \xrightarrow{\Delta} V^{-1} \otimes V^{-1} \xrightarrow{m} \bigwedge_2 V^{-1}.$$

The first is the multiplication by $1 + q^2$, while the second is multiplication by $1 + q^{-2}$. Similar facts on symmetric algebras and divided power algebra should be noted.

PROPOSITION 5.10. *Let $V = (V, \beta_V, \gamma_V)$ be Jimbo's YB triple of type $A_{N-1}^{(1)}$ and $E = (E, \beta_E, \gamma_E)$ be as in §4 Example (2). Then we have:*

1. *For any $k \geq 0$, the comultiplication $\Delta_{\bigwedge E}^{(1^k)}$ is a split injection. Hence, we have $\bigwedge E \simeq (\bigwedge' E)^\dagger$ as graded YB-bialgebras.*

2. *$\Delta_{SE}^{(1^k)}: S_k E \rightarrow T_k E$ is injective if $R = \mathbb{Z}[Q, Q^{-1}]$, where Q is an indeterminate.*

PROOF. 1. We define a map $\rho_k: E^{\otimes k} \rightarrow \bigwedge_k E$ by

$$\rho_k(x_{i_1 j_1} \otimes \cdots \otimes x_{i_k j_k}) = \begin{cases} x_{i_1 j_1} \wedge \cdots \wedge x_{i_k j_k} & (\forall l \ i_l < i_{l+1}, \text{ or } j_l > j_{l+1} \text{ and } i_l = i_{l+1}) \\ 0 & (\text{otherwise}). \end{cases}$$

Then $\rho_k \circ \Delta_{\bigwedge E}^{(1^k)} = \text{id}_{\bigwedge_k E}$. Hence, $\Delta_{\bigwedge E}^{(1^k)}$ splits. The second assertion is a consequence of Proposition 5.2.

2. We may localize, and assume that $R = \mathbb{Q}(Q)$. We define a map $\pi_k: E^{\otimes k} \rightarrow S_k E$ by

$$\pi_k(x_{i_1 j_1} \otimes \cdots \otimes x_{i_k j_k}) = \begin{cases} (i=1 \prod_{l'=1}^N [u_{l, l'}]_{Q^{-2}})^{-1} \cdot x_{i_1 j_1} \cdots x_{i_k j_k} & (\text{if } \forall h \ i_h < i_{h+1}, \text{ or } i_h = i_{h+1} \text{ and } j_h < j_{h+1}) \\ 0 & (\text{otherwise}), \end{cases}$$

where $u_{l, l'} = \#\{h \mid (i_h, j_h) = (l, l')\}$. Then $\pi_k \circ \Delta_{SE}^{(1^k)} = \text{id}_{S_k E}$. Hence $\Delta_{SE}^{(1^k)}$ is an injection. \square

Assume that R is a field. Since $(S_2 V)^* = (\text{Coker}(1 - \beta))^*$ is identified with $D_2 V = \text{Ker}(1 - \beta)$, the Koszul complex of the quadratic algebra SV defined in [26] is nothing but $SV \otimes (\bigwedge' V)^\dagger$. The boundary map is the composite map

$$\partial: SV \otimes (\bigwedge' V)^\dagger \xrightarrow{1 \otimes \Delta} SV \otimes V \otimes (\bigwedge' V)^\dagger \xrightarrow{m \otimes 1} SV \otimes (\bigwedge' V)^\dagger,$$

where Δ is the abbreviation for the map

$$(\bigwedge^t V)^\dagger \xrightarrow{\Delta} (\bigwedge^t V)^\dagger \otimes (\bigwedge^t V)^\dagger \xrightarrow{\text{proj} \otimes 1} (\bigwedge_1^t V)^* \otimes (\bigwedge^t V)^\dagger = V \otimes (\bigwedge^t V)^\dagger,$$

and m is the product in SV . If SV is a (homogeneous) Koszul algebra (see [28]), then $SV \otimes (\bigwedge^t V)^\dagger$ is a resolution of the left SV -module $R = SV/(V)$. If $V = (V, \beta, \gamma)$ is Jimbo's YB triple of type $A_{N-1}^{(1)}$, then $SV \otimes (\bigwedge^t V)^\dagger$ is rewritten as $SV \otimes \bigwedge V$, and the boundary map is described explicitly as

$$(5.5) \quad \partial(a \otimes u_{i_1} \wedge \cdots \wedge u_{i_k}) = \sum_{j=1}^k (-q)^{j-1} a u_{i_j} \otimes u_{i_1} \wedge \cdots \wedge \check{u}_{i_j} \wedge \cdots \wedge u_{i_k} \quad (1 \leq i_1 < \cdots < i_k \leq N).$$

In this case, SV is a Koszul algebra [32, Theorem 5.3], and the Koszul complex $SV \otimes \bigwedge V$ is acyclic.

REMARK 5.11. Let V be a YB pair. By the discussion in the proof of Proposition 5.6, the dual of the projection $T_k V \rightarrow S_k V$ is naturally identified with the inclusion $(T_k V^\vee)^{\mathfrak{B}_k} \rightarrow T_k V^\vee$.

6. Schur modules and Weyl modules. Throughout this section, (V, β_V) is Jimbo's YB pair of type $A_{N-1}^{(1)}$ (cf. (1.2)) over the base ring R determined by a unit $q \in R$.

For (infinite) sequences of integers $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$, we define the sum $\lambda + \mu$ to be the sequence $(\lambda_1 + \mu_1, \lambda_2 + \mu_2, \dots)$, and we define $k \cdot \lambda$ to be the sequence $(k \cdot \lambda_1, k \cdot \lambda_2, \dots)$ for $k \in \mathbb{Z}$. We also denote $\text{supp}(\lambda) = \{i \in N \mid \lambda_i \neq 0\}$. We will denote by Ω^+ the set $\{\lambda : \text{a sequence of integers } |^{\forall i \in N, \lambda_i \geq 0} \text{ and } \#(\text{supp}(\lambda)) < \infty\}$. For $\lambda \in \Omega^+$, we define the *length* of λ by $\ell(\lambda) = \max \text{supp}(\lambda)$, and the *degree* of λ by $|\lambda| = \sum_{i \in N} \lambda_i$. When $\lambda = (0, 0, \dots)$, we define $\ell(\lambda) = 0$. We set $\Omega^- = \{\lambda \in \Omega^+ \mid \forall i \in N \lambda_i \geq \lambda_{i+1}\}$ and call an element of Ω^- a *partition*. For any $\lambda \in \Omega^+$, its *transpose* $\tilde{\lambda}$ is defined to be the partition which satisfies $\tilde{\lambda}_i = \#\{j \in N \mid \lambda_j \geq i\}$ for any $i \in N$. It is easy to see that $(\tilde{\lambda})^\sim$ is the partition obtained by rearrangement of the sequence λ in non-increasing order.

We introduce a lexicographic order $<$ into Ω^+ . Namely, for two elements λ and $\mu \in \Omega^+$, we say that $\lambda > \mu$ when there exists some j such that $\lambda_j > \mu_j$ and $\lambda_i = \mu_i$ for any $i < j$. With this ordering, Ω^+ is a totally ordered set.

We will denote the partition $(0, 0, \dots)$ of degree zero by 0.

For $\lambda, \mu \in \Omega^+$, we say that $\lambda \rhd \mu$ if and only if $\lambda_i \geq \mu_i$ for each $i \in N$. It is clear that \rhd is an order and $\lambda \rhd \mu$ implies $\lambda \geq \mu$. It holds that $\lambda \rhd \mu$ if and only if $\tilde{\lambda} \rhd \tilde{\mu}$. For $\lambda, \mu \in \Omega^-$, we say that λ/μ is a *skew partition*, if $\lambda \rhd \mu$.

DEFINITION 6.1. The *diagram* (or *shape*) Δ_λ of an element $\lambda \in \Omega^+$ is the set $\{(i, j) \in N^2 \mid j \leq \lambda_i\}$. We use the convention as in the case of matrices, namely, the row index i increases as one goes downward, and the column index j increases from left to right (see Figure). The *skew shape* of a pair $\lambda, \mu \in \Omega^+$ with $\lambda \rhd \mu$ is $\Delta_{\lambda/\mu} := \Delta_\lambda \setminus \Delta_\mu$.




FIGURE.

Clearly, $\Delta_{\lambda/0} = \Delta_\lambda$. So we will always adapt the terminology for skew-shape (or skew partition) to the terminology for shape (or partition) by letting $\mu=0$.

DEFINITION 6.2. Let X be a set and let $\lambda, \mu \in \Omega^+$. We define $\text{Tab}_{\lambda/\mu}(X)$ to be the set $\text{Map}(\Delta_{\lambda/\mu}, X)$. An element of $\text{Tab}_{\lambda/\mu}(X)$ is called a *tableau of shape λ/μ with values in the set X* . For $S \in \text{Tab}_{\lambda/\mu}(X)$, we say that S is a *Young tableau*, if X is the set $[1, |\lambda| - |\mu|]$ and S is a bijection.

Pictorially, a tableau $S \in \text{Tab}_{\lambda/\mu} X$ is expressed as an array of elements in X in which $S(i, j)$ occupies the intersection of the i -th row and the j -th column. For example, if $\lambda=(4, 3, 3)$, $\mu=(2, 1)$ and

$$S = \begin{matrix} & 6 & 3 \\ & 1 & 4 \\ & 5 & 2 & 7 \end{matrix},$$

then S is a Young tableau, and $S(3, 2)=2$.

DEFINITION 6.3. Let Y be a totally ordered set. A tableau $S \in \text{Tab}_{\lambda/\mu}(Y)$ is said to be *row-standard* if the rows of S are strictly increasing, i.e., if we have $S(i, \mu_i+1) < S(i, \mu_i+2) < \dots < S(i, \lambda_i)$ for any i . The tableau S is said to be *column-standard* if the columns of S are non-decreasing, i.e., if for all $1 \leq j \leq \lambda_1$ we have $S(i, j) \leq S(i+1, j)$ whenever (i, j) and $(i+1, j)$ are both in $\Delta_{\lambda/\mu}$. S is said to be *standard* if it is both row- and column-standard. We denote by $\text{Row}_{\lambda/\mu} Y$ (resp. $\text{St}_{\lambda/\mu} Y$) the subset of $\text{Tab}_{\lambda/\mu} Y$ consisting of all row-standard (resp. standard) tableaux.

DEFINITION 6.4. A tableau $T \in \text{Tab}_{\lambda/\mu}(Y)$ is said to be *co-row-standard* if the rows of T are non-decreasing, and *co-column-standard* if the columns of T are strictly increasing. T is said to be *co-standard* if it is co-row- and co-column-standard. We denote by $\text{Corow}_{\lambda/\mu} Y$ (resp. $\text{Cost}_{\lambda/\mu} Y$) the subset of $\text{Tab}_{\lambda/\mu} Y$ consisting of all co-row-standard (resp. co-standard) tableaux.

Let $\lambda, \mu \in \Omega^+$ with $\lambda \supset \mu$. We define an element $\chi_{\lambda/\mu}$ in $\mathfrak{S}_{|\lambda|-|\mu|}$ as follows. First we introduce a lexicographic order into the set $\Delta_{\lambda/\mu}$. We define $(i, j) <_r (i', j')$ if $i < i'$, or $i = i'$ and $j < j'$. Since $<$ is a total ordering, we have a unique order-preserving bijection

$\alpha_{r, \lambda/\mu} : [1, |\lambda| - |\mu|] \rightarrow (\mathcal{A}_{\lambda/\mu}, <_r)$. Secondly, we introduce another lexicographic order into $\mathcal{A}_{\lambda/\mu}$. We define $(i, j) <_c (i', j')$ if $j < j'$, or $j = j'$ and $i < i'$. We also have a unique order-preserving bijection $\alpha_{c, \lambda/\mu} : [1, |\lambda| - |\mu|] \rightarrow (\mathcal{A}_{\lambda/\mu}, <_c)$. We define $\chi_{\lambda/\mu} := \alpha_{c, \lambda/\mu}^{-1} \circ \alpha_{r, \lambda/\mu}$. Here we show an example.

EXAMPLE 6.5. Let $\lambda = (5, 4, 2)$ and $\mu = (1, 1)$. The permutation $\chi_{\lambda/\mu}$ is described as follows:

$$\alpha_{r, \lambda/\mu}^{-1} = \begin{matrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 \\ 8 & 9 \end{matrix} \xrightarrow{\chi_{\lambda/\mu}} \begin{matrix} 2 & 5 & 7 & 9 \\ 3 & 6 & 8 \\ 1 & 4 \end{matrix} = \alpha_{c, \lambda/\mu}^{-1}.$$

The permutation w_i in Proposition 4.2 is nothing but $\chi_{(i,i)}$

DEFINITION 6.6. For any $\lambda, \mu \in \Omega^+$, we define

$$\begin{aligned}\bigwedge_{\lambda/\mu} V &= \bigwedge_{\lambda_1 - \mu_1} V \otimes \cdots \otimes \bigwedge_{\lambda_s - \mu_s} V, \\ S_{\lambda/\mu} V &= S_{\lambda_1 - \mu_1} V \otimes \cdots \otimes S_{\lambda_s - \mu_s} V, \\ D_{\lambda/\mu} V &= D_{\lambda_1 - \mu_1} V \otimes \cdots \otimes D_{\lambda_s - \mu_s} V, \\ T_{\lambda/\mu} V &= T_{\lambda_1 - \mu_1} V \otimes \cdots \otimes T_{\lambda_s - \mu_s} V,\end{aligned}$$

when $\lambda \supset \mu$, where s is a sufficiently large integer. If $\lambda \not\supset \mu$, they are all 0 by definition.

Note that $T_{\lambda/\mu}V$ is naturally isomorphic to $T_{|\lambda|-|\mu|}V$.

DEFINITION 6.7. Let λ/μ be a skew partition with $\ell(\lambda/\mu)=s$ and $\lambda_1=t$. We denote the composite map

from $\bigwedge_{\lambda/\mu} V$ to $S_{\tilde{\lambda}/\tilde{\mu}} V$ by $d_{\lambda/\mu}(V)$, and call it the *Schur map* (cf. [4]). The image $\text{Im } d_{\lambda/\mu}(V)$ of the Schur map is denoted by $L_{\lambda/\mu} V$, and is called the *Schur module of V with respect to the skew partition λ/μ* .

By definition, $d_{\lambda/\mu}(V)$ and $L_{\lambda/\mu}V$ are the usual Schur map and the Schur module, respectively, when $q=1$. Note that $(q^{-2}\beta)(\chi_{\lambda/\mu})$ in the definition can be replaced by $\beta(\chi_{\lambda/\mu})$, $(-\beta)(\chi_{\lambda/\mu})$, or $(-q^{-2}\beta)(\chi_{\lambda/\mu})$ for the purpose of defining $L_{\lambda/\mu}V$, since q is a unit

(using $\beta(\chi_{\lambda/\mu})$ might be the simplest; however we use $(q^{-2}\beta)(\chi_{\lambda/\mu})$, since it is more consistent with the notation in [14] (see §8)). It is easy to see that $d_{\lambda/\mu}V$ is an SE^\vee -homomorphism, and $L_{\lambda/\mu}V$ is an SE^\vee -comodule. We denote $d_{\lambda/0}(V)$ and $L_{\lambda/0}V$ by $d_\lambda(V)$ and $L_\lambda V$, respectively.

We set $Y(\sigma) = \{u_{\sigma_1} < \dots < u_{\sigma_N}\}$, the ordered basis of V for $\sigma \in \mathfrak{S}_N$, and $X = \{u_1, \dots, u_N\}$ its underlying set.

DEFINITION 6.8. Let λ/μ be a skew partition with $\ell = \ell(\lambda/\mu)$ and $\tilde{\ell} = \ell(\tilde{\lambda}/\tilde{\mu})$, and $S \in \text{Tab}_{\lambda/\mu}X$. We define

$$\begin{aligned}\xi_S &= S(1, \mu_1 + 1) \wedge \dots \wedge S(1, \lambda_1) \otimes \dots \otimes S(\ell, \mu_\ell + 1) \wedge \dots \wedge S(\ell, \lambda_\ell) \in \bigwedge_{\lambda/\mu} V, \\ \eta_S &= S(\tilde{\mu}_1 + 1, 1) \cdots S(\tilde{\lambda}_1, 1) \otimes \dots \otimes S(\tilde{\mu}_{\tilde{\ell}} + 1, \tilde{\ell}) \cdots S(\tilde{\lambda}_{\tilde{\ell}}, \tilde{\ell}) \in S_{\tilde{\lambda}/\tilde{\mu}} V, \\ \zeta_S &= S(1, \mu_1 + 1) \otimes \dots \otimes S(1, \lambda_1) \otimes \dots \otimes S(\ell, \mu_\ell + 1) \otimes \dots \otimes S(\ell, \lambda_\ell) \in T_{\lambda/\mu} V, \\ \tilde{\zeta}_S &= S(\tilde{\mu}_1 + 1, 1) \otimes \dots \otimes S(\tilde{\lambda}_1, 1) \otimes \dots \otimes S(\tilde{\mu}_{\tilde{\ell}} + 1, \tilde{\ell}) \otimes \dots \otimes S(\tilde{\lambda}_{\tilde{\ell}}, \tilde{\ell}) \in T_{\tilde{\lambda}/\tilde{\mu}} V.\end{aligned}$$

For $\sigma \in \mathfrak{S}_N$, we set

$$\begin{aligned}\bigwedge_{\lambda/\mu} Y(\sigma) &= \{\xi_S \mid S \in \text{Row}_{\lambda/\mu} Y(\sigma)\}, \\ L_{\lambda/\mu} Y(\sigma) &= \{d_{\lambda/\mu}(V)(\xi_S) \mid S \in \text{St}_{\lambda/\mu} Y(\sigma)\}.\end{aligned}$$

It is obvious that $\bigwedge_{\lambda/\mu} Y(\sigma)$ and the set $\{\eta_S \mid S \in \text{Tab}_{\lambda/\mu} Y(\sigma), S \text{ is column-standard}\}$ are bases of $\bigwedge_{\lambda/\mu} V$ and $S_{\tilde{\lambda}/\tilde{\mu}} V$, respectively, for any $\sigma \in \mathfrak{S}_N$. It follows that $\tilde{\zeta}_S = \tau(\chi_{\lambda/\mu}) \cdot \zeta_S$, where τ is the trivial twisting. In the rest of this section, we sometimes tacitly identify $T_{\lambda/\mu} V$ with $T_{|\lambda|-|\mu|} V$. In such an occasion, we also identify $S \in \text{Tab}_{\lambda/\mu} X$ with $S \circ \alpha_{r, \lambda/\mu} \in \text{Map}([1, N], X)$. For an arbitrary totally ordered set Y and skew partition λ/μ , we introduce two (pseudo-) orders into $\text{Tab}_{\lambda/\mu} Y$, as follows:

DEFINITION 6.9. For a skew partition λ/μ , $I \subset N$, totally ordered set Y , $Y_0 \subset Y$ and $S \in \text{Tab}_{\lambda/\mu} Y$, we define

$$n_I(S, Y_0) = \#\{(i, j) \in \Delta_{\lambda/\mu} \mid i \in I, S(i, j) \in Y_0\}.$$

DEFINITION 6.10. Let S and S' be elements in the set $\text{Tab}_{\lambda/\mu} Y$. We say that $S \leq S'$ when it holds

$$n_{\{1, \dots, i\}}(S, (-\infty, y]) \geq n_{\{1, \dots, i\}}(S', (-\infty, y])$$

for any $i \in [1, \ell(\lambda/\mu)]$ and $y \in Y$, where $(-\infty, y]$ denotes the subset $\{z \in Y \mid z \leq y\}$ of Y . We say that $S \leq_{\text{lex}} S'$ when $S = S'$ or $S \circ \alpha_{r, \lambda/\mu}(j) < S' \circ \alpha_{r, \lambda/\mu}(j)$, where j is the minimum of the numbers i such that $S \circ \alpha_{r, \lambda/\mu}(i) \neq S' \circ \alpha_{r, \lambda/\mu}(i)$. We say that $S < S'$ (resp. $S <_{\text{lex}} S'$) when $S \leq S'$ and $S \not\leq S'$ (resp. $S \leq_{\text{lex}} S'$ and $S \not\leq_{\text{lex}} S'$).

Note that \leq is a pseudo-order for $\text{Tab}_{\lambda/\mu} Y$, and \leq_{lex} is a total order of $\text{Tab}_{\lambda/\mu} Y$. The restriction of \leq to $\text{Row}_{\lambda/\mu} Y$ is an order, and $S, S' \in \text{Row}_{\lambda/\mu} Y$, $S \leq S'$ imply $S \leq_{\text{lex}} S'$.

EXAMPLE 6.11. Let $\lambda=(3, 2)$, $\mu=(1)$, $Y=\{1 < 2 < 3\}$, and S_1 , S_2 and S_3 be as follows:

$$S_1 = \begin{smallmatrix} 1 & 3 \\ 2 & 2 \end{smallmatrix}, \quad S_2 = \begin{smallmatrix} 2 & 2 \\ 1 & 3 \end{smallmatrix}, \quad S_3 = \begin{smallmatrix} 3 & 1 \\ 2 & 2 \end{smallmatrix}.$$

Then we have $S_1 \leq S_3 \leq S_1$ since S_3 is obtained by row-permutation on the entries of S_1 . We have $S_2 \not\leq S_1 \not\leq S_2 \not\leq S_3 \not\leq S_2$, since $n_{\{1\}}(S_1, \{1\})=n_{\{1\}}(S_3, \{1\})=1>0=n_{\{1\}}(S_2, \{1\})$ and $n_{\{1\}}(S_1, \{1, 2\})=n_{\{1\}}(S_3, \{1, 2\})=1<2=n_{\{1\}}(S_2, \{1, 2\})$. Since $S_1(1, 2)=1 < S_2(1, 2)=2 < S_3(1, 2)=3$, we have $S_1 <_{\text{lex}} S_2 <_{\text{lex}} S_3$.

Returning to the situation $X=\{u_1, \dots, u_N\}$, we denote $S \leq_{\sigma} S'$ (resp. $S \leq_{\text{lex}, \sigma} S'$) for $S, S' \in \text{Tab}_{\lambda/\mu} X$, when $S \leq S'$ (resp. $S \leq_{\text{lex}} S'$) with S, S' viewed as elements of $\text{Tab}_{\lambda/\mu} Y(\sigma)$. We will omit the subscript σ if $\sigma=\text{id}$. Hence, $S \leq S'$ (resp. $S \leq_{\text{lex}} S'$) and $S \geq_{\omega_0} S'$ (resp. $S \geq_{\text{lex}, \omega_0} S'$) are equivalent, where $\omega_0 \in \mathfrak{S}_N$ is the unique element with maximal length.

For $k \in \mathbb{N}$ and $\rho \in \mathfrak{S}_k$, we define the *decomposition* of ρ by induction on k . We let $\rho_1=(1 2 \cdots \rho^{-1} 1)$, the cyclic permutation. Since $\rho \rho_1^{-1}(1)=1$, we may assume that $\rho \rho_1^{-1} \in \mathfrak{S}_{k-1}$. Let $\rho \rho_1^{-1}=\rho_{k-1} \cdots \rho_2$ be the decomposition of $\rho \rho_1^{-1}$. We define the decomposition of ρ to be $\rho=\rho_{k-1} \cdots \rho_1$. Note that this decomposition is always a reduced expression.

LEMMA 6.12. Let λ/μ be a skew partition, and $S \in \text{Tab}_{\lambda/\mu} Y$. Then we have

$$\beta(\chi_{\lambda/\mu})\zeta_S = q^* \zeta_S + \sum_{S' > S} c_{S'} \zeta_{S'}, \quad (c_{S'} \in \mathbb{Z}[q, q^{-1}]),$$

where by q^* we mean some power of q .

PROOF. We set $k=|\lambda|-|\mu|$, and proceed by induction on k . We denote by μ' the unique partition such that $\Delta_{\lambda/\mu'}$ is obtained by removal of the box $\alpha_{c, \lambda/\mu}(1)$ from $\Delta_{\lambda/\mu}$. We denote $\chi_{\lambda/\mu}^{-1}(1)$ by d (in Example 6.5, $\alpha_{c, \lambda/\mu}(1)=(3, 1)$ and $d=8$). If the decomposition of $\chi_{\lambda/\mu}$ is $\chi_{\lambda/\mu}=\rho_{k-1} \cdots \rho_2 \rho_1$, then we have $\rho_{k-1} \cdots \rho_2=\chi_{\lambda/\mu'}$, with the natural identification of \mathfrak{S}_{k-1} with the isotropy group of 1 in \mathfrak{S}_k . Hence,

$$\tau(\chi_{\lambda/\mu}^{-1})\beta(\chi_{\lambda/\mu})=[1_V \otimes \tau(\chi_{\lambda/\mu}^{-1})\beta(\chi_{\lambda/\mu'})] \circ [\tau(\rho_1^{-1})\beta(\rho_1)],$$

where we identify $T_{\lambda/\mu} V$ with $V \otimes T_{\lambda/\mu'} V$ via the isomorphism $\tau(\rho_1)$. By the remark after Definition 6.8, it suffices to show that we have an expression

$$\tau(\chi_{\lambda/\mu}^{-1})\beta(\chi_{\lambda/\mu})\zeta_S = q^* \zeta_S + \sum_{S' > S} c_{S'} \zeta_{S'}.$$

By the induction hypothesis, it also suffices to prove that we have an expression

$$(*) \quad \tau(\rho_1^{-1})\beta(\rho_1)\zeta_S = q^* \zeta_S + \sum_{S' > S} c_{S'} \zeta_{S'}.$$

We set $\pi_i = \sigma_i \cdots \sigma_{d-1}$ for $1 \leq i \leq d-1$, where σ_j is the transposition $(j \ j+1)$. Note that $\rho_1 = \pi_1$. To prove that we have an expression $(*)$, we will show that

$$(**) \quad \beta(\pi_i)\zeta_S = \sum_I c_I^i \zeta_{S'_i(I)} \quad (c_I^i \in \mathbb{Z}[q, q^{-1}], \quad c_\emptyset^i = q^*)$$

for any $i \in [1, d-1]$, where the sum is taken over all subsequences $I = (i_1, i_2, \dots, i_h)$ of $(i, i+1, \dots, d-1)$ such that $S(i_1) < S(i_2) < \cdots < S(i_h) < S(d)$. The empty subsequence is denoted by \emptyset . For such a subsequence I , let $J_i = (j_1, \dots, j_{d-h-i})$ be the complementary subsequence of I in $(i, i+1, \dots, d-1)$. The tableau $S'_i(I)$ is defined by $S'_i(I) = \sigma_{j_1} \cdots \sigma_{j_{d-h-i}} \cdot S (= S \circ \sigma_{j_{d-h-i}} \cdots \sigma_{j_1})$. We shall prove $(**)$ by descending induction on i . For each subsequence $I = (i_1, \dots, i_h)$ of $(i+1, \dots, d-1)$, we have

$$\beta(\sigma_i) \cdot \zeta_{S'_{i+1}(I)} = \begin{cases} \zeta_{S'_i(I)} & (\text{if } S(i) = S(i_1)), \\ q\zeta_{S'_i(I)} & (\text{if } S(i) < S(i_1)), \\ q\zeta_{S'_i(I)} + (1-q^2)\zeta_{S'_{i+1}(I')} & (\text{if } S(i) > S(i_1)), \end{cases}$$

by (1.3), where we assume that $i_1 = d$ when I is empty, and I' is the sequence (i, i_1, \dots, i_h) . So $(**)$ holds by the induction hypothesis. Moreover, we have

$$\pi_i^{-1} \cdot S'_i(I) = \pi_{i+1}^{-1} \sigma_i \pi_{i+1} \pi_{i+1}^{-1} S'_i(I) = \pi_{i+1}^{-1} S'_{i+1}(I)$$

and

$$\pi_i^{-1} \cdot S'_i(I') = \pi_{i+1}^{-1} \sigma_i \pi_{i+1} \pi_{i+1}^{-1} S'_i(I') = (i \ d) \pi_{i+1}^{-1} S'_{i+1}(I).$$

Since $(\pi_{i+1}^{-1} \cdot S'_{i+1}(I))(i) = S(i) > S(i_1) = (\pi_{i+1}^{-1} S'_{i+1}(I))(d)$ and $\alpha_{r, \lambda/\mu}(i)$ lies in an earlier row than the row in which $\alpha_{r, \lambda/\mu}(d)$ lies, it is easy to see that $\pi_i^{-1} S'_i(I') < \pi_{i+1}^{-1} S'_{i+1}(I)$ (cf. [4, Lemma II. 2. 14]).

Using the descending induction again, we have $\rho_1^{-1} \cdot S'_1(\emptyset) = S > \rho_1^{-1} S'_1(I)$ for any non-empty I . By $(**)$, we have an expression

$$\tau(\rho_1^{-1})\beta(\rho_1)\zeta_S = \sum_I c_I^1 \zeta_{\rho_1^{-1} \cdot S'_1(I)} \quad (c_I^1 \in \mathbb{Z}[q, q^{-1}], \quad c_\emptyset^1 = q^*)$$

which is of the form $(*)$. □

Let $k \geq 0$ and (i_1, \dots, i_k) be a sequence with $1 \leq i_1 < \cdots < i_k \leq N$. Then by Lemma 5.3, we have

$$\begin{aligned} \Delta^{(1^k)} \nabla (u_{i_k} \wedge \cdots \wedge u_{i_1}) &= (-q)^{k(k-1)/2} \Delta^{(1^k)} (u_{i_1} \wedge \cdots \wedge u_{i_k}) \\ &= (-q)^{k(k-1)/2} \sum_{\sigma \in \mathfrak{S}_k} (-q)^{\ell(\sigma)} u_{i_{\sigma 1}} \otimes \cdots \otimes u_{i_{\sigma k}} \\ &= (-q)^{k(k-1)/2} \sum_{\sigma \in \mathfrak{S}_k} (-q)^{-\ell(\sigma)} u_{i_{\sigma k}} \otimes \cdots \otimes u_{i_{\sigma 1}}. \end{aligned}$$

For a skew partition λ/μ and $S \in \text{Row}_{\lambda/\mu}(\omega_0)$, the image of ζ_S by the coproduct

$\Delta: \bigwedge_{\lambda/\mu} V \rightarrow T_{\lambda/\mu} V$ is of the form $q^* \sum_{\sigma \in \mathfrak{S}_{\lambda-\mu}} (-q)^{-\ell(\sigma)} \zeta_{\sigma S}$, where $\omega_0 \in \mathfrak{S}_N$ is the permutation of maximal length. Note that $\sigma S \geq S$, $\sigma S \leq S$, and $\sigma S \neq S$ for any $\sigma \in \mathfrak{S}_{\lambda-\mu} \setminus \{\text{id}\}$.

PROPOSITION 6.13. *Let λ/μ be a skew partition. Then $L_{\lambda/\mu} Y(\omega_0) = \{d_{\lambda/\mu}(V)(\xi_S) \mid S \in \text{St}_{\lambda/\mu} Y(\omega_0)\}$ is linearly independent.*

PROOF. We denote by $\bigwedge'_{\lambda/\mu} V$ (resp. $S'_{\lambda/\mu} V$) the submodule of $\bigwedge_{\lambda/\mu} V$ (resp. $S_{\lambda/\mu} V$) generated by $\{\xi_S \mid S \in \text{St}_{\lambda/\mu} Y(\omega_0)\}$ (resp. $\{\eta_S \mid S \in \text{St}_{\lambda/\mu} Y(\omega_0)\}$). We denote the composite map

$$\bigwedge'_{\lambda/\mu} V \hookrightarrow \bigwedge_{\lambda/\mu} V \xrightarrow{d_{\lambda/\mu}(V)} S_{\tilde{\lambda}/\tilde{\mu}} V \xrightarrow{p} S'_{\tilde{\lambda}/\tilde{\mu}} V$$

by H , where p is the projection via the standard basis $\{\eta_S \mid S \in \text{Tab}_{\lambda/\mu} Y(\omega_0)\}$, S is column-standard} of $S_{\tilde{\lambda}/\tilde{\mu}} V$. Since both $\bigwedge'_{\lambda/\mu} V$ and $S'_{\lambda/\mu} V$ have bases indexed by the same totally order set $(\text{St}_{\lambda/\mu} Y(\omega_0), \leq_{\text{lex}, \omega_0})$, we can express H by means of a square matrix with respect to these bases. It suffices to show that this matrix is upper triangular and each diagonal entry is of the form q^* . By Lemma 6.12 and the preceding observation, we have an expression

$$d_{\lambda/\mu}(V)(\xi_S) = q^* \eta_S + \sum_{S' \geq S, S' \neq S} c_{S'} \eta_{S'}$$

for $S \in \text{St}_{\lambda/\mu} Y(\omega_0)$. For each S' , we have $\eta_{S'} = q^* \eta_{(S')^\circ}$, where $(S')^\circ$ is the column-standardization of S' with respect to the order of $Y(\omega_0)$. Hence we have

$$H(\xi_S) = q^* \eta_S + \sum_{S' \geq S, S' \neq S, (S')^\circ \in \text{St}_{\lambda/\mu} Y(\omega_0)} c'_{(S')} \cdot Y_{(S')^\circ}.$$

If S' appears in the sum and $S' \leq S$, then S' is not row-standard. Hence $S' \neq (S')^\circ$, since $(S')^\circ$ is row-standard. So we have $(S')^\circ <_\omega S' \leq_\omega S$. If $S' > S$, then we have $(S')^\circ <_\omega S$, too.

□

For $i \geq 1$, we denote the sequence $(0, \dots, 0, \overset{i}{1}, 0, \dots)$ by ε_i , and we define $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$. For a skew partition λ/μ , we define the subset $S_{\square}(\lambda/\mu)$ of Ω^+ by:

$$S_{\square}(\lambda/\mu) = \{v \in \Omega^+ \mid \exists t \in N, \exists k \in N_0 \text{ } k < \lambda_{t+1} - \mu_t \text{ and } v = \gamma + (\gamma_{t+1} - k) \cdot \alpha_t\},$$

where $\gamma = \lambda - \mu$.

DEFINITION 6.14. Let λ/μ be a skew partition, and $v \in S_{\square}(\lambda/\mu)$. We have $v = \lambda - \mu + r \cdot \alpha_t$ for some $r, t \geq 0$. Then we define the *box map* $\square_{\lambda/\mu}^v: \bigwedge_v V \rightarrow \bigwedge_{\lambda/\mu} V$ to be the composite map

$$\begin{aligned} \bigwedge_v V &= \bigwedge_{\gamma_1} V \otimes \cdots \otimes \bigwedge_{\gamma_{t-1}} V \otimes \bigwedge_{\gamma_t+r} V \otimes \bigwedge_{\gamma_{t+1}-r} V \otimes \bigwedge_{\gamma_{t+2}} V \otimes \cdots \otimes \bigwedge_{\gamma_s} V \\ &\quad \downarrow 1 \otimes \cdots \otimes 1 \otimes \square \otimes 1 \otimes \cdots \otimes 1 \\ \bigwedge_{\lambda/\mu} V &= \bigwedge_{\gamma_1} V \otimes \cdots \otimes \bigwedge_{\gamma_{t-1}} V \otimes \bigwedge_{\gamma_t} V \otimes \bigwedge_{\gamma_{t+1}} V \otimes \bigwedge_{\gamma_{t+2}} V \otimes \cdots \otimes \bigwedge_{\gamma_s} V, \end{aligned}$$

where $\gamma = \lambda - \mu$ and \square is the composite map

$$\bigwedge_{\gamma_t+r} V \otimes \bigwedge_{\gamma_{t+1}-r} V \xrightarrow{\Delta \otimes 1} \bigwedge_{\gamma_t} V \otimes \bigwedge_r V \otimes \bigwedge_{\gamma_{t+1}-r} V \xrightarrow{1 \otimes m} \bigwedge_{\gamma_t} V \otimes \bigwedge_{\gamma_{t+1}} V.$$

The sum of these box maps $\sum_{v \in S_{\square}(\lambda/\mu)} \bigwedge_v V \rightarrow \bigwedge_{\lambda/\mu} V$ is denoted by $\square_{\lambda/\mu}$.

LEMMA 6.15 (cf. [4, Lemma II.2.9]). *Let $\lambda = (\lambda_1, \lambda_2)$ and $\mu = (\mu_1, \mu_2)$ be partitions with $\lambda \supset \mu$. We set $\gamma = \lambda - \mu$. Let a and b be non-negative integers with $a + b < \lambda_2 - \mu_1$. Then the image of the composite map*

$$\begin{aligned} \bar{\square}_{(a, b)} : \bigwedge_a V \otimes \bigwedge_{|\gamma| - a - b} V \otimes \bigwedge_b V &\xrightarrow{1 \otimes \Delta \otimes 1} \bigwedge_a V \otimes \bigwedge_{\gamma_1 - a} V \otimes \bigwedge_{\gamma_2 - b} V \otimes \bigwedge_b V \\ &\xrightarrow{m \otimes m} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2} V = \bigwedge_{\lambda/\mu} V \end{aligned}$$

is contained in $\text{Im } \square_{\lambda/\mu}$.

PROOF. We use induction on a . If $a = 0$, then we have $(\gamma_1 + \gamma_2 - b, b) \in S_{\square}(\lambda/\mu)$, and $\bar{\square}_{(a, b)} = \square_{\lambda/\mu}^{(\gamma_1 + \gamma_2 - b, b)}$ by assumption. So the assertion is clear. We consider the case $a > 0$. For each $i \in [0, a]$, we denote by $\Delta(i)$ the composite map

$$\begin{aligned} \bigwedge_a V \otimes \bigwedge_{|\gamma| - a - b} V &\xrightarrow{\Delta \otimes \Delta} \bigwedge_i V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_1 - i} V \otimes \bigwedge_{\gamma_2 - a + i - b} V \\ &\xrightarrow{1 \otimes \varphi_{\Delta} \otimes 1} \bigwedge_i V \otimes \bigwedge_{\gamma_1 - i} V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_2 - a + i - b} V. \end{aligned}$$

Then, we have a commutative diagram

$$(*) \quad \begin{array}{ccc} \bigwedge_a V \otimes \bigwedge_{|\gamma| - a - b} V & \xrightarrow{m} & \bigwedge_{|\gamma| - b} V \\ \downarrow \sum_{i=0}^a \Delta(i) \otimes \Delta & & \downarrow \Delta \\ \sum_{i=0}^a \bigwedge_i V \otimes \bigwedge_{\gamma_1 - i} V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_2 - a + i - b} V & \xrightarrow{m \otimes m} & \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2 - b} V, \end{array}$$

since m is a homomorphism of coalgebras. Note that $\bar{\square}_{(a, b)}$ is nothing but the composite map

$$\begin{aligned} \bigwedge_a V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_b V &\xrightarrow{\Delta(a) \otimes 1} \bigwedge_a V \otimes \bigwedge_{\gamma_1-a} V \otimes \bigwedge_{\gamma_2-b} V \otimes \bigwedge_b V \\ &\xrightarrow{m \otimes m} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2} V = \bigwedge_{\lambda/\mu} V. \end{aligned}$$

On the other hand, the image of the composite map

$$\begin{aligned} \bigwedge_a V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_b V &\xrightarrow{m \otimes 1} \bigwedge_{|\gamma|-b} V \otimes \bigwedge_b V \xrightarrow{\Delta \otimes 1} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2-b} V \otimes \bigwedge_b V \\ &\xrightarrow{1 \otimes m} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2} V = \bigwedge_{\lambda/\mu} V \end{aligned}$$

is clearly contained in $\text{Im } \square_{\lambda/\mu}$. Hence, it suffices to show that the image of the composite map

$$\begin{aligned} f_i: \bigwedge_a V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_b V &\xrightarrow{\Delta(i) \otimes 1} \bigwedge_i V \otimes \bigwedge_{\gamma_1-i} V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_2-a+i-b} V \otimes \bigwedge_b V \\ &\xrightarrow{m \otimes m \otimes 1} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2-b} V \otimes \bigwedge_b V \xrightarrow{1 \otimes m} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2} V = \bigwedge_{\lambda/\mu} V \end{aligned}$$

is contained in $\text{Im } \square_{\lambda/\mu}$ for $i \in [0, a-1]$, because $\bar{\square}_{(a,b)} = f_a$ and $\text{Im } (\sum_{i=0}^a f_i) \subset \text{Im } \square_{\lambda/\mu}$ by the commutativity of (*). Consider the diagram

$$\begin{array}{ccc} \bigwedge_{a-i} V \otimes \bigwedge_{|\gamma|-a-b} V & \xrightarrow{\varphi_{\wedge V}} & \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_{a-i} V \\ \downarrow 1 \otimes \Delta & & \downarrow \Delta \otimes 1 \\ \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_1-i} V \otimes \bigwedge_{\gamma_2-a+i-b} V & & \\ \downarrow \varphi_{\wedge V} \otimes 1 & & \downarrow \\ \bigwedge_{\gamma_1-i} V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{\gamma_2-a+i-b} V & \xrightarrow{1 \otimes \varphi_{\wedge V}} & \bigwedge_{\gamma_1-i} V \otimes \bigwedge_{\gamma_2-a+i-b} V \otimes \bigwedge_{a-i} V \\ \searrow m \otimes 1 & & \swarrow 1 \otimes m \\ & \bigwedge_{\gamma_1-i} V \otimes \bigwedge_{\gamma_2-b} V. & \end{array}$$

The upper square commutes, since $(\bigwedge V, \varphi_{\wedge V})$ is a YB coalgebra. Since $\varphi_{\wedge V}$ equals $(-q^2)^* \psi_{\wedge V}$ on $\bigwedge_{a-i} V \otimes \bigwedge_{\gamma_2-a+i-b} V$, the lower triangle commutes up to unit thanks to Proposition 4.11 (1). Together with the associativity of the algebra $\bigwedge V$, these commutativity of the diagram above yields that f_i is a unit multiple of the composite map

$$\begin{aligned} \bigwedge_a V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_b V &\xrightarrow{\Delta \otimes 1} \bigwedge_i V \otimes \bigwedge_{a-i} V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_b V \\ &\xrightarrow{1 \otimes \varphi_{\wedge V} \otimes 1} \bigwedge_i V \otimes \bigwedge_{|\gamma|-a-b} V \otimes \bigwedge_{a-i} V \otimes \bigwedge_b V \end{aligned}$$

$$\xrightarrow{1 \otimes m} \bigwedge_i V \otimes \bigwedge_{|\gamma| - i - (a+b-i)} V \otimes \bigwedge_{a+b-i} V \xrightarrow{\square_{(i,a+b-i)}} \bigwedge_{\gamma_1} V \otimes \bigwedge_{\gamma_2} V = \bigwedge_{\lambda/\mu} V,$$

whose image is contained in $\text{Im } \square_{\lambda/\mu}$ by the induction hypothesis. \square

We emphasize that the proof above is a slight modification of the proof of [4, Lemma II.2.9]. Similarly, we can prove the following lemma using the idea of the proof of [4, Proposition II.2.8] and the properties of YB bialgebras $\bigwedge V$ and SV from §4.

LEMMA 6.16. *Let $\lambda/\mu = (\lambda_1, \lambda_2)/(\mu_1, \mu_2)$ be a skew shape consisting of two rows, and $v \in S_{\square}(\lambda/\mu)$. Then, the composite map*

$$\bigwedge_v V \xrightarrow{\square_{\lambda/\mu}^v} \bigwedge_{\lambda/\mu} V \xrightarrow{d_{\lambda/\mu}} S_{\tilde{\lambda}/\tilde{\mu}} V$$

is zero.

The proof is left to the reader.

LEMMA 6.17. *Let λ/μ be a skew partition, and $v \in S_{\square}(\lambda/\mu)$. Then the composite map*

$$\bigwedge_v V \xrightarrow{\square_{\lambda/\mu}^v} \bigwedge_{\lambda/\mu} V \xrightarrow{d_{\lambda/\mu}(V)} S_{\tilde{\lambda}/\tilde{\mu}} V$$

is zero.

PROOF. We set $v = \lambda - \mu + t \cdot \alpha_k$, $\lambda^+ = (\lambda_1, \dots, \lambda_{t-1})$, $\mu^+ = (\mu_1, \dots, \mu_{t-1})$, $\lambda^0 = (\lambda_t, \lambda_{t+1})$, $\mu^0 = (\mu_t, \mu_{t+1})$, $\lambda^- = (\lambda_{t+2}, \dots, \lambda_s)$, and $\mu^- = (\mu_{t+2}, \dots, \mu_s)$, where $s = \ell(\lambda/\mu)$. We have a reduced expression $\chi_{\lambda/\mu} = \rho \cdot \chi_{\lambda^+/mu^+} \cdot \chi_{\lambda^0/\mu^0} \cdot \chi_{\lambda^-/\mu^-}$ with unique $\rho \in \mathfrak{S}^{(|\lambda^+|-|\mu^+|, |\lambda^0|-|\mu^0|, |\lambda^-|-|\mu^-|)}$. Since $(SV, \varphi_{\bigwedge V})$ is a YB algebra, this Schur map $d_{\lambda/\mu}(V)$ coincides with the composite map

$$\begin{aligned} \bigwedge_{\lambda/\mu} V &= \bigwedge_{\lambda^+/mu^+} V \otimes \bigwedge_{\lambda^0/\mu^0} V \otimes \bigwedge_{\lambda^-/\mu^-} V \\ &\xrightarrow{d_{\lambda^+/mu^+} \otimes d_{\lambda^0/\mu^0} \otimes d_{\lambda^-/\mu^-}} S_{\tilde{\lambda}^+/mu^+} V \otimes S_{\tilde{\lambda}^0/\mu^0} V \otimes S_{\tilde{\lambda}^-/\mu^-} V \xrightarrow{m} S_{\tilde{\lambda}/\tilde{\mu}} V, \end{aligned}$$

where m is the restriction of the multiplication $T_{\lambda_1} SV \otimes T_{\lambda_1} SV \otimes T_{\lambda_1} SV \rightarrow T_{\lambda_1} SV$ defined in Proposition 4.2 (which agrees with $m_{SV}^{\otimes \lambda_1} \circ \varphi_{SV}(w)$, where m_{SV} is the multiplication $SV \otimes SV \otimes SV \rightarrow SV$ and $w \in \mathfrak{S}_{3\lambda_1}$ is given by $w(i\lambda_1 + j) = 3j + i - 2$ for $i = 0, 1, 2$ and $j = 1, \dots, \lambda_1$). Observe that the restriction of $\varphi_{SV}(w)$ to $S_{\tilde{\lambda}^+/mu^+} V \otimes S_{\tilde{\lambda}^0/\mu^0} V \otimes S_{\tilde{\lambda}^-/\mu^-} V$ is induced by $(q^{-2}\beta)(w) : T_{\tilde{\lambda}^+/mu^+} V \otimes T_{\tilde{\lambda}^0/\mu^0} V \otimes T_{\tilde{\lambda}^-/\mu^-} V \rightarrow T_{\tilde{\lambda}/\mu} V$. On the other hand, the box map $\square_{\lambda/\mu}^v$ is nothing but $1 \otimes \square_{\lambda^0/\mu^0}^{v^0} \otimes 1$, where $v^0 = (v_t, v_{t+1})$. Therefore, the lemma is clear by Lemma 6.16. \square

LEMMA 6.18. *Let λ/μ be a skew partition, $\sigma \in \mathfrak{S}_N$, and $S \in \text{Row}_{\lambda/\mu} Y(\sigma) \setminus \text{St}_{\lambda/\mu} Y(\sigma)$. Then, there exist $S'_1, \dots, S'_r \in \text{Row}_{\lambda/\mu} Y(\sigma)$ ($r \in N$) with $S'_i <_{\sigma} S$ for each $i \in [1, r]$ such that*

$$\xi_S - \sum_i c_i \xi_{S'_i} \in \text{Im } \square_{\lambda/\mu} \text{ for some } c_i \in \mathbb{Z}[q, q^{-1}].$$

PROOF. The proof is quite similar to that of [4, II. Lemma 2.15] (the case $q=1$), so we only point out some non-trivial differences. The proof in [4] depends on the fact corresponding to (2.3) and (2.4) for the case $q=1$, while we can use these without much problem, since $\pm q^*$ is a unit by our assumption. It also depends on the following fact: (*) Let $s, t \in N$ and i_1, \dots, i_{s+t} be distinct integers in $[1, N]$. Then, the image of $u_{i_1} \wedge \cdots \wedge u_{i_{s+t}}$ by the map $\Delta_{\bigwedge V}^{(s,t)}: \bigwedge_{s+t} V \rightarrow \bigwedge_s V \otimes \bigwedge_t V$ equals

$$\sum_{\pi \in \mathfrak{S}^{(s,t)}} \pm q^* u_{i_{\pi 1}} \wedge \cdots \wedge u_{i_{\pi s}} \otimes u_{i_{\pi(s+1)}} \wedge \cdots \wedge u_{i_{\pi t}}.$$

(for the case $q=1$).

Let us prove (*). If $i_1 < \cdots < i_{s+t}$ (this case corresponds to the case $\sigma = \text{id}$), then

$$\Delta_{\bigwedge V}^{(s,t)}(u_{i_1} \wedge \cdots \wedge u_{i_{s+t}}) = \sum_{\pi \in \mathfrak{S}^{(s,t)}} (-q)^{\ell(\pi)} u_{i_{\pi 1}} \wedge \cdots \wedge u_{i_{\pi s}} \otimes u_{i_{\pi(s+1)}} \wedge \cdots \wedge u_{i_{\pi t}}$$

by Proposition 4.8 and (5.4). The general case follows from (2.4). Since the coefficients in the sum are units, we can use (*). We have generalized [4, Lemma II.2.9], which is used in the proof, to Lemma 6.15, and we can use this generalized version. We leave the complete modifications to the interested reader. \square

THEOREM 6.19. *Let λ/μ be a skew partition. Then the Schur module $L_{\lambda/\mu} V$ is an R-free module with $L_{\lambda/\mu} Y(\sigma) = \{d_{\lambda/\mu}(V)(\xi_S) \mid S \in \text{St}_{\lambda/\mu} Y(\sigma)\}$ its free basis for any $\sigma \in \mathfrak{S}_N$. The sequence of SE^\vee -comodules*

$$(*) \quad \sum_{v \in S_{\square}(\lambda/\mu)} \bigwedge_v V \xrightarrow{\square_{\lambda/\mu}} \bigwedge_{\lambda/\mu} V \xrightarrow{d_{\lambda/\mu}(V)} L_{\lambda/\mu} V \longrightarrow 0$$

is exact. The Schur module $L_{\lambda/\mu} V$ is a direct summand of $S_{\lambda/\bar{\mu}} V$ via the natural inclusion, as an R-module.

PROOF. Clearly, $d_{\lambda/\mu}(V): \bigwedge_{\lambda/\mu} V \rightarrow L_{\lambda/\mu} V$ is surjective. With any $a = \sum_{S \in \text{Row}_{\lambda/\mu} Y(\sigma)} c_S \xi_S \in \bigwedge_{\lambda/\mu} V$, we associate a non-negative integer $N(a) = \sum_{c_S \neq 0} 2^{\text{ht}(S)}$, where $\text{ht}(S)$ is the height of S with respect to the order $\leq_{\text{lex}, \sigma}$ of $\text{Row}_{\lambda/\mu} Y(\sigma)$. We claim that a can be expressed in the form $a = a_0 + \sum_{S \in \text{St}_{\lambda/\mu} Y(\sigma)} c'_S \xi_S$ with $a_0 \in \text{Im } \square_{\lambda/\mu}$. We prove this by induction on $N(a)$. If each S appearing in the sum with $c_S \neq 0$ is standard, then there is nothing to prove. So we may assume that some S is not standard. For such an S , we have an expression $\xi_S = \sum_i c''_i \cdot S'_i + a''_0$ for some $a''_0 \in \text{Im } \square_{\lambda/\mu}$ and $S'_i <_{\sigma} S$ by Lemma 6.18. Then we have $a = c_S a''_0 + (a - c_S a''_0)$, and $N(a - c_S a''_0) < N(a)$. So the claim is now clear by the induction hypothesis. Since we have $d_{\lambda/\mu}(a) = \sum_S c'_S d_{\lambda/\mu}(\xi_S)$ by Lemma 6.17, $L_{\lambda/\mu} Y(\sigma)$ generates $L_{\lambda/\mu} V$ for any σ . By Proposition 6.13, $L_{\lambda/\mu} Y(\omega_0)$ is a free basis of $L_{\lambda/\mu} V$. Since $\#L_{\lambda/\mu} Y(\sigma) = \#L_{\lambda/\mu} Y(\omega_0)$, the first assertion is now clear. If $a \in \text{Ker } d_{\lambda/\mu}$, then we have $0 = d_{\lambda/\mu}(a) = \sum_S c'_S \cdot d_{\lambda/\mu}(\xi_S)$. Since $d_{\lambda/\mu}(\xi_S)$ are linearly independent, we have $c'_S = 0$ for all S . Hence, $a = a_0 \in \text{Im } \square_{\lambda/\mu}$, and the second assertion is proved. As we have seen in the

proof of Proposition 6.13, $L_{\lambda/\mu}V + S''_{\tilde{\lambda}/\tilde{\mu}}V = S_{\tilde{\lambda}/\tilde{\mu}}V$, where $S''_{\tilde{\lambda}/\tilde{\mu}}V$ is the R -span of the set $\{\eta_S \mid S \in \text{Tab}_{\lambda/\mu} Y(\text{id})$, S is column-standard but is not row standard}. This sum must be a direct sum by the first assertion, so the last assertion is now clear. \square

COROLLARY 6.20. *Let $f: R \rightarrow R'$ be a homomorphism of commutative rings. Then we have an isomorphism of $S(E')^\vee$ -comodules $L_\lambda(R' \otimes_R V) \simeq R' \otimes_R L_\lambda V$, where $E' = R' \otimes_R E$, and $R' \otimes_R V$ is Jimbo's YB-pair obtained by base change.*

PROOF. Note that $\bigwedge_{\lambda/\mu}$ and $\sum_v \bigwedge_v$ are compatible with base changes, and that $\square_{\lambda/\mu}$ is defined over $\mathbb{Z}[Q, Q^{-1}]$, where Q is an indeterminate. By Theorem 6.19, we have an isomorphism $\text{Coker } \square_{\lambda/\mu} \simeq L_{\lambda/\mu}V$. Since cokernels are compatible with base change, the assertion is clear. \square

We shall now define the Weyl modules of V .

DEFINITION 6.21. Let λ/μ be a skew partition. Let $d'_{\lambda/\mu}(V)$ be the composite map

$$\begin{aligned} D_{\lambda/\mu}V &\xrightarrow{A \otimes \cdots \otimes A} T_{\lambda/\mu}V = T_{|\lambda|-|\mu|}V \xrightarrow{(q^{-2}\beta)(\chi_{\lambda/\mu})} T_{|\lambda|-|\mu|}V \\ &= T_{\tilde{\lambda}/\tilde{\mu}}V \xrightarrow{p \otimes \cdots \otimes p} \bigwedge_{\tilde{\lambda}/\tilde{\mu}}V. \end{aligned}$$

It is called the *Weyl map*. We denote the image $\text{Im } d'_{\lambda/\mu}(V)$ of $d'_{\lambda/\mu}(V)$ by $K_{\lambda/\mu}V$ and called it the *Weyl module of V with respect to the skew partition λ/μ* .

With the natural identifications $\bigwedge_{\tilde{\lambda}/\tilde{\mu}}V = (\bigwedge_{\tilde{\lambda}/\tilde{\mu}}^t V)^*$ and $D_{\lambda/\mu}V = (S_{\lambda/\mu}^t V)^*$, we can identify $d'_{\lambda/\mu}(V)$ with $(d_{\tilde{\lambda}/\tilde{\mu}}(V))^*$, since $\chi_{\tilde{\lambda}/\tilde{\mu}}^{-1} = \chi_{\lambda/\mu}$ so that $((q^{-2}\beta)(\chi_{\tilde{\lambda}/\tilde{\mu}}))^* = (q^{-2}\beta)(\chi_{\lambda/\mu})$. Since $L_{\tilde{\lambda}/\tilde{\mu}}V$ is a direct summand of $S_{\lambda/\mu}^t V$, we have a standard isomorphism $K_{\lambda/\mu}V \simeq (L_{\tilde{\lambda}/\tilde{\mu}}V)^*$. In particular, $K_{\lambda/\mu}V$ is a free R -module with the same rank as $L_{\tilde{\lambda}/\tilde{\mu}}V$.

To describe a presentation and the standard basis of $K_{\lambda/\mu}V$, we need additional notation. We define $\Omega_k^+ = \{\alpha \in \Omega^+ \mid |\alpha| = k\}$ for $k \geq 0$. We denote the dual basis of X by ' $X = \{v_1, \dots, v_N\}$ '. So we have a basis $\{v^\alpha \mid \alpha \in \Omega_k^+, \ell(\alpha) \leq N\}$ of $S_k^t V$ for $k \geq 0$, where $v^\alpha = v_1^{\alpha_1} \cdots v_N^{\alpha_N}$. The dual basis $\{u^{(\alpha)} \mid \alpha \in \Omega_k^+, \ell(\alpha) \leq N\}$ is a basis of $D_k V$, where $u^{(\alpha)}$ is the basis corresponding to v^α . For $i \in [1, N]$ and $k \geq 0$, we denote $u_i^{k-\epsilon_i}$ by $u_i^{(k)}$. Since the product of DV is the dual of the coproduct of $S^t V$, we have $u_i^k = [k]_{q-2}! u_i^{(k)}$ in $D_k V$ by Lemma 5.4 applied to $S^t V$, where u_i on the left hand side is $u_i^{(1)} \in D_1 V = V$. We call $u_i^{(k)}$ the k -th divided power of u_i .

LEMMA 6.22. *Let k, α and v be as in Lemma 5.4. Then we have $u^{(\alpha)} = u_1^{(\alpha_1)} \cdots u_N^{(\alpha_N)}$. If $\sigma \in \mathfrak{S}^\alpha$, then we have*

$$u_{v\sigma^{-1}1} \cdots u_{v\sigma^{-1}k} = q^{-\ell(\sigma)} u^\alpha = \left(\prod_{i=1}^N [\alpha_i]_{q-2}! \right) q^{-\ell(\sigma)} u^{(\alpha)}$$

in $D_k V$.

PROOF. The second assertion follows immediately by Lemma 5.4, since the product of DV is the dual of the coproduct of $S'V$. To prove the first assertion, we may assume that the base ring R is $\mathbb{Z}[q, q^{-1}]$ with q transcendental over \mathbb{Z} . By the observation above, we have

$$u^\alpha = u_1^{\alpha_1} \cdots u_N^{\alpha_N} = \left(\prod_{i=1}^N [\alpha_i]_{q^{-2}}^! \right) u_1^{(\alpha_1)} \cdots u_N^{(\alpha_N)}.$$

On the other hand, we have $u^\alpha = (\prod_i [\alpha_i]_{q^{-2}}^!) u^{(\alpha)}$ by the first assertion. Since $\prod_i [\alpha_i]_{q^{-2}}^!$ is a non-zero-divisor, we have $u^{(\alpha)} = u_1^{(\alpha_1)} \cdots u_N^{(\alpha_N)}$ as desired.

For a skew partition λ/μ and $S \in \text{Tab}_{\lambda/\mu} X$, we set

$$\xi'(S) = q^{-(\ell(\sigma^{(1)}) + \cdots + \ell(\sigma^{(s)}))} u^{(\alpha^{(1)})} \otimes \cdots \otimes u^{(\alpha^{(s)})} \in D_{\lambda/\mu} V,$$

where $s = \ell(\lambda - \mu)$, $\alpha^{(i)} = (\alpha_1^{(i)}, \dots, \alpha_N^{(i)})$ is given by $\alpha_t^{(i)} = \#\{j \mid S(i, j) = u_t\}$ for $t \in [1, N]$, and $\sigma^{(i)}$ is the unique element in $\mathfrak{S}^{\alpha^{(i)}}$ such that $S(i, \mu_i + \sigma^{(i)} j)$ is the j -th element in the sequence $(1^{\alpha_1^{(i)}}, \dots, N^{\alpha_N^{(i)}})$. It is clear that $\{\xi'(S) \mid S \in \text{Corow}_{\lambda/\mu} Y(\rho)\}$ is a free basis of $D_{\lambda/\mu} V$ for any $\rho \in \mathfrak{S}_N$.

THEOREM 6.23. *Let λ/μ be a skew partition. Then the Weyl module $K_{\lambda/\mu} V$ is an R -free module with $\{d'_{\lambda/\mu}(V)(\xi_S) \mid S \in \text{Cost}_{\lambda/\mu} Y(\rho)\}$ as its free basis for any $\rho \in \mathfrak{S}_N$. The sequence of SE^\vee -comodules*

$$\bullet \sum_{v \in S \square(\lambda/\mu)} D_v V \xrightarrow{\square'_{\lambda/\mu}} D_{\lambda/\mu} V \xrightarrow{d'_{\lambda/\mu}(V)} K_{\lambda/\mu} V \longrightarrow 0$$

is exact, where $\square'_{\lambda/\mu}$ is defined similarly to $\square_{\lambda/\mu}$. The Weyl module $K_{\lambda/\mu}$ is a direct summand of $\bigwedge_{\tilde{\lambda}/\tilde{\mu}} V$ as an R -module via the natural inclusion.

PROOF. Since the inclusion $K_{\lambda/\mu} V \subset \bigwedge_{\tilde{\lambda}/\tilde{\mu}} V$ can be identified with the dual of the Schur map $d_{\tilde{\lambda}/\tilde{\mu}}(V) : \bigwedge_{\tilde{\lambda}/\tilde{\mu}} V \rightarrow L_{\tilde{\lambda}/\tilde{\mu}} V$, the last assertion is trivial. The rest of the theorem is proved similarly to Theorem 6.19 (see also the proof of [4; Theorem II.3.16]). To translate this proof into our context, we need Lemma 6.22). We only remark that there is no need to prove the fact corresponding to Proposition 6.13, since we already know that $K_{\lambda/\mu} V$ is free of rank $\#\text{Cost}_{\lambda/\mu} Y(\rho) = \#\text{St}_{\tilde{\lambda}/\tilde{\mu}} Y(\rho)$. \square

COROLLARY 6.24. *Let λ/μ be a skew partition, and $R \rightarrow R'$ a homomorphism of commutative rings. Then we have an SE^\vee -comodule isomorphism $K_\lambda(R' \otimes V) \simeq R' \otimes K_\lambda V$, where $E' = R' \otimes E$.*

7. Universality of Weyl modules. Throughout this section, $V = (V, \beta_V, \gamma_V)$ denotes Jimbo's YB triple determined by a basis $\{u_1, \dots, u_N\}$ and $q \in R^\times$, and $E = (E, \beta, \gamma)$ is as in Example (2) in §4.

Let A be an R -algebra (resp. R -coalgebra). We denote by $A\text{-mff}$ (resp. $\text{mff-}A$, resp. $A\text{-cff}$, resp. $\text{cff-}A$) the category of left A -modules (resp. right A -modules, resp. left

A -comodules, resp. right A -comodules) which are free of finite rank as R -modules. Let C be an R -coalgebra which is free of finite rank as an R -module, and M a C -comodule. The map $C^* \otimes M \rightarrow M$ given by $a \otimes u \mapsto \sum_i \text{ev}(a \otimes u_i^{(2)})u_i^{(1)}$ is a left action of the dual algebra C^* of C on M , where $\omega(u) = \sum_i u_i^{(1)} \otimes u_i^{(2)}$. This correspondence is an equivalence between the category of right C -comodules and the category of left C^* -modules, and we have an equivalence $\text{cff-}C \cong C^*\text{-mff}$. We will sometimes identify $\text{cff-}C$ with $C^*\text{-mff}$ by this correspondence. The corresponding notion of the R -dual $(\)^* = \text{Hom}_R(-, R)$: $C^*\text{-mff} \rightarrow \text{mff-}C^*$ is the R -dual $\text{cff-}C \rightarrow C\text{-cff}$ (see §5).

Since the coalgebra SE^\vee is decomposed into the direct sum $SE^\vee = \bigoplus_{k \geq 0} S_k E^\vee$, we have a decomposition $M = \bigoplus_{k \geq 0} M_k$ for any SE^\vee -comodule M , where M_k is the image of the map

$$M \xrightarrow{\omega} M \otimes SE^\vee \xrightarrow{\text{proj}} M \otimes S_k E^\vee \xrightarrow{1 \otimes e} M.$$

Clearly, M_k is an $S_k E^\vee$ -comodule. We call M_k the *degree k component* of M . The comodule M is said to be *locally free of finite rank* when M_k is free of finite rank as an R -module for each $k \geq 0$. We denote the category of locally finite free right (resp. left) SE^\vee -comodules by \mathcal{P} (resp. \mathcal{Q}). It is clear that $\mathcal{P} \cong \prod_{k \geq 0} \text{cff-}S_k E^\vee$ and $\mathcal{Q} \cong \prod_{k \geq 0} S_k E^\vee\text{-cff}$. The categories \mathcal{P} and \mathcal{Q} are additive, and are closed under extensions and tensor products. Taking the product of the dual $(\)^* : \text{cff-}S_k E^\vee \rightarrow S_k E^\vee\text{-cff}$, we obtain the *graded dual* $(\)^\dagger := \prod_{k \geq 0} (\)^* : \mathcal{P} \rightarrow \mathcal{Q}$. The graded dual is a contravariant equivalence of categories, and preserves tensor products. Similarly, the graded dual $\mathcal{Q} \rightarrow \mathcal{P}$ is also defined, and is also denoted by $(\)^\dagger$. It holds that $(\)^\dagger \circ (\)^\dagger = \text{Id}$.

The dual algebra of $S_k E^\vee$ is $(S_k E^\vee)^* = D_k E$. Since the projection $(E^\vee)^{\otimes k} \rightarrow S_k E^\vee$ is nothing but the product, $D_k E$ is a subalgebra of $E^{\otimes k} \cong \text{End}_R V^{\otimes k}$ via the coproduct $A_{D^* E}^{(1,k)}$. By Proposition 5.6, we have $D_k E \cong (E^{\otimes k})^{\mathfrak{B}_k} \cong \text{End}_{\mathfrak{B}_k}(V^{\otimes k})$.

DEFINITION 7.1. The dual algebra $D_k E = \text{End}_{\mathfrak{B}_k} V^{\otimes k}$ of $S_k E^\vee$ is called the *Schur algebra* of degree k .

We denote the dual basis of $\{u_1, \dots, u_N\}$ by $\{v_1, \dots, v_N\}$. The R -linear map $\rho : V \rightarrow V^*$ given by $\rho(u_i) = v_i$ is an element of $\mathcal{U}\mathcal{B}_R(V, {}^t V) \cong \mathcal{U}\mathcal{B}_R(V^{-1}, V^\vee)$. Clearly, we have $\rho^* = \rho^{-1}$. Since SE^\vee and $S^t E$ are the quotients of the same bialgebra $TE^\vee = T^t E$ defined by the same biideal $(\text{Im}(1 - \beta_E^\vee)) = (\text{Im}(1 - {}^t \beta_E))$, they are isomorphic as bialgebras. The composite map

$$SE^\vee \xrightarrow{\sim} S^t E \xrightarrow{S\tau_{v^*, v}} SE^{-1} \xrightarrow{S(\rho \otimes \rho^{-1})} SE^\vee$$

is an anticoalgebra algebra isomorphism, and is denoted by tr . Since $((\rho \otimes \rho^{-1}) \circ \tau_{v^*, v})^2 = \text{id}$, we have $\text{tr}^2 = \text{id}$.

DEFINITION 7.2. The composite functor $\text{tr}_* \circ (\)^\dagger : \mathcal{P} \rightarrow \mathcal{P}$ is denoted by $(\)^\circ$, and

is called the *contravariant dual*, where $\text{tr}_\sharp: \mathcal{P} \rightarrow \mathcal{P}$ is the push-forward with respect to tr .

LEMMA 7.3. (1) *The contravariant dual $(\)^\circ: \mathcal{P} \rightarrow \mathcal{P}$ is a contravariant additive functor, and preserves tensor products.*

$$(2) ((\)^\circ)^2 = \text{id}_{\mathcal{P}}.$$

$$(3) \rho: V \rightarrow V^\circ \text{ is an isomorphism of } SE^\vee\text{-comodules. } \beta_V^\circ \text{ is the composite map}$$

$$V^\circ \otimes V^\circ \xrightarrow{\rho^{-1} \otimes \rho^{-1}} V \otimes V \xrightarrow{\beta_V} V \otimes V \xrightarrow{\rho \otimes \rho} V^\circ \otimes V^\circ.$$

PROOF. Part (1) is obvious. Part (2) follows easily from the fact $\text{tr}^2 = 1$. (3) The action of $D_1 E = \text{End } V$ on $V^* = V^\circ$ is given by $(u_i \otimes v_j) \cdot v_k = v_k (\text{tr}^*(u_i \otimes v_j)) = v_k \cdot (v_j \otimes u_i) = \delta_{jk} v_i$. Hence ρ is an isomorphism. The map β_V° is nothing but β_V by definition. Since $\rho \in \mathcal{YB}_R(V, {}^t V)$, we have completed the proof. \square

PROPOSITION 7.4. *The maps $T\rho: TV \rightarrow \text{tr}_\sharp T^t V \simeq (TV)^\circ$, $D\rho: DV \rightarrow \text{tr}_\sharp D^t V \simeq (SV)^\circ$, $S\rho: SV \rightarrow \text{tr}_\sharp S^t V \simeq (DV)^\circ$, $\wedge\rho: \bigwedge V \rightarrow \text{tr}_\sharp \bigwedge^t V \simeq (\bigwedge V)^\circ$ are isomorphisms of SE^\vee -equivariant graded YB bialgebras, where \simeq 's are the usual identifications (see §5).*

PROOF. The identifications \simeq 's are isomorphisms of SE^\vee -equivariant graded YB bialgebras, thanks to the observation after Lemma 5.8. Since $\rho: V \rightarrow {}^t V$ is an isomorphism of YB pairs, it suffices to show that $T\rho$, $D\rho$, $S\rho$ and $\wedge\rho$ are SE^\vee -homomorphisms. This is clear for $T\rho$ by Lemma 7.3. So the induced maps $S\rho$ and $\wedge\rho$ are SE^\vee -homomorphisms. It is easy to see that $D\rho = (S\rho)^\circ$, so it is also an SE^\vee -homomorphism. \square

COROLLARY 7.5. *For a partition λ , $D_\lambda\rho: D_\lambda V \rightarrow \text{tr}_\sharp D_\lambda {}^t V \simeq (S_\lambda V)^\circ$, $\wedge_\lambda\rho: \bigwedge_\lambda V \rightarrow \text{tr}_\sharp \bigwedge_\lambda {}^t V \simeq (\bigwedge_\lambda V)^\circ$, $L_\lambda\rho: L'_\lambda V \rightarrow \text{tr}_\sharp L'_\lambda {}^t V \simeq (K_\lambda V)^\circ$ are isomorphisms of SE^\vee -comodules. We have $(K_\lambda\rho)^\circ = L_\lambda\rho$ and $D_\lambda\rho = (S_\lambda\rho)^\circ$.*

REMARK 7.6. It is easy to see that $\beta_E^\pm, \gamma_E^\pm \in \mathcal{P}((E^\vee)^{\otimes 2}, (E^\vee)^{\otimes 2})$, where $(E^\vee)^{\otimes 2}$ is the subcomodule of the right regular representation $SE_{SE^\vee}^\vee$. It follows that SE^\vee and $S^t E$ are SE^\vee -equivariant YB-bialgebras. The direct sum of Schur algebras $DE = \bigoplus_{k \geq 0} D_k E$ is an R -algebra without unit. The graded duals of the left and the right regular representations $({}_{DE}DE)^\dagger$ and $(DE_{DE})^\dagger$ are identified with the regular representations ${}_{SE^\vee}SE^\vee$ and $SE_{SE^\vee}^\vee$ respectively. Since

$$\text{tr}: SE_{SE^\vee}^\vee \longrightarrow \text{tr}_\sharp({}_{SE^\vee}SE^\vee)$$

is an isomorphism in \mathcal{P} , we have $({}_{DE}DE)^\circ \simeq {}_{SE^\vee}SE^\vee$. It follows that ${}_{DE}DE$, viewed as SE^\vee -comodule, is also an SE^\vee -equivariant YB bialgebra. The left DE -module DE^{-1} is identified with $(S^t E_{SE^\vee})^\circ$, and is also an SE^\vee -equivariant YB bialgebra.

Let I be the ideal of SE^\vee generated by $\{x_{ij} \mid i \neq j\}$. It is easy to see that I is a homogeneous bi-ideal of SE^\vee . By (3.6), we have $SE^\vee/I = S(\bigoplus_{i=1}^N Rt_i, \tau)$, where t_i is the image of x_{ii} in SE^\vee/I , and τ is the trivial twisting. Let us denote by T the YB pair

$(\bigoplus_{i=1}^N R t_i, \tau)$. The structure of ST is independent of q , and is commutative and cocommutative. In fact, $ST = R[t_1, \dots, t_N]$ is a polynomial ring, and each monomial $t^\lambda = t^{\lambda_1} \cdots t_N^{\lambda_N}$ is group-like for $\lambda \in \Omega^+$ with $\ell(\lambda) \leq N$. Taking the dual, we see that $D_k T = D_k t^\lambda T$ is a subalgebra of the Schur algebra $D_k E$, and $\{t^{*(\lambda)} = t_1^{*(\lambda_1)} \cdots t_N^{*(\lambda_N)} \mid \lambda \in \Omega_k^+, \ell(\lambda) \leq N\}$ is a set of mutually orthogonal primitive idempotents of $D_k T$. It follows that $1 = \sum_\lambda t^{*(\lambda)}$, and that $D_k T \simeq \prod_\lambda t^{*(\lambda)} (D_k T) t^{*(\lambda)} \simeq \prod_\lambda R$.

DEFINITION 7.7. Let $M \in \mathcal{P}$ and $\lambda \in \Omega_k^+$ with $\ell(\lambda) \leq N$. We call the R -module $t^{*(\lambda)} M_k$ the weight λ component of M , and denote it by M_λ .

Clearly, M_λ is a finitely generated projective R -module. By [6, p. 375], we have

$$(7.1) \quad M_\lambda = t^{*(\lambda)} M_k \simeq \text{Hom}_{D_k T}((D_k T) t^{*(\lambda)}, M) \simeq \text{Hom}_{D_k E}((D_k E) t^{*(\lambda)}, M),$$

so that $(\)_\lambda$ is an exact functor $((D_k E) t^{*(\lambda)})$ is projective). Since $1 = \sum_\lambda t^{*(\lambda)}$, we have $M = \bigoplus_\lambda M_\lambda$. It is easy to see that $M_\lambda = \{m \in M \mid \omega(m) = m \otimes t_\lambda\}$, where ω is the coaction as an ST -comodule. For a base change $R \rightarrow R'$, we have $(R' \otimes_R M)_\lambda \simeq R' \otimes_R M_\lambda$. If the prime spectrum $\text{Spec } R$ is connected, then $\dim_K(K \otimes_R M)_\lambda$ does not depend on the field K or the map $R \rightarrow K$.

LEMMA 7.8. Let $M \in \mathcal{P}$ and $\lambda \in \Omega_k^+$ with $\ell(\lambda) \leq N$. Then we have an isomorphism of R -modules $(M^\circ)_\lambda \simeq M_\lambda^*$.

PROOF. We may replace M by M_k . Clearly, we have $M^* t^{*(\lambda)} \simeq (M_\lambda)^*$. On the other hand, the restriction of tr^\dagger to DT is the identity map. \square

PROPOSITION 7.9. For $k \geq 0$ and $\lambda \in \Omega_k^+$ with $\ell(\lambda) \leq N$, we have:

(1) $(D_k E) t^{*(\lambda)} \simeq D_\lambda V$ as SE^\vee -comodules.

(2) $D_\lambda V$ is projective, and is generated by $c(\lambda) := u_1^{(\lambda_1)} \otimes \cdots \otimes u_N^{(\lambda_N)}$ as a left $D_k E$ -module.

PROOF. First note that the identification $SE^\vee \simeq S'E$ maps a monomial M to M . Hence, the identification $D_k E \simeq D_k E^{-1}$ maps $t^{*(\lambda)}$ to $t^{*(\lambda)}$. For $i \in [1, N]$, we define $\varphi_i: V \rightarrow E = V \otimes V^*$ by $\varphi_i(u_j) = u_j \otimes v_i$. It is clear that $\varphi_i \in \mathcal{YB}_R(V, E)$, and that φ_i is a homomorphism of left $\text{End}(V) = E$ -module. Hence, the composite map

$$\varphi_\lambda: D_\lambda V = D_{\lambda_1} V \otimes \cdots \otimes D_{\lambda_N} V \xrightarrow{D_{\lambda_1} \varphi_1 \otimes \cdots \otimes D_{\lambda_N} \varphi_N} D_{\lambda_1} E \otimes \cdots \otimes D_{\lambda_N} E \xrightarrow{m} D_k E$$

is a homomorphism of left $D_k E$ -modules. Observe that $\varphi_\lambda(c(\lambda)) = t^{*(\lambda)}$ (cf. Lemma 6.22). The sum $\sum_\lambda \varphi_\lambda: \bigoplus_\lambda D_\lambda V \rightarrow D_k E$ is surjective, since $\sum_\lambda \varphi_\lambda(c(\lambda)) = 1$. On the other hand, we have $\text{rank}_R(\bigoplus_\lambda D_\lambda V) = \text{rank}_R(D_k E)$ (since the rank in question is stable under base change, we may assume that $R = C$ and $q = 1$). In this case, see e.g. [2]). It follows that $\sum_\lambda \varphi_\lambda$ is an isomorphism. Since $\sum_\lambda \varphi_\lambda$ is injective, and $\text{Im } \varphi_\lambda \supset (D_k E) t^{*(\lambda)}$ for each λ , we have $\text{Im } \varphi_\lambda = (D_k E) t^{*(\lambda)}$, and $\varphi_\lambda: D_\lambda V \rightarrow (D_k E) t^{*(\lambda)}$ is an isomorphism. Hence, we have (1). Now (2) is obvious. \square

COROLLARY 7.10. *Let $\lambda \in \Omega_k^+$ and $\ell(\lambda) \leq N$. For $M \in \mathcal{P}$, we have an isomorphism $\Psi_\lambda: M_\lambda \xrightarrow{\sim} \mathcal{P}(D_\lambda, M)$ given by $(\Psi_\lambda(a))(c(\lambda)) = a$ for $a \in M_\lambda$.*

LEMMA 7.11. *Let $\lambda \in \Omega^+$ with $\ell(\lambda) \leq N$. For any $\sigma \in \mathfrak{S}_N$, we have an isomorphism of SE^\vee -comodules $D_\lambda V \simeq D_{\sigma\lambda}V$, where $\sigma\lambda = (\lambda_{\sigma^{-1}1}, \dots, \lambda_{\sigma^{-1}N})$. In particular, we have $D_\lambda V \simeq D_{(\tilde{\lambda})^\sim}V$. For $M \in \text{ob } \mathcal{P}$, we have $M_\lambda \simeq M_{\sigma\lambda}$.*

PROOF. The SE^\vee -isomorphism $\psi_{DV}(\sigma): (DV)^{\otimes N} \rightarrow (DV)^{\otimes N}$ maps $D_\lambda V$ onto $D_{\sigma\lambda}V$. The last assertion follows from Corollary 7.10. \square

Let λ and μ be elements in Ω^+ such that $\ell(\lambda), \ell(\mu) \leq N$. We say that $\mu \leqq \lambda$ when $\lambda - \mu$ is a non-negative linear combination of $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ ($i \in [1, N-1]$). It is easy to see that $\mu \leqq \lambda$ implies $|\mu| = |\lambda|$ and $\mu \leq \lambda$. For $\lambda \in \Omega^+$ with $\ell(\lambda) \leq N$, we have $\lambda \leqq \tilde{\lambda}^\sim$. For $\mu \in \Omega^-$ with $\ell(\mu) \leq N$ and $\lambda \in S_\square(\mu)$, we have $\mu < \lambda$ by the definition of $S_\square(\lambda)$ (see §6).

THEOREM 7.12. *Let $k \geq 0$, and $\lambda \in \Omega_k^-$ with $\ell(\lambda) \leq N$. Then we have:*

- (1) $\mu \in \Omega_k^-, \ell(\mu) \leq N, (K_\lambda V)_\mu \neq 0 \Rightarrow \mu \leqq \lambda$.
- (2) $(K_\lambda V)_\lambda \simeq R$.
- (3) $K_\lambda V$ is generated by $(K_\lambda V)_\lambda$ as a $D_k E$ -module.

Let M be a $D_k E$ -module which satisfies:

- (1') $\mu \in \Omega_k^-, \ell(\mu) \leq N, \mu > \lambda \Rightarrow M_\mu = 0$.
- (3') M is generated by some $a \in M_\lambda$.

Then M is a homomorphic image of $K_\lambda V$. Moreover, we have:

- (1'') $\mu \in \Omega^-, \ell(\mu) \leq N, M_\mu \neq 0 \Rightarrow \mu \leqq \lambda$.
- (2') $\text{Hom}_{D_k E}(K_\lambda V, M) \simeq M_\lambda \simeq Ra$.

Conversely, any homomorphic image M of $K_\lambda V$ satisfies (1') and (3').

PROOF. By Proposition 7.9 (2), $K_\lambda V$ is generated by $d'_\lambda(V)(c(\lambda)) \in (K_\lambda V)_\lambda$. So we have (3). By Corollary 6.24, $\text{rank}_R(K_\lambda V)_\mu$ does not depend on R nor q . Parts (1) and (2) are well-known in the case $R = \mathbb{C}$ and $q = 1$ (see [4]). So the general case immediately follows, since any rank one projective module over Laurent polynomial ring $\mathbb{Z}[Q, Q^{-1}]$ is free. Now the last assertion is obvious, since $(\)_\mu$ is an exact functor for any μ . Let M be a $D_k E$ -module which satisfies (1') and (3'). By (3') and Corollary 7.10, we have a surjective map $\Psi_\lambda(a): D_\lambda V \rightarrow M$. By Theorem 6.23, we have an exact sequence

$$0 \rightarrow \text{Hom}_{D_k E}(K_\lambda V, M) \rightarrow \text{Hom}_{D_k E}(D_\lambda V, M) \rightarrow \bigoplus_{\mu \in S_\square(\lambda)} \text{Hom}_{D_k E}(D_\mu V, M).$$

For $\mu \in S_\square(\lambda)$, we have $\text{Hom}_{D_k E}(D_\mu V, M) \simeq M_\mu \simeq M_{\tilde{\mu}^\sim}$ by Lemma 7.11. Since $\tilde{\mu}^\sim \geqq \mu > \lambda$ and $\tilde{\mu}^\sim \in \Omega_k^-$, we have $M_{\tilde{\mu}^\sim} = 0$ by (1'). It follows that $\text{Hom}_{D_k E}(d'_\lambda(V), M): \text{Hom}_{D_k E}(K_\lambda V, M) \rightarrow \text{Hom}_{D_k E}(D_\lambda V, M) \simeq M_\lambda$ is an isomorphism. In particular, we have an induced surjective map $\overline{\Psi_\lambda(a)}: K_\lambda V \rightarrow M$. Now (1'') and (2') are obvious. \square

REMARK 7.13. Since DV and DV^{-1} are isomorphic as SE^\vee -equivariant YB coalgebras, we have $D_\lambda V \simeq D_\lambda V^{-1}$ for $\lambda \in \Omega^+$. So it is easy to see that $K_\lambda V^{-1}$ satisfies

(1') and (3') in the theorem for $\lambda \in \Omega^-$ with $\ell(\lambda) \leq N$. We have a surjective map $K_\lambda V \rightarrow K_\lambda V^{-1}$. This map is an isomorphism, since $K_\lambda V$ and $K_\lambda V^{-1}$ have the same rank. Since $\bigwedge V$ and $\bigwedge V^{-1}$ are isomorphic as SE^\vee -equivariant YB algebras, the image $K'_\lambda V$ of the composite map

$$d''_\lambda(V): D_\lambda V \xrightarrow{A \otimes \cdots \otimes A} T_\lambda V \xrightarrow{(-\gamma_V^{-1})(\chi_\lambda)} T_{\tilde{\lambda}} V \xrightarrow{m \otimes \cdots \otimes m} \bigwedge_{\tilde{\lambda}} V$$

is isomorphic to $K_\lambda V^{-1}$. Hence, we have $K'_\lambda V \simeq K_\lambda V$. Taking the contravariant dual, we have $L_{\tilde{\lambda}} V \simeq L_{\tilde{\lambda}} V^{-1} \simeq (K'_\lambda V)^\circ$.

As we will see in the next section, the study of SE^\vee -comodules is closely related to the representations of the *Iwahori Hecke algebra*. The Iwahori Hecke algebra \mathfrak{H}_k is defined to be the quotient algebra

$$(7.2) \quad \mathfrak{H}_k \simeq R[\mathfrak{B}_k]/((1 - b_i)(1 + q^{-2}b_i) \mid i = 1, \dots, k-1).$$

We set $t = q^{-2}$. By Iwahori's theorem, $h(\sigma) := t^{\ell(\sigma)} \bar{b}_{i_1} \cdots \bar{b}_{i_{\ell(\sigma)}}$ is independent of the choice of the reduced expression $\sigma = \sigma_{i_1} \cdots \sigma_{i_{\ell(\sigma)}}$ of $\sigma \in \mathfrak{S}_k$, where \bar{b}_j ($1 \leq j < k$) is the image of b_j in \mathfrak{H}_k . It is known and easily shown that \mathfrak{H}_k is R -free of rank $k!$, and $\{h(\sigma) \mid \sigma \in \mathfrak{S}_k\}$ is a free basis of \mathfrak{H}_k . By (1.4), the representation of $R[\mathfrak{B}_k]$ on $V^{\otimes k}$ given by (1.8) yields the representation $B: \mathfrak{H}_k \rightarrow \text{End}_R V^{\otimes k}$ given by $B(\bar{b}_i) = (\beta_V)_i$ ($1 \leq i < k$). Since $D_k E = \text{End}_{\mathfrak{H}_k} V^{\otimes k}$, we have $\text{Im } B \subset \text{End}_{D_k E} V^{\otimes k}$. Note that $B(h(\sigma)) = (-\gamma_V)(\sigma)$ for $\sigma \in \mathfrak{S}_k$.

LEMMA 7.14. *Assume that $N \geq k$. Then, $B: \mathfrak{H}_k \rightarrow \text{End}_{D_k E} V^{\otimes k}$ is an isomorphism.*

PROOF. Since $V^{\otimes k} = D_{(1^k)} V$, we have an isomorphism $\Psi_{(1^k)}: (V^{\otimes k})_{(1^k)} \cong \text{End}_{D_k E} V^{\otimes k}$ by Corollary 7.10. Under this identification, $B: \mathfrak{H}_k \rightarrow V_{(1^k)}^{\otimes k}$ is given by $B(h(\sigma)) = (-\gamma_V)(\sigma)(u_1 \otimes \cdots \otimes u_k)$. By (5.4), we have $(-\gamma_V)(\sigma)(u_1 \otimes \cdots \otimes u_k) = q^{-\ell(\sigma)}(u_{\sigma^{-1}1} \otimes \cdots \otimes u_{\sigma^{-1}k})$. Hence, B is bijective. \square

8. Polynomial representations of GL_q over a field. In this section, we work on a base field K , instead of R . Throughout this section, $V = (V, \beta_V, \gamma_V)$ and $E = (E, \beta_E, \gamma_E)$ are as in §7. A *polynomial representation* of $GL_q(N, K)$ is a finite-dimensional right SE^\vee -comodule by definition. This section is devoted to studying polynomial representations of $GL_q(N, K)$ over a field K . Note that the categories $\text{cfl-}S_k E$ and \mathcal{P} (§7) are abelian, and that the contravariant dual $()^\circ$ is exact.

DEFINITION 8.1. For a polynomial representation M of $GL_q(N, K)$, we define a polynomial

$$(8.1) \quad \chi(M) = \sum_{\lambda \in \Omega^+, \ell(\lambda) \leq N} (\dim_K M_\lambda) t_1^{\lambda_1} \cdots t_N^{\lambda_N} \in \mathbb{Z}[t_1, \dots, t_N]$$

in the variables t_1, \dots, t_N , and call it the *formal character* of M .

By Lemma 7.11, $\chi(M)$ is a symmetric polynomial in t_1, \dots, t_N . For polynomial representations M and N of $GL_q(N, K)$ and $\lambda \in \Omega^+$ with $\ell(\lambda) \leq N$, we have $(M \otimes N)_\lambda \simeq \bigoplus_{\mu + \mu' = \lambda} M_\mu \otimes N_{\mu'}$. Since $(\cdot)_\lambda$ is an exact functor for $\lambda \in \Omega^+$ with $\ell(\lambda) \leq N$, the formal character χ is a homomorphism from the Grothendieck ring POL of polynomial representations of $GL_q(N, K)$ to the ring $Z[t_1, \dots, t_N]^{\mathfrak{S}_k}$ of symmetric polynomials. If \tilde{M} is an SE^\vee -comodule over the Laurent polynomial ring $Z[Q, Q^{-1}]$ such that it is $Z[Q, Q^{-1}]$ -free of finite rank, then $\chi(K \otimes_{Z[Q, Q^{-1}]} \tilde{M})$ does not depend on the field K nor the value of q by the remark before Lemma 7.8. Hence, for $k \geq 0$ and $\lambda \in \Omega^-$ with $\ell(\lambda) \leq N$, we obtain the following formulas:

$$(8.2) \quad \chi(\bigwedge_k V) = e_k := \sum_{1 \leq i_1 < \dots < i_k \leq N} t_{i_1} \cdots t_{i_k}.$$

$$(8.3) \quad \chi(S_k V) = h_k := \sum_M M, \text{ where } M \text{ runs through the monomials of degree } k.$$

$$(8.4) \quad \chi(K_\lambda V) = s_\lambda := \det(h_{\lambda_i - i + j})_{1 \leq i, j \leq N} = \det(e_{\tilde{\lambda}_i - i + j})_{1 \leq i, j \leq N}.$$

LEMMA 8.2. *For a polynomial representation M of $GL_q(N, K)$, we have $\chi(M^\circ) = \chi(M)$.*

This follows immediately from Lemma 7.8.

In the rest of this section, we fix $k \geq 0$, and set $A = D_k E$. We denote the Jacobson radical of A by J . For $M \in A\text{-mff}$, $(M/JM)^\circ$ is the unique maximal semisimple submodule of M° . Namely, we have $(M/JM)^\circ \simeq \text{Soc } M^\circ$, where Soc denotes the socle.

LEMMA 8.3. *Let $\lambda \in \Omega_k^-$ and $\ell(\lambda) \leq N$. Then, $K_\lambda V/J(K_\lambda V) \simeq (\text{Soc}(L_{\tilde{\lambda}} V))^\circ$ is simple (irreducible). In particular, $K_\lambda V$ and $L_{\tilde{\lambda}} V$ are indecomposable.*

PROOF. Let M be a proper submodule of $K_\lambda V$. Then, we have $M_\lambda = 0$ by (2) and (3) of Theorem 7.12. We define $(K_\lambda V)_{\max}$ to be the sum of all proper submodules of $K_\lambda V$. We have $(K_\lambda V)_{\max} \neq K_\lambda V$, since $((K_\lambda V)_{\max})_\lambda = 0$. It follows that $(K_\lambda V)_{\max}$ is the unique maximal proper submodule of $K_\lambda V$. Hence, we have $(K_\lambda V)_{\max} = J(K_\lambda V)$, and $K_\lambda V/J(K_\lambda V)$ is simple. By Corollary 7.5, we have $(\text{Soc}(L_{\tilde{\lambda}} V))^\circ \simeq K_\lambda V/J(K_\lambda V)$. \square

For $\lambda \in \Omega_k^-$ with $\ell(\lambda) \leq N$, we denote by F_λ the simple module $K_\lambda V/J(K_\lambda V)$.

PROPOSITION 8.4. *Each simple A -module is isomorphic to F_λ for exactly one $\lambda \in \Omega_k^-$ such that $\ell(\lambda) \leq N$.*

PROOF. In this proof, we use the lexicographic order \leq of the set $\{\lambda \in \Omega_k^- \mid \ell(\lambda) \leq N\}$. For $0 \neq M \in A\text{-mff}$, we define

$$(8.5) \quad h(M) := \max\{\lambda \in \Omega_k^- \mid \ell(\lambda) \leq N, M_\lambda \neq 0\}.$$

Since $0 \neq M = \bigoplus_\lambda M_\lambda$ and $M_\lambda \simeq M_{\tilde{\lambda}}$, the partition $h(M)$ is well-defined. Since F_λ is a non-zero homomorphic image of $K_\lambda V$, we have $h(F_\lambda) = \lambda$ by Theorem 7.12. Hence,

$F_\lambda \neq F_\mu$ for $\mu \neq \lambda$. Let M be a simple A -module. By the definition of $h(M)$, the condition (1') of Theorem 7.12 is satisfied for $\lambda = h(M)$. Since M is simple, the condition (3') of Theorem 7.12 is satisfied for any non-zero vector $a \in M_{h(M)}$. Hence, M is a homomorphic image of $K_{h(M)}$, and we have $M \simeq F_{h(M)}$. \square

COROLLARY 8.5. *The formal character $\chi: \text{POL} \rightarrow \mathbb{Z}[t_1, \dots, t_N]^{\otimes N}$ is a ring isomorphism, where POL is the Grothendieck ring of polynomial representations of $GL_q(N, K)$.*

PROOF. It suffices to show that $\chi: [A\text{-mff}] \rightarrow \text{Sym}(k)$ is a bijection, where $[A\text{-mff}]$ denotes the Grothendieck group of A -mff and $\text{Sym}(k)$ is the module of symmetric polynomials of degree k in t_1, \dots, t_N . By [25], $\text{Sym}(k)$ is a \mathbb{Z} -free module with a free basis $\{s_\lambda \mid \lambda \in \Omega_k^-, \ell(\lambda) \leq N\}$. By (8.4), $\chi: [A\text{-mff}] \rightarrow \text{Sym}(k)$ is surjective. By the proposition, we have $\text{rank}[A\text{-mff}] = \text{rank Sym}(k)$. \square

COROLLARY 8.6. *Let $M \in A\text{-mff}$. Then, M and M° have the same composition factors. In particular, We have $F_\lambda \simeq F_\lambda^\circ \simeq \text{Soc}(L_{\tilde{\lambda}} V)$ for $\lambda \in \Omega_k^-$ with $\ell(\lambda) \leq N$.*

This follows immediately from Lemma 8.2 and Corollary 8.5.

LEMMA 8.7. *Let $\lambda \in \Omega_k^-$ with $\ell(\lambda) \leq N$. Then, the following are equivalent:*

- (1) $K_\lambda V$ is simple.
- (2) $L_{\tilde{\lambda}} V$ is simple.
- (3) $K_\lambda V \simeq L_{\tilde{\lambda}} V$.

PROOF. $K_\lambda V \simeq F_\lambda \Leftrightarrow (K_\lambda V)^\circ \simeq F_\lambda^\circ \Leftrightarrow L_{\tilde{\lambda}} V \simeq F_\lambda^\circ$. Hence, we have (1) \Leftrightarrow (2). By Corollary 8.6, (1) (or (2)) also implies (3). We shall show that (3) \Rightarrow (2). By Theorem 7.12, $L_{\tilde{\lambda}} V$ is generated by $(L_{\tilde{\lambda}} V)_\lambda$. Since both $(L_{\tilde{\lambda}} V)_\lambda$ and $(F_\lambda)_\lambda$ are one-dimensional, $L_{\tilde{\lambda}} V$ is generated by its simple socle F_λ . Hence, we have $L_{\tilde{\lambda}} V \simeq F_\lambda$. \square

LEMMA 8.8. *Let $\lambda \in \Omega_k^-$ and $\ell(\lambda) \leq N$. Then, F_λ is absolutely irreducible.*

PROOF. Since $0 \neq \text{End}_A F_\lambda \subset \text{Hom}_A(K_\lambda, F_\lambda) \simeq K$ by Theorem 7.12, $\text{End}_A F_\lambda$ is one-dimensional. The lemma follows from [6, (29.13)]. \square

The rest of this sections is devoted to the proof of the following theorem.

THEOREM 8.9. *Let K be a field, (V, β_V) be Jimbo's YB pair of type $A_{N-1}^{(1)}$ defined by $q \in K^\times$ and (1.3), and let λ be a partition with $\ell(\lambda) \leq N$. If the product $\prod_{(i,j) \in A_\lambda} h_q(i,j)$ of (q -analogues of) hook lengths is non-zero, then the Weyl module $K_\lambda V$ is simple (irreducible) and projective, where $h_q(i,j) = [\lambda_i + \tilde{\lambda}_j - i - j + 1]_{q-2}$.*

To prove the theorem, we use Gyoja's q -analogues of Young symmetrizers [14]. For this purpose, let us clarify the relationship between the notation in [14] and that of ours. Our size k of the Iwahori Hecke algebra, is denoted by n in [14]. The notation $h(\sigma)$ agrees, while q in [14] corresponds to our $t = q^{-2}$. To avoid confusion, we will

stick to our notation k , (our) q and $t=q^{-2}$. For $\lambda \in \Omega_k^-$, our \mathfrak{S}_λ , $\mathfrak{S}_{\tilde{\lambda}}$, $\alpha_{r,\lambda}^{-1}$, $\alpha_{c,\lambda}^{-1}$, χ_λ and $\chi_{\tilde{\lambda}}=\chi_\lambda^{-1}$ correspond to $W_+(\lambda)$, $W_-(\lambda)$, $T_+(\lambda)$, $T_-(\lambda)$, $[-, +]$ and $[+, -]$ in [14], respectively. Hence, we have

$$(8.6) \quad Be_+ = \sum_{\sigma \in \mathfrak{S}_\lambda} Bh(\sigma) = \sum_{\sigma \in \mathfrak{S}_\lambda} (-\gamma_V)(\sigma),$$

$$(8.7) \quad Be_- = \sum_{\sigma \in \mathfrak{S}_{\tilde{\lambda}}} (-t)^{-\ell(\sigma)} Bh(\sigma) = \sum_{\sigma \in \mathfrak{S}_{\tilde{\lambda}}} (-\beta_V)(\sigma),$$

where $e_+ = e_+(\lambda)$ and $e_- = e_-(\lambda)$ are those in [14], and B is as defined in §7.

PROOF OF THEOREM 8.9. We set $k=|\lambda|$. It suffices to show that $K'_\lambda V$ (see Remark 7.13) is simple and projective. We define the map $\Phi_\lambda: S_\lambda V \rightarrow D_\lambda V$ to be the restriction of $T_N \Phi: (SV)^{\otimes N} \rightarrow (DV)^{\otimes N}$ to $S_\lambda V$, where $\Phi=\Phi^{SV}$ is the map which appears in Proposition 5.2. By the remark below Lemma 5.8, Φ_λ is a homomorphism of SE^\vee -comodules. We denote by $\kappa_\lambda(V)$ the composite map

$$K'_\lambda V \hookrightarrow \bigwedge_{\tilde{\lambda}} V \xrightarrow{d_\lambda(V)} L_{\tilde{\lambda}} V \hookrightarrow S_\lambda V \xrightarrow{\Phi_\lambda} D_\lambda V \xrightarrow{d''_\lambda(V)} K'_\lambda V,$$

where $d''_\lambda(V)$ is the map defined in Remark 7.13. The map $\kappa_\lambda(V)$ is defined over the Laurent polynomial ring $\mathbb{Z}[Q, Q^{-1}]$, and is a multiplication by a scalar, say c , by Theorem 7.12. We claim that $c \in \mathbb{Z}[Q, Q^{-1}]$ is expressed as $c = Q^* \prod_{(i,j) \in \Delta_\lambda} h_Q(i,j)$. Assume that the claim is true. Then, $\kappa_\lambda(V) = q^* \prod h_q(i,j) \neq 0$ is an isomorphism over K . Since the isomorphism $\kappa_\lambda(V)$ factors through $L_{\tilde{\lambda}} V$ and $\dim_K K'_\lambda V = \dim_K L_{\tilde{\lambda}} V$, we have $K'_\lambda V \simeq L_{\tilde{\lambda}} V$. By Lemma 8.7, $K'_\lambda V$ is simple. Since the scalar multiplication $\kappa_\lambda(V)$ factors through $D_\lambda V$, the epimorphism $d''_\lambda(V): D_\lambda V \rightarrow K'_\lambda V$ splits. Since $D_\lambda V$ is projective by Proposition 7.9, (2), the direct summand $K'_\lambda V$ is also projective, and the proof is complete. To prove the claim, we may extend the base ring and work over the base field $K=\mathbf{C}(Q)$. We still use the notation $t=Q^{-2}$. We denote the maps $m \otimes \cdots \otimes m: V^{\otimes k} \simeq T_\lambda V \rightarrow D_\lambda V$, $\Delta \otimes \cdots \otimes \Delta: D_\lambda V \rightarrow T_\lambda V \simeq V^{\otimes k}$, $m \otimes \cdots \otimes m: V^{\otimes k} \simeq T_{\tilde{\lambda}} V \rightarrow \bigwedge_{\tilde{\lambda}} V$, $\Delta \otimes \cdots \otimes \Delta: \bigwedge_{\tilde{\lambda}} V \rightarrow T_{\tilde{\lambda}} V \simeq V^{\otimes k}$, $m \otimes \cdots \otimes m: V^{\otimes k} \simeq T_\lambda V \rightarrow S_\lambda V$ and $\Delta \otimes \cdots \otimes \Delta: S_\lambda V \rightarrow T_\lambda V \simeq V^{\otimes k}$ by m_D , Δ_D , m_Λ , Δ_Λ , m_S and Δ_S , respectively. Since Φ is a homomorphism of YB bialgebras, it holds that $\Phi_\lambda \circ m_S = m_D$ and $\Delta_D \circ \Phi_\lambda = \Delta_S$ ($\Phi_{(1k)}$ is identified with $\text{id}_{V^{\otimes k}}$ and omitted). By (5.1) and (8.7), we have $Be_- = \Delta_\Lambda \circ m_\Lambda$. By (5.2) and (8.6), we have $Be_+ = \Delta_S \circ m_S = \Delta_D \circ \Phi_\lambda \circ m_S = \Delta_D \circ m_D$. Now consider the map

$$(8.8) \quad V^{\otimes k} \xrightarrow{m_D} D_\lambda V \xrightarrow{d''_\lambda} K'_\lambda V \xrightarrow{\kappa_\lambda} K'_\lambda V \hookrightarrow \bigwedge_{\tilde{\lambda}} V \xrightarrow{\Delta_\Lambda} V^{\otimes k} \xrightarrow{h_-} V^{\otimes k},$$

where $h_- = h([- , +])$. Since the diagram

(8.9)

$$\begin{array}{ccccccccccccc}
V^{\otimes k} & \xrightarrow{e_+} & V^{\otimes k} & \xrightarrow{h_-^{-1}} & V^{\otimes k} & \xrightarrow{e_-} & V^{\otimes k} & \xrightarrow{h_-} & V^{\otimes k} & \xrightarrow{e_+} & V^{\otimes k} & \xrightarrow{h_-^{-1}} & V^{\otimes k} & \xrightarrow{e_-} & V^{\otimes k} & \xrightarrow{h_-} & V^{\otimes k} \\
& \searrow m & \uparrow \Delta & \downarrow m & \nearrow m & \Delta & \searrow m & \nearrow m & \downarrow m & \nearrow m & \Delta & \downarrow m & \nearrow m & \Delta & \downarrow m & \nearrow m & \Delta & \downarrow m & \nearrow m \\
& D_\lambda V & \longrightarrow & \bigwedge_{\tilde{\lambda}} V & \longrightarrow & S_\lambda V & \longrightarrow & D_\lambda V & \longrightarrow & \bigwedge_{\tilde{\lambda}} V & \longrightarrow & K'_\lambda V & \longrightarrow & L_{\tilde{\lambda}} V & \longrightarrow & K'_\lambda V & \longrightarrow & K'_\lambda V
\end{array}$$

is commutative and the bottom row map is $\kappa_\lambda(V)$, the map (8.8) is $B(h_- e_- h_-^{-1} e_+)^2$. On the other hand, we have $\kappa_\lambda(V) = c$. So the map (8.8) equals $cB(h_- e_- h_-^{-1} e_+)$ by the same diagram. Hence, we have

$$(8.10) \quad B(h_- e_- h_-^{-1} e_+)^2 = cB(h_- e_- h_-^{-1} e_+).$$

Since $K = C(Q)$, we see that $\Delta: S_i^t V \rightarrow T_i^t V$ ($i \geq 0$) is injective by Lemma 5.4. It follows that m_D is surjective. Again by (8.9), we have $(h_- e_- h_-^{-1} e_+)V^{\otimes k} \simeq K'_\lambda V \neq 0$. Hence, c is uniquely determined by (8.10).

It is now clear that the scalar c is nothing but $c(T_+)$ in [14]. Here we recall the value of c [14, (3.9)].

$$(8.11) \quad c = \frac{\prod_{i=1}^m \sum_{l=1}^{\lambda_i + m - i} (t^l - 1)}{\prod_{1 \leq i < j \leq m} (t^{\lambda_i + m - i} - t^{\lambda_j + m - j})} t^{(m)} \cdot (t-1)^{-k},$$

where m is any integer such that $m \geq N$. By [25, p. 9], we have

$$c = \frac{\prod_i \prod_{l=1}^{\mu_i} (t^l - 1)}{\prod_{i < j} (\mu_i - \mu_j - 1)} t^*(t-1)^{-k} = t^* \prod_{(i,j) \in \Delta_\lambda} \frac{t^{h_1(i,j)} - 1}{t-1} = t^* \prod_{(i,j) \in \Delta_\lambda} h_Q(i,j),$$

as desired, where $\mu = (\mu_1, \dots, \mu_m)$ is given by $\mu_i = \lambda_i + m - i$ for $i \in [1, m]$. \square

COROLLARY 8.10. *Assume that $[a]_t \neq 0$ for $a \in [1, k]$. Then the Schur algebra $A = \text{End}_{\mathfrak{H}_k} V^{\otimes k}$ is semisimple, and $\{K_\lambda \mid \lambda \in \Omega_k^-, \ell(\lambda) \leq N\}$ is a complete set of non-isomorphic simple A -modules. In this case, the Iwahori Hecke algebra \mathfrak{H}_k is also semisimple.*

PROOF. By the theorem, $K_\lambda V$ is simple and projective for any $\lambda \in \Omega_k^-$ with $\ell(\lambda) \leq N$. By Proposition 8.4, $\{K_\lambda \mid \lambda \in \Omega_k^-, \ell(\lambda) \leq N\}$ is a complete set of non-isomorphic simple A -modules. Since each simple module is projective, $A = \text{End}_{\mathfrak{H}_k} V^{\otimes k}$ is semisimple. Applying this for $N \geq k$, we see that $\mathfrak{H}_k = \text{End}_A V^{\otimes k}$ is also semisimple, by the double centralizer theorem. \square

NOTE. Gyoja and Uno [15] proved the semisimplicity of \mathfrak{H}_k under the same assumption as in the corollary above (and some extra condition on the base field, which can be dropped easily). The semisimplicity of A also follows from their result using the double centralizer theorem.

COROLLARY 8.11. *In the situation of Corollary 8.10, $A = \text{End}_{\mathfrak{H}_k} V^{\otimes k}$ is isomorphic to $\prod_{\lambda \in \Omega_k, \ell(\lambda) \leq N} \text{End}_K(K_\lambda V)$ as a K -algebra. We have an irreducible decomposition*

$$(8.12) \quad A \simeq S_k E^\vee \simeq \bigoplus_{\lambda \in \Omega_k, \ell(\lambda) \leq N} K_\lambda V \otimes (K_\lambda V)^*$$

of A -bicomodules of A and $S_k E^\vee$.

PROOF. The first assertion is a consequence of Lemma 8.8 and Corollary 8.10. So we have $S_k E^\vee \simeq \bigoplus_{\lambda} (K_\lambda V)^* \otimes K_\lambda V$ as a K -coalgebra, and the isomorphism (8.12) is established. The category of finite dimensional A -bimodules can be identified with $(A \otimes A^{\text{op}})\text{-mff}$, where $(\cdot)^{\text{op}}$ denotes the opposite algebra. By the first assertion, we have $A \otimes A^{\text{op}} \simeq \prod_{\lambda} \text{End}_K(K_\lambda V \otimes (K_\lambda V)^*)$, and the decomposition (8.12) is an irreducible decomposition. \square

In the next section, we will discuss the A -bimodule (or SE^\vee -bicomodule) structure of $S_k E^\vee$ over an arbitrary commutative ring R and $q \in R^\times$.

9. Quantum determinants and the straightening formulas. Throughout this section, V and E are as in §7. Let $M \in \text{cff-}SE^\vee$. Then we have the coefficient map $\text{cf}_M : M^* \otimes M \rightarrow SE^\vee$ (see §3). Since cf_M is a homomorphism of coalgebras, $M^* \otimes M$ is an SE^\vee -bicomodule and cf_M is a homomorphism of SE^\vee -bicomodules. The SE^\vee -bicomodule $M^* \otimes M$ agrees with the tensor product of the left SE^\vee -comodule M^* and the right SE^\vee -comodule M . Since the counit \bar{e} of $M^* \otimes M$ is nothing but $\text{ev} : M^* \otimes M \rightarrow R$, the map $(1 \otimes \text{ev}) \circ (\omega \otimes 1) : M^* \otimes M \rightarrow SE^\vee$ agrees with cf_M . It follows that the diagrams

$$(9.1) \quad \begin{array}{ccc} N^* \otimes M & \xrightarrow{1 \otimes \varphi} & N^* \otimes N \\ \downarrow \varphi^* \otimes 1 & & \downarrow \text{cf}_N \\ M^* \otimes M & \xrightarrow{\text{cf}_M} & SE^\vee \end{array}$$

and

$$(9.2) \quad \begin{array}{ccc} M_1^* \otimes \cdots \otimes M_s^* \otimes M_1 \otimes \cdots \otimes M_s & \xrightarrow{\sim} & (M_1 \otimes \cdots \otimes M_s)^* \otimes (M_1 \otimes \cdots \otimes M_s) \\ \downarrow \tau & & \downarrow \text{cf}_{M_1 \otimes \cdots \otimes M_s} \\ M_1^* \otimes M_1 \otimes \cdots \otimes M_s^* \otimes M_s & \xrightarrow{\text{cf}_{M_1} \otimes \cdots \otimes \text{cf}_{M_s}} & SE^\vee \otimes \cdots \otimes SE^\vee \xrightarrow{m} SE^\vee \end{array}$$

are commutative for $M, N, M_1, \dots, M_s \in \text{cff-}SE^\vee$ and $\varphi \in \text{cff-}SE^\vee(M, N)$, where τ is an appropriate trivial twisting.

In what follows, we identify $\bigwedge V$ with $(\bigwedge V)^\dagger$ via the left SE^\vee -equivariant graded YB bialgebra isomorphism $\Phi^\dagger: \bigwedge V \rightarrow (\bigwedge V)^\dagger$, where $\Phi: \bigwedge V \rightarrow (\bigwedge V)^\dagger$ is the map which appears in Proposition 5.2. So $\bigwedge_k V$ ($k \geq 0$) is identified with $(\bigwedge_k V)^*$. More generally, we identify $\bigwedge_\lambda V$ with $(\bigwedge_\lambda V)^*$ via the isomorphism

$$\bigwedge_\lambda V \xrightarrow{\Phi_{\lambda_1}^* \otimes \cdots \otimes \Phi_{\lambda_s}^*} (\bigwedge_{\lambda_1} V)^* \otimes \cdots \otimes (\bigwedge_{\lambda_s} V)^* \cong (\bigwedge_\lambda V)^*$$

for $\lambda = (\lambda_1, \dots, \lambda_s) \in \Omega^+$.

DEFINITION 9.1. For $k \geq 0$, the coefficient map $\text{cf}_{\bigwedge_k V}: \bigwedge_k V \otimes \bigwedge_k V \rightarrow S_k E^\vee$ is called the *determinant map*, and is denoted by \det_k . For $\lambda = (\lambda_1, \dots, \lambda_s) \in \Omega_k^+$, $\text{cf}_{\bigwedge_\lambda V}: \bigwedge_\lambda V \otimes \bigwedge_\lambda V \rightarrow S_k E^\vee$ is denoted by \det_λ .

By the commutativity of (9.2), \det_λ is the composite map

$$(9.3) \quad \begin{aligned} \bigwedge_\lambda V \otimes \bigwedge_\lambda V &\xrightarrow{\tau} \bigwedge_{\lambda_1} V \otimes \bigwedge_{\lambda_1} V \otimes \cdots \otimes \bigwedge_{\lambda_s} V \otimes \bigwedge_{\lambda_s} V \\ &\xrightarrow{\det_{\lambda_1} \otimes \cdots \otimes \det_{\lambda_s}} S_\lambda E^\vee \xrightarrow{m} S_k E^\vee. \end{aligned}$$

Note that \det_λ is an SE^\vee -bicomodule homomorphism.

LEMMA 9.2. Let $k \geq 0$. Then, \det_k is given by

$$(9.4) \quad \begin{aligned} \det_k(v_{i_1} \wedge \cdots \wedge v_{i_k} \otimes u_{j_1} \wedge \cdots \wedge u_{j_k}) &= \sum_{\sigma \in \mathfrak{S}_k} (-q)^{\epsilon(\sigma)} x_{i_{\sigma(1)} j_1} \cdots x_{i_{\sigma(k)} j_k} \\ &= \sum_{\sigma \in \mathfrak{S}_k} (-q)^{\epsilon(\sigma)} x_{i_{1,j_{\sigma(1)}}} \cdots x_{i_{k,j_{\sigma(k)}}} \end{aligned}$$

for $1 \leq i_1 < \cdots < i_k \leq N$ and $1 \leq j_1 < \cdots < j_k \leq N$. The map $\det_{(1^k)}: (V) \otimes V \rightarrow S_k E^\vee$ is surjective.

PROOF. It is easy to see that $\det_1: V \otimes V \rightarrow E^\vee$ is nothing but the identification $V^* \otimes V \cong E^\vee$. By (9.3), $\det_{(1^k)}$ is surjective, and is given by

$$(9.5) \quad \det_{(1^k)}((v_{a_1} \otimes \cdots \otimes v_{a_k}) \otimes (u_{b_1} \otimes \cdots \otimes u_{b_k})) = x_{a_1 b_1} \cdots x_{a_k b_k}$$

for $1 \leq a_1, \dots, a_k, b_1, \dots, b_k \leq N$. Since $m: (V) \otimes V \rightarrow \bigwedge_k V$ (resp. $\Delta: \bigwedge_k V \rightarrow (V) \otimes V$) is the dual of $\Lambda: \bigwedge_k V \rightarrow V \otimes V$ (resp. $m: V \otimes V \rightarrow \bigwedge_k V$), the diagram

$$(9.6) \quad \begin{array}{ccccc} ({'}V)^{\otimes k} \otimes \bigwedge_k V & \xrightarrow{m \otimes 1} & \bigwedge_k {}^t V \otimes \bigwedge_k V & \xleftarrow{1 \otimes m} & \bigwedge_k {}^t V \otimes V^{\otimes k} \\ \downarrow 1 \otimes \Delta & & \downarrow \det_k & & \downarrow \Delta \otimes 1 \\ ({'}V)^{\otimes k} \otimes V^{\otimes k} & \xrightarrow{\det_{(1^k)}} & S_k E^\vee & \xleftarrow{\det_{(1^k)}} & ({'}V)^{\otimes k} \otimes V^{\otimes k} \end{array}$$

is commutative, thanks to the commutativity of the diagram (9.1). By the commutativity of the left (resp. right) square in (9.6) and Lemma 5.3, the first (resp. the second) equality in (9.4) holds. \square

LEMMA 9.3 (The Laplace expansion). *Let $i, j \geq 0$. Then, the following diagram commutes.*

$$\begin{array}{ccccc} \bigwedge_i {}^t V \otimes \bigwedge_j {}^t V \otimes \bigwedge_{i+j} V & \xrightarrow{m \otimes 1} & \bigwedge_{i+j} {}^t V \otimes \bigwedge_{i+j} V & \xleftarrow{1 \otimes m} & \bigwedge_{i+j} {}^t V \otimes \bigwedge_i V \otimes \bigwedge_j V \\ \downarrow 1 \otimes \Delta & & \downarrow \det_{i+j} & & \downarrow \Delta \otimes 1 \\ \bigwedge_i {}^t V \otimes \bigwedge_j {}^t V \otimes \bigwedge_i V \otimes \bigwedge_j V & \xrightarrow{\det_{(i,j)}} & S_{i+j} E^\vee & \xleftarrow{\det_{(i,j)}} & \bigwedge_i {}^t V \otimes \bigwedge_j {}^t V \otimes \bigwedge_i V \otimes \bigwedge_j V. \end{array}$$

PROOF. Since $m: \bigwedge_i {}^t V \otimes \bigwedge_j {}^t V \rightarrow \bigwedge_{i+j} {}^t V$ (resp. $\Delta: \bigwedge_{i+j} {}^t V \rightarrow \bigwedge_i {}^t V \otimes \bigwedge_j {}^t V$) is identified with the dual of $\Delta: \bigwedge_{i+j} V \rightarrow \bigwedge_i V \otimes \bigwedge_j V$ (resp. $m: \bigwedge_i V \otimes \bigwedge_j V \rightarrow \bigwedge_{i+j} V$), the lemma follows from the commutativity of (9.1). \square

LEMMA 9.4 (cf. [4, Proposition III.1.1]). *Let $k \geq 0$ and $\lambda \in \Omega_k^-$. For $\mu \in S_\square(\lambda)$, the following diagram commutes:*

$$(9.7) \quad \begin{array}{ccccc} \bigwedge_\mu {}^t V \otimes \bigwedge_\lambda V & \xrightarrow{1 \otimes \square_\lambda^\mu({'}V)^*} & \bigwedge_\mu {}^t V \otimes \bigwedge_\mu V & \xleftarrow{\square_\lambda^\mu(V)^* \otimes 1} & \bigwedge_\lambda {}^t V \otimes \bigwedge_\mu V \\ \downarrow \square_\lambda^\mu(V) \otimes 1 & & \downarrow \det_\mu & & \downarrow 1 \otimes \square_\lambda^\mu(V) \\ \bigwedge_\lambda {}^t V \otimes \bigwedge_\lambda V & \xrightarrow{\det_\lambda} & S_k E^\vee & \xleftarrow{\det_\lambda} & \bigwedge_\lambda {}^t V \otimes \bigwedge_\lambda V. \end{array}$$

This follows immediately from the commutativity of (9.1).

Let $M, N \in \text{cff-SE}^\vee$, and $\varphi: M \rightarrow N$ be an SE^\vee -isomorphism. Then, we have $\text{Im } \text{cf}_M = \text{Im } \text{cf}_N$ by the commutativity of (9.1).

LEMMA 9.5. *Let $\lambda = (\lambda_1, \dots, \lambda_s) \in \Omega^+$ and $\sigma \in \mathfrak{S}_s$. Then, we have $\bigwedge_\lambda V \simeq \bigwedge_{\sigma\lambda} V$ and $\text{Im } \det_\lambda = \text{Im } \det_{\sigma\lambda}$, where $\sigma\lambda = (\lambda_{\sigma^{-1}1}, \dots, \lambda_{\sigma^{-1}s})$. In particular, we have $\text{Im } \det_\lambda = \text{Im } \det_{\tilde{\lambda}}$.*

PROOF. It suffices to show that $\bigwedge_\lambda V \simeq \bigwedge_{\sigma\lambda} V$ by the observation above. But this is obvious, since the isomorphism $\psi_{\bigwedge V}(\sigma): (\bigwedge V)^{\otimes s} \rightarrow (\bigwedge V)^{\otimes s}$ maps $\bigwedge_\lambda V$ onto $\bigwedge_{\sigma\lambda} V$. \square

Let $\lambda \geq 0$ and $\lambda \in \Omega_k^-$. We define:

$$(9.8) \quad M(\lambda) = \sum_{\mu \in \Omega_k^-, \mu \geq \lambda} \text{Im } \det_\mu \subset S_k E^\vee$$

and

$$(9.9) \quad \dot{M}(\lambda) = \sum_{\mu \in \Omega_k^-, \mu > \lambda} \text{Im } \det_\mu \subset M(\lambda).$$

Note that $M(\lambda)$ and $\dot{M}(\lambda)$ are subbicomodules of $S_k E^\vee$, and that $\lambda, \mu \in \Omega_k^-$, $\mu \geq \lambda$ imply $M(\mu) \subset M(\lambda)$. By Lemma 9.2, we have $S_k E^\vee = M((1^k))$. Hence, we have a filtration

$$S_k E^\vee = M((1^k)) \supset M((2, 1^{k-2})) \supset \cdots \supset M((k)) \supset 0$$

of $S_k E^\vee$.

THEOREM 9.6 (Straightening formula). *Let $k \geq 0$ and $\lambda \in \Omega_k^-$. Then we have a unique isomorphism of SE^\vee -bicomodules $\Theta_\lambda: L_\lambda^t V \otimes L_\lambda V \rightarrow M(\lambda)/\dot{M}(\lambda)$ which makes the following diagram commutative:*

$$(9.10) \quad \begin{array}{ccc} \bigwedge_\lambda V \otimes \bigwedge_\lambda V & \xrightarrow{\det_\lambda} & M(\lambda) \\ \downarrow d_\lambda \otimes d_\lambda & & \downarrow \text{proj.} \\ L_\lambda^t V \otimes L_\lambda V & \xrightarrow{\Theta_\lambda} & M(\lambda)/\dot{M}(\lambda). \end{array}$$

Hence, the associated graded object of the filtration $\{M(\lambda)\}_{\lambda \in \Omega_k^-}$ is $\bigoplus_{\lambda \in \Omega_k^-} L_\lambda^t V \otimes L_\lambda V$. The set $\{\det_\lambda(\xi_S \otimes \xi_T) \mid \lambda \in \Omega_k^-, S \in \text{St}_\lambda Y, T \in \text{St}_\lambda^t Y\}$ is a free basis of $S_k E^\vee$, where $Y = \{v_1 < \cdots < v_N\}$ is the dual basis of $Y = \{u_1 < \cdots < u_N\}$.

PROOF. Let $v \in S_{\square}(\lambda)$. Then, we have $\text{Im } \det_v = \text{Im } \det_{\tilde{v}}$ by Lemma 9.5. Since $\tilde{v} \sim \lambda$, we have $\text{Im } \det_v \subset \dot{M}(\lambda)$. By Lemma 9.4, we have $\det_\lambda(\text{Im}(1 \otimes \square_\lambda^*)) \subset \dot{M}(\lambda)$ and $\det_\lambda(\text{Im}(\square_\lambda^* \otimes 1)) \subset \dot{M}(\lambda)$. Hence, we have $\det_\lambda(\text{Im}(\square_\lambda^* \otimes 1) + \text{Im}(1 \otimes \square_\lambda)) \subset \dot{M}(\lambda)$. By Theorem 6.19, we have $\text{Ker}(d_\lambda \otimes d_\lambda) = \text{Im}(\square_\lambda^* \otimes 1) + \text{Im}(1 \otimes \square_\lambda)$. Hence, we have a unique induced map $\Theta_\lambda: L_\lambda^t V \otimes L_\lambda V \rightarrow M(\lambda)/\dot{M}(\lambda)$ which makes (9.10) commutative. By the definition of $M(\lambda)$ and $\dot{M}(\lambda)$, the composite map $\text{proj} \circ \det_\lambda$ is surjective, and so is Θ_λ . We set $\Gamma(\lambda) = \{\det_\mu(\xi_S \otimes \xi_T) \mid \mu \in \Omega_k^-, \mu \geq \lambda, S \in \text{St}_\mu Y, T \in \text{St}_\mu^t Y\}$ for $\lambda \in \Omega_k^-$. By induction on λ , it is easy to see that $\Gamma(\lambda)$ generates $M(\lambda)$, thanks to Theorem 6.19. On the other hand, we have $\text{rank } S_k E^\vee = \sum_{\lambda \in \Omega_k^-} (\#(\text{St}_\lambda Y))^2 = \#\Gamma(1^k)$. To prove this, we may assume that $R = C(q)$, and this case is a consequence of Corollary 8.11. It follows that $\Gamma(1^k)$ is linearly independent, and $\Gamma(\lambda)$ is a free basis of $M(\lambda)$ for $\lambda \in \Omega_k^-$. Hence, $\{\Theta_\lambda(d_\lambda \xi_S \otimes d_\lambda \xi_T) \mid S \in \text{St}_\lambda Y, T \in \text{St}_\lambda^t Y\}$ is linearly independent, and Θ_λ is injective for $\lambda \in \Omega_k^-$. Since \det_λ , proj and $d_\lambda \otimes d_\lambda$ are SE^\vee -bicomodule homomorphisms, so is Θ_λ . The theorem is now clear. \square

Let $N \in \mathcal{D}$, and $v \in \Omega^+$ with $\ell(v) \leq N$. We define ${}_v N := N t^{*(v)} \subset N$. By Lemma 7.8 and its proof, we have ${}_v N \simeq (N_v^*)^* \simeq (\text{tr}_\# N)_v$. For $\lambda \in \Omega_k^-$ with $\ell(\tilde{\lambda}) = \lambda_1 \leq N$, we set $c'(\lambda) := v_1 \wedge \cdots \wedge v_{\lambda_1} \otimes \cdots \otimes v_1 \wedge \cdots \wedge v_{\lambda_N} \in \bigwedge_\lambda V$. It is easy to see that $c'(\lambda) \in {}_{\tilde{\lambda}}(\bigwedge_\lambda V)$, and $d_\lambda(c'(\lambda))$ generates ${}_{\tilde{\lambda}}(L_\lambda^t V) = (K_{\tilde{\lambda}} V)_{\tilde{\lambda}}^* \simeq R$. We set

$$(9.11) \quad L_q^\lambda := \det_\lambda(c(\lambda) \cdot R \otimes \bigwedge_\lambda V) \subset S_k E^\vee.$$

This definition is due to Taft and Towber [37, §3]. Clearly, L_q^λ is a quotient $S_k E^\vee$ -comodule of $\bigwedge_\lambda V$ via the surjection

$$\bigwedge_\lambda V \xrightarrow{\sim} c'(\lambda)R \otimes \bigwedge_\lambda V \xrightarrow{\det_\lambda} L_q^\lambda V,$$

and is a subcomodule of $S_k E$.

PROPOSITION 9.7. *Let $\lambda \in \Omega_k^-$ and $\ell(\lambda) \leq N$. Then we have $L_q^\lambda \simeq L_\lambda V$ as an $S_k E^\vee$ -comodule.*

PROOF. For $\mu \in \Omega_k^-$ with $\ell(\lambda) \leq N$, we have $\mu > \lambda \Leftrightarrow \tilde{\mu} < \tilde{\lambda}$ (see [25, (1.11)]). Hence, $\mu > \lambda$ implies $\tilde{\mu} \not\leq \tilde{\lambda}$. By Theorem 7.12, we have ${}_{\tilde{\lambda}}(L_\mu^{-1}V) = (K_{\tilde{\mu}}V)_{\tilde{\lambda}}^* = 0$ for $\mu \in \Omega_k^-$ with $\ell(\mu) \leq N$ and $\mu > \lambda$. By Theorem 9.6, we have ${}_{\tilde{\lambda}}M(\lambda) = 0$. Hence, the projection $\text{proj}: {}_{\tilde{\lambda}}M(\lambda) \rightarrow {}_{\tilde{\lambda}}(M(\lambda)/\dot{M}(\lambda))$ is an isomorphism. Since L_q^λ is contained in ${}_{\tilde{\lambda}}M(\lambda)$, we have

$$\begin{aligned} L_q^\lambda &= \text{proj} \circ \det_\lambda(c'(\lambda) \cdot R \otimes \bigwedge_\lambda V) = \Theta_\lambda \circ (d_\lambda \otimes d_\lambda)(c'(\lambda) \cdot R \otimes \bigwedge_\lambda V) \\ &= d_\lambda(c'(\lambda)) \cdot R \otimes L_\lambda V \simeq L_\lambda V. \end{aligned}$$

□

REFERENCES

- [1] E. ABE, Hopf algebras, Cambridge Tracts in Math. 74, Cambridge Univ. Press, 1980.
- [2] K. AKIN AND D. A. BUCHSBAUM, Characteristic-free representation theory of the general linear group, Adv. in Math. 58 (1985), 149–200.
- [3] K. AKIN AND D. A. BUCHSBAUM, Characteristic-free representation theory of the general linear group II Homological considerations, Adv. in Math. 72 (1988), 171–210.
- [4] K. AKIN, D. A. BUCHSBAUM AND J. WEYMAN, Schur functors and Schur complexes, Adv. in Math. 44 (1982), 207–278.
- [5] G. M. BERGMAN, The diamond lemma for ring theory, Adv. in Math. 29 (1978), 178–218.
- [6] C. CURTIS AND I. REINER, Representation theory of finite groups and associative algebras, Pure and Appl. Math., vol. 11, Interscience, New York, 1962; 2nd ed., 1966.
- [7] V. G. DRINFELD, Quantum groups, Proc. ICM, Berkeley, 1986.
- [8] E. DATE, M. Jimbo and T. Miwa, Representations of $U_q(\mathfrak{gl}(n, C))$ at $q=0$ and the Robinson-Schensted correspondence, Memorial Volume for Vadim Kniznik, Physics and Mathematics of Strings, World Scientific, Singapore, 1990, 185–211.
- [9] P. DELIGNE AND J. MILNE, Tannakian categories, Lecture Notes in Math. 900, Springer-Verlag, Berlin Heidelberg New York, 1982, 101–228.
- [10] R. DIPPER AND G. JAMES, q -tensor space and q -Weyl modules, preprint.
- [11] P. DOOBILET, G.-C. ROTA AND J. STEIN, Foundations of combinatorics. IX. Combinatorial methods in invariant theory, Stud. Appl. Math. 43 (1971), 1020–1058.
- [12] L. D. FADDEEV, N. YU. RESHETIKHIN AND L. A. TAKHTAJAN, Quantization of Lie groups and Lie algebras, Algebra and Analysis 1 (1989), 178–206 (English translation: Leningrad Math. J. 1 (1990), 193–225).

- [13] J. A. GREEN, Polynomial representations of GL_n , Lecture Notes in Math. 830, Springer-Verlag, Berlin Heidelberg New York, 1980.
- [14] A. GYOJA, A q -analogue of Young symmetrizer, Osaka J. Math. 23 (1986), 841–852.
- [15] A. GYOJA AND K. UNO, On the semisimplicity of Hecke algebras, J. Math. Soc. Japan 41 (1989), 75–79.
- [16] T. HIBI, Distributive lattices, affine semigroup rings and algebras with straightening laws, in Advanced Studies in Pure Math. 11, Kinokuniya, Tokyo and North-Holland, Amsterdam, 1987, 93–109.
- [17] T. HAYASHI, Quantum deformation of classical groups, Publ. Res. Inst. Math. Sci., Kyoto Univ. 28 (1992), 57–81.
- [18] M. JIMBO, A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation, Lett. in Math. Phys. 10 (1985), 63–69.
- [19] M. JIMBO, Quantum R matrix related to the generalized Toda system: an algebraic approach, Lecture Notes in Phys. 246, Springer-Verlag, Berlin Heidelberg New York, 1986, 335–361.
- [20] M. JIMBO, A q -analogue of $U(\mathfrak{gl}(N+1))$, Hecke algebra and the Yang-Baxter equation, Lett. in Math. Phys. 11 (1986), 247–252.
- [21] P. P. KULISH, N. YU. RESHETIKHIN AND E. K. SKLYANIN, Yang-Baxter equation and representation theory. I, Lett. Math. Phys. 5 (1981), 393–403.
- [22] P. P. KULISH AND E. K. SKLYANIN, Quantum spectral transform method. Recent developments, Integrable quantum field theories, Lecture Notes in Physics 151, Springer-Verlag, Berlin Heidelberg New York, 1981, 61–119.
- [23] G. LUSZTIG, Modular representations and quantum groups, Contemp. Math. 82, Amer. Math. Soc., Providence, R. I., 1989, 59–77.
- [24] V. V. LYUBASHENKO, Hopf algebras and vector-symmetries, Uspekhi Mat. Nauk, 41–5 (1986), 185–186 (in Russian).
- [25] I. G. MACDONALD, Symmetric functions and Hall polynomials, Oxford Univ. Press, 1979.
- [26] YU. I. MANIN, Some remarks on Koszul algebras and quantum groups, Ann. Inst. Fourier, Grenoble 37 (1987), 191–205.
- [27] YU. I. MANIN, Multiparametric quantum deformation of the general linear supergroup, Commun. Math. Phys. 123 (1989), 163–175.
- [28] D. G. NORTHCOTT, Multilinear algebra, Cambridge University Press, 1984.
- [29] M. NOUMI, H. YAMADA AND K. MIMACHI, Finite dimensional representations of the quantum group $GL_q(n)$ and the zonal spherical functions on $U_q(n-1) \setminus U_q(n)$, preprint.
- [30] B. PARSHALL AND J. WANG, Quantum linear groups, Memoirs of the Amer. Math. Soc. 439 (1991).
- [31] P. PODLES, Quantum spheres, Lett. in Math. Phys. 14 (1987), 193–202.
- [32] S. B. PRIDDY, Koszul resolutions, Trans. Amer. Math. Soc. 152–1 (1970), 39–60.
- [33] N. YU. RESHETIKHIN, Quantum universal enveloping algebras, the Yang-Baxter equation and invariants of links. I, preprint.
- [34] M. SWEEDLER, Hopf algebras, W. A. Benjamin, New York, 1969.
- [35] M. TAKEUCHI, Matric bialgebras and quantum groups, Israel J. of Math. 72 (1990), 232–251.
- [36] T. TANISAKI, Finite dimensional representations of quantum groups, Osaka J. of Math. 28 (1991), 37–54.
- [37] E. TAFT AND J. TOWBER, Quantum deformation of flag schemes and Grassmann schemes I A q -deformation of the shape-algebra for $GL(n)$, J. Algebra 142 (1991), 1–36.
- [38] V. G. TURAEV, The Yang-Baxter equation and invariant of links, Invent. Math. 92 (1988), 527–553.
- [39] K. UENO, T. Takebayashi, Y. Shibukawa, Gelfand-Tsetlin basis for representations of the quantum group $GL_q(N+1)$ and the basic quantum affine space, preprint.
- [40] S. L. WORONOWICZ, Compact matrix pseudogroups, Comm. Math. Phys. 111 (1987), 613–665.

DEPARTMENT OF MATHEMATICS
SCHOOL OF SCIENCE
NAGOYA UNIVERSITY
NAGOYA 464
JAPAN

