### Quasi m-Cayley strongly regular graphs

### Klavdija Kutnar

#### University of Primorska, Koper, Slovenia

Feb, 2012

# Joint work with Luis Martinez Fernandez, Aleksander Malnič and Dragan Marušič.

A regular graph X of valency k and order v is called strongly regular graph (SRG) with parameters  $(v, k, \lambda, \mu)$  if any two adjacent vertices have  $\lambda$  common vertices and any two distinct non-adjacent vertices have  $\mu$  common vertices. For a prime power q such that  $q \equiv 1 \pmod{4}$  the Paley graph P(q) is a graph with vertex set  $\mathbb{F}_q$  in which two vertices are adjacent if their difference is non zero square.

P(q) is a strongly regular graph with parameters

$$(v, k, \lambda, \mu) = (q, \frac{q-1}{2}, \frac{q-5}{4}, \frac{q-1}{4}).$$

伺 とう きょう とう とう

A graph X is an *m*-Cayley graph on a group H if its automorphism group admits a semiregular subgroup H having *m* orbits, all of equal length.

If H is cyclic and

- m = 1 then X is said to be circulant;
- m = 2 then X is said to be bicirculant;
- m = 3 then X is said to be tricirculant.

A non-identity automorphism of a graph with m cycles of equal length n in its cycle decomposition is said to be (m, n)-semiregular.

(4) (3) (4) (3) (4)

## Example



æ

▲□→ ▲圖→ ▲厘→ ▲厘→

## Example



a (2,5)-semiregular automorphism  $\rho = (12345)(1'2'3'4'5')$ 

문 🛌 문

## Example



a (2,5)-semiregular automorphism  $\rho = (12345)(1'2'3'4'5')$ 



< 注 → 注

< E.

A particular attention has been given to questions regarding strong regularity for various classes of graphs satisfying certain special symmetry conditions.

strongly regular Cayley graphs, strongly regular 2-Cayley graphs, strongly regular 3-Cayley graphs (Bridges, Mena, Leung, Ma, Marušič, Miklavič, Šparl, de Resmini, Jungnickel, ...)

For example, by a classical result of Bridges and Mena (1979) it is known that the Paley graphs are the only SRGs among Cayley graphs on cyclic groups.

・ 回 と ・ ヨ と ・ ヨ と

A group G acts quasi-semiregularly on a set V if there exists an element  $\infty$  in V such that the stabilizer  $G_{\infty}$  of the element  $\infty$  in G is equal to G, and the stabilizer  $G_{v}$  of any element  $v \in V - \{\infty\}$  in G is trivial. The element  $\infty$  is called the point at infinity.

• E • • E • •

A group G acts quasi-semiregularly on a set V if there exists an element  $\infty$  in V such that the stabilizer  $G_{\infty}$  of the element  $\infty$  in G is equal to G, and the stabilizer  $G_{v}$  of any element  $v \in V - \{\infty\}$  in G is trivial. The element  $\infty$  is called the point at infinity.

A graph X is a quasi m-Cayley graph on a group H if its automorphism group admits a semiregular subgroup H having morbits, all of equal length.

If H is cyclic and

- m = 1 then X is said to be quasi circulant;
- m = 2 then X is said to be quasi bicirculant;
- m = 3 then X is said to be quasi tricirculant;

• ...

・ 同 ト ・ ヨ ト ・ ヨ ト

 If n = mk + 1 ≥ 2 then the complete graph K<sub>n</sub> is a quasi m-Cayley graph on a cyclic group Z<sub>k</sub>.

▲圖▶ ▲屋▶ ▲屋▶

æ

- If n = mk + 1 ≥ 2 then the complete graph K<sub>n</sub> is a quasi m-Cayley graph on a cyclic group Z<sub>k</sub>.
- Paley graphs P(q) are quasi bicirculants.

(4回) (4回) (4回)

2

The symbol of a quasi *m*-Cayley graph in the following way:

Let X be a quasi *m*-Cayley graph on a group G and let  $\{U_0, \ldots, U_{m-1}\}$ be the set of *m* orbits of G on  $V(X) - \{\infty\}$ . Let  $u_i \in U_i$ ,  $i \in \mathbb{Z}_m$ , let  $S_{i,j}$ ,  $i,j \in \mathbb{Z}_m$  be defined by  $S_{i,j} = \{\rho \in G \mid u_i \to \rho(u_j)\}$ , and let  $S_{\infty} \subseteq \mathbb{Z}_m$  be defined by

$$S_{\infty} = \{i \in \mathbb{Z}_m \mid U_i \subseteq N(\infty)\}.$$

Then the family  $(S_{i,j})$  together with  $S_{\infty}$  is called the symbol of X relative to  $(G; u_0, \ldots, u_{m-1}, S_{\infty})$ , and determines the adjacencies in the graph. We can assume by renumbering the orbits if necessary that  $S_{\infty} = \{0, \ldots, s-1\}$  for some s.

▲圖 ▶ ▲ 国 ▶ ▲ 国 ▶

## Quasi-partial difference family

Let G be a group of order n and m, s positive integers with  $s \leq m$ . A family  $\{S_{i,j}\}$  of subsets of G, with  $0 \leq i, j \leq m-1$  that satisfy  $0 \notin S_{i,i} \forall i$  and  $S_{j,i} = -S_{i,j} \forall i, j$  with  $i \neq j$ , is said to be an  $(m, n, s, \lambda, \mu)$  quasi-partial difference family if

$$\sum_{j=1}^{m} |S_{i,j}| = \begin{cases} ns-1 & \text{if } i \leq s \\ ns & \text{otherwise} \end{cases}, \sum_{j=1}^{s} |S_{i,j}| = \begin{cases} \lambda & \text{if } i \leq s \\ \mu & \text{otherwise} \end{cases}$$
(1)

and if the following identities hold in the group ring  $\mathbb{Z}[G]$ :

$$\sum_{k=0}^{m-1} S_{i,k} S_{k,j} = \delta_{i,j} \gamma\{0\} + \beta S_{i,j} + \mu' G,$$
(2)

where  $\delta_{i,j}$  is the Kronecker delta,  $\gamma = ns - \mu$ ,  $\beta = \lambda - \mu$  and

$$\mu' = egin{cases} \mu - 1 & ext{if } i,j \leq s \ \mu & ext{otherwise.} \end{cases}$$

御 と く き と く き と

#### Proposition

The quasi *m*-Cayley graph defined by the symbol  $(S_{i,j})$  with  $S_{\infty} = \{0, \ldots, s-1\}$  is an  $(mn + 1, ns, \lambda, \mu)$ -SRG iff  $(S_{i,j})$  forms an  $(m, n, s, \lambda, \mu)$  quasi-partial difference family.

イロン イ部ン イヨン イヨン 三日

An example of a (3, 3, 1, 0, 1) quasi-partial difference family on the cyclic group  $C_3$ :

$$S_{0,0} = \emptyset$$
,  $S_{0,1} = \{0\}$ ,  $S_{0,2} = \{0\}$ ,  
 $S_{1,1} = \emptyset$ ,  $S_{1,2} = \{1,2\}$ ,  $S_{2,2} = \emptyset$ .

個 と く ヨ と く ヨ と …

æ

An example of a (3, 3, 1, 0, 1) quasi-partial difference family on the cyclic group  $C_3$ :

$$S_{0,0} = \emptyset$$
,  $S_{0,1} = \{0\}$ ,  $S_{0,2} = \{0\}$ ,  
 $S_{1,1} = \emptyset$ ,  $S_{1,2} = \{1,2\}$ ,  $S_{2,2} = \emptyset$ .

This quasi-partial difference family generates the Petersen graph.

個 と く ヨ と く ヨ と …

#### Theorem

Let G be a cyclic group. If  $\{S_{i,j}\}$  is a (m, n)-circulant  $(m, n, s, \lambda, \mu)$  quasi-partial difference family and if one of the following three conditions is satisfied:

- 1 n is a prime
- 2 *n* is coprime to  $(m!)\Delta$

then  $\{S_{i,i}\}_{0 \le i \le m-1}$  covers all the elements of  $G - \{0\}$  the same number of times.

・ 回 ト ・ ヨ ト ・ ヨ ト

#### Theorem

If dm + 1 is a prime power, then a uniform

$$(m, d(md + 2), 1, d^2 - md + 3d - 1, d(d + 1))$$

QPDF exists.

・ロト ・回ト ・ヨト ・ヨト

æ

#### Theorem

If dm + 1 is a prime power, then a uniform

$$(m, d(md + 2), 1, d^2 - md + 3d - 1, d(d + 1))$$

QPDF exists.

An  $(m, n, 1, \lambda, \mu)$  QPDF  $\{S_{i,j}\}$  on a cyclic group  $C_n$  is uniform if the following three conditions hold:

- $\cup_{i=0}^{m-1}S_{i,i} = G \{0\}.$
- all the cardinalities  $|S_{i,i}|$  with  $i \ge 1$  are equal.
- all the cardinalities  $|S_{i,j}|$  with  $i,j \ge 1$  and  $i \ne j$  are equal.

(ロ) (同) (E) (E) (E)

Thank you!

æ