

 Open access • Journal Article • DOI:10.2298/FIL0702243K

Quasi-nearly subharmonic functions and conformal mappings — [Source link](#)

Vesna Kojić

Published on: 01 Jan 2007 - Filomat (National Library of Serbia)

Topics: Subharmonic function and Conformal map

Related papers:

- [Classes of Quasi-nearly Subharmonic Functions](#)
- [H_p spaces of several variables](#)
- [A generalized mean value inequality for subharmonic functions](#)
- [Subharmonic functions, generalizations and separately subharmonic functions](#)
- [Quasi-nearly subharmonic functions and quasiconformal mappings](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/quasi-nearly-subharmonic-functions-and-conformal-mappings-4xxz73wr97>

QUASI-NEARLY SUBHARMONIC FUNCTIONS AND CONFORMAL MAPPINGS

Vesna Kojić

Abstract

If φ is a conformal mapping and u is a quasi-nearly subharmonic function, then $u \circ \varphi$ is quasi-nearly subharmonic. A similar fact for “regularly oscillating” functions holds.

Introduction

If u is a nonnegative subharmonic function on a domain $\Omega \subset \mathbb{C}$ and $p \geq 1$, then

$$u(z)^p \leq \frac{1}{r^2} \int_{B(z,r)} u^p dm \quad (1)$$

for all $B(x, r) \subset \Omega$. Here $B(z, r)$ is the Euclidean disk with center z and radius r , and dm is the Lebesgue measure in \mathbb{C} normalized so that the measure of the unit ball equals one. If $0 < p < 1$, then (1) need not hold but there is a constant $C = C(p) \geq 1$ such that

$$u(z)^p \leq \frac{C}{r^2} \int_{B(z,r)} u^p dm \quad (2)$$

This fact, essentially due to Hardy and Littlewood [5], was first proved by Fefferman and Stein [3, Lemma 2, p. 172]. Fefferman and Stein’s proof is reproduced in Garnett [4, Lemma 3.7, pp. 121–123]. Although Fefferman

2000 *Mathematics Subject Classification*. Primary: 31B05, 31B25; Secondary: 31C05.

Key words and phrases. Subharmonic, quasi-nearly subharmonic, and regularly oscillating functions, Koebe’s theorems.

Received: August 20, 2007

The author is supported by MNZŽS Grant ON144010, Serbia

and Stein considered only the case when $u = |v|$ and v is harmonic their proof applies also in the case of nonnegative subharmonic functions. Perhaps, the first simple proof of (2) was given in [10, p. 64], although it depended on the hypothesis that u was subharmonic. For other proofs see [13, p. 18, and Theorem 1, p. 19], (see also [14, Theorem A, p. 15]), [18, Lemma 2.1, p. 233], and [19, Theorem, p. 188]).

That the validity of (2) for some p implies its validity for all p was observed in [1, p. 132], [13, Theorem 1], and [18, Lemma 2.1]. See also [11], where the case of “ M -subharmonic” functions was considered, and [8, 9], where some extensions of [11] were made.

For various applications of (2) we refer to [24, p. 191], [6, Theorems 1 and 2, pp. 117-118], [21, Theorems 1, 2 and 3, pp. 301, 307], [18, Theorem, p. 233], [22, Theorem 2, p. 271], [23, Theorem, p. 113], [15], [12], and [17]. Further information can be found in [19] and [17].

Quasi-nearly subharmonic functions

Let Ω be a subdomain of the complex plane \mathbb{C} . Following [20] and [17], we call a Borel measurable function $u : \Omega \rightarrow [0, \infty]$ *quasi-nearly subharmonic*, if $u \in L^1_{\text{loc}}(\Omega)$ and if there is a constant $K = K(u, \Omega) \geq 1$ such that

$$u(z) \leq \frac{K}{r^2} \int_{B(z,r)} u(w) dm(w) \quad (3)$$

for any disk $B(z, r) \subset \Omega$.

In [18], the term *pseudoharmonic functions* is used, while in [13], condition (3) is called *sh $_K$ -condition*. If $K = 1$ (and u takes its values in $[-\infty, \infty]$), then u is called *nearly subharmonic* (see [7]).

We will denote by $QNS_K(\Omega)$ the class of all functions satisfying (3) (for a fixed K) and by $QNS(\Omega)$ the class of all quasi-nearly subharmonic functions defined in Ω ; so

$$QNS(\Omega) = \bigcup_{K \geq 1} QNS_K(\Omega).$$

One of the most important properties of QNS is the following fact, which generalizes the above mentioned result of Fefferman and Stein [3].

Theorem A. [1, 13, 18] *If $u \in QNS_K(\Omega)$, and $p > 0$, then $u^p \in QNS_C(\Omega)$, where C is a constant depending only on p, K . In particular, if u^p is quasi-nearly subharmonic for some $p > 0$, then so is for every $p > 0$.*

In this paper we prove the following:

Theorem 1. *Let $u \in QNS_K(\Omega)$ and φ a conformal mapping from a domain G onto Ω , then the composition $u \circ \varphi$ belongs $QNS_C(G)$, where C depends only on K .*

Regularly oscillating functions

A function f defined in Ω is called regularly oscillating (see [17]) if f is of class $C^1(\Omega)$ and

$$|\nabla f(z)| \leq Kr^{-1} \sup_{B(z,r)} |f - f(z)|, \quad B(z,r) \subset \Omega. \quad (4)$$

The class of such functions is denoted in [13] and [16] by $OC_K^1(\Omega)$ (O = oscillation). The class of all regularly oscillating functions will be denoted by $RO(\Omega)$.

Theorem B. *[13, Theorem 3] If f is regularly oscillating, then $|f|$ and $|\nabla f|$ are quasi-nearly subharmonic. Moreover, if $f \in OC_K^1(\Omega)$, then $|f|$ and $|\nabla f|$ are in $QNS_C(\Omega)$, where C depends only on K .*

Example 1. Harmonic functions are regularly oscillating.

Example 2. [13] Convex functions are regularly oscillating. It follows that the modulus of the gradient of a convex function is quasi-nearly subharmonic.

Example 3. [14] If f is an eigenfunction of Δ , i.e., $\Delta f = \lambda f$ for some constant λ , and if Ω is bounded, then $f \in OC^2(\Omega)$.

Example 4 (Polyharmonic functions). A function $f \in C^{2k}(\Omega)$ is said to be polyharmonic (of degree k) if it is annihilated by the k -th power of the Laplacian. It is proved in [14, Corollary 5] (see also [15]) that every polyharmonic function is regularly oscillating, and therefore $|f|$ and $|\nabla f|$ are quasi-nearly subharmonic.

Here we prove the following:

Theorem 2. *If $f \in RO(\Omega)$, and φ is a conformal mapping from G onto Ω , then $f \circ \varphi \in G$. Moreover if $f \in OC_K^1(\Omega)$, then $f \circ \varphi$ is in $OC_C^1(G)$, where C depends only on K .*

Proofs

Our proofs are based on two theorems of Koebe (see [2, Theorem 2.3 and Theorem 2.5]).

Theorem C (Koebe one-quarter theorem). *Let φ be a conformal mapping from the disk $B(z_0, R)$ into \mathbb{C} , then the image $\varphi(B(z_0, R))$ contains the disk $B(\varphi(z_0), \rho)$, where $\rho = R|\varphi'(z_0)|/4$.*

Theorem D (Koebe distortion theorem). *Let φ be a conformal mapping from the disk $B(z_0, R)$ into \mathbb{C} , then there holds the inequalities*

$$\frac{R^2(R - |z - z_0|)}{(R + |z - z_0|)^3} \leq \frac{|\varphi'(z)|}{|\varphi'(z_0)|} \leq \frac{R^2(R + |z - z_0|)}{(R - |z - z_0|)^3}, \quad z \in B(z_0, R).$$

Consequently if $|z - z_0| < R/2$, then

$$\frac{|\varphi'(z)|}{|\varphi'(z_0)|} \geq \frac{4}{27}.$$

Proof of Theorem 1.

Let $u \in QNS_K(\Omega)$ and φ a conformal mapping from G onto Ω . We have to find a constant C such that

$$\int_{B(z_0, r)} u(\varphi(z)) \, dm(z) \geq u(\varphi(z_0))r^2/C, \quad (\dagger)$$

whenever $r < \text{dist}(z, \partial G)$. Let $w_0 = \varphi(z_0)$ and $\rho = r|\varphi'(z_0)|/4$, and let $\psi : \Omega \mapsto G$ denote the inverse of φ . Then

$$\begin{aligned} \int_{B(z_0, r)} u(\varphi(z)) \, dm(z) &= \int_{\varphi(B(z_0, r))} u(w)|\psi'(w)|^2 \, dm(w) \\ &\geq \int_{B(w_0, \rho/2)} u(w)|\psi'(w)|^2 \, dm(w), \end{aligned} \quad (5)$$

where we have applied the one-quarter theorem. Now we apply the distortion theorem to the function ψ to get $|\psi'(w)| \geq (4/27)|\psi'(w_0)|$, for $|w - w_0| < \rho/2$. It follows that

$$\begin{aligned} \int_{B(z_0, r)} u(\varphi(z)) \, dm(z) &\geq (4/27)^2 |\psi'(w_0)|^2 \int_{B(w_0, \rho/2)} u(w) \, dm(w) \\ &\geq (4/27)^2 |\psi'(w_0)|^2 (\rho/2)^2 u(w_0)/K \\ &= (4/27)^2 |\psi'(w_0)|^2 |\varphi'(z_0)|^2 u(w_0)r^2/16K. \end{aligned} \quad (6)$$

Now we use the identity $\psi'(w_0)\varphi'(z_0) = 1$ to get (†) with $C = 27^2K$. This concludes the proof of Theorem 1.

Proof of Theorem 2.

Let $u \in OC_K^1(\Omega)$ and φ a conformal mapping from G onto Ω . We have to find a constant C_1 such that

$$|\nabla u(\varphi(z_0))| \cdot |\varphi'(z_0)| \leq \frac{C_1}{\varepsilon} \sup_{z \in B(z_0, \varepsilon)} |u(\varphi(z)) - u(\varphi(z_0))|, \quad B(z_0, \varepsilon) \subset G. \quad (7)$$

Let $w_0 = \varphi(z_0)$ and $\rho = \varepsilon|\varphi'(z_0)|/4$, and let ψ be the inverse of φ . Then, the definition of OC_K^1 and by the one-quarter theorem,

$$\begin{aligned} & \sup\{|u(\varphi(z)) - u(\varphi(z_0))| : z \in B(z_0, \varepsilon)\} \\ &= \sup\{|u(w) - u(w_0)| : w \in \varphi(B(z_0, \varepsilon))\} \\ &\geq \sup\{|u(w) - u(w_0)| : w \in B(w_0, \rho)\} \\ &\geq |\nabla u(w_0)|\rho/K \\ &= |\nabla u(w_0)| \cdot |\varphi'(z_0)|\varepsilon/4K. \end{aligned}$$

This gives (7) with $C_1 = 4K$, concluding the proof.

References

- [1] P. Ahern and J. Bruna, *Maximal and area integral characterizations of Hardy-Sobolev spaces in the unit ball of \mathbb{C}^n* , Revista Mat. Iberoamericana 4(1988), 123–153.
- [2] P. L. Duren, *Univalent functions*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 259. Springer-Verlag, New York, 1983. xiv+382 pp.
- [3] C. Fefferman and E.M. Stein, *H^p spaces of several variables*, Acta Math. 129(1972), 137–193.
- [4] J.B. Garnett, *Bounded analytic functions*, Academic Press, New York, 1981.
- [5] G.H. Hardy and J.E. Littlewood, *Some properties of conjugate functions*, J. Reine Angew. Math. 167(1931), 405–423.

- [6] D.J. Hallenbeck, *Radial growth of subharmonic functions*, Pitman Research Notes 262, 1992, pp. 113–121.
- [7] M. Hervé, *Analytic and Plurisubharmonic Functions in Finite and Infinite Dimensional Spaces*, Lecture Notes in Mathematics 198, Springer-Verlag, Berlin, 1971.
- [8] M. Jevtić and M. Pavlović, *M-Besov p -classes and Hankel operators in the Bergman spaces on the unit ball*, Arch. Math. 61(1993), 367–376.
- [9] M. Jevtić and M. Pavlović, *Subharmonic behavior of generalized (α, β) -harmonic functions and their derivatives*, Indian J. Pure Appl. Math. 30(1999), 407–418.
- [10] M. Pavlović, *Mean values of harmonic conjugates in the unit disc*, Complex Variables 10(1988), 53–65.
- [11] M. Pavlović, *Inequalities for the gradient of eigenfunctions of the invariant Laplacian on the unit ball*, Indag. Math. (N.S.) 2(1991), 89–98.
- [12] M. Pavlović, *A proof of the Hardy–Littlewood theorem on fractional integration and a generalization*, Publ. Inst. Math. (Belgrade) 59(1996), 31–38.
- [13] M. Pavlović, *On subharmonic behavior and oscillation of functions on balls in \mathbb{R}^n* , Publ. Inst. Math. (Beograd) 55(69)(1994), 18–22.
- [14] M. Pavlović, *Subharmonic behavior of smooth functions*, Math. Vesnik 48(1996), 15–21.
- [15] M. Pavlović, *Decompositions of L^p and Hardy spaces of polyharmonic functions*, J. Math. Anal. Appl. 216(1997), 499–509.
- [16] M. Pavlović, *Introduction to function spaces on the disk*, Posebna izdanja 20, Matematički Institut SANU, 2004.
- [17] M. Pavlović and J. Riihenta, *Classes of quasi-nearly subharmonic functions*, preprint
- [18] J. Riihenta, *Subharmonic functions: non-tangential and tangential boundary behavior*, Function Spaces, Differential Operators and Non-linear Analysis, Proceedings of the Syöte Conference 1999 (eds. V. Mustonen and J. Rákosnik), Math. Inst., Czech Acad. Science, Praha, 2000, pp. 229–238.

- [19] J. Riihenta, *A generalized mean value inequality for subharmonic functions*, Expo. Math. 19(2001), 187–190.
- [20] J. Riihenta, *A generalized mean value inequality for subharmonic functions and applications*, arXiv:math.CA/0302261 v1 21 Feb 2003.
- [21] M. Stoll, *Weighted tangential boundary limits of subharmonic functions on domains in \mathbb{R}^n ($n \geq 2$)*, Math. Scand. 83(1998), 300–308.
- [22] N. Suzuki, *Nonintegrability of harmonic functions in a domain*, Japan. J. Math. 16(1990), 269–278.
- [23] N. Suzuki, *Nonintegrability of superharmonic functions*, Proc. Amer. Math. Soc. 113(1991), 113–115.
- [24] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Academic Press, London, 1986.

Fakultet organizacionih nauka, Jove Ilića 154, Belgrade, Serbia

E-mail: vesnak@fon.bg.ac.yu