

Published in final edited form as:

*J Am Chem Soc.* 2011 December 7; 133(48): 19350–19353. doi:10.1021/ja2093579.

## Radical Cation Diels–Alder Cycloadditions by Visible Light Photocatalysis

Shishi Lin, Michael A. Ischay, Charles G. Fry, and Tehshik P. Yoon\*


Department of Chemistry, University of Wisconsin–Madison, 1101 University Avenue, Madison, WI 53706.

### Abstract

Ruthenium(II) polypyridyl complexes promote the efficient radical cation Diels–Alder cycloaddition of electron-rich dienophiles upon irradiation with visible light. These reactions enable facile [4+2] cycloadditions that would be electronically mismatched under thermal conditions. Key to the success of this methodology is the availability of ligand-modified ruthenium complexes that enable the rational tuning of electrochemical properties of the catalyst without significantly perturbing the overall photophysical properties of the system.

The Diels–Alder reaction ranks among the most important carbon-carbon bond forming reactions in the repertoire of synthetic organic chemistry.<sup>1</sup> Factors governing the rate, stereoselectivity, and catalysis of this powerful transformation have been extensively studied and are well understood. In particular, the reaction rate is generally synthetically useful only when an electron-rich component (typically the diene) engages an electron-deficient reaction partner (typically the dienophile). Electronically mismatched Diels–Alder reactions between two electron-rich components require more forcing conditions and significantly longer reaction times.

On the other hand, the radical cations of electron-rich olefins undergo exceptionally facile [4+2] cycloadditions with electron-rich dienes. These radical cation Diels–Alder reactions can occur with high regio-, stereo-, and chemoselectivity and often occur at rates several orders of magnitude greater than thermal cycloaddition of the corresponding neutral species.<sup>2</sup> The requisite radical cations are most commonly generated either using one-electron chemical oxidants such as aminium salts<sup>2a–c,3</sup> or using photoinitiated electron transfer (PET) with an organic photosensitizer.<sup>4</sup> Both approaches require somewhat high loadings of the oxidant or photosensitizer, and the photochemical methods generally call for the use of high-intensity xenon lamps. In this communication, we report a highly efficient and operationally facile protocol for radical cation Diels–Alder cycloaddition that utilizes low loadings of a ruthenium photosensitizer and irradiation with visible light.




\*Corresponding Author tyoon@chem.wisc.edu.

Supporting Information Placeholder

ASSOCIATED CONTENT

**Supporting Information.** Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.



(2)

We recently reported that the [2+2] photocycloadditions of electron-rich olefins can be conducted upon visible light irradiation in the presence of catalytic  $\text{Ru}(\text{bpy})_3^{2+}$  and methyl viologen ( $\text{MV}^{2+}$ ).<sup>5,6,7</sup> This reaction involves oxidative quenching of the  $\text{Ru}^*(\text{bpy})_3^{2+}$  photoexcited state with  $\text{MV}^{2+}$  to afford an oxidized  $\text{Ru}(\text{bpy})_3^{3+}$  complex that is sufficiently oxidizing (+1.3 V vs SCE) to convert a variety of electron-rich styrenes to the corresponding radical cations. Given the facility of this approach, we speculated that other reactions known to involve radical cation intermediates, including the Diels–Alder cycloaddition, might also be amenable to visible light photocatalysis under similar conditions.

Our investigation commenced with an exploration of the reaction between anethole (**1**) and isoprene (**2**). As expected, this electronically mismatched Diels–Alder cycloaddition does not proceed upon thermal activation (200 °C, 24 h, eq 1). Nevertheless, we found that the photocatalytic conditions we had developed for intramolecular [2+2] cycloaddition of electron-rich styrenes<sup>5</sup> translated smoothly to this [4+2] process. Cycloadduct **3** was formed in 98% yield after just 1 h of irradiation with a household 23 W compact fluorescent light bulb at ambient temperature (Table 1, entry 1).

Despite this promising preliminary result, the scope of the  $\text{Ru}(\text{bpy})_3^{2+}/\text{MV}^{2+}$  catalyst system proved to be quite limited. Reactions involving sterically demanding dienes or less electron-rich dienophiles required much longer reaction times or higher catalyst loadings to achieve synthetically useful yields (*vide infra*). In addition, the loadings of the photocatalyst and  $\text{MV}^{2+}$  in this process (5 mol% and 15 mol%, respectively) are relatively high, and lower loadings result in significant loss of efficiency (Table 1, entry 2). Thus, it became clear that the development of a robust and efficient protocol for the radical cation Diels–Alder reaction would require a more reactive catalyst system for oxidative photocatalysis.

We speculated that rapid back electron transfer from the reduced viologen quencher ( $\text{MV}^+$ ) to the dienophile radical cation might be reducing the efficiency of the desired cycloaddition.<sup>8</sup> In order to slow the rate back electron transfer, we first replaced  $\text{MV}^{2+}$  with air as an irreversible co-oxidant for this process (Table 1, entry 3), without success. The observed lack of reactivity is consistent with the quenching of  $\text{Ru}^*(\text{bpy})_3^{2+}$  by net energy transfer rather than by oxidative quenching;<sup>9</sup> the key  $\text{Ru}(\text{bpy})_3^{3+}$  photooxidant required for one-electron oxidation of **1**<sup>10</sup> would not be expected to be efficiently formed under air. On the other hand, numerous ligand-modified variants of  $\text{Ru}(\text{bpy})_3^{2+}$  are known,<sup>11</sup> and the effect of these modifications on the photophysical and electrochemical properties of the complex have been studied in detail. In particular, the tris(bipyrazyl) analog,  $\text{Ru}(\text{bpz})_3^{2+}$  (**6**),<sup>12</sup> possesses an excited state oxidation potential (+1.4 V vs. SCE) sufficient to directly oxidize **1** without the need for a co-oxidant.

Indeed, irradiation of **1** and **2** under air in the presence of 1 mol%  $\text{Ru}(\text{bpz})_3(\text{PF}_6)_2$  (**6**·( $\text{PF}_6$ )<sub>2</sub>) produces [4+2] cycloadduct **3** in good yield (entry 4). Using this catalyst, we found that the concentration of the diene could be lowered to 3 equiv relative to **1** without adverse effect on yield (entry 5). We also investigated the effect of various counteranions. The BArF complex of the catalyst afforded marginally higher yields, and its greater solubility enabled us to conduct the cycloaddition in less polar solvents such as  $\text{CH}_2\text{Cl}_2$  (entries 6 and 7). We

discovered under these conditions that  $\text{MgSO}_4$  was having a deleterious effect, and that the yield improved when no desiccant was added (entry 8). Finally, we observed that the Diels–Alder cycloaddition remained high-yielding when the loading of  $6\bullet(\text{BARF})_2$  was reduced to 0.5 mol% (entry 9).

The scope studies outlined in Chart 1 demonstrate that this optimized protocol for the radical cation Diels–Alder reaction is quite general. A variety of substituted dienes are successful reaction partners, including sterically bulky (**7**), aryl- and heteroatom-substituted (**9** and **10**), and cyclic dienes (**12**), and in cases where diastereoselectivity is an issue, the cycloaddition exhibits a high *endo* selectivity (**9**, **10**, and **12**). Electron-deficient dienes such as **24** failed to produce cycloadduct, demonstrating that moderately electron-rich dienes are required due to the electron-deficiency of the radical cationic dienophile. Danishefsky's diene (**25**) also failed to undergo [4+2] cycloaddition, presumably due to competitive, unproductive oxidation of this highly electron-rich diene. The dienophile also tolerates significant structural modification, including multiple alkoxy groups (**13** and **14**) or other protected oxygen substituents (**15** and **16**). Substitution at the allylic position is tolerated (**17** and **18**), and a variety of cyclic electron-rich styrenes are suitable dienophiles in the cycloaddition (**19–21**). We also found that the scope of the reaction is not limited to styrenes; other electron-rich olefins such as enamines (**22**) and vinyl ethers (**23**) also undergo cycloaddition under these conditions. However, several key limitations remain to be addressed. Styrenes lacking electron-donating *para* or *ortho* substituents (**26**) do not react, presumably due to the greater difficulty of oxidizing these substrates. Disubstituted styrenes (**28**) and stilbenes (**29**) are also not suitable dienophiles. Finally, unprotected phenols (**30**) undergo competitive oxidative decomposition rather than productive cycloaddition. Nevertheless, within these constraints, the scope of the radical cation Diels–Alder cycloaddition under these conditions is quite broad.

Control and comparison experiments probing the reaction conditions are summarized in Table 2. Both  $\text{Ru}(\text{bpz})_3^{2+}$  and a light source are required for cycloaddition (entries 2 and 3), consistent with a photocatalytic process. However, a variety of visible light sources are suitable, including ambient sunlight (entry 4). Reactions conducted in degassed solvents in the absence of air produce the cycloadduct but do not proceed to completion (entry 5). We reason from this observation that a chain propagation mechanism is operative, as suggested by Ledwith's studies of radical cation cycloadditions,<sup>13</sup> and that the role of oxygen is to facilitate catalyst turnover from the photogenerated  $\text{Ru}(\text{bpz})_3^+$  complex to the resting  $\text{Ru}(\text{bpz})_3^{2+}$  state. Reactions conducted with Rose Bengal in place of the ruthenium photocatalyst (entry 6) did not produce cycloadduct, suggesting that the radical cation Diels–Alder reaction is not mediated by singlet oxygen. Finally, we also observed significantly diminished reactivity using the organic photosensitizers most commonly reported for promoting radical cation Diels–Alder reactions. No conversion occurs when the photocatalyst was replaced with 9,10-dicyanoanthracene (entry 7), and only 28% yield was obtained using triphenylpyrilium $\bullet\text{BF}_4$  (entry 8). Thus, the conditions reported in this paper are uniquely well suited to the promotion of this radical cation cycloaddition under visible light irradiation.

Based upon these observations, we propose the mechanism outlined in Scheme 1. The  $\text{Ru}(\text{bpz})_3^{2+}$  chromophore undergoes an MLCT transition in the visible range ( $\lambda_{\text{max}} = 440$  nm), which produces a redox-active photoexcited state ( $\text{Ru}^*(\text{bpz})_3^{2+}$ ) whose oxidation potential (+1.4 V vs SCE) is sufficiently positive to oxidize anethole (**1**, +1.1 V). The resulting radical cation (**1 $\bullet^+$** ) is activated towards [4+2] cycloaddition to afford the radical cation Diels–Alder product **3 $\bullet^+$** , which can abstract an electron from another equivalent of **1** in a chain-propagation step. Oxygen turns over the reduced  $\text{Ru}(\text{bpz})_3^+$  complex to the photoactive  $\text{Ru}(\text{bpz})_3^{2+}$  state, which may then initiate additional radical cation chains.

Synthetically, the radical cation Diels–Alder process is best understood as an umpolung process that reverses both the intrinsic electronic character of the electron-rich dienophile and the overall regiochemical preference of the cycloaddition.<sup>14</sup> Given this analysis, we became intrigued by the structure of heitziamide A (**31**), an amide natural product isolated as a racemate from the medicinal shrub *Fagara heitzii*.<sup>15</sup> This compound almost certainly arises from a Diels–Alder cycloaddition between *trans*-fagaramide (**33**), another amide endogenous to *F. heitzii*,<sup>15</sup> and the monoterpene myrcene (**34**). However, the polarization of the diene and dienophile  $\pi$  bonds suggests that the regiochemistry expected from the thermal Diels–Alder reaction would afford the isomeric cycloadduct **32**. Indeed, heating **33** with 20 equiv of neat diene **34** at 150 °C for 72 h afforded 60% of the unnatural regioisomer **32** without any trace of natural heitziamide A.

On the other hand, due to its umpolung reactivity, the intrinsic regiochemical preference of a radical cation in the Diels–Alder reaction is opposite that of the corresponding neutral dienophile. Indeed, the natural regiochemistry of heitziamide A is produced upon irradiation of styrene **35** and myrcene in the presence of 2 mol% Ru(bpz)<sub>3</sub>(PF<sub>6</sub>)<sub>2</sub>.<sup>16</sup> No trace of the unnatural regioisomer could be observed by <sup>1</sup>H NMR analysis of the unpurified reaction mixture. The regiochemistry was confirmed by deprotection of the silyl ether and oxidation<sup>17</sup> followed by a EDC coupling of the resulting carboxylic acid (**37**) with isobutylamine to afford a compound whose spectral properties were identical to those of natural heitziamide A.

In summary, we have developed a robust protocol for radical cation Diels–Alder cycloadditions using visible light photocatalysis. These reactions feature low catalyst loadings and short reaction times and are conducted open to the atmosphere. In contrast to previously reported photocatalytic radical cation Diels–Alder processes, this method does not require the use of specialized photochemical equipment and can be conducted using a variety of convenient visible light sources including ambient sunlight. Thus, we have developed an extremely convenient strategy for promoting electronically mismatched Diels–Alder cycloadditions between electron-rich coupling partners that would otherwise be difficult to accomplish. Further demonstrations of the applicability of this reaction in synthesis are under development in our laboratory.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

Dedicated to Prof. Howard Zimmerman in celebration of his career and his many contributions to photochemistry. This research was conducted using funds from the ACS PRF (49817-ND1), NIH (GM095666), Sloan Foundation, Beckman Foundation, and Research Corporation. The NMR facilities at UW-Madison are funded by the NSF (CHE-9208463, CHE-9629688) and NIH (RR08389-01, RR13866-01).

## REFERENCES


1. a Bear BR, Sparks SM, Shea KJ. *Angew. Chem. Int. Ed.* 2001; 40:820–849. b Nicolaou KC, Snyder SA, Montagnon T, Vassilikogiannakis G. *Angew. Chem. Int. Ed.* 2002; 41:1668–1698. c Stocking EM, Williams RM. *Angew. Chem. Int. Ed.* 2003; 42:3078–3115. d Takao K-I, Munakata R, Tadano K-I. *Chem. Rev.* 2005; 105:4779–4807. [PubMed: 16351062]
2. a Bauld NL, Bellville DJ, Pabon R, Chelsky R, Green G. *J. Am. Chem. Soc.* 1983; 105:2378–2382. b Bellville DJ, Bauld NL, Pabon R, Gardner SA. *J. Am. Chem. Soc.* 1983; 105:3584–3588. c Pabon RA, Bauld NL. *J. Am. Chem. Soc.* 1984; 106:1145–1146. d Wiest O, Steckhan E, Grein F. *J. Org. Chem.* 1992; 57:4034–4037. e Bauld NL, Yang J. *Org. Lett.* 1999; 1:773–774. [PubMed: 16118883]

- f Valley NA, Wiest O. *J. Org. Chem.* 2007; 72:559–566. [PubMed: 17221974] g Sevov CS, Wiest O. *J. Org. Chem.* 2008; 73:7909–7915. [PubMed: 18785777]
3. a Bellville DJ, Wirth DW, Bauld NL. *J. Am. Chem. Soc.* 1981; 103:718–720. b Yueh W, Bauld NL. *J. Phys. Org. Chem.* 1996; 9:529–538.
4. For seminal reports of organic PET sensitization of radical cation Diels–Alder, see: a Maroulis AJ, Arnold DR. *J. Chem. Soc., Chem. Commun.* 1979:351–352.; b Jones CR, Allman BJ, Mooring A, Spahic B. *J. Am. Chem. Soc.* 1983; 105:652–654.; c Pabon RA, Bellville DJ, Bauld NL. *J. Am. Chem. Soc.* 1983; 105:5158–5159.; d Calhoun GC, Schuster GB. *J. Am. Chem. Soc.* 1984; 106:6870–6871.; e Mlcoch J, Steckhan E. *Angew. Chem. Int. Ed. Engl.* 1985; 24:412–414.; e Gieseler A, Steckhan E, Wiest O. *Synlett.* 1990:275–277.; f Gieseler A, Steckhan E, Wiest O, Knoch F. *J. Org. Chem.* 1991; 56:1405–1411.
5. Ischay MA, Lu Z, Yoon TP. *J. Am. Chem. Soc.* 2010; 132:8572–8574. [PubMed: 20527886]
6. For reviews on recent developments in transition metal photoredox catalysis in organic synthesis, see: a Zeitler K. *Angew. Chem., Int. Ed.* 2009; 48:9785–9789.; b Narayanam JMR, Stephenson CRJ. *Chem. Soc. Rev.* 2011; 40:102–113. [PubMed: 20532341] ; c Yoon TP, Ischay MA, Du J. *Nat. Chem.* 2010; 2:527–532. [PubMed: 20571569]
7. For leading references on transition metal photocatalysis from other groups, see: a Nicewicz DA, MacMillan DWC. *Science.* 2008; 322:77–80. [PubMed: 18772399] ; b Pham PV, Nagib DA, MacMillan DWC. *Angew. Chem. Int. Ed.* 2011; 50:6119–6122.; c Narayanam JMR, Tucker JW, Stephenson CRJ. *J. Am. Chem. Soc.* 2009; 131:8756–8757. [PubMed: 19552447] ; d Furst L, Narayanam JMR, Stephenson CRJ. *Angew. Chem. Int. Ed.* 2011; 50:9655–9659.; e Andrews RS, Becker JJ, Gagné MR. *Angew. Chem. Int. Ed.* 2010; 49:7274–7276.; f Chen Y, Kamlet AS, Steinman JB, Liu DR. *Nat. Chem.* 2011; 3:146–153. [PubMed: 21258388] ; g Larraufie M-H, Pellet R, Fensterbank L, Goddard J-P, Lacôte E, Malacria M, Ollivier C. *Angew. Chem. Int. Ed.* 2011; 50:4463–4466.
8. The effect of back electron transfer on the efficiency of PET sensitized radical cation chemistry is well appreciated. For instance, see: a Reynolds DW, Lorenz KT, Chiou HS, Bellville DJ, Pabon RA, Bauld NL. *J. Am. Chem. Soc.* 1987; 109:4960–4968.; b Mattay J. *Angew. Chem. Int. Ed. Engl.* 1987; 26:825–845.
9. a Winterle JS, Kliger DS, Hammond GS. *J. Am. Chem. Soc.* 1976; 98:3719–3721. b Demas JN, Harris EW, McBride RP. *J. Am. Chem. Soc.* 1977; 99:3547–3551.
10. The half-wave potential of 1 is +1.1 V vs. SCE. If Ru\*(bpy)<sub>3</sub><sup>2+</sup> is not oxidatively quenched to the more strongly oxidizing Ru(III) state (+1.3 V), the excited state itself is insufficiently oxidizing (+0.8 V) to produce the alkene radical cation. These values are from ref 3b and Kalyanasundaram K. *Coord. Chem. Rev.* 1982; 46:159–244.
11. Juris A, Balzani V, Barigelli F, Campagna S, Belser P, von Zelewsky A. *Coord. Chem. Rev.* 1988; 84:85–277.
12. Crutchley RJ, Lever ABP. *J. Am. Chem. Soc.* 1980; 102:7128–7129.
13. a Crellin RA, Lambert MC, Ledwith A. *J. Chem. Soc. D, Chem. Commun.* 1970:682–683. b Ledwith A. *Acc. Chem. Res.* 1972; 5:133–139.
14. The regiochemistry of the cycloadducts in Chart 1 is consistent with polarization of the dienophile as shown for structure 1\*<sup>+</sup> (Scheme 1). The umpolung polarization of alkene radical cations has been exploited in synthetic applications by Moeller. For an excellent, concise account, see: Moeller KD. *Synlett.* 2009:1208–1218.
15. Mbaze LM, Lado JA, Wansi JD, Shiao TC, Chiozem DD, Mesaik MA, Choudhary MI, Lacaille-Dubois M-A, Wandji J, Roy R, Sewald N. *Phytochemistry.* 2009; 70:1442–1447. [PubMed: 19747699]
16. Aminium cation initiators (ref 4c) failed to produce any trace of the cycloadduct in the same reaction time. Similarly, we also replaced Ru(bpz)<sub>3</sub>(PF<sub>6</sub>)<sub>2</sub> with the photosensitizer triphenylpyrilium tetrafluoroborate (ref 4d) and failed to observed significant formation of 35. See the Supporting Information for experimental details.
17. a Haga Y, Okazaki M, Shuto Y. *Biosci. Biotech. Biochem.* 2003; 67:2183–2193. b Lopes ECS, Coelho F. *J. Braz. Chem. Soc.* 2007; 18:1415–1438.




<sup>a</sup> Conditions, unless noted: 0.5 mol% Ru(bpz)<sub>3</sub>(BArF)<sub>2</sub>, 3 equiv diene, CH<sub>2</sub>Cl<sub>2</sub>, irradiated with a 23 W compact fluorescent bulb. <sup>b</sup> Yields in brackets are for the same time point using the following conditions: 5 mol% Ru(bpy)<sub>3</sub>(PF<sub>6</sub>)<sub>2</sub>, 15 mol% MV(PF<sub>6</sub>)<sub>2</sub>, MgSO<sub>4</sub>, MeNO<sub>2</sub>. <sup>c</sup> Using 2 equiv of diene. <sup>d</sup> Using 3 mol% Ru(bpz)<sub>3</sub>(BArF)<sub>2</sub>. <sup>e</sup> Yield determined by <sup>1</sup>H NMR analysis using CH<sub>2</sub>Br<sub>2</sub> as an internal standard.

**Chart 1.**  
Scope of the radical cation Diels–Alder cycloaddition<sup>a</sup>



**Scheme 1.**  
Proposed mechanism



**Scheme 2.**  
 Synthetic studies on heitziamide A



Table 1

Optimization of conditions for the radical cation Diels–Alder cycloaddition.

entry	catalyst (mol%)	co-oxidant (amount)	diene (equiv)	solvent	time	yield (%) <sup>a</sup>
1 <sup>b</sup>	Ru(bpy) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub> (5)	MV(PF <sub>6</sub> ) (1.5 mol%)	10	MeNO <sub>2</sub>	1 h	98%
2 <sup>b</sup>	Ru(bpy) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub> (1)	MV(PF <sub>6</sub> ) (3 mol%)	10	MeNO <sub>2</sub>	1 h	25%
3 <sup>b</sup>	Ru(bpy) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub> (1)	air (1 atm)	10	MeNO <sub>2</sub>	1 h	0%
4 <sup>b</sup>	Ru(bpz) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub> (1)	air (1 atm)	10	MeNO <sub>2</sub>	1 h	78%
5 <sup>b</sup>	Ru(bpz) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub> (1)	air (1 atm)	3	MeNO <sub>2</sub>	1 h	76%
6 <sup>b</sup>	Ru(bpz) <sub>3</sub> (BArF) <sub>2</sub> (1)	air (1 atm)	3	MeNO <sub>2</sub>	30 min	83%
7 <sup>b</sup>	Ru(bpz) <sub>3</sub> (BArF) <sub>2</sub> (1)	air (1 atm)	3	CH <sub>2</sub> Cl <sub>2</sub>	30 min	85%
8	Ru(bpz) <sub>3</sub> (BArF) <sub>2</sub> (1)	air (1 atm)	3	CH <sub>2</sub> Cl <sub>2</sub>	30 min	92%
9	Ru(bpz) <sub>3</sub> (BArF) <sub>2</sub> (0.5)	air (1 atm)	3	CH <sub>2</sub> Cl <sub>2</sub>	1 h	98% <sup>c</sup>

<sup>a</sup>Yields determined by <sup>1</sup>H NMR analysis using CH<sub>2</sub>Br<sub>2</sub> as an internal standard unless otherwise noted.

<sup>b</sup>Conducted in the presence of 200 wt% MgSO<sub>4</sub>.

<sup>c</sup>Isolated yield.

**Table 2**Control and comparison studies for cycloaddition of **1** and **2** (eq 2).

entry	Conditions	time	yield (%)
1	standard conditions	1 h	98%
2	no Ru(bpz) <sub>3</sub> (PF <sub>6</sub> ) <sub>2</sub>	1 h	0%
3	no light	1 h	0%
4	ambient sunlight instead of fluorescent lamp	1 h	98%
5	no air (under N <sub>2</sub> )	1 h	46%
6	Rose Bengal instead of Ru(bpz) <sub>3</sub> <sup>2+</sup>	1 h	0%
7	9,10-Dicyanoanthracene instead of Ru(bpz) <sub>3</sub> <sup>2+</sup>	1 h	0%
8	Triphenylpyrilium•BF <sub>4</sub> instead of Ru(bpz) <sub>3</sub> <sup>2+</sup>	1 h	28%