

RADIO-AGN IN THE AKARI-NEP FIELD AND THEIR ROLE IN THE EVOLUTION OF GALAXIES

M. KAROUZOS, M. IM, AND THE AKARI-NEP TEAM

Department of Physics and Astronomy, Seoul National University, Seoul 151-747, Korea

Center for the Exploration of the Origin of the Universe

E-mail: mkarouzos@astro.snu.ac.kr

(Received June 28, 2012; Accepted July 27, 2012)

ABSTRACT

Radio-loud active galaxies have been found to exhibit a close connection to galactic mergers and host galaxy star-formation quenching. We present preliminary results of an optical spectroscopic investigation of the AKARI NEP field. We focus on the population of radio-loud AGN and use photometric and spectroscopic information to study both their star-formation and nuclear activity components. Preliminary results show that radio-AGN are associated with early type, massive galaxies with relatively old stellar populations.

Key words: infrared: telescope; conferences: proceedings

1. INTRODUCTION

In the context of galaxy evolution in the Universe, the role of nuclear activity, in particular radio-loud active galactic nuclei (AGN), is still under debate. Are radio-AGN a phase of a galaxy's evolution? How are they triggered and what is their effect on their host galaxy? We identify radio-AGN within the AKARI-NEP field and study their host galaxy properties in terms of an hierarchical evolutionary scheme.

2. CROSS-IDENTIFICATION

We cross-identify all AKARI-NEP (wide and deep) sources detected in the N2 band of AKARI with the sources from the WSRT catalog at 1.5 GHz ([WH10]), following [DO86] (also see [WH12]). In total 401 and 168 radio sources are matched for NEP-wide and -deep, respectively. Photo- z for NEP-deep cross matched sources range between 0.37 and 2.2, with most sources having z between 0.37 and 1.

We also cross-identify 1.5 GHz WSRT sources with the optical spectroscopy catalogs available (Shim et al., Takagi et al., private communication). For a matching radius of 3 arcsec, 48 radio sources are matched (spec- z between 0.03 and 4, with a few above 1). Radio-samples are defined in Table 1.

TABLE 1.
Radio Samples Selection Criteria

Sample	Description	Selection
(1)	All	-
(2)	Luminosity	$L_{1.5GHz} > 10^{23} \text{W/Hz}^*$
(3)	Flat-spectrum	$\alpha_{radio} < 0.5^{**}$

*limit definition following [CO92][MA07]
** α_{radio} calculated using 2 or 3 bands

3. OPTICAL SPECTROSCOPY

We are in the process of analyzing all the available optical spectra using IDL routines (emission line fluxes, equivalent widths, 4,000 Å break, etc.). Using the BPT emission line classification diagram (Fig. 1) we find 10 AGN/LINERs and 21 transitional objects in a total of 84 sources. 3 radio-sources are classified as AGN/LINERs and 3 as transitional.

4. OPTICAL COLORS AND STELLAR AGES

We find that luminosity-selected AGN are predominantly associated with elliptical galaxies ($u-r > 2.22$; [ST01]), with a similar trend for our flat-spectrum sample. A two-sample KS test gives a 99.8%




Fig. 1. Baldwin-Phillips-Terlevich (BPT) emission line ratios diagnostic diagram to separate star-forming galaxies from active AGN and LINERs. The dashed line is from [n], the dotted line from [l], and the continuous lines are from [m] (horizontal) and [n] (diagonal). They separate star-forming galaxies, LINERs, and Seyferts, respectively.

probability that sample (2) is drawn from a different population than its parent sample. For the comparison with sample (3) the KS test does not provide a significant result. Both samples (1) and (2) show C_{4000A} characteristic of old stellar populations and early-type galaxies (e.g., [GA05]). A small fraction of sample (2) shows low values of C_{4000} indicative of a strong power-law non-thermal continuum (Fig. 2). Assuming that rest-frame N2 luminosity is a good proxy for the stellar mass of a galaxy, both samples (1) and (2) inhabit more massive galaxies compared to the non-radio sample.

ACKNOWLEDGEMENTS

MK acknowledges the support from the Creative Research Initiative program, No. 2010-0000712, of the National Research Foundation of Korea (NRFK) funded by the Korea government(MEST).

REFERENCES

- Condon, J. J., 1992, Radio Emission from Normal Galaxies, *ARA&A*, 30, 575
 Downes, A. J., Peacock, J. A., Savage, A., et al., 1986, The Parkes Selected Regions - Powerful Radio Galaxies and Quasars at High Redshifts, *MNRAS*, 218, 31
 Gallazzi, A., Charlot, S., Brinchmann, J., et al., 2005,




Fig. 2. Normalized distributions of the 4000A break index C_{4000A} and rest-frame N2 luminosity for all non-radio sources (black), all radio sources (dashed black), and luminosity-selected radio-AGN (shaded turquoise). Only sources with spectroscopic redshifts are included here. The dotted line denotes the L_* .

- The Ages and Metallicities of Galaxies in the Local Universe, *MNRAS*, 362, 41
 Ho, L. C., Filippenko, A. V., & Sargent, W. L., 1997, A Search for “Dwarf” Seyfert Nuclei. III. Spectroscopic Parameters and Properties of the Host Galaxies, *ApJS*, 112, 315
 Kauffmann, G., Heckman, T. M., Tremonti, C., et al., 2003, The Host Galaxies of Active Galactic Nuclei, *MNRAS*, 346, 1055
 Kewley, L. J. & Dopita, M. A., 2002, Using Strong Lines to Estimate Abundances in Extragalactic H II Regions and Starburst Galaxies, *ApJS*, 142, 35
 Mauch, T. & Sadler, E. M., 2007, Radio Sources in the 6dFGS: Local Luminosity Functions at 1.4 GHz for Star-Forming Galaxies and Radio-Loud AGN, *MNRAS*, 375, 931
 Strateva, I., Ivezić, Ž., Knapp, G. R., et al., 2001, Color Separation of Galaxy Types in the Sloan Digital Sky Survey Imaging Data, *AJ*, 122, 1861
 White, G. J., Pearson, C., Braun, R., et al., 2010, A Deep Survey of the AKARI North Ecliptic Pole Field. I. WSRT 20 cm Radio Survey Description, Observations and Data Reduction, *A&A*, 517A, 54
 White, G. J., Hatsukade, B., Pearson, C., et al., 2012, A deep ATCA 20cm Radio Survey of the AKARI Deep Field South near the South Ecliptic Pole, arXiv:1207.2262