

RESEARCH ARTICLE

Open Access

Radioactive ^{125}I seeds inhibit cell growth and epithelial-mesenchymal transition in human glioblastoma multiforme via a ROS-mediated signaling pathway

Yunhong Tian^{1,2†}, Qiang Xie^{3†}, Jie He¹, Xiaojun Luo¹, Tao Zhou³, Ying Liu³, Zuoping Huang³, Yunming Tian¹, Dan Sun³ and Kaitai Yao^{1*}

Abstract

Background: Glioblastoma multiforme (GBM) is the most common primary central nervous system neoplasm in adults. Radioactive ^{125}I seed implantation has been widely applied in the treatment of cancers. Moreover, previous clinical trials have confirmed that ^{125}I seeds treatment was an effective therapy in GBM. We sought to investigate the effect of ^{125}I seed on GBM cell growth and Epithelial-mesenchymal transition (EMT).

Methods: Cells were exposed to irradiation at different doses. Colony-formation assay, EdU assay, cell cycle analysis, and TUNEL assay were performed to investigate the radiation sensitivity. The effects of ^{125}I seeds irradiation on EMT were measured by transwell, Boyden and wound-healing assays. The levels of reactive oxygen species (ROS) were measured by DCF-DA assay. Moreover, the radiation sensitivity and EMT were investigated with or without pretreatment with glutathione. Additionally, nude mice with tumors were measured after treated with radiation.

Results: Radioactive ^{125}I seeds are more effective than X-ray irradiation in inhibiting GBM cell growth. Moreover, EMT was effectively inhibited by ^{125}I seed irradiation. A mechanism study indicated that GBM cell growth and EMT inhibition were induced by ^{125}I seeds with the involvement of a ROS-mediated signaling pathway.

Conclusions: Radioactive ^{125}I seeds exhibit novel anticancer activity via a ROS-mediated signaling pathway. These findings have clinical implications for the treatment of patients with GBM by ^{125}I seeds.

Keywords: Irradiation, Radioactive ^{125}I seeds, Glioblastoma multiforme, Epithelial-mesenchymal transition

Background

Glioblastoma multiforme (GBM) is the most common and lethal type of primary central nervous system neoplasm in adults [1]. Unlike most other tumors that metastasize to distant organs, malignant glioma very rarely metastasizes outside the central nervous system. In this sense, GBM may be regarded as a “local” tumor [2]. In GBM, standard treatment involves maximal resection followed by concomitant and adjuvant chemoradiotherapy with temozolomide. Even with this

comprehensive treatment strategy, outcomes for patients with this malignancy remain very poor. Thus, in order to improve the current therapeutic regimens, it is important to explore effective new modalities for GBM patients. Radioactive ^{125}I seed implantation has been widely applied in the treatment of cancers [3-6]. It has been shown to be an effective adjuvant therapy in recurrent GBM [7,8] and Low-grade (WHO grades I and II) gliomas (LGGs) [9,10]. Furthermore, several studies have shown that ^{125}I seed irradiation directly causes more cell death by comparing with ^{60}Co - γ or X-ray irradiation [11-14]. However, few studies of the biological effects of ^{125}I seed irradiation on GBM cells are available.

Epithelial-mesenchymal transition (EMT) is a key developmental program that is often activated during

* Correspondence: yao.kaitai@yahoo.cn

†Equal contributors

¹Cancer Research Institute, Southern Medical University, Guangzhou 510, 515 Guangdong Province, People's Republic of China

Full list of author information is available at the end of the article

cancer invasion and metastasis [15]. Cells that have undergone EMT are resistant to many of the chemotherapeutic and adjuvant drugs that are used to treat epithelial tumors, and may therefore drive tumor recurrence [16,17]. For the lack of E-cadherin expression in GBM cells suggesting a non-classic EMT, only very few recent reports described an EMT phenomenon in GBMs and its association with the poor prognostic mesenchymal subgroup of GBMs [18]. Reactive oxygen species (ROS) play an important role in cellular metabolism and cancer therapy [19,20]. The absorption of ionizing radiation by living cells can act indirectly through radiolysis of water, thereby generating ROS [21]. Moreover, in the past few years, nuclear DNA damage-sensing mechanisms activated by ionizing radiation have been identified, including ataxia-telangiectasia mutated (ATM)/ATM-and Rad3-related (ATR) and the DNA-dependent protein kinase [22,23].

Therefore, in this study, we evaluated the effect of radioactive ^{125}I seeds on GBM cell growth and EMT. The results showed that radioactive ^{125}I seeds were more effective than X-ray irradiation in inhibiting GBM cell growth. Moreover, EMT in GBM cells was effectively inhibited by ^{125}I seed irradiation. A mechanism study indicated that GBM cell growth and EMT were inhibited by ^{125}I seeds with the involvement of a ROS-mediated signaling pathway. Pretreatment of cells with glutathione (GSH) significantly blocked ^{125}I seed irradiation-induced inhibition of cell migration and growth by recovering the expression levels of ROS. Meanwhile, the results of an *in vivo* study confirmed that ^{125}I seed irradiation inhibits tumor growth and EMT via a ROS-mediated signaling pathway. Taken together, these results suggest that radioactive ^{125}I seeds exhibit novel anticancer activity via a ROS-mediated signaling pathway. These findings have clinical implications for the treatment of patients with GBM by ^{125}I seeds.

Methods

Cell culture and reagents

U251 and U87 human GBM cell lines were available at the Cancer Institute of Southern Medical University (Guangzhou, China) and were originally purchased from the American Type Culture Collection (ATCC). Cells were maintained in Dulbecco's Modified of Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and antibiotics (100 IU/ml penicillin and 100 mg/ml streptomycin) at 37°C under a humidified atmosphere of 95% air and 5% CO₂. To investigate the effect of ROS on migration, 5 mM GSH (Sigma-Aldrich, MO, USA) was added 2 hours before irradiation.

Treatment of GBM cells with ^{125}I seeds and X-ray irradiation

^{125}I seeds were obtained from Beijing Atom and High Technique Industries Inc. (Beijing, China). The *in vitro*

irradiation was carried out as previously described [13]. The absorbed doses were calculated as follows: 44, 92, 144, and 204 hours were required for doses of 2, 4, 6, and 8 Gy, respectively [14]. X-ray irradiation with a clinically calibrated irradiation field of 10 × 10 cm was performed at the Department of Radiotherapy, Armed Police Corps Hospital of Guangdong Province, using the Elekta precise treatment system (Stockholm, Sweden).

Colony-formation and thiazolyl blue tetrazolium bromide (MTT) assay

According to a previous study, the plating efficiency (PE) of unirradiated controls was calculated using the following formula: number of colonies/number of seeded cells × 100%. U87 and U251 cells were exposed to radiation and then seeded using a cell-dilution assay. Surviving fractions (SFs) were calculated as following formula: SF = number of colonies/number of seeded cells × PE. The dose-survival curve was fitted based on the single-hit multi-target theory formula: $\text{SF} = 1 - (1 - e^{-D/D_0})^N$; $\log N = D_q/D_0$. Cell viability was determined by MTT assay as previously described [24].

Annexin V-PI apoptosis and Caspase-3 activity assay

Cells in exponential growth were irradiated and harvested 24 hours after irradiation. Then cells were assessed according to the protocol of the Alexa Fluor® 488 annexin V/Dead Cell Apoptosis kit (Invitrogen, CA, USA). For caspase-3 activity, cells incubated 48 hours after irradiation at different doses were lysed with lysis buffer (100 μl per 2 × 10⁶ cells) for 15 minutes on ice following washing with D-Hank's medium. Then cell extracts mixed with Ac-DEVD-pNA substrate were incubated at 37°C for 2 hours. The values measured by colorimetric measurement of p-nitroanilide product at 405 nm were normalized to untreated controls allowing determination of the fold change in caspase-3 activity.

Cell cycle measured by flow cytometry

Cells in exponential growth were irradiated and harvested 24 hours after irradiation. Then they were washed with cold phosphate-buffered saline (PBS) and fixed overnight in cold 70% ethanol. Fixed cells washed with PBS were resuspended in 100 μl RNaseA (250 μg/ml), incubated for 30 minutes at 37°C. Then, 50 μg/ml PI was added and incubated at room temperature in the dark for 30 minutes followed by PI-detection with BD FACSCaria™ (BD Biosciences, CA, USA).

Analysis of apoptosis by terminal deoxynucleotidyl transferase (TdT)-mediated dUTP-digoxigenin nick-end labeling (TUNEL) assay

We applied a TUNEL assay according to the manufacturer's instructions (Beyotime Institute of Biotechnology, Jiangsu, China) to evaluate the apoptotic response in

tumor cells. Briefly, cells cultured on chamber slides were fixed with 3.7% formaldehyde and permeabilized with 0.1% Triton X-100 in PBS. Then, the cells were incubated with TUNEL reaction mixture for 1 hour and cell nuclei were stained with 4', 6-diamino-2-phenylindole (DAPI; Invitrogen). The cells were then washed with PBS and examined.

Transwell and Boyden chamber assays

Cells (10^6 cells/100 μ l) in serum-free DMEM were added to the upper chamber and 500 μ l of the DMEM with 10% FBS was added to the lower chamber with permeable supports (Corning, NY, USA). Then, cells on the upper surface which were incubated for 24 hours at 37°C were removed using a cotton-tipped applicator. Finally, cells on the lower surface of the filter were stained with crystal violet to calculate the average number of migrated cells [25].

Wound-healing assay

Cells exposed to irradiation at a dose of 4 Gy were scraped with a conventional 10 μ l micropipette tip across the monolayer. The distance between the wound edges was measured immediately and again 24 hours later. The total distance migrated by wounded U251 and U87 cells was evaluated using Adobe Photoshop and is expressed as a percentage of the initial wound distance.

Immunofluorescence assay

Cells seeded on slides were fixed in 4% paraformaldehyde and permeabilized in 0.5% Triton X-100. Primary antibody (1:200; Santa Cruz Biotechnology, CA, USA) and Alexa Fluor 488-conjugated secondary antibody (1:500; Invitrogen) were used to detect the location and expression of E-cadherin and vimentin. The cell nuclei were stained with DAPI. Finally, the images were recorded by fluorescence microscopy with a Nikon eclipse 80i microscope.

Detection of ROS in intracellular

For intracellular ROS analysis, cells were loaded with 10 μ M DCF-DA (Sigma-Aldrich), incubated at 37°C for 30 minutes, and immediately analyzed by microscope and flow cytometry (BD Biosciences).

Western blotting analysis

Cells and tissues were lysed in RIPA buffer. Tumors were ground in liquid nitrogen and lysed. Protein concentration was determined using the BCA Kit (Beyotime Institute of Biotechnology). Proteins were mixed with loading buffer and heated at 70°C for 10 minutes on sodium dodecyl sulfate (SDS)-polyacrylamide gels at 30 μ g per lane. The proteins were transferred to polyvinylidene fluoride (PVDF, Millipore, MA, USA) after electrophoresis. Membranes

were blocked for 2 hours in 5% BSA and incubated overnight at 4°C with antibodies against γ -H2AX, ATM, ATR, Chk1, cell-cycle controller-2 (Cdc2), E-cadherin, vimentin, caspase-3, and caveolin-1 (Cav-1). The blots were then incubated with HRP-conjugated secondary antibody (1:1000; Santa Cruz Biotechnology). Finally, bands were visualized by enhanced chemiluminescence (Thermo Scientific Pierce, IL, USA).

In vivo experiments

Female BALB/c nude mice (age 4–5 weeks) were purchased from the Model Animal Research Center of Nanjing University (Nanjing, China). This study was approved by the ethics committee of Southern Medical University. Animals were injected subcutaneously (s.c.) with U251 cells into the right hind limb (5×10^6 cells/100 μ l). Two weeks later, mice whose tumor volumes had reached approximately 200 mm³ were randomly divided into three groups with 4 mice in every group. The three groups were: (1) irradiation at 20 Gy (2 Gy/day \times 10 F, 5 fractions/week for X-ray irradiation); (2) implanted with ¹²⁵I seeds at a total dose of 20 Gy, the number of which calculated by the treatment planning system (TPS) (RT-RSI, Beijing Atom and High Technique Industries Inc., Beijing, China); and (3) untreated group. The dimensions of xenograft nodules were callipered every 3 days for 2 successive weeks. The animals were euthanized day 15 after treatment. Finally, immunohistochemistry (IHC) and western blotting for E-cadherin and vimentin were performed in xenograft tumor samples.


Statistical analysis

Statistical analysis was performed with the SPSS statistical package (v15.0). *In vitro* experiments were usually performed in triplicate and repeated three times. The data are presented as mean \pm standard deviation (SD). Statistical differences among groups were examined using one-way analysis of variance (ANOVA), with *p* values of less than 0.05 considered statistically significant. Multiple comparisons of the means were done by the least significance difference (LSD) test.

Results

Radioactive ¹²⁵I seeds are more effective than X-ray in inhibiting GBM cell growth

In this study, the effects of irradiation on the growth of GBM cells were measured by colony-formation, MTT, and apoptosis assay. The results showed that the colony-formation ability was significantly reduced by irradiation in a dose-dependent manner (Figure 1A). The SFs of cells exposed to ¹²⁵I seed irradiation were significantly lower than that of cells exposed to X-ray irradiation at the same doses, both in U251 and U87 cells. Based on the dose-survival curve fitted with the



single-hit multi-target theory formula, the D_0 and D_q of ¹²⁵I seeds and X-ray were 1.703 versus 0.722 and 19.245 versus 5.736 in U87. Moreover, the MTT assay also confirmed that the viability of cells exposed to ¹²⁵I seeds was lower than that of cells exposed to X-ray irradiation (Figure 1B), especially at a dose of 4 Gy; the SF decreased from 77.63% to 47.57% in U251 cells

and from 84.42% to 61.69% in U87 cells. Finally, cell proliferation was measured and the results indicated that the cell proliferation of U251 and U87 was significantly inhibited by irradiation in a dose-dependent manner. However, compared with the X-ray group, the cell proliferation was significantly inhibited by ¹²⁵I seeds at the same doses (Figure 1C). Taken together, these

assays indicate that GBM cells are more sensitive to ^{125}I seed irradiation than to X-ray irradiation.

An annexin V- PI apoptosis assay was performed to examine the effect of ^{125}I seed irradiation on cell death. The results showed that apoptotic cell death was markedly induced by X-ray and ^{125}I seed irradiation in a dose-dependent manner. After X-ray irradiation, apoptosis significantly increased in GBM cells. However, compared with X-ray irradiation, ^{125}I seed irradiation led to a higher percentage of apoptosis under the same doses. For example, the apoptotic rate after ^{125}I seed irradiation at a dose of 4 Gy increased from 2.12% (control) to 15.21% in U251 cells and to 14.26% in U87 cells (Figure 2A and B). We further investigated whether

the observed irradiation-induced apoptosis was related to caspase-3. As expected, the results showed that caspase-3 activity increased after irradiation, with ^{125}I seed irradiation exhibiting a greater effect on caspase-3 activity compared with X-ray (Figure 2C). To gain a better understanding of the apoptosis induced by the ionizing irradiation, DNA distribution histograms of GBM cells were acquired. The results indicated that dose-dependent increases in the G2/M cell population were observed in cells exposed to X-ray and ^{125}I seed irradiation, without significant differences in the S and G0/G1 phases (Figure 2D). As we know, TUNEL-positive cells showed typical apoptosis. The results of our study indicate that the number of TUNEL-positive cells was significantly




Figure 2 Apoptosis and G2/M arrest in glioblastoma multiforme (GBM) cells is induced by ^{125}I seed irradiation. Apoptosis of GBM cells was examined by Annexin V-PI co-staining flow cytometric analysis (A, B) and caspase-3 activity assay (C). (D) Effects of ^{125}I seed on the cell-cycle distribution of GBM cells were examined by flow cytometric analysis. (E) TUNEL assay was used to evaluate the apoptotic response in U251 (left panel) and U87 (right panel) cells. (F) Treatment of cells with ^{125}I seeds caused obvious apoptosis in a dose-dependent manner, accompanied by up-regulation of cleaved caspase-3, cleaved PARP, and phosphorylated Cdc2 (Tyr 15). Data are presented as mean \pm SD (n = 3). Significant differences between the ^{125}I seed and X-ray groups under the same dose are indicated by * $P < 0.05$ and ** $P < 0.01$.

increased with ^{125}I seed irradiation as compared with X-ray irradiation at the same doses (Figure 2E). With irradiation at a dose of 4 Gy, the percentage of apoptotic cells increased to 19.18% with ^{125}I seed treatment, compared with 12.32% with X-ray. The enzyme poly-ADP-ribosepolymerase (PARP), the expression of which is triggered by DNA-strand breaks, acts as a substrate for caspases. In cells undergoing apoptosis, it is cleaved by caspase-3 during the degradation of cellular DNA, thus preventing DNA damage repair. It is therefore critical for the repair of some DNA lesions [26,27]. Historically, mitotic entry and exit were thought to be a direct consequence of Cdc2 activation and inactivation, respectively [28]. Therefore, PARP, caspase-3, and Cdc2 in U251 were measured by western blotting to further confirm the apoptosis and G2/M arrest induced by ^{125}I seeds. As expected, treatment of cells with ^{125}I seeds caused obvious apoptosis and G2/M arrest in a dose-dependent manner, as accompanied by up-regulation of cleaved caspase-3, cleaved PARP, and phosphorylated Cdc2 (Tyr 15). Interestingly, results indicated that O6-methylguanine DNA methyltransferase (MGMT) which was an independent favorable prognostic factor in patients with GBM [29] was also up-regulated by irradiation (Figure 2F). Altogether, these results suggest a higher potency of ^{125}I seed irradiation in inducing cancer-cell apoptosis and G2/M arrest.

Effects of irradiation on EMT in GBM cells


GBM cells are characterized by migration and invasion into the surrounding normal brain tissue. This study compared the EMT in GBM cells with X-ray and ^{125}I seed irradiation. As shown in Figure 3A, transwell and Boyden chamber assays were performed to investigate the effects of these treatments on cell invasion and migration. After X-ray irradiation, the number of migrating cells per high-power field (HPF) in U251 cells increased from 42.3 to 59.4, whereas it reduced to 22.9 after ^{125}I seed irradiation. Similar findings were seen with U87 cells: the number of migrating U87 cells decreased significantly after ^{125}I seed irradiation. When the potential involvement of ^{125}I seeds in reducing the invasion of GBM cells was determined by Boyden chamber assay, there was a two-fold decrease in the number of invading cells of both U251 and U87 compared with the control group. As shown in Figure 3B, the migration index of ^{125}I seed irradiation measured by wound-healing assay reduced from 40.9% to 22.1% compared with control. However, a higher migration index of U251 cells was observed 24 hours after X-ray irradiation. Moreover, dramatic morphological changes, in which the spindle-like, fibroblastic morphology was replaced by a typical cobblestone-like appearance of normal epithelium, indicating inhibition of EMT by ^{125}I seeds, were observed

in ^{125}I seed-treated GBM cells (Figure 3C). In GBM cell lines, the capabilities of motility and invasion have been ascribed to Snail and ZEB1 expression [30]. Thus, our studies investigated the levels of ZEB1 and Snail. Our results indicated that the levels of ZEB1 were involved in EMT inhibited by ^{125}I seeds, with no significant changes in Snail in U251 and U87. As anticipated, decreased vimentin were also observed in ^{125}I seed-treated U87 and U251 cells (Figure 3D). To further test the hypothesis that ^{125}I seeds inhibit EMT, the results of immunofluorescence assay indicated that ^{125}I seeds down-regulated ZEB1 in the nuclear of U251 cells, suggesting that ^{125}I seeds inhibit EMT in GBM cells (Figure 3E). Taken together, these results demonstrate that ^{125}I seed irradiation inhibits the nonclassic EMT in GBM cells.

Radioactive ^{125}I seeds suppress cell migration and growth via a ROS-mediated signaling pathway


Studies have shown that ROS play an important role in cancer therapy; in particular, low-linear energy transfer (LET) radiation exposure is more dependent on ROS generation. Therefore, we hypothesized that ^{125}I seeds may inhibit cell growth and EMT via increased ROS generation in GBM cells. To test this hypothesis, we sought to determine whether ^{125}I seeds were more effective than X-ray in generating ROS. DCF-DA staining and flow cytometric assays showed that the levels of ROS were markedly increased in U251 and U87 cells after 24 hours of ^{125}I seed irradiation. As shown in Figure 4A, the degree of ROS was measured under a microscope, with green fluorescence indicating the creation of ROS. Results indicated that ^{125}I seeds stimulated higher level of ROS in both U87 and U251 cells, as reflected by the intensity of green fluorescence and the percentage of cells carrying ROS. In addition, flow cytometry analysis revealed that ^{125}I seeds were more effective in stimulating ROS generation than X-ray irradiation at the same doses (Figure 4B). For example, the percentage of U251 cells carrying ROS increased from 79.3% to 97.1%. Moreover, the mean fluorescence intensity measured by flow cytometry significantly increased by 1.5-fold in U251 cells and 1.8-fold in U87 cells, compared with the control group. Taken together, these results suggest that the inhibition of GBM cell growth and EMT induced by ^{125}I seeds may be associated with ROS generation.

Increased ROS is also known to produce DNA double-strand breaks (DSBs) with accumulation of the known marker γH2AX . ATM and its downstream kinase Chk1 phosphorylate several targets that regulate DNA repair, cell-cycle checkpoints, and apoptosis [31,32]. Other studies have implicated ROS in the aggressive behavior of cancer [33]. Moreover, we have found that ^{125}I seeds can induce ROS production. Therefore, we



hypothesized that ^{125}I seeds may lead to inhibition of cell growth and invasion via a ROS-mediated signaling pathway. To test this hypothesis, western blotting was

performed and GSH, which can scavenge ROS, was used. The results indicated that pretreatment of cells with GSH obviously decreased DNA damage, G2/M arrest, and




apoptosis (Figure 4C). Interestingly, ^{125}I seed-inhibited EMT associated protein recovered when GSH was applied. To further test whether EMT and cell growth were inhibited via a ROS-mediated signaling pathway, Invasion and growth of GBM cells were measured with and without GSH. The results indicated that pretreatment of cells with GSH recovered the cell invasion suppressed by ^{125}I seeds (Figure 4D). The inhibited colony-formation in GBM cells was also rescued by GSH (Figure 4E). Moreover, when cells were pretreated with GSH, ^{125}I seed-induced cell apoptosis, as reflected by sub-G1, decreased from 13.46% to 1.72%, and the percentage of G2/M cells decreased from 22.85% to 8.23% in U251 (Figure 4F). Taken together, these results suggest that radioactive ^{125}I seeds suppress cell migration and cell growth by activating a ROS-mediated

signaling pathway, and that GSH blocks the ^{125}I seed irradiation-induced inhibition of cell migration and cell growth.

Radioactive ^{125}I seeds exhibit greater *in vivo* anticancer activity than X-ray

An *in vivo* experiment was performed to further evaluate the effect of ^{125}I seed irradiation. This study was performed according to TPS to ensure consistency with clinical therapy. As anticipated, both X-ray and ^{125}I seed irradiation at a cumulative dose of 20 Gy significantly inhibited xenograft tumor growth *in vivo* (Figure 5A and B). Measurement of xenograft tumor volumes showed that the sizes of tumors deriving from the ^{125}I seed and X-ray groups were significantly smaller than those from the



control group by day 12 (Figure 5C). Interestingly, compared with X-ray irradiation, ¹²⁵I seed irradiation more effectively inhibits tumor growth. Moreover, the expression of proteins associated with DNA damage, G2/M, apoptosis and EMT were detected in xenograft tumors by western blotting and IHC. The *in vivo* results indicate that ¹²⁵I seeds caused up-regulation of γH2AX and of the phosphorylation levels of ATM (Ser 1981), ATR (Ser 428), and Cdc2 (Tyr 15), indicating that ¹²⁵I seeds are more effective than X-ray in inhibiting cell growth. Moreover, EMT was inhibited by ¹²⁵I seeds, as reflected by decreased vimentin and ZEB1 (Figure 5D and E). Interestingly, the body weight of the nude mice exposed to X-ray irradiation decreased more significantly than that of the ¹²⁵I irradiation group (data not shown). Taken together, these findings indicate that exposure to ¹²⁵I seed radiation is more effective than X-ray in inhibiting GBM cell growth and EMT *in vivo*.

Discussion and conclusions

GBM is the most common primary central nervous system neoplasm in adults. The annual incidence of malignant gliomas in Western countries is approximately 5 cases per 100,000 people [1]. Unlike most other tumors,

extracranial metastases from GBM are extremely rare [2]. In GBM, standard treatment involves maximal resection followed by concomitant and adjuvant chemoradiotherapy with temozolomide. Even with complete resection, tumors will recur because of the infiltrative and diffuse nature of GBM and its intrinsic radioresistance [34]. Although the comprehensive treatment strategy for GBM has continuously progressed over recent decades, patients with GBM still have an extremely poor prognosis with median survival of 14.6 months [35]. Thus, the discovery of novel treatments that can overcome the diffuse nature of GBM and its intrinsic radioresistance has been an aspiration of scientists in the fields of oncology. Interestingly, previous clinical trials have indicated that ¹²⁵I seeds treatment is an adjuvant therapy to be effective in recurrent GBM [8,36]. Stereotactic brachytherapy (SBT) with ¹²⁵I seeds was confirmed to be an effective and safe treatment for patients with small unresectable or incompletely resected LGG. Moreover, it has been confirmed that radioactive ¹²⁵I seed inhibited the cell growth, migration, and invasion of NPC cells [12]. These encouraging results compel us to embark on testing larger numbers of cancer with this treatment modality. Thus, the current study was performed to investigate the biological effects of ¹²⁵I seeds on GBM. Results


confirmed that radioactive ^{125}I seeds are more effective than X-ray irradiation in inhibiting GBM cell growth. Moreover, EMT in GBM cells was effectively inhibited by ^{125}I seed irradiation *in vitro* and *in vivo*. Further studies indicated that the anticancer effect of ^{125}I seed irradiation was achieved via a ROS-mediated signaling pathway.

Previous studies have reported that ^{125}I seed irradiation is more effective than ^{60}Co - γ ray irradiation in inhibiting cell growth in PANC-1, PC3 prostate cancer, and CL187 colonic cells [12,14]. Therefore, we evaluated the radiosensitivity of GBM cells. The results showed that GBM cells are more sensitive to ^{125}I seed irradiation than to X-ray irradiation. Moreover, a significant increase in the proportion of apoptosis with the involvement caspase-3 and G2/M arrest was observed after ^{125}I seeds irradiation. Similarly to our results, a prolonged accumulation in the G2/M phase and inhibited growth after exposure to ^{125}I seed irradiation has been previously described [12,14]. EMT is a tightly regulated process that occurs during tumor cell metastasis. Without knowledge of the underlying mechanisms, a previous study reported that a sublethal dose of X-ray irradiation promoted the migration and invasion of glioma cells in the border area during postoperative radiotherapy [37]. In contrast, some studies have shown that both proton and carbon ion irradiation significantly decreased cell migration and invasion [38,39]. Therefore, in our study, the effects of ^{125}I seed and X-ray irradiation on GBM cell migration were measured. Interestingly, we showed that X-ray irradiation promoted the migration and invasion of GBM cells, while inhibitory effects were observed in the ^{125}I seed irradiation group. Consistently, EMT of NPC cell lines were inhibited by ^{125}I seeds [12]. Currently, Kim et al. found that activation of the Snail pathway by X-ray was important for radioresistance primarily via EMT induction [40]. Mahabir et al. also described that GBM post irradiation undergo EMT and stemness features throughout the early to late phases by upregulating Snail [30]. Gomez et al. confirmed that non-small cell lung cancer cells survived X-ray irradiation treatment display cancer stem cell and EMT phenotypes, with enhancement of motility and invasiveness and elevated resistance to apoptosis [41]. On the basis of these findings that EMT which can be induced by X-ray irradiation contributes to radioresistance, though a classic EMT is not involved in GBMs progression, we indicate that ^{125}I seeds might be an effective treatment for GBM.

What could be the mechanisms behind the inhibition of GBM cell growth and invasion induced by ^{125}I seed irradiation? Previous studies confirmed that ^{125}I seed irradiation inhibited NPC cell line migration by inactivating VEGF-A/ERK signaling [11]. Kahlert et al. indicated that the activity of the canonical WNT/ β -catenin signaling pathway was directly associated with increased motility of the

GBM cells [18]. However, decreased WNT signaling or VEGF-A were not observed in GBM cells treated with ^{125}I seeds (data not shown). As we all know, the absorption of ionizing radiation by living cells can act indirectly through radiolysis of water, thereby generating ROS that can damage nucleic acids, proteins, and lipids [21]. ROS play an important role in cancer therapy, particularly with low-LET irradiation [19,42]. Therefore, we hypothesized that cell growth and invasion inhibition were induced by ^{125}I seeds with the involvement of increasing ROS. The results indicated that ^{125}I seed irradiation led to a higher level of ROS than X-ray under the same doses. Moreover, pretreatment of cells with GSH rescued the EMT and cell growth suppressed by ^{125}I seeds. Taken together, these results indicate that radioactive ^{125}I seeds inhibit GBM cell invasion and growth via a ROS signaling pathway, and that ROS inhibition can block the ^{125}I seed irradiation-induced inhibition of cell migration and cell growth. Following this, the anticancer action of ^{125}I seed and X-ray irradiation *in vivo* was investigated. As with the *in vitro* findings, the results indicate that ^{125}I seeds exhibit greater anticancer activity than X-ray irradiation *in vivo*.

Increasing evidence suggests that up-regulated ROS induced by chemotherapy or radiotherapy is associated with increased EMT of cancer cells. In our study, however, the results indicate that ROS generated by ^{125}I seeds can prevent GBM cells migration. This may be due to the type of ROS, and the dose of irradiation. Luanpitpong et al. have confirmed that superoxide anions and hydrogen peroxide down-regulate Cav-1 expression and inhibit cell migration and invasion, whereas hydroxyl radicals up-regulate Cav-1 expression and promote cell migration and invasion [33]. Interestingly, Cav-1 was confirmed to be down-regulated by ^{125}I seeds in our study, and the inhibited EMT could be reversed by scavenger of ROS by GSH. Moreover, Urbich et al. have indicated that CD40 ligand inhibits endothelial cell migration by increasing the production of endothelial ROS, and that H_2O_2 -prevented endothelial cell migration can be reversed vitamin C [43]. Although further studies need to perform to confirm the type of ROS generated by ^{125}I seeds, we can conclude that ^{125}I seeds can inhibit EMT and growth GBM cells through increased ROS. Effective treatment options are limited for patients with GBM, especially for recurrent GBM, thus, novel treatment approaches are needed. ^{125}I seeds treatment is an adjuvant therapy that has been shown to be effective in recurrent GBM [7,44]. Permanent placement of ^{125}I seeds for recurrent GBM may prolong survival in patients with recurrent GBM [36]. Darakchiev et al. have indicated that the use of adjuvant therapy combining carmustine wafers and permanent ^{125}I seeds was a good treatment option for patients with recurrent GBM who have undergone previous surgery and



Scheme 1 Proposed signaling pathway by which radioactive ¹²⁵I seeds inhibit epithelial-mesenchymal transition (EMT) and cell growth in glioblastoma multiforme (GBM) cells. ¹²⁵I seeds act indirectly through radiolysis of water, thereby generating ROS and producing DNA damage. DNA damage activates the sensory ATM/ATR kinases and finally results in cell growth inhibition and G2/M arrest. Moreover, ROS induced by ¹²⁵I seeds can inhibit EMT via Cav-1 in GBM cells.

radiation therapy [8]. Overall, previous clinical trials have confirmed that ¹²⁵I was an effective treatment for patients with GBM. However, our study was performed to investigate the biological effects of ¹²⁵I seeds on GBM. Our results indicated that ¹²⁵I seed irradiation was more effective than X-ray irradiation in inhibiting GBM cells via the ROS pathway. Obviously, our data are in line with the majority of published clinical trials studies. These results reported here also confirm and extend previous findings. Though GBM cell growth and invasion were inhibited by radioactive ¹²⁵I seeds in this study, however the signaling pathway via which radioactive ¹²⁵I seeds inhibited GBM cell growth and EMT in human GBM was ROS, which was different from previous study [11].

In summary, we have demonstrated, for the first time, that radioactive ¹²⁵I seeds are more effective than X-ray irradiation in inhibiting GBM cell growth. Moreover, EMT of GBM cells was effectively inhibited by ¹²⁵I seed irradiation. A mechanism study indicated that the cell growth and EMT inhibition in GBM cells was induced by ¹²⁵I seeds with the involvement of a ROS signaling pathway. To our knowledge, this is the first attempt to study the effect of ¹²⁵I seeds on GBM cells. These results suggest that radioactive ¹²⁵I seeds exhibit novel anticancer activity via a ROS signaling pathway (Scheme 1). Furthermore, EMT of GBMs was inhibited by ¹²⁵I seeds via ROS

pathway. These findings have clinical implications for the treatment of GBM patients with ¹²⁵I seeds. Based on the current report, ¹²⁵I seeds either alone or in combination with other treatment may be a better choice for GBM patients. Future studies are needed to determine whether the biological effects of ¹²⁵I seed irradiation identified in GBM cells can be reproduced in other cancer-cell types, thus broadening the significance of the data reported here.

Abbreviations

Cav-1: Caveolin-1; Cdc2: Cell-cycle controller-2; CT: Computed tomography; DAPI: 4'-6-diamino-2-phenylindole; EdU: 5-ethynyl-2'-deoxyuridine; EMT: Epithelial-mesenchymal transition; FBS: Fetal bovine serum; GBM: Glioblastoma multiforme; GSH: Glutathione; MTT: Thiazolyl blue tetrazolium bromide; PARP: Poly-ADP-ribose polymerase; PBS: Phosphate buffered saline; ROS: Reactive oxygen species; SD: Standard deviation; SF: Surviving fraction; TPS: Treatment planning system; TUNEL: Deoxynucleotidyl transferase (TdT)-mediated dUTP- digoxigenin nick-end labeling.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KY carried out the design of experiment and helped to alter the manuscript. YT performed all of the *in vivo* and *in vitro* work and wrote the manuscript. QX carried out the flow cytometry, *in vivo* experiment and western blotting, and helped to draft the manuscript. He Jie carried out the western blotting and IF, XL participated in the MTT assay, colony-formation assay, and EdU assay. TZ carried out the TUNEL assay. YT carried out the annexin V-PI apoptosis and Caspase-3 activity assay. DS performed the statistical analysis and participated in the *in vivo* experiment. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to thank Huanxing, Liu, Ph.D. for their technical assistance and revised the thesis. This study was supported by the National Natural Science Foundation of China-Guangdong Joint Fund (K1060006).

Author details

¹Cancer Research Institute, Southern Medical University, Guangzhou 510, 515 Guangdong Province, People's Republic of China. ²Department of Radiation Oncology, Cancer Center of Guangzhou Medical University, Guangzhou, Guangdong Province, People's Republic of China. ³Department of Oncology, Armed Police Corps Hospital of Guangdong Province, Guangzhou, People's Republic of China.

Received: 17 August 2013 Accepted: 4 November 2014

Published: 19 February 2015

References

- Meyer MA. Malignant gliomas in adults. *N Engl J Med.* 2008;359(5):492–507.
- Wang Y, Jiang T. Understanding high grade glioma: molecular mechanism, therapy and comprehensive management. *Cancer Lett.* 2013;331(2):139–46.
- Wang JJ, Yuan HS, Li JN, Jiang WJ, Jiang YL, Tian SQ. Interstitial permanent implantation of 125I seeds as salvage therapy for re-recurrent rectal carcinoma. *Int J Colorectal Dis.* 2009;24(4):391–9.
- Coatmeur O, Truc G, Barillot I, Horiot JC, Maingon P. Treatment of T1-T2 rectal tumors by contact therapy and interstitial brachytherapy. *Radiother Oncol.* 2004;70(2):177–82.
- Jiang YL, Meng N, Wang JJ, Ran WQ, Yuan HS, Qu A, et al. Percutaneous computed tomography/ultrasonography-guided permanent iodine-125 implantation as salvage therapy for recurrent squamous cell cancers of head and neck. *Cancer Biol Ther.* 2010;9(12):959–66.
- Meng N, Jiang YL, Wang JJ, Ran WQ, Yuan HS, Qu A, et al. Permanent implantation of iodine-125 seeds as a salvage therapy for recurrent head and neck carcinoma after radiotherapy. *Cancer Invest.* 2012;30(3):236–42.
- Chan TA, Weingart JD, Parisi M, Hughes MA, Olivi A, Borzillary S, et al. Treatment of recurrent glioblastoma multiforme with GliSite brachytherapy. *Int J Radiat Oncol Biol Phys.* 2005;62(4):1133–9.
- Darakchiev BJ, Albright RE, Breneman JC, Warnick RE. Safety and efficacy of permanent iodine-125 seed implants and carmustine wafers in patients with recurrent glioblastoma multiforme. *J Neurosurg.* 2008;108(2):236–42.
- Ruge MI, Simon T, Suchorska B, Lehrke R, Hamisch C, Koerber F, et al. Stereotactic brachytherapy with iodine-125 seeds for the treatment of inoperable low-grade gliomas in children: long-term outcome. *J Clin Oncol.* 2011;29(31):4151–9.
- Lopez WO, Trippel M, Doostkam S, Reithmeier T. Interstitial brachytherapy with iodine-125 seeds for low grade brain stem gliomas in adults: diagnostic and therapeutic intervention in a one-step procedure. *Clin Neurol Neurosurg.* 2013;115(8):1451–6.
- Tian Y, Xie Q, Tian Y, Liu Y, Huang Z, Fan C, et al. Radioactive (125I) Seed Inhibits the Cell Growth, Migration, and Invasion of Nasopharyngeal Carcinoma by Triggering DNA Damage and Inactivating VEGF-A/ERK Signaling. *PLoS One.* 2013;8(9):e74038.
- Wang J, Liao A, Zhuang H, Zhao Y. The direct biologic effects of radioactive 125I seeds on pancreatic cancer cells PANC-1, at continuous low-dose rates. *Cancer Biother Radiopharm.* 2009;24(4):409–16.
- Zhuang HQ, Wang JJ, Liao AY, Wang JD, Zhao Y. The biological effect of 125I seed continuous low dose rate irradiation in CL187 cells. *J Exp Clin Cancer Res.* 2009;28:12.
- Liao A, Wang J, Zhuang H, Zhao Y. Relative biological effectiveness and cell-killing efficacy of continuous low-dose-rate 125I seeds on prostate carcinoma cells in vitro. *Integr Cancer Ther.* 2010;9(1):59–65.
- Wu F, Ye X, Wang P, Jung K, Wu C, Douglas D, et al. Sox2 suppresses the invasiveness of breast cancer cells via a mechanism that is dependent on Twist1 and the status of Sox2 transcription activity. *BMC Cancer.* 2013;13:317.
- Haslehurst AM, Koti M, Dharsee M, Nuin P, Evans K, Geraci J, et al. EMT transcription factors snail and slug directly contribute to cisplatin resistance in ovarian cancer. *BMC Cancer.* 2012;12:91.
- Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *J Clin Invest.* 2009;119(6):1420–8.
- Kahlert UD, Nikkhah G, Maciaczyk J. Epithelial-to-mesenchymal(like) transition as a relevant molecular event in malignant gliomas. *Cancer Lett.* 2013;331(2):131–8.
- Trachootham D, Alexandre J, Huang P. Targeting cancer cells by ROS-mediated mechanisms: a radical therapeutic approach? *Nat Rev Drug Discov.* 2009;8(7):579–91.
- Nathan FM, Singh VA, Dhanoa A, Palanisamy UD. Oxidative stress and antioxidant status in primary bone and soft tissue sarcoma. *BMC Cancer.* 2011;11:382.
- Azzam EI, Jay-Gerin JP, Pain D. Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Cancer Lett.* 2012;327(1–2):48–60.
- Romilda C, Marika P, Alessandro S, Enrico L, Marina B, Andromachi K, et al. Oxidative DNA damage correlates with cell immortalization and mir-92 expression in hepatocellular carcinoma. *BMC Cancer.* 2012;12:177.
- Mikkelsen RB, Wardman P. Biological chemistry of reactive oxygen and nitrogen and radiation-induced signal transduction mechanisms. *Oncogene.* 2003;22(37):5734–54.
- Chen T, Wong YS. Selenocystine induces reactive oxygen species-mediated apoptosis in human cancer cells. *Biomed Pharmacother.* 2009;63(2):105–13.
- Cheng C, Kong X, Wang H, Gan H, Hao Y, Zou W, et al. Trihydrophobin 1 interacts with PAK1 and Regulates ERK/MAPK Activation and Cell Migration. *J Biol Chem.* 2009;284(13):8786–96.
- De Vos M, Schreiber V, Dantzer F. The diverse roles and clinical relevance of PARPs in DNA damage repair: current state of the art. *Biochem Pharmacol.* 2012;84(2):137–46.
- Riffell JL, Lord CJ, Ashworth A. Tankyrase-targeted therapeutics: expanding opportunities in the PARP family. *Nat Rev Drug Discov.* 2012;11(12):923–36.
- Burgess A, Vigneron S, Brioudes E, Labbe JC, Lorca T, Castro A. Loss of human Greatwall results in G2 arrest and multiple mitotic defects due to deregulation of the cyclin B-Cdc2/PP2A balance. *Proc Natl Acad Sci U S A.* 2010;107(28):12564–9.
- Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med.* 2005;352(10):997–1003.
- Mahabir R, Tanino M, Elmansuri A, Wang L, Kimura T, Itoh T, et al. Sustained elevation of Snail promotes glial-mesenchymal transition after irradiation in malignant glioma. *Neuro-Oncol.* 2014;16(5):671–85.
- Yamamori T, Yasui H, Yamazumi M, Wada Y, Nakamura Y, Nakamura H, et al. Ionizing radiation induces mitochondrial reactive oxygen species production accompanied by upregulation of mitochondrial electron transport chain function and mitochondrial content under control of the cell cycle checkpoint. *Free Radic Biol Med.* 2012;53(2):260–70.
- Lemon JA, Rollo CD, Boreham DR. Elevated DNA damage in a mouse model of oxidative stress: impacts of ionizing radiation and a protective dietary supplement. *Mutagenesis.* 2008;23(6):473–82.
- Luanpitpong S, Talbott SJ, Rojanasakul Y, Nimmannit U, Pongrakhananon V, Wang L, et al. Regulation of lung cancer cell migration and invasion by reactive oxygen species and caveolin-1. *J Biol Chem.* 2010;285(50):38832–40.
- Stupp R, Tonn JC, Brada M, Pentheroudakis G. High-grade malignant glioma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2010;21 Suppl 5:v190–3.
- Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med.* 2005;352(10):987–96.
- Patel S, Breneman JC, Warnick RE, Albright Jr RE, Tobler WD, van Loveren HR, et al. Permanent iodine-125 interstitial implants for the treatment of recurrent glioblastoma multiforme. *Neurosurgery.* 2000;46(5):1123–8. discussion 1128–1130.
- Wild-Bode C, Weller M, Rimmer A, Dichgans J, Wick W. Sublethal irradiation promotes migration and invasiveness of glioma cells: implications for radiotherapy of human glioblastoma. *Cancer Res.* 2001;61(6):2744–50.
- Ogata T, Teshima T, Kagawa K, Hishikawa Y, Takahashi Y, Kawaguchi A, et al. Particle irradiation suppresses metastatic potential of cancer cells. *Cancer Res.* 2005;65(1):113–20.
- Liu C, Lin J, Zhao L, Yang Y, Gao F, Li B, et al. Gamma-ray irradiation impairs dendritic cell migration to CCL19 by down-regulation of CCR7 and induction of cell apoptosis. *Int J Biol Sci.* 2011;7(2):168–79.
- Kim E, Youn H, Kwon T, Son B, Kang J, Yang HJ, et al. PAK1 Tyrosine Phosphorylation Is Required to Induce Epithelial-Mesenchymal Transition and Radioresistance in Lung Cancer Cells. *Cancer Res.* 2014;74(19):5520–31.
- Gomez-Casal R, Bhattacharya C, Ganesh N, Bailey L, Basse P, Gibson M, et al. Non-small cell lung cancer cells survived ionizing radiation treatment

display cancer stem cell and epithelial-mesenchymal transition phenotypes. *Mol Cancer*. 2013;12(1):94.

42. Shao J, Dai Y, Zhao W, Xie J, Xue J, Ye J, et al. Intracellular distribution and mechanisms of actions of photosensitizer Zinc(II)-phthalocyanine solubilized in Cremophor EL against human hepatocellular carcinoma HepG2 cells. *Cancer Lett*. 2013;330(1):49–56.
43. Urbich C, Dernbach E, Aicher A, Zeiher AM, Dimmeler S. CD40 ligand inhibits endothelial cell migration by increasing production of endothelial reactive oxygen species. *Circulation*. 2002;106(8):981–6.
44. Wernicke AG, Sherr DL, Schwartz TH, Pannullo SC, Stieg PE, Boockvar JA, et al. The role of dose escalation with intracavitary brachytherapy in the treatment of localized CNS malignancies: outcomes and toxicities of a prospective study. *Brachytherapy*. 2010;9(1):91–9.

doi:10.1186/1471-2407-15-1

Cite this article as: Tian *et al.*: Radioactive ^{125}I seeds inhibit cell growth and epithelial-mesenchymal transition in human glioblastoma multiforme via a ROS-mediated signaling pathway. *BMC Cancer* 2015 **15**:1.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

