

REVIEW ARTICLE

Radioprotection by Plant Products: Present Status and Future Prospects

Rajesh Arora, Damodar Gupta†, Raman Chawla, Ravinder Sagar, Ashok Sharma, Raj Kumar, Jagdish Prasad, Surinder Singh, Namita Samanta and Rakesh Kumar Sharma*

Division of Radiopharmaceuticals and Radiation Biology, Institute of Nuclear Medicine and Allied Sciences, Defence Research and Development Organization, Brig. SK Mazumdar Road, Delhi-110 054, India

The development of radioprotective agents has been the subject of intense research in view of their potential for use within a radiation environment, such as space exploration, radiotherapy and even nuclear war. However, no ideal, safe synthetic radioprotectors are available to date, so the search for alternative sources, including plants, has been on going for several decades. In Ayurveda, the traditional Indian system of medicine, several plants have been used to treat free radical-mediated ailments and, therefore, it is logical to expect that such plants may also render some protection against radiation damage. A systematic screening approach can provide leads to identifying potential new candidate drugs from plant sources, for mitigation of radiation injury. This article reviews some of the most promising plants, and their bioactive principles, that are widely used in traditional systems of medicine, and which have rendered significant radioprotection in both *in vitro* and *in vivo* model systems. Plants and their constituents with pharmacological activities that may be relevant to amelioration of radiation-mediated damage, including antiemetic, antiinflammatory, antioxidant, cell proliferative, wound healing and haemopoietic stimulatories are also discussed. Copyright © 2005 John Wiley & Sons, Ltd.

Keywords: herbal radioprotection; ionizing radiation; plants; traditional medicine; radioprotectors; bioactive principles.

INTRODUCTION

The development of effective radioprotectors and radiorecovery drugs is of great importance in view of their potential application during both planned radiation exposure (e.g. radiotherapy) and unplanned radiation exposure (e.g. in the nuclear industry, natural background radiation emanating from the earth or other sources) (Arora and Goel, 2000; Bump and Malaker, 1998; Coleman *et al.*, 2003; Moulder, 2002; Nair *et al.*, 2001). These drugs are also likely to be useful in nuclear warfare to provide protection to personnel (Giambaressi and Jacobs, 1987). Over the past 50 years, research in the development of radioprotectors worldwide has focused on screening a plethora of chemical and biological compounds (Maisin, 1998; Sweeney, 1979; Weiss *et al.*, 1990; Weiss and Landauer, 2003). Numerous drugs of both synthetic and natural origin, e.g. antioxidants (Hahn *et al.*, 1994, 1999; Kumar *et al.*, 2002; Mitchell *et al.*, 1991, 2000; Vijaylaxmi *et al.*, 1996), cytoprotective agents (Links and Lewis, 1999), angiotensin-converting enzyme (ACE) inhibitors (Molteni *et al.*, 2000; Moulder *et al.*, 1998a,b), or angiotensin-II type-1 (AT1) receptor antagonists (e.g.

losartan) (Moulder *et al.*, 1998c), metalloelements (Matsubara *et al.*, 1987; Miko *et al.*, 1998; Satoh *et al.*, 1989), immunomodulators (Furuse *et al.*, 1997; Guenechea *et al.*, 1997; Kalechman *et al.*, 1995a,b; Landauer *et al.*, 1997; Real *et al.*, 1992; Weiss and Simic, 1988), sulphhydryl compounds (Capizzi and Oster, 1995; Livesey and Reed, 1987; Ramnath *et al.*, 1997; Spencer and Goa, 1995; Tannehill and Mehta, 1996; Wasserman, 1994; Weiss, 1997), lipopolysaccharides and prostaglandins (Hanson *et al.*, 1988; Joshima *et al.*, 1992; Riehl *et al.*, 2000; Van Buul *et al.*, 1999), vitamin A, C and E (Haranpanhalli *et al.*, 1994) and DNA binding ligands (Denison *et al.*, 1992; Martin *et al.*, 1996; Martin and Anderson, 1999) have been tested in both *in vitro* and *in vivo* models, and in human clinical trials to mitigate injuries caused by ionizing radiation exposure in the sublethal to supralethal range. Combinations of agents have also been tested with little success (Weiss *et al.*, 1990; Kumar and Gupta, 2002). The potential combination of differential radiomodifiers with metabolic modulators has been demonstrated in cell culture and animal models (Dwarakanath *et al.*, 1999; Sharma *et al.*, 2000a,b; Sharma and Jain, 2002).

Among the molecular radioprotectors, WR-2721 [S-2-(3-aminopropyl-amino) ethyl phosphorothioic acid], also known as amifostine, ethiophos (USA) or gammaphos (former USSR), is the most thoroughly investigated radioprotective drug, initially developed at the Walter Reed Army Research Institute, USA under the Antiradiation Drug Development Program of the US Army Medical Research and Development Command (Schuchter and Glick, 1993; Sweeney, 1979). However, the radioprotective effects

* Correspondence to: Dr R. K. Sharma, Division of Radiopharmaceuticals and Radiation Biology, Institute of Nuclear Medicine and Allied Sciences, Brig. SK Mazumdar Road, Delhi-110 054, India.
E-mail: rks@inmas.org

† Present address: Department of Cancer Biology, Cleveland Clinic Foundation, Lerner Research Institute, 9500 Euclid Avenue, Ohio, 44195, USA.

of phosphorothioate compounds, including amifostine, are short term, and associated with severe side effects (e.g. nausea, vomiting, diarrhoea, hypotension, hypocalcaemia, nephro- and neuro-toxicity) at clinically effective doses (Cairnie, 1983; Kligerman *et al.*, 1984; Glover *et al.*, 1983; Landauer *et al.*, 1987). These limitations have greatly restricted their clinical use. Despite its drawbacks, amifostine (Ethyol®) is the only radioprotector that has been approved by the Food and Drug Administration (FDA), USA. Amifostine is being used clinically for ameliorating the incidence of xerostomia (dry mouth) in patients undergoing radiotherapy for the treatment of head and neck cancer (Brizel *et al.*, 2000).

In recent years, an array of haemopoietic growth factors and cytokines such as interleukin-7 (Bolotin *et al.*, 1996), interleukin-11 (Van der Meeren *et al.*, 2002), granulocyte-colony stimulating factor (G-CSF) (Russel *et al.*, 2000), granulocyte, macrophage-colony stimulating factor (GM-CSF) (Mettler and Guskova, 2001; Vose and Armitage, 1995), stem cell factor (SCF) (Zsebo *et al.*, 1992), antiapoptotic cytokine combinations (Herodin *et al.*, 2003), and epithelial cell-specific growth factors such as keratinocyte growth factor (Dorr *et al.*, 2001) have been used to mitigate radiation-induced damage and to augment recovery of stem cells and their precursors after radiation exposure. However, the success with these compounds has also been limited.

The fact remains that to date there is not a single radioprotective agent available which meets all the prerequisites of an ideal radioprotector, i.e. produces no cumulative or irreversible toxicity, offers effective long-term protection, possesses a shelf life of 2–5 years, and can be easily administered (Maisin, 1998; Coleman *et al.*, 2003). In view of this, the search for newer, less toxic and more effective radioprotector drugs continues.

Plants have been utilized since time immemorial for curing diseases. Even today, nearly 70% of the world's population is dependent on plants for handling their health related problems (Fabricant and Farnsworth, 2001).

A number of plants have been utilized successfully for the treatment of free radical-mediated diseases in humans such as rheumatoid arthritis, atherosclerosis,

cancer, Alzheimer's disease, Parkinson's disease, aging and several other conditions including inflammatory diseases (Singh *et al.*, 2000; Das, 2002). It is, therefore, reasonable to expect that plants may contain certain compounds that can protect against radiation-induced reactive oxygen species (ROS)-mediated damage.

A number of medicinal plants evaluated for their radioprotective efficacy have shown protective effects against the damaging effects of ionizing radiation (Arora and Goel, 2000; Arora *et al.*, 2003a,b, 2004; Ben-Hur and Fulder, 1981; Gupta *et al.*, 2003a; Jagetia *et al.*, 2002; Jagetia and Baliga, 2002a,b; Kamat *et al.*, 1999; Maharwal *et al.*, 2003; Shen *et al.*, 1989; Shimoi *et al.*, 1994, 1996; Uma Devi *et al.*, 1999). Plant extracts eliciting radioprotective efficacy contain a plethora of compounds including antioxidants, immunostimulants, cell proliferation stimulators, antiinflammatory and antimicrobial agents, some of which may act in isolation as well as in combination with other constituents from the same plant. They may also augment the efficacy of compounds present in other plant species, to provide protection against radiation-induced damage (Fig. 1). In traditional Ayurveda, Chinese, Japanese, Korean, Siddha, European, Tibetan and Unani systems of medicine it is a common practice to use a multi-plant formulation for treating diseases. Synergistic effects may be present, and some of the toxic effects generated by active constituents of one plant may be countered by other constituents present. Most studies using natural plant products have focused on evaluation of radioprotective efficacy of whole extracts or polyherbal formulations, and in some cases fractionated extracts and isolated constituents (Abraham *et al.*, 1993; Agarwal and Nagaratnum, 1981; Arora and Goel, 2000; Arora *et al.*, 2003a,b, 2004; Hsu *et al.*, 1993; Gupta *et al.*, 2003a; Hsu *et al.*, 1997; Kim *et al.*, 2003; Kumar *et al.*, 1996; Kure, 1992; Narimanov 1993; Ohara *et al.*, 2001; Pande *et al.*, 1998b; Wang *et al.*, 1992; Wang, 1996; William *et al.*, 1996; Yang *et al.*, 1997) (Table 1; Fig. 1).

There is a need to critically review the radioprotective properties of these plant products. The present review attempts to provide an overview of plants and their constituents with pharmacological activities relevant to radioprotection, radiorecovery, and treatment of radiation injuries, including antioxidant, antiinflammatory,




Figure 1. Some mechanisms by which natural plant products render radioprotection.

Table 1. Traditional herbal plants showing therapeutic activities relevant to radioprotection

Plant	Family	Traditional uses and radioprotective efficacy	Reference
<i>Aegle marmelos</i> Corr. Ex Roxb	Rutaceae	To promote digestion, treat colic, diarrhoea and dysentery, intermittent fever, melancholia and heart palpitation <i>A. marmelos</i> provided protection against radiation-induced sickness and mortality in mice. Optimum protective dose was 15 mg/kg b.w. [1/82 of the LD ₅₀ dose (2250 mg/kg b.w.)]	Jagetia <i>et al.</i> , 2003e
<i>Acanthopanax senticosus</i> Harms (<i>Shigoka</i>)	Araliaceae	To restore normal functioning of spleen and kidneys. Also used as a remedy for bronchitis, heart ailments and rheumatism Pre-irradiation administration of Shigoka extract (5 mg/kg b.w.; –24 h; i.p.) rendered maximum survival (80%), while post-irradiation administration (+12 h; 9.5 Gy) exhibited 30% survival. The extract also increased leukocyte counts and diminished cerebral haemorrhage	Yonezawa <i>et al.</i> , 1989 Miyanomae and Frindel, 1988.
<i>Ageratum conyzoides</i> L.	Asteraceae	In India <i>A. conyzoides</i> leaves are applied to cuts and sores, while the juice is considered as antilithic. An alcoholic extract of <i>A. conyzoides</i> (75 mg/kg b.w.; –1 h) effectively protected mice against 10 Gy-induced gastrointestinal and bone marrow related death (DMF: 1.3)	Jagetia <i>et al.</i> , 2003d; Satyavati <i>et al.</i> , 1987
<i>Allium cepa</i> L.	Alliaceae	Dried bulb has been used orally for treating diabetes, dropsy, colic, diabetes, dysentery, fever, chronic bronchitis, body heat and as an emmenagogue Administration of the dried bulb of <i>Allium cepa</i> at a concentration of 20 mg/kg was active against x-irradiation	Bakina <i>et al.</i> , 1967
<i>Allium sativum</i> L. <i>Gaertn</i>	Alliaceae	The plant has been reported to possess antioxidant, antimicrobial, antitumour, antimutagenic, antiinflammatory, antiviral and antiulcer properties. Radioprotective efficacy of aged garlic extract (produced by storing sliced raw garlic in 15–20% ethanol for 20 months and containing compounds such as S-allylcysteine, S-allylmercaptocysteine, allixin and selenium which are stable, highly bioavailable and possess significant antioxidant and anticarcinogenic properties) has been reported	Reeve <i>et al.</i> , 1993; Singh <i>et al.</i> , 1995a; Gupta, 1996
<i>Aloe arborescens</i>	Liliaceae	Acts as a cell proliferant, healer, demuculent and allergy reducer. Topically it is used for skin ulcers, burns, irritations and bites An extract of <i>Aloe arborescens</i> (AA S6-3-b) provided protection to mouse skin against soft x-irradiation by scavenging hydroxyl radicals and reducing alterations in enzyme activity (SOD and glutathione peroxidase activity)	Sato <i>et al.</i> , 1990
<i>Archangelica officinalis</i> Hoffm.	Umbelliferae	Used in traditional Chinese medicine (TCM) to promote fertility, relieve fatigue, migraine and dysmenorrhoea, and ease anxiety and nervous tension Administration of a combination of <i>Archangelica officinalis</i> and <i>Ledum palustre</i> extracts to mice 5–15 min before irradiation [7.5 Gy (LD _{90/30})] rendered 70% survival (DMF: 1.48)	Narimanov <i>et al.</i> , 1991; Narimanov, 1993.
<i>Angelica sinensis</i> (Oliver) Diels	Apiaceae	Used in TCM to replenish blood, improve the rhythmicity and tonicity of uterine muscles, as an emollient and laxative for chronic constipation in aged and debilitated humans. <i>Angelica sinensis</i> is used externally for burns <i>Angelica sinensis</i> root extract, administered to mice via i.v. route (post-irradiation), restored 80% of pregnancy rate vs none in controls. The polysaccharide fraction, containing a ferulic acid, increased survival in irradiated mice (>30 days) by promoting haemopoietic stem cell proliferation	Liu and Xiao, 1993; Mei <i>et al.</i> , 1988, 1991
<i>Amaranthus paniculatus</i> Linn.	Amaranthaceae	It is used for purifying blood and treating scrofulous sores <i>A. paniculatus</i> leaf extract (600 mg/kg b.w./ day for 2 wks) protected mice against 5 Gy by reducing lipid peroxidation, glycogen and cholesterol levels in brain	Verma <i>et al.</i> , 2003

Table 1. (continued)

Plant	Family	Traditional uses and radioprotective efficacy	Reference
<i>Curcuma longa</i> Linn.	Zingiberaceae	<i>Curcuma longa</i> rhizome is widely used in Indian cuisine as well as in traditional medicine. Pharmacological activities include antiinflammatory, anti-HIV, antibacteria, antitumour, antioxidant and nematocidal effects. Curcumin (diferuloylmethane) has been reported to render radioprotective effect	Ammon, 1993; Ammon and Wahl, 1991; Chaudhary <i>et al.</i> , 1999
<i>Glycyrrhiza glabra</i> L.	Fabaceae	<i>Glycyrrhiza glabra</i> finds use in both Indian and TCM due to its antibacterial, antiinflammatory, antiviral, antimutagenic, antioxidant free radical scavenging and immunomodulating activities <i>G. glabra</i> (70% methanol extract; 100 µg/mL) protected rat microsomal membranes from γ -radiation (up to 500 Gy) induced lipid peroxidation, while 20 µg/mL could provide almost complete (99%) protection to plasmid (pBR322) DNA from radiation-induced strand breaks	Agarwal and Singh, 1999; Belinky <i>et al.</i> , 1998; Kovalenko <i>et al.</i> , 2003; Shetty <i>et al.</i> , 2002
<i>Hypericum perforatum</i> Linn.	Hypericaceae	Traditionally used in several European countries and USA for healing of wounds, depression, nervous disorders, insomnia, Herpes infections, inflammations and menstrual complaints. In India, it is used as an antihelmintic and emmenagogue In murine model, <i>Hypericum perforatum</i> aqueous extract protected bone marrow and intestinal mucosa against x-ray in a concentration and time-dependent manner	Smyshlieva and Kudriashov, 1992
<i>Lycium chinense</i>	Solanaceae	Used in TCM as a tonic for general debility, replenishment of blood, eyesight improvement, vertigo, lumbago and impotence Administration of root extract (500 mg/kg body wt) prior to x-irradiation significantly improved the recovery of leukocyte, erythrocyte and thrombocyte counts and haematocrit in ICR strain mice	Hsu <i>et al.</i> , 1999
<i>Mentha arvensis</i> Linn.	Lamiaceae	It has carminative, antiseptic, refrigerant, stimulant, emmenagogue and diuretic properties Pre-irradiation treatment with chloroform extract protected mice against gastrointestinal and bone marrow death (DMF: 1.2)	Jagetia and Baliga, 2002a
<i>Moringa oleifera</i> Lam.	Moringaceae	<i>M. oleifera</i> is used in Ayurveda to treat asthma, gout, rheumatism, inflammation, epilepsy, cardiac and circulatory disorders, nervous debility and healing of wounds. Pre-treatment (i.p.) with a leaf extract (150 mg/kg; single dose) significantly reduced the percent aberrant cells in metaphase chromosomes to normal range by day 7 post-irradiation in mice	Rao <i>et al.</i> , 2001
<i>Syzygium cumini</i> L. Skeels	Myrtaceae	In Ayurveda, <i>S. cumini</i> is used to treat bronchitis, asthma, dyspepsia, diabetes, ulcers and blood impurities. Treatment of human peripheral blood lymphocytes with <i>S. cumini</i> leaf extract (0–100 µg/mL) before γ -radiation (3 Gy) significantly reduced micronuclei-induction	Jagetia <i>et al.</i> , 2002
<i>Tephrosia purpurea</i> (L.) Pers.	Fabaceae	<i>T. purpurea</i> roots are used to treat snake bite; diarrhoea, liver and spleen disorders, inflammation, boils and pimples. <i>Tephrosia</i> extract (200 mg/kg b.wt) protected Swiss albino mice against radiation (5 Gy)-induced haemopoietic injury	Taraphdar <i>et al.</i> , 2002

immunostimulatory, wound healing and antimicrobial effects, are discussed.

BIOLOGICAL ACTIVITIES RELEVANT TO RADIATION PROTECTION

Antiemetic activity

Exposure to ionizing radiation induces nausea and vomiting, particularly in scenarios relevant to radiation disaster. Plants with antiemetic activity, e.g. *Centella asiatica*, *Mentha piperita*, *Zingiber officinale* have been evaluated for their ability to provide radiation protection (Srivastava *et al.*, 1997; Meyer *et al.*, 1995; Phillips *et al.*, 1993). *C. asiatica* rendered protection against radiation-induced conditioned taste aversion in rats (Shobi and Goel, 2001). An ethanol extract of *Ginkgo biloba*, along with ginger extract, has been shown to have antiemetic activity as observed by conditioned taste aversion (CTA) studies (Frisch *et al.*, 1995), and these workers opined that it might be useful in achieving radioprotection too. In human clinical trials, *Zingiber officinale* (ginger) has been reported to ameliorate chemotherapy-induced nausea and vomiting and to protect against gastrointestinal haemorrhage (Meyer *et al.*, 1995). Recently, Jagetia *et al.* (2003b) have also shown the radioprotective property of *Zingiber officinale*. The antiemetic principles present in *Zingiber officinale* include shogaols (6-, 8-, 10-shogaol) and gingerols (6-, 8-, 10-gingerol) (Akita *et al.*, 1998; Yang *et al.*, 2002). However, the role of *Zingiber officinale* and its bioactive compounds in mitigation of radiation-induced emesis has not been worked out so far, and the same remains true for several other plants.

With the growing need for behavioural radioprotectors, in view of their potential applications during space exploration and rescue operations, there is a need to search for more effective antiemetic drugs from plants.

Antiinflammatory activity

A number of plants and their bioactive constituents, including flavonoids, exhibit antiinflammatory properties (Handa *et al.*, 1992; Middleton *et al.*, 2000) and the radioprotective response in several cases is mediated by this effect. Novel antiinflammatory drugs could, therefore, be useful in ameliorating radiation damage with less severe side effects than the synthetic drugs currently available. *Glycyrrhiza glabra*, liquorice, a plant with antiinflammatory and chemopreventive properties (Jo *et al.*, 2004), is known to possess radioprotective properties. Topical application of glycyrrhizin, an active constituent of the rhizome, has been shown to exhibit antiinflammatory properties (Ozaki and Ono, 2002). A number of plants, e.g. *Allium sativum*, *Aloe vera*, *Tinospora cordifolia*, *Hippophae rhamnoides*, *Curcuma longa*, *Centella asiatica*, *Stephania tetrantra*, *Spirulina platensis*, *Syzygium cumini*, *Ocimum sanctum*, *Moringa oleifera*, *Zingiber officinale*, *Eleutherococcus senticosus* (Ammon and Wahl, 1991; Ben-Hur and Fulder, 1981; Chen *et al.*, 1999; Ezeamuzie *et al.*, 1996; Jaiswal and Bordia, 1996; Park *et al.*, 1998; Penna

et al., 2003; Thomson *et al.*, 2002; Udupa *et al.*, 1994; Vazquez *et al.*, 1996) and some of their bioactive constituents such as quercetin, curcumin, c-phycoyanin, allicin, gingerol, caffeine exhibit antiinflammatory properties (Hebbar *et al.*, 2002; Romay *et al.*, 1998; Sidhu *et al.*, 1999; Zhang *et al.*, 1997) and may have potential in the management of radiation injury. Future developments may head towards evaluation of more herbal products that either stimulate the production of antiinflammatory cytokines (e.g. IL-10, IL-13) or selectively inhibit radiation-induced signal transduction pathways.

Antimicrobial activity

One of the major causes of death following radiation-induced myelosuppression is infection arising primarily from translocated endogenous Gram-negative gastrointestinal bacteria (Gordon *et al.*, 1955). A number of antibiotics e.g. aminoglycosides, such as gentamycin, netilmycin, tobramycin and amikacin, ureido-penicillins and monobactams have been used for therapeutic purposes post-irradiation to suppress infection. Plants such as *Ocimum sanctum*, *Podophyllum hexandrum*, *Mentha arvensis*, *Stephania tetrantra*, *Hypericum perforatum*, *Syzygium cumini*, *Moringa oleifera*, *Terminalia chebula*, *Curcuma longa* that exhibit antimicrobial activity have been reported to be radioprotective (Agarwal and Singh, 1999; Eilert *et al.*, 1981; Jagetia and Baliga, 2002a; Singh and Shah, 1994; Sakar and Tamer, 1990; Schapoval *et al.*, 1988). Compounds reported to have antimicrobial activity include benzopyrans, xanthenes, flavonoids and tannins (Ishiguro *et al.*, 1986). The coming years are likely to witness the application of plant products which particularly can stimulate radiorecovery via their antimicrobial effect.

Antioxidant activity

Antioxidants protect the cell against reactive oxygen/nitrogen species (ROS/RNS) by scavenging the free radicals in the cellular milieu. The increased levels of ROS/RNS generated during irradiation have been shown to be effectively scavenged by some antioxidants present in plants (Gupta *et al.*, 2003a; Uma Devi *et al.*, 2000). Antioxidants delay the oxidation of biomolecules by inhibiting the initiation and propagation of oxidizing chain reactions thereby interfering with the initiation of apoptosis (Halliwell and Gutteridge, 1989, 1990). Antioxidants can be phenolic compounds (tocopherols, flavonoids and phenolic acids), nitrogen compounds (alkaloids, amines, amino acids and chlorophyll derivatives) or carotenoids as well as vitamins including ascorbic acid (Beckman and Ames, 1998; Bloor *et al.*, 2000; Burton *et al.*, 1982; Choi *et al.*, 2002; Diplock *et al.*, 1998; Duthie and Crozier, 2000; Duthie *et al.*, 1997; Havsteen, 1984; Jadhav and Bhutani, 2002; Kandaswami and Middleton, 1994; Parshad *et al.*, 1998; Scarterzzini and Speroni, 2000). A relation between the antioxidant property and radiation protection by plant flavonoids and phenols was suggested by Shimoi *et al.* (1994, 1996) and Emerit *et al.* (1997a,b). Antioxidant molecules have been shown to be present in plant species belonging to different levels of organization (from simple to

complex): algae (Karpov *et al.*, 2000; Qishen *et al.*, 1989; Sarma *et al.*, 1993; Singh *et al.*, 1995b; Upasani and Balaraman, 2003; Zozulia and Iurchenko, 2000), gymnosperms (Alaoui-Youssefi *et al.*, 1999) and angiosperms and are known to provide protection from oxidative damage in biological systems. The radioprotective property of plants including *Asparagus racemosus*, *Potentilla alba*, *Ocimum sanctum*, *Podophyllum hexandrum*, *Stephania tetrantra*, *Tinospora cordifolia*, *Hippophae rhamnoides*, *Zingiber officinalis*, *Centella asiatica*, *Ginkgo biloba*, *Syzygium cumini*, *Ligusticum wallichii*, *Vitis vinifera*, *Portulaca oleracea*, *Panax ginseng*, and the ability to extend life span, has been attributed to the presence of antioxidant molecules in these plants (Arora and Goel, 2000; Arora *et al.*, 2003a,b, 2004; Bestwick and Milne, 2001; Cao *et al.*, 1993; Castillo *et al.*, 2000; Ganasoundari *et al.*, 1997a,b, 1998; Goel *et al.*, 2002a,b; Gupta *et al.*, 2003a; Shetty *et al.*, 2002; Gohil *et al.*, 2000; Jagetia *et al.*, 2003b; Jagetia and Baliga, 2003; Jadhav and Bhutani, 2002; Kamat *et al.*, 2000a,b; Landauer *et al.*, 2000; Pincemail *et al.*, 1989; Shimoi *et al.*, 1994; Uma Devi *et al.*, 2000; Yoshioka, 1997; Zhang and Zhang, 1990).

Plants with radioprotective properties have been shown almost invariably to possess antioxidant biomolecules. The radioprotective effect of antioxidant molecules such as eugenol from *Zingiber officinalis*, genistein from *Glycine max*; orientin, vicenin and ursolic acid from *Ocimum sanctum*, curcumin from *Curcuma longa*, bixin from *Bixa orellana*, quercetin from *Podophyllum hexandrum*, β -carotene from the heat-tolerant algae *Dunaliella baradwiil*, luteolin from *Aspalanthus linearis*, allicin from *Allium sativum*, glycyrrhizin from *Glycyrrhiza glabra*, caffeine from *Coffea arabica*, flavan-3-ols (procyanidins) from *Vitis vinifera*, flavone glycosides and terpenes from *Ginkgo biloba*, silymarin from *Silybum marianum*, epigallocatechin from 'Thea viridis' (green tea percolate) and melatonin (N-acetyl-5-methoxytryptamine) from *Hypericum perforatum*, *Silybum marianum*, *Lycium spp.* has largely been attributed to the antioxidative properties of these compounds (Abraham *et al.*, 1993; Alaoui-Youssefi *et al.*, 1999; Ben Amotz *et al.*, 1996, 1998; Booth *et al.*, 1999; Chaudhary *et al.*, 1999; Devasagayam and Kesavan, 1996; George *et al.*, 1999; Inano and Onado, 2002; Kamat *et al.*, 2000a; Kumar *et al.*, 2001; Kim *et al.*, 2003; Kropacova *et al.*, 1998; Landauer *et al.*, 2000, 2003; Ramadan *et al.*, 2002; Stelzer *et al.*, 1994; Reiter and Tan, 2002; Shimoi *et al.*, 1994; Theresamma *et al.*, 1996; Yoshioka, 1997). Evidently, the use of plants and their bioactive constituents with antioxidant activity is highly relevant in mitigation of radiation-induced oxidative damage.

Haemopoietic stimulation

Exposure of mammals to ionizing radiation leads to the development of a complex dose dependent cascade of changes including injury to the lymphoid and haemopoietic system, which can result in septicaemia and death (Prasad, 1999). Agents capable of enhancing survival in the radiation dose inducing the haemopoietic syndrome have typically been associated with accelerated haemopoietic regeneration. An accelerated ability to regenerate new haemopoietic elements, especi-

ally those that are important in controlling microbial infections, such as granulocytes, allows the host to resist opportunistic infections better and, hence, enhances survival. *Acanthopanax senticosus*, *Ginkgo biloba*, *Hippophae rhamnoides*, *Panax ginseng*, *Podophyllum hexandrum*, *Tinospora cordifolia*, *Boerhaavia diffusa*, *Spirullina* provide total-body radiation protection by stimulating haemopoiesis (Goel *et al.*, 2002a,b,c; Kapoor and Mehta, 1998; Miyanomae and Frindel, 1988; Song *et al.*, 2003; Takeda *et al.*, 1982; Thali *et al.*, 1998; Yonezawa *et al.*, 1981). Both isolated compounds (e.g. ginsan, a purified polysaccharide isolated from *Panax ginseng* (Song *et al.*, 2003; Kim *et al.*, 1998b) and glycyrrhizic acid (Lin *et al.*, 1996) and complex herbal preparations used in Ayurveda [e.g. Liv 52 (Ganapathi and Jagetia, 1995; Saini *et al.*, 1985), Triphala (Jagetia *et al.*, 2002), Abana (Jagetia *et al.*, 2003c), Mentat (Jagetia and Baliga, 2003) Chayawanprash (Agarwal *et al.*, 2003)], Chinese medicine [Si-Jun-Zi-Tang (Hsu *et al.*, 1996a), Si-Wu-Tang (Hsu *et al.*, 1996b), Kuei-Pi-Tang (Hsu *et al.*, 1991; Jeng-Sheng-Yang-Yung-Tang (Hsu *et al.*, 1992), Lifukang (Kim *et al.*, 1998a)] and Japanese medicine [Juzen-Taiho-Toh (Ohnishi *et al.*, 1990)] have been evaluated for their radioprotective effects particularly for attenuating damage to the haemopoietic system. In coming years, it can be expected that plant products that stimulate the haemopoietic system will find a use in mitigating radiation induced-injury and enhancing radiorecovery.

Immunostimulant activity

Numerous plants have been reported to exert radioprotective effect via immunostimulatory activity in *in vitro* and *in vivo* models (Agarwal and Singh, 1999; Song *et al.*, 2003). Plants possessing immunostimulatory activity, e.g. *Podophyllum hexandrum*, *Hippophae rhamnoides*, *Viscum album*, *Ocimum sanctum*, *Tinospora cordifolia*, have been reported to provide protection by increasing spleen colony forming units (Ganasoundari *et al.*, 1997b, 1998; Goel *et al.*, 2002b; Narimanov *et al.*, 1992). Several triterpenoids such as glycyrrhizic acids, ursolic acid and oleanolic acid possess immunopotentiating activity (Raphael and Kuttan, 2003). The radioprotective effects of glycyrrhizin and glycyrrhizic acid on cellular immunocompetence has been reported by Lin *et al.* (1996). *Panax ginseng*, a known radioprotective plant, is known to augment natural killer (NK) cell activity (Kim *et al.*, 1990), production of interleukin-1 (IL-1) (Kim *et al.*, 1998b), interleukin-2 (IL-2) (Ma *et al.*, 1995), tumor necrosis factor-alpha (TNF- α) (Gao *et al.*, 1996), granulocyte macrophage-colony stimulating factor (GM-CSF; Kim *et al.*, 1998b), increase in population of CD3, CD4, CD8 cells (Mizuno *et al.*, 1994). An acidic polysaccharide (ginsan) isolated from *Panax ginseng* has been reported to activate multiple effector pathways of the immune system thereby rendering radioprotection (Kim *et al.*, 1998b).

Thatte *et al.* (1988) suggested that *Tinospora cordifolia* activates macrophages to release GM-CSF activity, while Kapil and Sharma (1997) have reported enhanced humoral and cell mediated immunity by syringin (TC-4) and cordiol (TC-7), the active constituents of *Tinospora cordifolia*. Other plants with

immunostimulant properties such as *Allium sativum*, *Ocimum sanctum* (which augments NK cells, stimulates T cells and IL-2 production), *Aloe vera* (which stimulates IL-1 and TNF- α (Agarwal and Singh, 1999) and *Eleutherococcus senticosus*, have been shown to provide radioprotection (Ben-Hur and Fulder, 1981).

Metal chelation activity

Iron is considered to be an important contributor in the generation of reactive oxygen species (Stevens *et al.*, 2000). Agents that chelate free iron can reduce ROS-mediated damage including radiation-induced damage (Morel *et al.*, 1993; Ramadan *et al.*, 2002). Polyphenolic compounds, including flavonoids, present in a number of medicinal plants have been reported to possess metal chelating properties (Bars *et al.*, 1994; Sgaragli *et al.*, 1993). Flavonoids (orientin and vicenin) from *Ocimum sanctum*, catechin present in *Camellia sinensis*, and quercetin present in *Podophyllum hexandrum* extract are known to play a pivotal role in metal (iron) chelation (Ganasoundari *et al.*, 1997a; Prem Kumar and Goel, 2000; Sestili *et al.*, 1998; Uma Devi *et al.*, 2000). Other phytoconstituents such as quercetin, catechin, silymarin and luteolin are also known to chelate metal ions, thereby rendering radioprotection (Duthie *et al.*, 2000; Gebhardt, 2002; Korina and Afanas'ev, 1997; Morel *et al.*, 1993; Ramadan *et al.*, 2002). Afanas'ev and coworkers (1989) have reported that rutin and quercetin can inhibit iron ion-dependent lipid peroxidation by chelating iron ions. Rutin has been shown to protect against tert-butyl hydroperoxide-induced oxidative damage to DNA by acting as a metal ion chelator (Aherne and O'Brien, 2000). Another compound, mimosine, can chelate iron and reversibly block cell cycle progression in mammalian cells (Kulp and Vulliet, 1996) via inhibition of cyclin-dependent kinase (cdk) activity (Kulp *et al.*, 1996). One of the mechanisms whereby *Podophyllum* provides radioprotection, apart from iron chelation (Prem Kumar and Goel, 2000), could include its ability to arrest the cell cycle in G1 phase.

Wound healing activity

In the aftermath of a nuclear accident, a number of patients will present combined injury wounds (i.e. wounds from burns and/or trauma with local or systemic radiation exposure before, during or after the injuries). The wound healing response in these patients is likely to be delayed due to their compromised immunological state and the synergistic characteristic of combined injury. The mortality of radiation injury is greatly increased by concomitant trauma and thermal injuries. Topical application of steroidal or non-steroidal antiinflammatory agents is the most common treatment for radiation injury of the skin, however, the results are usually not satisfactory and local toxicity has been observed (Chen *et al.*, 1999). Therefore, plant products with wound healing activity could be useful for treating combined injury wounds.

A number of plants containing triterpenes, alkaloids and other constituents are known to promote wound healing (Fleischner, 1985; Sarma *et al.*, 1990). These bioactive constituents influence different phases of

the healing process namely, inflammation that leads to haemostasis and clot formation, fibroplasias and neovascularization, formation of a granulation tissue, reepithelialization and finally the formation of new extracellular matrix and tissue remodelling (Chitra *et al.*, 1998).

Several plants, e.g. *Aloe vera*, *Angelica sinensis*, *Centella asiatica*, *Terminalia chebula*, *Hippophae rhamnoides*, *Ocimum sanctum* and *Curcuma longa*, with radioprotective properties, possess antibacterial, anti-fungal, antioxidant, antiinflammatory and cell proliferative activity (Ahmad *et al.*, 1998; Dutta *et al.*, 1998; Klein and Penneys, 1988; Sakar and Tamer, 1990; Udupa *et al.*, 1994) and are useful for the treatment of burns and wounds (Davis *et al.*, 1987; Ianev *et al.*, 1995; Suguna *et al.*, 2002; Shetty *et al.*, 1999; Rodrigues-Bigas *et al.*, 1988). Ascorbic acid has been reported to alter glutathione, superoxide dismutase and lipid peroxidation in mouse skin exposed to fractionated γ -radiation, thereby protecting the skin (Jagetia *et al.*, 2003a). *Aloe arborescens* and *Aloe vera* have been shown to be effective for treating radiation-induced burns (Chitra *et al.*, 1998; Sato, 1990, 1991) and acute radiation dermatitis (Chen *et al.*, 1999). Chamomile cream and almond ointment has also been reported to mitigate acute radiation skin reactions (Maiche, 1991). *Centella asiatica*, *Curcuma longa*, *Hypericum perforatum* and *Hippophae rhamnoides* extracts have been traditionally used to accelerate wound healing (Ianev *et al.*, 1995), and in recent years particularly in cases of chronic post-surgical and post-traumatic wounds (Jadhav and Bhutani, 2002). *Curcuma longa* has been used for healing of corneal wounds (Mehra *et al.*, 1984), while *C. asiatica* extracts have been used as a therapy in the treatment of second- and third-degree burns (WHO Monographs, 1999). *Centella asiatica* extracts (marketed under the trade name Madecassol[®]) and tetrandrine (a bisbenzylisoquinoline alkaloid isolated from *Stephania tetrantra* S. Moore) have been shown to reduce acute radiation reactions via their antiinflammatory activity (Chen *et al.*, 1999). Shukla and coworkers (1999) have shown that asiaticoside obtained from *Centella asiatica* helps in wound healing. The search for promising plant extracts that can stimulate wound healing, while decreasing pain, is therefore warranted.

PLANTS WITH RADIOPROTECTIVE EFFICACY: THEIR PHARMACOLOGICAL BASIS

Centella asiatica Linn

Centella asiatica (Apiaceae), is prescribed in the Indian system of medicine for the treatment of various diseases (Diwan *et al.*, 1991; Zafar and Naaz, 2002). It has traditionally been used to improve mental ability (Sarma *et al.*, 1995; Sanjay, 2000), healing of ulcers, wounds and skin lesions (Srivastava *et al.*, 1997; Tan *et al.*, 1997), and in an attempt to restore youth, memory and longevity (Kapoor, 1990). *Centella* has also been reported to have CNS depressant effects and improve maze learning capability in rats (Rao *et al.*, 1999). *Centella asiatica* contains several antioxidant molecules such as carotenoids, ascorbic acid, terpenoids and other biologically

active components such as asiaticoside, asiatic acid, brahmoside, brahminoside, brahmie acid, centoic acid, centellic acid, isobrahmic acid and thankunic acid (Srivastava *et al.*, 1997; Brinkhaus *et al.*, 2000).

There are several reports regarding the neuromodulatory effect of *Centella asiatica*, but only a few studies document the protective effects against radiation-induced behavioural changes and performance deficits (Goel *et al.*, 2000d; Shobi and Goel, 2001). Studies related to behavioural radioprotection assume importance since radiation is known to cause severe behavioural perturbations such as conditioned taste aversion (CTA), performance decrement and learning. Total-body exposure to even very low doses (0.1 cGy) of electron beam can induce retrograde amnesia (Wheeler and Hardy, 1985), while higher dose of radiation (10–100 Gy) induce emesis, nausea, taste aversion and diarrhoea, besides behavioural degradation in terms of coordination, performance, learning and memory (Brogo, 1984; Burghardt and Hunt, 1985; Franz, 1985). An aqueous extract of *Centella asiatica* (100 mg/kg body weight; i.p.; single dose; –1 h) when used to protect Sprague Dawley rats against the adverse effects of low-dose ionizing radiation (2 Gy) rendered significant protection against radiation-induced body weight loss and conditioned taste aversion (since there is no emesis in rodents, taste aversion is an equivalent manifestation), suggesting that *Centella asiatica* could be useful in preventing radiation-induced behavioural changes during clinical radiotherapy (Shobi and Goel, 2001).

Centella asiatica extract (100 mg/kg body wt.) administered orally has recently been shown to provide total-body protection in mice against sublethal (8 Gy) ⁶⁰Co gamma radiation (Sharma and Sharma, 2002). These workers have reported significantly less radiation-induced body weight loss in drug treated animals. Though the precise reason for radioprotection has not been elucidated, some of the reasons put forth include reduction of lipid peroxidation and membrane peroxidation, increase in haemoglobin percentage, and increased secretion of serotonin (a known radioprotective agent). The detoxification effect of *Centella asiatica* against combined toxicity of γ -radiation and cadmium chloride has been reported by Agarwal and coworkers (2001), while the effect on cognition and markers of oxidative stress in rats has been reported by Kumar and Gupta (2002).

***Ginkgo biloba* Linn.**

Ginkgo biloba (Cycadaceae), a plant indigenous to China, Japan and Korea, has been reported to stimulate endogenous antioxidants such as glutathione and attenuate oxidative stress (Rong *et al.*, 1996a,b). It has also been used in the modern system of medicine for the treatment of circulatory (Heinjen and Knipschild, 1992) and equilibrium disorders (Haguenauer *et al.*, 1986), asthma and senility. A *Ginkgo biloba* extract (Egb761), which is a mixture of flavonoids, heterosides and terpenes with antioxidant properties (De Feudis, 1991; Huguet *et al.*, 1994), has been shown to prevent mitochondrial aging by reducing oxidative damage (Sastre *et al.*, 1998). *Ginkgo biloba* extract is also useful in the treatment of cerebral disorders due to aging and hypoxia (Duche *et al.*, 1988).

G. biloba contains nearly 300 compounds including ascorbic acid, α -carotene, β -carotene, flavonoids (kaempferol, quercetin, myricetin, ginkgetin, isoginkgetin etc.), coumarins, catechins, ginkgolides, bilobalide, rhamnetin, γ -tocopherol to name a few (De Feudis, 1991), many of which individually in isolated form render radioprotective effects.

An ethanol (30%) extract of the dried leaf at a concentration of 100 μ g/mL was effective when tested on a culture exposed to clastogenic factors from plasma of human subjects exposed to irradiation (Emerit *et al.*, 1995a). Treatment of recovery workers from the Chernobyl accident site was found to be effective when an oral dose of 40 mg/day was given 3 times daily for 2 months (Emerit *et al.*, 1995b).

An intravenous infusion of an ethanol extract of *G. biloba* leaves, at a dose of 100 mg/person was found to be effective on patients with vasogenic oedema observed after irradiation of the brain (Hannequin *et al.*, 1986).

G. biloba extract provided protection to brain neurons against oxidative stress (Oyama *et al.*, 1996; Smith *et al.*, 1996). *G. biloba* leaf extract (30%) at a concentration of 100 μ g/mL assayed in rat cerebellar neuronal cell culture, was active on neurons against hydroxyl radical-induced apoptosis (Ni *et al.*, 1996). Agents that inhibit free radical-mediated apoptosis are known to provide radioprotection, which could help to explain the radioprotective effect of *G. biloba*.

***Hippophae rhamnoides* Linn.**

Hippophae rhamnoides (Sea Buckthorn; Family: Elaeagnaceae) has been used in traditional Tibetan and Indian systems of medicine for centuries. In Tibet, the plant was used as early as 900 AD. The plant has been extensively exploited for treatment of sluggish digestion, stomach malfunctioning (Nikitin *et al.*, 1989; Xiao *et al.*, 1992), burn and wound healing (Ianev *et al.*, 1995; Nikulin *et al.*, 1992), circulatory disorders, ischaemic heart disease (Liu *et al.*, 1998; Zhang, 1987), hepatic injury (Cheng *et al.*, 1994) and neoplasia (Nikitin *et al.*, 1989). Sea Buckthorn oil is used as a treatment of oral mucositis, vaginal mucositis, cervical erosion, burns, scalds, duodenal ulcers, gastric cancers and skin ulcers (Li, 1999). The plant has been well documented to have antioxidative, antiinflammatory, antimicrobial, pain-relieving, immunostimulative and regenerative properties.

The berries of *H. rhamnoides* contain polyphenolic compounds (namely, isorhamnetin, rhamnetin, quercetin, kaempferol), carotenes (α , β , γ), vitamins (A, E, C, K), riboflavin, folic acid, tannins, glycerides of palmitic, stearic and oleic acids, and some essential amino acids (Chan *et al.*, 1990), which play a major role in contributing towards bioactivities such as free radical scavenging, chromatin compaction and hypoxia induction (Goel *et al.*, 2000c, 2001c, 2003b; Prem Kumar *et al.*, 2002), all of which have been demonstrated to play a vital role in radioprotection.

Gileva and Lukin (1984) reported the radioprotective efficacy of compounds from *Hippophae*, while Mizina and Sitnikova (1999) showed that oral administration of a *H. rhamnoides* fruit juice concentrate to rats before or after irradiation (x-ray; 1 Gy) was accompanied by an increase in life span, restoration of the

11-oxycorticosteroid level in the blood and weight of isolated adrenals, and also normalization of their basal activity and response to ACTH (corticotropin) under *in vitro* conditions.

An aqueous-alcohol (50% ethanol) extract of berries of *H. rhamnoides* (30 mg/kg body weight) when administered to Strain 'A' mice 30 min before whole-body ^{60}Co γ -irradiation (10 Gy) increased life span and rendered 82% survival (at 30 days) compared with 100% mortality (within 15 days) in irradiated controls (Goel *et al.*, 2002a; Sharma *et al.*, 2004). The endogenous colony forming unit (CFU) counts in mouse spleen on post-irradiation day 10, and various other haematological parameters, clearly demonstrated the radioprotective effects of the extract. In addition, the herbal extract of *H. rhamnoides* also inhibited the Fenton reaction and radiation-induced generation of hydroxyl radicals *in vitro*, superoxide anion-mediated nitroblue tetrazolium reduction and ferrous sulphate-mediated lipid peroxidation in mouse liver.

The hydroalcohol extract of *H. rhamnoides* has also been shown to protect mice against gamma radiation-induced genotoxicity (Agarwal and Goel, 2002). Pre-irradiation administration of the herbal extract to mice reduced the radiation-induced micronuclei frequency in a dose-dependent manner, suggesting its radioprotective efficacy. Treatment with *H. rhamnoides* extract enhanced DNA synthesis (S-phase) in unirradiated controls and also countered the radiation-mediated depression of S-phase to facilitate replenishment of cells lost as a result of radiation injury.

Irradiation causes damage to mitochondria, leading to a bio-energetic catastrophe in the cell (Kroemer and Reed, 2000). Since energy is essentially required for repair and restoration of normal cellular (metabolic) functions, protecting mitochondria against the deleterious effects of ionizing radiation can help to provide radiation protection and radiorecovery.

A hydro-alcohol extract of fruits (berries) of *H. rhamnoides* has been shown to protect the functional integrity of mouse liver mitochondria against lethal γ -radiation (10 Gy) under *in vivo* conditions (Gupta *et al.*, 2003c). It was found that pre-irradiation treatment of mice with the extract (30 mg/kg body weight; i.p.; single dose; -30 min) significantly inhibited the radiation-induced increase of superoxide anions, oxidized glutathione (GSSG), thiobarbituric acid reactive substances (TBARS), mitochondrial complex I and complex I/III activity and protein oxidation. The mitochondrial complex II/III activity and mitochondrial membrane potential (MMP; which was reduced during irradiation) was significantly enhanced.

H. rhamnoides has been shown to provide protection to the gastrointestinal system against lethal whole-body γ -radiation (Goel *et al.*, 2003a). Administration of a hydroethanol (50:50 v/v) extract 30 min before irradiation increased the number of surviving crypts in the jejunum by a factor of 2.02 and villi cellularity by 2.5 fold in comparison with the irradiated control. The extract also reduced the incidence of apoptotic bodies in the crypts in a time-dependent manner and increased cellularity in the crypts and villi (84 h post-irradiation) compared with the control. Caspase-3 activity was also found to be significantly lower in the mice administered *Hippophae* extract before irradiation compared with irradiated control, thereby implying that by

reducing the caspase activity, *Hippophae* extract might play a pivotal role in the protection of crypts from apoptosis. The study showed that reduction in the radiation-induced loss of cellularity of crypts and villi, and also the decrease in frequency of apoptosis by *H. rhamnoides* extract could have contributed to the overall radioprotective effect.

The mode of action of the alcoholic extract (50%) of whole berries of *H. rhamnoides* at molecular and cellular level has been partially elucidated (Goel *et al.*, 2001c, 2003b; Prem Kumar *et al.*, 2002). Single cell gel electrophoresis (comet assay) revealed that the extract inhibits radiation-induced DNA strand breaks in mouse thymocytes in a dose-dependent manner (Goel *et al.*, 2001c; Prem Kumar *et al.*, 2002). *Hippophae* extract under *ex vivo* conditions induced a strong compaction of chromatin making the nuclei resistant to a radiation dose as high as 1000 Gy (Goel *et al.*, 2000c, 2001c). The compaction of chromatin was not reversed even by relaxation buffer, indicating that the salt concentration, did not have any role to play in the herbal extract-induced chromatin compaction. The alkaline halo assay also corroborated the results of the comet assay. A thermal denaturation assay was used to show that the extract interacts with DNA.

Further investigations on *H. rhamnoides*, done in cultured thymocytes, revealed an ability to bring about a concentration-dependent compaction of both a reversible (<100 $\mu\text{g}/\text{mL}$) and irreversible (>100 $\mu\text{g}/\text{mL}$) nature, which was further correlated to the magnitude of DNA-protein crosslinks formed (Goel *et al.*, 2003b). *H. rhamnoides* extracts can maintain chromatin organization and block the cell cycle at G2-M phase by interfering with topoisomerase I activity (Goel *et al.*, 2003b), thereby contributing towards the radioprotective efficacy of the extract. Thus, *H. rhamnoides* appears to be a promising herb in the prophylactic treatment of radiation-induced damage but further research is necessary to identify appropriate dosing regimes and to characterize the active constituents.

Mentha piperita Linn.

Mentha piperita (Lamiaceae) is an aromatic plant with a diverse array of medicinal properties. Its stimulative and carminative properties have been used for allaying nausea, flatulence and vomiting for over a thousand years. The antioxidant and antiperoxidant properties have been attributed to the presence of caffeic acid, eugenol, rosmarinic acid and α -tocopherol (Rastogi and Mehrotra, 1991). *Mentha* extract and mint oil have been shown to possess antibacterial and antifungal activities. Antimutagenic properties (ability to enhance error-free repair of DNA damage) of *Mentha* has also been reported (Vokovic-Gacis and Simic, 1993). The chemopreventive effect against shamma (a complex mixture of powdered tobacco used in Saudi Arabia which has been linked to oral cancer) induced carcinogenesis may be largely due to the antimutagenic properties of *Mentha* (Samman *et al.*, 1998).

Leaves of *Mentha piperita* contain 7-O-rutinosides of eriodictyol and luteolin, while *Mentha* oil contains menthol, menthone, neomenthone, cineole, menthyl acetate, isomenthol, limonene and pinene (Rastogi and Mehrotra, 1995).

Pre-treatment with a *Mentha piperita* extract protected haematological constituents and serum phosphatases activity in Swiss albino mice against γ -radiation (Samarth *et al.*, 2001, 2002b). *M. piperita* administration elevated the counts of endogenous spleen colonies and spleen weight significantly (Samarth *et al.*, 2001). The leaf extract of *M. piperita* was shown to provide protection against radiation-induced alterations (reduction in villus height, mucosal, total cells and mitotic figures/crypt section) in the intestinal mucosa of mice (Samarth *et al.*, 2002a). *M. piperita* pre-treatment also protected against the radiation-induced increase in goblet cells/villus section and dead cells/crypt section in the jejunum of mice.

Oral administration of *M. piperita* (1 g/kg body weight/day) prior to sublethal radiation exposure (8 Gy) was found to be effective against the chromosomal damage in bone marrow of Swiss albino mice (Samarth and Kumar, 2003). Irradiated animals exhibited chromosomal aberrations in the form of chromatid and chromosome breaks, centric rings, dicentric exchanges and acentric fragments, while animals pre-treated with *M. piperita* extract showed a significantly lesser number of aberrant cells. It also significantly increased GSH levels and decreased the lipid peroxidation level in irradiated mice. The radioprotective effect of *M. piperita* was also demonstrated by determining dose modification factor, which was 1.78 (DMF refers to the ratio of the radiation dose required to elicit the same effect in the presence and in the absence of the radioprotectant) (Samarth and Kumar, 2003).

Jagetia and Baliga (2002a) have reported the protective effect (DMF: 1.27) of a chloroform extract of *Mentha arvensis*, a related species, in mice exposed to 10 Gy radiation.

Ocimum sanctum Linn.

Ocimum sanctum (Holy Basil; Tulasi; Family: Lamiaceae) is an Indian medicinal herb widely distributed in the semi-tropical and tropical regions of the country. The medicinal value of the plant has been well documented in various ancient Indian texts, and virtually every part of this plant is used in the traditional, Ayurvedic and Siddha systems of medicine for treating a plethora of human ailments. In particular, *O. sanctum* is used for treating infections, skin diseases, hepatic disorders, common cold and cough, malarial fever and as an antidote for bites by snakes and poisonous insects (Satyavati *et al.*, 1987). During the past few decades, the plant has been investigated extensively and has been shown to possess a range of biological activities, many of which are relevant to radiation protection, e.g. antibacterial (Phadke and Kulkarni, 1989), antifungal (Rai, 1996), hypoglycaemic (Chattopadhyay, 1999), antiinflammatory (Godhwani *et al.*, 1987; Singh and Agarwal, 1991; Singh *et al.*, 1996), antiviral (Kumar *et al.*, 1997), antioxidant (Uma Devi, 2001), antiulcer (Dharmani *et al.*, 2002), anticarcinogenic (Uma Devi, 2001), hepatoprotective (Chattopadhyay *et al.*, 1992), analgesic (Godhwani *et al.*, 1987), immunostimulatory (Godhwani *et al.*, 1988) and wound healing activities (Shetty *et al.*, 1999). The antistress effect of *O. sanctum* has also been reported in a rodent model (Bhargava and Singh, 1981).

The leaves and stem of *O. sanctum* contain a number of constituents including apigenin, carvacrol, cirsilineol, cirsimaritin, eugenol, isothymonin, luteolin, methyl eugenol, orientin, rosmarinic acid, sesquiterpene hydrocarbon caryophyllene, ursolic acid, vicenin, etc. (Nair *et al.*, 1982). The presence of vicenin-2, rosmarinic acid, galuteolin, cirsilineol gallic acid, gallic acid methylester, gallic acid ethylester, protocatechic acid, vanillic acid, vanillin, caffeic acid has been reported in the ethanol extract of *O. sanctum* (Norr and Wagner, 1982).

The radioprotective property of *O. sanctum* was first reported by Uma Devi and Ganasoundari (1995). Thirty-day lethality studies in Swiss albino mice were carried out following treatment with single graded doses of aqueous and ethanol extracts from dried leaves of *O. sanctum* (Krishna Tulasi; the dark-leaved variety of *O. sanctum*) and it was found that the aqueous extract was more effective in increasing survival, compared with the ethanol extract. The optimal dose for protection was reported to be 50 mg/kg b.w (intraperitoneal administration), while the acute LD₅₀ was 6 g/kg b.w. Administration of a fractionated dose of the herbal extract via the i.p. route (10 mg/kg/day for 5 consecutive days to mice prior to irradiation) was more effective compared with a single dose (50 mg/kg b.w.). The optimum dose (fractionated dose of 10 mg/kg/day of aqueous extract of *O. sanctum* for 5 consecutive days via the intraperitoneal route) administered to mice prior to irradiation gave a dose modification factor of 1.28. It was also found that the i.p. route of drug administration was more effective than the oral route.

Ganasoundari *et al.* (1997a) studied the effect of *O. sanctum* on the survival of mice after whole-body lethal irradiation and compared it with WR-2721, a standard radioprotector. Their results indicated that *O. sanctum* promotes recovery and regeneration of haemopoietic progenitor cells in mice bone marrow. An intraperitoneal (i.p.) injection of an optimum dose (10 mg/kg daily for 5 days) of leaf extract of *O. sanctum* to mice before delivering sub-lethal (2 Gy) total-body γ -radiation produced a significantly higher bone marrow stem cell survival than a pre-treatment with 300 mg/kg (approx. 40% of its LD₅₀) of WR-2721 (amifostine), suggesting that in terms of the protective dose and toxicity, the herbal extract is a better radioprotector than the synthetic drug. Analysis of chromosomal aberrations in mouse bone marrow exposed to γ -radiation showed that the *O. sanctum* extract could significantly reduce the percentage of aberrant metaphases and other chromosomal aberrations, including dicentric and rings, induced by sub-lethal whole-body radiation doses (3–5 Gy). The decline in the percent aberrant metaphases by *O. sanctum* pre-treatment was comparable to that provided by 400 mg/kg of WR-2721. *O. sanctum* pre-treatment did not manifest any toxic side effects, while WR-2721 (300–400 mg/kg b.w.) administration prior to irradiation resulted in an increase in the percent aberrant cells at 14 days post-irradiation. Administration of a combination of *O. sanctum* and WR-2721 to mice prior to γ -irradiation considerably enhanced the chromosome protection by nearly two-fold, compared with individual administration, and also eliminated the delayed chromosome toxicity associated with the treatment of WR-2721. In addition, *O. sanctum* extract also protected mouse liver against radiation-induced lipid peroxidation.

The responses of single or sequential doses of *O. sanctum* were not substantially different. The anti-lipid peroxidative effect was attributed to increased levels of cellular antioxidants such as reduced glutathione (GSH), GSH-transferase, GSH-peroxidase and reductase as well as superoxide dismutase (SOD).

The work of Ganasoundari *et al.* (1997b) showed that the aqueous extract of leaves of *O. sanctum* significantly inhibited the OH radical-induced deoxyribose degeneration. A combination of WR-2721 and *O. sanctum* extract produced a significantly higher inhibition of the OH radical activity compared with either agent individually (Ganasoundari *et al.*, 1998).

Orientin (8-C- β -D-glucopyranosyl-luteolin) and vicenin-1 (6-C- β -D-xylopyranosyl-8-C- β -D-glucopyranosyl apigenin), water-soluble compounds from *O. sanctum* did not exhibit any systemic toxicity in mice even at a dose of 100 mg/kg b.w. Both compounds significantly increased mouse survival when administered 30 min prior to lethal whole-body γ -irradiation. The optimum dose for protection was found to be 50 μ g/kg b.w. via the i.p. route. Other routes of administration, e.g. oral and intravenous route were also found to be effective, but to a lesser extent. Vicenin provided a slightly higher protection (DMF: 1.37), compared with orientin (DMF: 1.30) in murine model system (Uma Devi *et al.*, 1999) and also reduced the chromosomal aberrations better than amifostine in the bone marrow of mice exposed to 2 Gy γ -irradiation (Uma Devi *et al.*, 1998). Both the flavonoids, orientin and vicenin, were found to be equally effective in rendering protection against γ -radiation-induced lipid peroxidation in mouse liver. These compounds also significantly inhibited the Fenton reaction-induced OH radical activity under *in vitro* conditions (Uma Devi *et al.*, 2000) and protected human lymphocyte chromosomes (Vrinda and Uma Devi, 2001). Though the role of orientin and vicenin in radiation protection has been established, it is plausible that other constituents present in *O. sanctum*, may also be involved in the observed radioprotective effects, since radioprotection of the whole organism usually requires multifarious activities, i.e. simultaneous protection of various target tissues and organs.

***Panax ginseng* CA Meyer**

Ginseng is one of the most extensively used medicinal plants, particularly in traditional oriental medicine for the treatment of various diseases. The plant has been thoroughly exploited for its adaptogenic, antistress, antitumor, antioxidant, antiaging, antifungal and rejuvenating properties (Keum *et al.*, 2000; Kim *et al.*, 1993, 1998b, 2002; Lam and Ng, 2002). Ginseng root and its major bioactive constituents have complex and multiple pharmacological actions (Attele *et al.*, 2002).

The stem and leaves of ginseng contain ginsenosides-Rb-1, Rb-2, Rc, Rd, Re, Rf, Rg-1, Rg-2, Rg-3, kaempferol, triofolin, salicylic, p-coumaric, gentisic and caffeic acids, while roots have been reported to contain panaxan A, B, C, D, E, F, G, H, acidic polysaccharides, tripalmatin, panxynol, panaxytriol, linolein, palmitic acid, β -sitosterol (Dewick, 2002).

The radioprotective efficacy of ginseng has been reported by several workers (Kim *et al.*, 1993, 2001; Kumar *et al.*, 2003; Pande *et al.*, 1998a; Takeda *et al.*,

1982). Yonezawa and coworkers (1985) showed that recovery of thrombocyte and erythrocyte counts in blood after irradiation was the major factor responsible for radiation protection. The whole extract of ginseng and the relative protective effects of various fractions (carbohydrate, protein and saponins) have been evaluated. The results showed that the water-soluble whole extract of ginseng provided the best protection against ^{60}Co gamma radiation in C3H mice, while isolated protein and carbohydrate fractions were less effective, the saponin fraction was ineffective (Zhang *et al.*, 1987). Similar results were obtained by Kim and coworkers (2001), who found that whole ginseng extract and its fractions increased endogenous spleen colony formation in irradiated mice and also reduced apoptosis in jejunal crypt cells. Ginsan, a purified polysaccharide (molecular weight 2000 kD) isolated from the ethanol-soluble fraction of *P. ginseng* aqueous extract, possesses immunological activities such as induction of several cytokines (IL-1, IL-2, IL-8, interferon- γ and GM-CSF) (Sonoda 1998), mitogenic activity, stimulation of natural killer (NK) cells, lymphokine-activated killer (LAK) cells and macrophages (Song *et al.*, 2002; Ma *et al.*, 1995).

Song and coworkers (2003) have studied the protective effect of ginsan on irradiated mice, due to its ability to induce proliferation of murine bone marrow as well as the release of haematopoietic growth factors. Ginsan was found significantly to induce production of bone marrow, spleen and granulocyte-macrophage colony-forming cells (GM-CFU), and also circulating neutrophils, lymphocytes and platelets in irradiated mice. Ginsan also induced a battery of cytokines in spleen cells including Th1-type (IL-2, IL-12, IFN- γ), Th2-type (IL-4, IL-5, IL-10), and proinflammatory (IL-1, IL-6, TNF- α) cytokines under *in vitro* conditions. The expression patterns of cytokines after ginsan treatment *in vitro* and *in vivo* were different. Ginsan induced the endogenous production of cytokines such as IL-1, IL-6, IFN- γ and IL-12, which are required for haematopoietic recovery, and also enhanced TH-1 function while interfering with the TR-2 response in irradiated mice. Pre-treatment of BALB/c mice, 24 h prior to irradiation, with ginsan (100 mg/kg body weight; i.p.; single dose) protected mice from the lethal effects of ionizing radiation (8 Gy) more effectively than when it was administered immediately (within 15 min) or at various times after irradiation (+3 h and +24 h) (Lee *et al.*, 1997).

The radioprotective effect of *P. ginseng* root extract on testicular enzymes (acid and alkaline phosphatases and lipid peroxidation) has been recently reported by Kumar *et al.* (2003). These workers administered 10 mg/kg b.w. of *P. ginseng* extract continuously for 4 days to Swiss albino mice, and on the day 4 they were given 8 Gy γ -radiation 30 min after extract administration and a protective effect was reported.

***Podophyllum hexandrum* Royale (Syn. *P. emodi* Wall.)**

Podophyllum hexandrum (Himalayan Mayapple; Berberidaceae) is a perennial herb, thriving in the Himalayan region at altitudes ranging between 2500 to 4000 m. The root and rhizome of the plant have been extensively used in India, for over 2000 years to treat a

number of ailments such as cold, constipation, biliary fever, septic wounds, erysipelas, insect bites, mental disorders and rheumatism (Singh and Shah, 1994). *P. hexandrum* has been used to provide symptomatic relief in some allergic and inflammatory conditions of skin. This herb, and its constituents, have also been used in the treatment of cancer (Singh and Shah, 1994), venereal warts (Beutner and Von Krogh, 1990), monocytoid leukaemia, Hodgkin's disease, non-Hodgkin's lymphoma and cancer of the brain, bladder and lung (Blasko and Cordell, 1988; Singh and Shah, 1994). Antiviral and anti-HIV properties of this plant have also been reported (Gowdey *et al.*, 1995).

P. hexandrum contains a number of bioactive constituents including lignans (podophyllotoxin, podophyllotoxone, podophyllin, peltatins α and β) and flavonoids including quercetin, kaempferol, astragaloside and kaempferol-3-glucoside (Singh and Shah, 1994; Wong *et al.*, 2000).

The radioprotective effect of *P. hexandrum* has been reported in Swiss albino mice by Goel and co-workers (Goel *et al.*, 2000a,b, 2001a). Pre-irradiation administration of *P. hexandrum* rhizome extract protected mice in a dose-dependent manner (optimal dose being 200 mg/kg body weight rendering 80% survival for 30 days) against whole-body lethal γ -irradiation (10 Gy). The extract was also found to provide cytogenetic protection, as observed by a decrease in radiation (2 Gy)-induced micronuclei frequency upon pre-irradiation treatment. Radioprotective properties of *P. hexandrum* were found to be comparable and in some cases even better than synthetic radioprotectors such as diltiazem and WR-2721. Subsequent 30-day survival studies by our group revealed over 80% protection against 10 Gy and a DMF of 1.33, was observed (unpublished data).

P. hexandrum chelated Fe^{2+} more efficiently than Fe^{3+} (in a dose dependent manner; measured using chelating agents 2,2-bipyridyl and potassium thiocyanate, respectively), and also modulated the $\text{Fe}^{2+}/\text{Fe}^{3+}$ ratio (Prem Kumar and Goel, 2000).

Radiation doses in the range 7–12 Gy cause a gastrointestinal syndrome (Coleman *et al.*, 2003), which is characterized by denuded, eroded and shrunken villi, stromal cores, elongated and dilated crypts, depopulated crypts, decrease in crypt cell numbers and a decrease in the mitotic index and apoptotic bodies in the crypts. The effect of an aqueous extract of *P. hexandrum* rhizome in ameliorating radiation (10 Gy)-induced gastrointestinal damage, resulting from destruction of clonogenic crypt cells and eventual depopulation and denudation of villi, has been reported (Salin *et al.*, 2001). Using an *in vivo* micro colony survival assay these workers demonstrated that pre-irradiation administration of *P. hexandrum* extract (200 mg/kg b.w. per mouse by i.p. route 2 h prior to gamma irradiation) increased (3-fold) the number of surviving crypts in the jejunum and villi cellularity (2.7-fold). The herbal extract induced cell division arrest and also reduced radiation-induced apoptosis in crypt cells, thereby rendering protection against lethal γ -radiation.

P. hexandrum extract has been shown to render radioprotection to the developing nervous system (Goel *et al.*, 2002c). Since cells of the embryonic nervous system are radiosensitive and actively differentiating and maturing during the fetal and early postnatal periods (Reyners *et al.*, 1992), the effect of sublethal (2 Gy)

γ -radiation was studied in rats *in utero* on day 17 of gestation to monitor radiation-induced retardation of neurophysiological development in new-borns. Treatment of rats with *P. hexandrum* extract (200 mg/kg b.w.; single dose; i.p. route) 120 min before irradiation mitigated radiation-induced physiological alterations (Goel *et al.*, 2002c).

P. hexandrum has been shown to modulate antioxidant enzyme levels. The effect of an aqueous extract of *P. hexandrum* in modulating the antioxidant defence system in murine liver, jejunum and ileum with special reference to enzymes such as glutathione-S-transferase (GST), which neutralizes electrophiles by conjugation with glutathione thereby making them readily excretable from the body, superoxide dismutase (SOD), which scavenges superoxide anion to form H_2O_2 and catalase, which inactivates H_2O_2 to form water has been investigated (Mittal *et al.*, 2002). Pre-irradiation treatment with *P. hexandrum* extract significantly enhanced liver GST and SOD at 12 h post-irradiation and intestinal SOD at 84 h post-irradiation intervals, while no significant change was manifested in hepatic catalase activity.

P. hexandrum has also been demonstrated to provide protection to the male reproductive system. The suitability of the aqueous extract of rhizome of *P. hexandrum* in rendering radioprotection, was evaluated in male germinal tissue in mice (Samanta and Goel, 2002). Administration of *P. hexandrum* via the i.p. route 2 h prior to irradiation produced a significant increase in the testis weight, repopulating the seminiferous tubules and increasing resting primary spermatocytes, stem cell survival index, sperm counts and reduction in abnormalities of sperm morphology. This suggests that *Podophyllum* extract, if put to clinical application, will not be harmful to the testicular system.

P. hexandrum has been shown to protect mammalian mitochondrial systems against radiation damage (Gupta *et al.*, 2002, 2003a,b). Radioprotection by an aqueous extract of *P. hexandrum* has been investigated in HepG2 cells (human hepatoma) by evaluating colony forming efficiency, redox status of mitochondria, reactive oxygen species (ROS), nitric oxide (NO) generation, peroxidation of lipids and intracellular glutathione (Gupta *et al.*, 2003b). Lower concentrations of the extract (0.1 and 1.0 $\mu\text{g}/\text{mL}$) rendered maximum radioprotection when administered 1 or 2 h before irradiation, while higher concentrations (5 and 10 $\mu\text{g}/\text{mL}$) were more effective at greater time intervals (4 or 8 h). The time of drug administration is very important for achieving radioprotection, as shown by several workers (Coleman *et al.*, 2003; Weiss and Landauer, 2003). Pre-treatment with *P. hexandrum* extract significantly inhibited radiation-induced MTT reduction in a concentration and time-dependent manner by decreasing gamma radiation-induced leakage of electrons from the mitochondrial electron transport chain. Pre-irradiation administration of the extract significantly reduced both ROS and NO generation and enhanced glutathione levels, and inhibited lipid peroxidation, thereby rendering radiation protection.

Since mitochondria play a vital role in cell metabolism, they are crucial in post-irradiation repair and recovery processes, being the site of ATP production (Haraguchi *et al.*, 2000). The role of *P. hexandrum* in affording radiation protection at the whole-body level, and the crucial role played by mitochondria therein,

was investigated at *in vivo* level (Gupta *et al.*, 2004). Exposure of γ -radiation (10 Gy) to mice significantly increased the generation of mitochondrial superoxide anions O_2^- , while pre-irradiation treatment of mice with *P. hexandrum* extract significantly reduced the formation of radiation-induced superoxide anions. Increased levels of mitochondrial glutathione were observed upon irradiation, whereas in the case of pre-irradiation treatment, levels were found to be significantly higher at all time periods studied, indicative of stimulation of endogenous antioxidants. The radiation-mediated increase in complex I (NADH: UQ oxidoreductase) activity was significantly ameliorated upon pre-irradiation administration with the herbal extract. The radiation-mediated alteration in the flow and leakage of electrons from complex I/III (NADH: cytochrome c oxidoreductase) and complex II/III (succinate: cytochrome c oxidoreductase) were also properly maintained by pre-irradiation administration of the extract. Pre-irradiation administration of *P. hexandrum* extract to mice inhibited the radiation mediated decrease in mitochondrial membrane potential (MMP). Administration of the extract to mice prior to irradiation also lowered radiation-induced oxidative damage to lipids and proteins, increased antioxidant levels, and simultaneously inhibited radiation-induced alterations in mitochondrial membrane potential (MMP), lipid peroxidation, protein oxidation and electron transport chain (ETC) activity, thereby affording protection to the hepatic mitochondria against lethal γ -radiation. *P. hexandrum* has been shown to act in a multifaceted manner and provide protection to haematopoietic, gastrointestinal, reproductive and central nervous system (CNS) (Goel *et al.*, 2000a,b, 2001a; Samanta and Goel, 2002; Sajikumar and Goel, 2003). It is apparent that *P. hexandrum* is a promising radioprotector and may be useful in providing protection during both planned and unplanned radiation exposures. Fractionation of the herbal extract(s), isolation of the individual active components, and evaluation of their bioactivities has not yet been reported. Identification of the bioconstituents, determination of appropriate dosing regimens, and potential effects of long-term usage also need to be considered before undertaking clinical trials, especially in view of the cytotoxic nature of podophyllotoxin.

Tinospora cordifolia (Willd.) Miers

Tinospora cordifolia (Menispermaceae) is considered to be a Rasayana, Medha and anti-aging drug in Ayurveda. It is considered as a tonic, vitalizer, anti-diabetic, hepatoprotective, antipyretic, antistress, anti-ulcer, anticancer, antioxidant and immunomodulatory agent, and has been widely used for the treatment of various ailments including jaundice, skin diseases, anaemia, emaciation and infections (Stanley *et al.*, 1999a,b). The aqueous stem extract of *Tinospora cordifolia* has been reported to prevent abdominal infections and sepsis, to improve Kupffer cell function and poly-morphonuclear cell (PMN)-mediated phagocytosis in rats with chronic liver damage and patients of surgical jaundice (Subramaniam *et al.*, 2003). The antioxidant properties of *Tinospora cordifolia* may be substantially responsible for these medicinal effects.

Tinospora cordifolia contains syringin and cordiol, which reduce immunohaemolysis and significantly increase IgG antibodies (Kapil and Sharma, 1997). Cordioside, cordiofolioside, cordiofolioside B and clerodane furanoditerpene glucosides have been reported to have immunostimulatory properties (Wazir *et al.*, 1995).

The radioprotective property of *Tinospora* was first reported by Goel and co-workers (2001b). The aqueous stem extract of *Tinospora cordifolia* exhibited free radical scavenging and metal chelation properties and thereby provided protection against ionizing radiation (Goel *et al.*, 2002b). The extract quenched radiation-mediated 2-deoxyribose degradation in a dose-dependent fashion (IC_{50} value: 700 μ g/mL), and inhibited the formation of Fe^{2+} -bipyridyl complex and formation of the comet tail in irradiated thymocytes by chelating Fe^{2+} ions. It also inhibited ferrous sulphate mediated lipid peroxidation in liver homogenate. The authors opined that the direct antioxidant mechanisms (chelation of metal ions) of action of *Tinospora cordifolia* aqueous extract contributed in a comprehensive and competitive manner to ameliorate the radiation-induced oxidative stress. The antioxidant properties (both direct and indirect mechanisms), along with other properties such as immunomodulation and cell proliferative capacity, could also be responsible for radioprotective manifestation, as was also revealed by survival against radiation (10 Gy)-induced mortality in Strain 'A' mice (Goel *et al.*, 2004).

The radioprotective activity of a pure arabinogalactan polysaccharide, isolated from the aqueous extract of the dried stem of *Tinospora cordifolia*, has been established using *Saccharomyces cerevisiae* X 2180 strain as the *in vivo* test model (Subramanian *et al.*, 2003). The radioprotective activity was attributed entirely to the superior radical scavenging capacity of the preparation, since it did not enhance expression of the protective endogenous antioxidant enzymes, namely, catalase and superoxide dismutase in the yeast cells. The preparation, even at a low concentration of 6.9 mg/mL, provided nearly 39% protection to plasmid (pBR322) DNA against γ -ray induced single strand breaks as was revealed by an increase in the supercoiled form of the plasmid. Exposure of pBR322 to γ -radiation led to the complete conversion of the supercoiled form to the open circular form.

Oral administration of an aqueous extract of *Tinospora cordifolia* (5 mg/kg body weight per day) to Swiss albino mice 1 h prior to sublethal whole-body radiation exposure (8 Gy) provided 33% survival (at 30 days). The same dose, when given for 15 consecutive days, produced 100% survival until day 9 and 50% of the animals survived until day 24. All the animals died within 30 days (Pahadiya and Sharma, 2003) suggesting that *Tinospora cordifolia* is partially effective against sublethal radiation dose.

CONCLUSIONS

It is apparent that many plants exhibit, or have the potential to show, a diverse array of biological activities that may be relevant to the mitigation of ionizing radiation-induced damage in mammalian systems.

However, so far, only a fraction of these plants have been investigated. There is an urgent need to develop newer, more efficient and reliable bioassays for large-scale rapid evaluation of radioprotective efficacy of plant extracts.

Further studies are necessary systematically to evaluate efficacy using standardized extracts, and to identify the bioactive compounds responsible for the radioprotective manifestation. Isolation of the bioactive constituents, and subsequent combination in appropriate proportions along with bio-enhancers may further potentiate the effects of herbal radioprotective drugs.

Most radioprotectors are prophylactic in nature rather than therapeutic, restricting their usage. There is usually a 'window' of time (in most cases 30 min to 2 h) prior to irradiation, when the administration of the relevant herbal preparations renders maximum survival.

There is, therefore, a need to develop simultaneously therapeutic formulations that can also be of use in a post-irradiation scenario.

Several plants found to render radioprotection, e.g. *Ginkgo biloba* and *Podophyllum hexandrum* already fall in the category of endangered plants, and efforts need to be made to save these plants from overexploitation. Biotechnological interventions for large-scale production of high-value, low-volume radioprotective constituents could also reduce the reliance on field-grown plants, and lead to the production of novel biomolecules with substantially improved radioprotective efficacy.

Clinical trials have not yet been undertaken with most herbal radioprotectors. If these are performed, herbal radioprotective drugs for human use from several of these plants may soon be available.

REFERENCES

- Abraham SK, Sarma L, Kesavan PC. 1993. Protective effects of chlorogenic acid, curcumin and beta-carotene against gamma-radiation-induced *in vivo* chromosomal damage. *Mutat Res* **303**: 109–112.
- Afanas'ev IB, Dorozhko AI, Brodskii AV, Kostyuk VA, Potapovitch AI. 1989. Chelating and free radical scavenging mechanisms of inhibitory action of rutin and quercetin in lipid peroxidation. *Biochem Pharmacol* **38**: 1763–1769.
- Agarwal GN, Katiyar GK, Arora D, Kachroo P. 2003. Usefulness of Dabur Chyawanprash special Ayurvedic medicine in prevention of early reactions during radiotherapy. *Antiseptic* **100**: 189–192.
- Agarwal OP, Nagaratnam A. 1981. Radioprotective property of flavonoids in mice. *Toxicol* **19**: 201–204.
- Agrawal PK, Goel HC. 2002. Protective effect of RH-3 with special reference to radiation-induced micronuclei in mice bone marrow. *Indian J Exp Biol* **40**: 525–530.
- Agarwal SS, Singh VK. 1999. Immunomodulation: A review of studies on Indian medicinal plants and synthetic peptides. *Proc Indian Natl Acad Sci* **B65**: 79–204.
- Agarwal V, Sharma J, Jaimala, Pahadiya S, Sharma R. 2001. Use of *Centella asiatica* extract as detoxicating agent against combined toxicity of gamma radiation and cadmium chloride. *J Phytol Res* **14**: 25–28.
- Aherne SA, O'Brien NM. 2000. Mechanism of protection by the flavonoids, quercetin and rutin against tert-butylhydroperoxide and menadione-induced DNA single strand breaks in Caco2 cells. *Free Rad Biol Med* **29**: 507–514.
- Ahmad I, Mehmood Z, Mohammed F. 1998. Screening of some Indian medicinal plants for their antimicrobial properties. *J Ethnopharmacol* **62**: 183–193.
- Akita Y, Yang Y, Kawai T, et al. 1998. New assay method for surveying anti-emetic compounds from natural sources. *Nat Prod Sci* **4**: 72–77.
- Alaoui-Youssefi A, Lamproglou I, Driou K, Emerit I. 1999. Anticlastogenic effects of *Ginkgo biloba* (EGB761) extract and some of its constituents in irradiated rats. *Mutat Res* **445**: 99–104.
- Ammon HP. 1993. Mechanism of antiinflammatory actions of curcumin and boswellic acids. *J Ethnopharmacol* **38**: 113–119.
- Ammon HPT, Wahl MA. 1991. Pharmacology of *Curcuma longa*. *Planta Med* **57**: 1–7.
- Arora R, Goel HC. 2000. Herbal radioprotectors. In *Proceedings of the International Conference on Radiation Biology, Radiobiology 2000*. Cancer Research Institute: Thiruvananthapuram, India, 87.
- Arora R, Gupta D, Prasad J, et al. 2003a. Bioprospection of Himalayan high-altitude region flora for radioprotective biomolecules. In *Proceedings of the National Seminar on New Millenium Strategies for Quality, Safety and GMPs for Herbal Drugs/Products*, NBRI, Lucknow, India.
- Arora R, Gupta D, Prasad J, et al. 2003b. Screening of indigenous medicinal and aromatic plants for their radioprotective potential: some considerations. In *Proceedings of the XXIV Annual Conference of Indian Association of Biomedical Scientists, Institute of Nuclear Medicine and Allied Sciences, Delhi, India*, 86–87.
- Arora R, Gupta D, Sharma AK, et al. 2004. Modification of radiation-induced damage in mammalian model systems by natural plant products: implications for radiation protection. In *Proceedings of the IUPAC International Conference on Biodiversity and Natural Products, Delhi University, Delhi, India*, 33.
- Attele AS, Zhou YP, Xie JT, et al. 2002. Anti-diabetic and antiobese effect of *Panax ginseng* berry extract and identification of its active component. *Diabetes* **51**: 1851–1858.
- Bakina EE, Rodina Y, Kinzburskii, Kopytin BM. 1967. Use of P vitamins, quercetin and flavallicept during radiation sickness in rats. *Vliyanie Organizm Fiz Khim Faktorov Vnesh Sredy Sb Rab Mater Nauch Konf* **1967**: 57–58.
- Bars W, Michael C, Saran M. 1994. Flavonoids antioxidants: Rate constants for reactions with oxygen radicals. *Methods Enzymol* **234**: 426–429.
- Beckman KB, Ames BN. 1998. The free radical theory of aging matures. *Physiol Rev* **78**: 547–581.
- Belinky PA, Aviram M, Mahmood S, Vaya J. 1998. Structural aspects of the inhibitory effects of glabridin on LDL oxidation. *Free Radic Biol Med* **24**: 1419–1429.
- Ben-Amotz A, Rachmilechiv B, Greenberg S, Sela M, Weshler Z. 1996. Natural beta-carotene and whole body irradiation in rats. *Radiat Environ Biophys* **35**: 285–288.
- Ben-Amotz A, Yatziv S, Sela M, et al. 1998. Effect of beta-carotene supplementation in children exposed to radiation from the Chernobyl accident. *Radiat Environ Biophys* **37**: 187–193.
- Ben-Hur E, Fulder S. 1981. Effect of *Panax ginseng* saponins and *Eleutherococcus senticosus* on survival of cultured mammalian cells after ionizing radiation. *Am J Chin Med* **9**: 48–56.
- Bestwick CS, Milne L. 2001. Quercetin modifies reactive oxygen levels but exerts only partial protection against oxidative stress within HL-60 cells. *Biochim Biophys Acta* **1528**: 49–59.
- Beutner KR, Von Krogh G. 1990. Current status of podophyllotoxin for the treatment of genital warts. *Semin Dermatol* **9**: 148.
- Bhargava KP, Singh N. 1981. Antistress activity of *Ocimum sanctum* Linn. *Indian J Med Res* **73**: 443.
- Blasko G, Cordell GA. 1988. *Recent Developments in the Chemistry of Plant derived Anticancer Agents*. In *Economic and Medicinal Plant Research*, Wagner H, Hiroshi H, Farnsworth NR (eds). Academic Press: London.
- Bolloor KK, Kamat JP, Devasagayam TPA. 2000. Chlorophyllin as a protector of mitochondrial membranes against gamma radiation and photosensitization. *Toxicol* **155**: 63–71.

- Bolotin E, Smogorzewska M, Smith M, Widmer M, Weinberg K. 1996. Enhancement of thymopoiesis after bone marrow transplant by *in vivo* interleukin-7. *Blood* **88**: 1887–1894.
- Booth C, Hargreaves DF, O'Shea JA, Potten CS. 1999. *In vivo* administration of genistein has no effect on small intestinal epithelial proliferation and apoptosis, but a modest effect on clonogen survival. *Cancer Lett* **144**: 169–175.
- Brinkhaus B, Linder M, Schuppan D, Hahn EG. 2000. Chemical, pharmacological and clinical profile of the East Asian medicinal plant *Centella asiatica*. *Phytomedicine* **7**: 427–448.
- Brizel DM, Wasserman TH, Henke M, et al. 2000. Phase III randomized trial of amifostine as a radioprotector in head and neck cancer. *J Clin Oncol* **18**: 3339–3345.
- Brogo V. 1984. Effects of bremsstrahlung and electron radiation on rat motor performance. *Radiat Res* **100**: 313–320.
- Bump EA, Malaker K (eds). 1998. *Radioprotectors: Chemical, Biological and Clinical Perspective*. CRC Press, Boca Raton, FL, 431.
- Burghardt WB, Hunt WA. 1985. Characterization of radiation-induced performance decrement using a two-lever shock-avoidance task. *Radiat Res* **103**: 149–157.
- Burton GW, Joyce A, Ingold KU. 1982. β -carotene is an unusual type of lipid antioxidant. *Science* **224**: 569–573.
- Cairnie A. 1983. Adverse effects of WR-2721. *Radiat Res* **94**: 221–226.
- Cao ZF, Chen ZG, Guo P, et al. 1993. Scavenging effects of ginger on superoxide anion and hydroxyl radical. *Chung-Kuo-Chung-Yao-Tsa-Chih* **18**: 750–764.
- Capizzi RL, Oster W. 1995. Protection of normal tissue from the cytotoxic effects of chemotherapy and radiation by amifostine: clinical experiences. *Eur J Cancer* **31**: S8–S13.
- Castillo J, Benavente-Garcia O, Lorente J, et al. 2000. Antioxidant activity and radioprotective effects against chromosomal damage induced *in vivo* by x-rays of flavan-3-ols (Procyanidins) from grape seeds (*Vitis vinefera*): comparative study versus other phenolic and organic compounds. *J Agric Food Chem* **48**: 1738–1745.
- Chan Y, Jiang Z, Qin W, Ni M, Li X, He Y. 1990. Chemical composition and characteristics of sea buckthorn fruit and its oil. *Chem Ind Forest Prod* **10**: 163–175.
- Chattopadhyay RR. 1999. A comparative evaluation of some blood sugar lowering agents of plant origin. *J Ethnopharmacol* **67**: 367.
- Chattopadhyay RR, Sarkar SK, Ganguly S, Medda C, Basu TK. 1992. Hepatoprotective activity of *Ocimum sanctum* leaf extract against paracetamol induced hepatic damage in rats. *Indian J Pharmacol* **24**: 163.
- Chaudhary D, Chandra D, Kale RK. 1999. Modulation of radioreponse of glyoxalase system by curcumin. *J Ethnopharmacol* **64**: 1–7.
- Chen YJ, Dai YS, Chen BF, et al. 1999. The effect of tetrandrine and extracts of *Centella asiatica* on acute radiation dermatitis in rats. *Biol Pharm Bull* **22**: 703–706.
- Cheng TJ, Pu JK, Wu LW, Ma ZR, Cao Z, Li TJ. 1994. A preliminary study on hepatoprotective action of seed oil of *Hippophae rhamnoides* L. (HR) and its mechanism of action. *Chung Kuo Chung Yao Tsa Chih* **19**: 367–370.
- Chitra P, Sajithlal GB, Chandrakasan G. 1998. Influence of *Aloe vera* on collagen characteristics in healing dermal wounds in rats. *Mol Cell Biochem* **181**: 71–76.
- Choi JS, Chung HY, Kang SS, et al. 2002. The structure-activity relationship of flavonoids as scavengers of peroxy nitrite. *Phytother Res* **16**: 232–235.
- Coleman NE, Blakely WF, Fike JR, et al. 2003. Molecular and cellular biology of moderate-dose (1–10 Gy) radiation and potential mechanisms of radiation protection: Report of a workshop (Bethesda, Maryland, USA, December, 17–18, 2001). *Radiat Res* **159**: 812–834.
- Das UN. 2002. A radical approach to cancer. *Med Sci Monit* **8**: 79–82.
- Davis RH, Kabbani JM, Maro NP. 1987. *Aloe vera* and wound healing. *J Am Pediatr Med Assoc* **77**: 165–169.
- De Feudis FV. 1991. *Ginkgo biloba Extract (Egb 761): Pharmacological Activities and Clinical Applications*. Elsevier: Paris.
- Denison L, Haigh A, D' Cunha G, Martin F. 1992. DNA ligands as radioprotectors: molecular studies with Hoechst 33342 and Hoechst 33258. *Int J Radiat Biol* **61**: 69–81.
- Devasagayam TPA, Kesavan PC. 1996. Radioprotective and antioxidant action of caffeine: mechanistic considerations. *Indian J Exp Biol* **34**: 291–297.
- Devasagayam TPA, Sainis KB. 2002. Immune system and antioxidants, especially those derived from Indian medicinal plants. *Indian J Exp Biol* **40**: 639–655.
- Dewick PM. 2002. *Medicinal Natural Products: A Biosynthetic Approach*, 2nd edn. Wiley: Chichester, 507.
- Dharmani P, Kulshreshta DK, Palit Z. 2002. Evaluation of anticancerogenic activity of crude drug extract of *Ocimum sanctum* Linn. *Indian J Pharmacol* **34**: 296.
- Diplock AT, Charleux JL, Crozier-Willi G, et al. 1998. Functional food science and defence against reactive oxygen species. *Br J Nutr* **80**: 77–112.
- Diwan PV, Karwande I, Singh AK. 1991. Anti-anxiety profile of Mandukparni *Centella asiatica* Linn. in animals. *Fitoterapia* **62**: 255–257.
- Dorr WF, Noack R, Spekl K, Farell CL. 2001. Modification of oral mucositis by keratinocyte growth factor: single radiation exposure. *Int J Radiat Res* **77**: 341–347.
- Duche JC, Barre J, Guinot P, Duchier J, Cournot A, Tillement JP. 1988. Effect of *Ginkgo biloba* constituents related to protection against brain damage caused by hypoxia. *Pharmacol Res Commun* **20**: 349–368.
- Duthie G, Crozier A. 2000. Plant-derived phenolic antioxidants. *Curr Opin Lipidol* **11**: 43–47.
- Duthie GG, Duthie SJ, Kyle JAM. 2000. Plant polyphenols in cancer and heart disease: implications as nutritional antioxidants. *Nutr Res Rev* **13**: 79–106.
- Duthie SJ, Collins AR, Duthie GG, Dobson VL. 1997. Quercetin and myricetin protect against hydrogen peroxide-induced DNA damage (strand breaks and oxidized pyrimidines) in human lymphocytes. *Mutat Res* **248**: 223–231.
- Dutta BK, Rahman I, Das TK. 1998. Antifungal activity of Indian medicinal plant extracts. *Mycoses* **41**: 535–536.
- Dwarakanath BS, Adhikari JS, Jain V. 1999. Hematoporphyrin derivatives potentiate the radiosensitizing effects of 2-deoxy-D-glucose in cancer cells. *Int J Radiat Oncol Biol Phys* **43**: 1125–1133.
- Eilert U, Wolter SB, Nahrstedt A. 1981. The antibiotic principle of seeds of *Moringa oleifera* and *Moringa stenopetala*. *Planta Med* **42**: 55–61.
- Emerit I, Arutyunyan R, Oganessian N, et al. 1995a. Radiation-induced clastogenic factors: Anticlastogenic effects of *Ginkgo biloba* extract. *Free Radic Biol Med* **18**: 985–991.
- Emerit I, Oganessian N, Pogossian A, et al. 1997a. Oxidative stress-related clastogenic factors in plasma of Chernobyl liquidators: protective effects of antioxidant plant phenols, vitamins and oligoelements. *Mutat Res* **377**: 239–246.
- Emerit I, Oganessian N, Sarkisian T, et al. 1995b. Clastogenic factors in the plasma of Chernobyl recovery workers: Anticlastogenic effects of *Ginkgo biloba* extract. *Radiat Res* **144**: 198–205.
- Emerit I, Quastel M, Goldsmith J, et al. 1997b. Clastogenic factors in the plasma of children exposed at Chernobyl. *Mutat Res* **373**: 47–54.
- Ezeamuzie IC, Ambakederemo AW, Shode FO, Ekwebelem SC. 1996. Antiinflammatory effects of *Moringa oleifera* root extract. *Int J Pharmacog* **34**: 207–212.
- Fabricant DS, Farnsworth NR. 2001. The value of plants used in traditional medicine for drug discovery. *Environ Health Perspect* **109** (Suppl 1): 69–75.
- Fleischner AM. 1985. Plant extracts to accelerate healing and reduce inflammation. *Cosmet Toilet* **100**: 45–58.
- Franz CG. 1985. Effects of mixed neutron-gamma total-body irradiation on physical activity performance of rhesus monkeys. *Behav Neural Biol* **40**: 114–118.
- Frisch C, Asenohri RU, Mattern CM, Hacker R, Huston JP. 1995. Blockade of lithium chloride-induced conditioned taste aversion as a test for antiemetic agents: Comparison of metoclopramide with combined extracts of *Zingiber officinale* and *Ginkgo biloba*. *Pharmacol Biochem Behav* **52**: 321–327.
- Furuse M, Tsuneoka K, Uchida K, Nomoto K. 1997. Acceleration of granulocytic cell recovery in irradiated mice by a single subcutaneous injection of a heat-killed *Lactobacillus casei* preparation. *J Radiat Res* **38**: 111–120.
- Ganapathi NG, Jagetia GC. 1995. Liv 52 pretreatment inhibits radiation-induced lipid peroxidation in mouse liver. *Curr Sci* **68**: 601–603.
- Ganasoundari A, Uma Devi P, Rao BSS. 1998. Enhancement of bone marrow radiation protection and reduction in WR-2721 toxicity by *Ocimum sanctum*. *Mutat Res* **397**: 303–312.

- Ganasoundari A, Uma Devi P, Rao MNA. 1997a. Protection against radiation-induced chromosome damage in mouse bone marrow by *Ocimum sanctum*. *Mutat Res* **373**: 271–276.
- Ganasoundari A, Zare SM, Uma Devi P 1997b. Modification of normal tissue sensitivity by some medicinal plants. *Br J Radiol* **70**: 599–602.
- Gao H, Wang F, Lien F-J, Trousdale MD. 1996. Immunostimulating polysaccharides from *Panax notoginseng*. *Pharm Res* **13**: 1198–1200.
- Gebhardt R. 2002. Oxidative stress, plant-derived antioxidants and liver fibrosis. *Planta Med* **68**: 289–296.
- George KC, Hebbar SA, Kale SP, Kesavan PC. 1999. Caffeine protects mice against whole-body lethal dose of gamma-irradiation. *J Radiol Prot* **19**: 171–176.
- Giambarresi L, Jacobs AJ. 1987. Radioprotectants. In *Military Radiobiology*, Conklin JJ, Walker RI (eds). Academic Press: London, 265–301.
- Gileva TG, Lukin AV. 1984. Assessment of the efficacy of radioprotective compounds in radiation therapy of larynx carcinoma. *Kazanskii Meditsinskii Zhurnal* **75**: 374–375.
- Glover D, Riley L, Carmichael K, et al. 1983. Hypocalcemia and inhibition of parathyroid hormone secretion after administration of WR-2721 (radioprotective and chemopreventive agent). *N Engl J Med* **309**: 1137–1141.
- Godhwani S, Godhwani JL, Vyas DS. 1987. *Ocimum sanctum*: an experimental study evaluating its anti-inflammatory, analgesic, and antipyretic activity in animals. *J Ethnopharmacol* **21**: 153.
- Godhwani S, Godhwani JL, Vyas DS. 1988. *Ocimum sanctum* – a preliminary study evaluating its immunoregulatory profile in albino rats. *J Ethnopharmacol* **24**: 193.
- Goel HC, Arora R, Prasad J, et al. 2000a,b. *A Process for Preparation of a Radioprotective Herbal Extract-I and II*. Indian Patent filed, Patent Office, New Delhi, India.
- Goel HC, Arora R, Prasad J, et al. 2001a. *A Radioprotective Herbal Formulation*. Indian Patent filed, Patent Office, New Delhi, India.
- Goel HC, Arora R, Shobi V, Mathew TL. 2000d. *A Process for Preparation of a Behavioural Radioprotective Herbal Extract*. Indian Patent filed, Patent Office, New Delhi, India.
- Goel HC, Kumar IP, Samanta N, Arora R. 2001c. *A Process for Preparation of an Anticancer and Radioprotective Herbal Extract*. Indian Patent filed, Patent Office, New Delhi, India.
- Goel HC, Prasad J, Singh S, et al. 2004. Radioprotective potential of a herbal extract of *Tinospora cordifolia*. *J. Radiat. Res* **45**: 61–68.
- Goel HC, Prasad J, Singh S, Sagar RK, Prem Kumar I, Sinha AK. 2002a. Radioprotection by a herbal preparation of *Hippophae rhamnoides*, RH-3, against whole body lethal irradiation in mice. *Phytomedicine* **9**: 15–25.
- Goel HC, Prem Kumar I, Rana SVS. 2002b. Free radical scavenging potential of *Tinospora cordifolia*, a possible role in radioprotection. *Indian J Exp Biol* **40**: 727–734.
- Goel HC, Prem Kumar I, Samanta N, Rana SVS. 2003b. Induction of DNA-protein cross-links by *Hippophae rhamnoides*, implications in radioprotection and cytotoxicity. *Mol Cell Biochem* **245**: 57–67.
- Goel HC, Saji Kumar S, Sharma AK. 2002c. Effects of *Podophyllum hexandrum* on radiation-induced delay of post-natal appearance of reflexes and physiological markers in rats irradiated *in utero*. *Phytomedicine* **9**: 447–454.
- Goel HC, Salin CA, Prakash H, 2003a. Protection of jejunal crypts by RH-3 (a preparation of *Hippophae rhamnoides*) against lethal whole body gamma irradiation. *Phytother Res* **17**: 222–226.
- Goel HC, Sinha AK, Dogra R, et al. 2001b. *A Process for Preparation of a Radioprotective Herbal Extract from Tinospora sps*. Indian Patent filed, Patent Office, New Delhi, India.
- Goel HC, Sinha AK, Joshi BP, et al. 2000c. *A Process for Preparation of a Radioprotective Herbal Extract from Hippophae rhamnoides*. Indian Patent filed, Patent Office, New Delhi, India.
- Gohil K, Moy RK, Farzin S, Maguire JJ, Packer L. 2000. mRNA expression profile of a human cancer cell line in response to *Ginkgo biloba* extract: induction of antioxidant response and the Golgi system. *Free Radic Res* **33**: 831–849.
- Gordon L, Ruml D, Hahne H, Miller P. 1955. Studies on susceptibility to infection following ionizing radiation. *J Exp Med* **102**: 403.
- Gowdey G, Lee RK, Carpenter WM. 1995. Treatment of HIV-related hairy leukoplakia with podophyllum resin 25% solution. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **79**: 64–67.
- Guenechea G, Albella B, Bueren JA, et al. 1997. AM218, a new polyanionic polysaccharide, induces radioprotection in mice when administered shortly before irradiation. *Int J Radiat Biol* **71**: 101–108.
- Gupta D, Arora R, Garg AP, Bala M, Goel HC. 2004. Modification of radiation-damage to mitochondrial system *in vivo* by *Podophyllum hexandrum*: mechanistic aspects. *Mol Cell Biochem* **266**: 65–77.
- Gupta D, Arora R, Garg AP, Goel HC. 2003b. Radiation-induced damage to mitochondria and its modification by *Podophyllum hexandrum* Royale in HepG2 cells. *Mol Cell Biochem* **247**: 25–40.
- Gupta D, Arora R, Goel HC. 2002. Effect of Himalayan Mayapple on mitochondria: implications in radiation protection. *Indian J Nuclear Med* **17**: 19–20.
- Gupta D, Arora R, Gupta S, Garg AP, Goel HC. 2003a. Radiation-induced oxidative damage to mitochondria and its modification by endogenous antioxidants *in vivo*: role of some herbal preparations. *Indian J Nuclear Med* **18**: 14.
- Gupta D, Arora R, Garg AP, Goel HC. 2003c. Radioprotection of mice liver mitochondria by *Hippophae rhamnoides* L. *in vivo*. In *Proceedings of 3rd Biennial Conference on Hyperthermic Oncology and Medicine, Nanavati Hospital, Mumbai, India*.
- Gupta NK. 1996. Modification of radiation-induced changes in murine hepatic lipid profiles by garlic (*Allium sativum* Linn.) unsaturated oils. *Indian J Exp Biol* **34**: 851–853.
- Haguenauer JP, Cantenot F, Koskas H, Pierart H. 1986. Treatment of equilibrium disorders with *Ginkgo biloba* extract. A multicenter double-blind drug vs. placebo study. *Presse Med* **25**: 1569–1572.
- Hahn SM, Krishna MC, Samuni A, et al. 1994. Potential use of nitroxides in radiation oncology. *Cancer Res* **54**: 2006S–2010S.
- Hahn SM, Sullivan FJ, De Luca AM, et al. 1999. Hemodynamic effect of the nitroxide superoxide mimics. *Free Rad Biol Med* **27**: 529–535.
- Halliwell B, Gutteridge JMC. 1989. Lipid peroxidation: A radical chain reaction. In *Free Radicals in Biology and Medicine*. Clarendon Press: Oxford.
- Halliwell B, Gutteridge JMC. 1990. Role of free radicals and catalytic metal ions in human disease: An overview. *Methods Enzymol* **186**: 1–85.
- Handa SS, Chawla AS, Sharma AK. 1992. Plants with anti-inflammatory activity. *Fitoterapia* **63**: 3–31.
- Hannequin D, Thibert A, Vaschalde Y. 1986. Development of a model to study the anti-oedema properties of *Ginkgo biloba* extract. *Presse Med* **15**: 1575–1576.
- Hanson WR, Houseman KA, Colins PW. 1988. Radiation protection *in vivo* by prostaglandins and related compounds of the arachidonic acid cascade. *Pharmacol Ther* **39**: 347–356.
- Haraguchi H, Hoshida N, Ishikawa H, Tamura Y, Mizutani K, Kinoshita T. 2000. Protection of mitochondrial functions against oxidative stresses by isoflavones from *Glycyrrhiza glabra*. *J Pharm Pharmacol* **52**: 219–223.
- Harapanalli RS, Nara VR, Yaghamai V, et al. 1994. Vitamins as radioprotectors *in vivo*: II Protection by vitamin A and soybean oil against radiation damage caused by internal radionuclides. *Radiat Res* **139**: 115–122.
- Havsteen B. 1984. Flavonoids: A class of natural products of high pharmacological potency. *Biochem Pharmacol* **32**: 1141–1148.
- Hebbar SA, Mitra AK, George KC, Verma NC. 2002. Caffeine ameliorates radiation-induced skin reactions in mice but does not influence tumor radiation response. *J Radiol Prot* **22**: 63–69.
- Heinjnjen J, Knipschild P. 1992. *Ginkgo biloba*. *Lancet* **340**: 1136–1139.
- Herodin FJ, Bourin P, Mayol JF, Lataillade JJ, Drouet M. 2003. Short-term injection of antiapoptotic cytokine combinations soon after lethal gamma irradiation promotes survival. *Blood* **101**: 2109–2116.
- Hsu HY, Ho YH, Lian SL, Lin CC. 1991. Preliminary study on antiradiation effect of Kuei-Pi-Tang. *Am J Chin Med* **19**: 275–284.

- Hsu HY, Ho YH, Lian SL, Lin CC. 1993. Preliminary study on the anti-radiation effect of Jeng-Sheng-Yang-Yung-Tang. *Am J Chin Med* **21**: 187–195.
- Hsu HY, Lin CC, Hau DM. 1992. Restoration of radiation injury in mice by two Chinese medicinal prescriptions Kuei-Pi-Tang and Jeng-Sheng-Yang-Yung-Tang. *Phytother Res* **6**: 294–299.
- Hsu HY, Yang JJ, Lian SL, Ho YH, Lin CC. 1996a. Recovery of the haematopoietic system by Si-Jun-Zi-Tang in whole body irradiated mice. *J Ethnopharmacol* **54**: 69–75.
- Hsu HY, Yang JJ, Lin SY, Lin CC. 1997. Comparisons of geniposidic acid and geniposide on antitumor and radioprotection after sublethal irradiation. *Cancer Lett* **113**: 31–37.
- Hsu HY, Yang JJ, Ho YJ, Lin CC. 1999. Difference in the effects of radioprotection between aerial and root parts of *Lycium chinese*. *J Ethnopharmacol* **64**: 101–108.
- Hsu JY, Ho YH, Lin CC. 1996b. Protection of mouse bone marrow by Si-Wu-Tang against whole body irradiation. *J Ethnopharmacol* **52**: 113–117.
- Huguot F, Drieu K, Piriou A. 1994. Decreased cerebral 5-HT_{1A} receptors during ageing reversal by *Ginkgo biloba* extract (Egb 761). *J Pharm Pharmacol* **46**: 316–318.
- Ianev E, Radiv S, Balutsov M, Klouchek E, Popov A. 1995. The effect of an extract of Sea Buckthorn (*Hippophae rhamnoides* L.) on the healing of experimental skin wounds in rats. *Khirurginia (Sofia)* **48**: 30–33.
- Inano H, Onoda M. 2002. Radioprotective action of curcumin extracted from *Curcuma longa* Linn.: inhibitory effect of formation of urinary 8-hydroxy-2'-deoxyguanosine, tumorigenesis, but not mortality, induced by gamma-ray irradiation 8-hydroxy-2'-deoxyguanosine, tumorigenesis, but not mortality, induced by gamma-ray irradiation. *Int J Radiat Oncol Biol Phys* **53**: 735–743.
- Ishiguro K, Yamaki M, Kashihara M, Takagi S. 1986. Sarothralen A and B, new antibiotic compounds from *Hypericum japonicum*. *Planta Med* **53**: 415–417.
- Jadhav HR, Bhutani KK. 2002. Antioxidant properties of Indian medicinal plants. *Phytother Res* **16**: 771–773.
- Jagetia GC, Baliga MS. 2002a. Influence of the leaf extract of *Mentha arvensis* Linn. (mint) on the survival of mice exposed to different doses of gamma radiation. *Strahlenther Onkol* **178**: 91–98.
- Jagetia GC, Baliga MS. 2002b. *Syzygium cumini* (Jamun) reduces the radiation-induced damage in the cultured human peripheral blood lymphocytes: a preliminary study. *Toxicol Lett* **132**: 19–25.
- Jagetia GC, Baliga MS. 2003. Treatment of mice with a herbal preparation (Mentat) protects against radiation-induced mortality. *Phytother Res* **17**: 876–881.
- Jagetia GC, Baliga MS, Aruna R, Rajanikant GK, Jain V. 2003a. Effect of abana (a herbal preparation) on the radiation-induced mortality in mice. *J Ethnopharmacol* **2857**: 1–7.
- Jagetia GC, Baliga MS, Malagi KJ, Kamath SM. 2002. The evaluation of radioprotective effect of triphala (an Ayurvedic rejuvenating drug) in the mice exposed to gamma-radiation. *Phytomedicine* **9**: 99–108.
- Jagetia GC, Baliga MS, Venkatesh P, Ulloor JN. 2003b. Influence of ginger rhizome (*Zingiber officinale* Rosc) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to gamma radiation. *Radiat Res* **160**: 584–592.
- Jagetia GC, Rajnikant GK, Rao SK, Shrinath BM. 2003c. Alteration of the glutathione, glutathione peroxidase, superoxide dismutase and lipid peroxidation by ascorbic acid in the skin of mice exposed to fractionated gamma radiation. *Clin Chim Acta* **332**: 111–121.
- Jagetia GC, Shirwaikar A, Rao SK, Bhilegaonkar PM. 2003d. Evaluation of the radioprotective effect of *Ageratum conyzoides* Linn. Extract in mice exposed to different doses of gamma radiation. *J Pharm Pharmacol* **55**: 51–58.
- Jagetia GC, Venkatesh P, Baliga MS. 2003e. Evaluation of radioprotective effect of *Aegle marmelos* (L.) Correa in the cultured human peripheral blood lymphocytes exposed to different doses of γ -radiation: a micronucleus study. *Mutagenesis* **18**: 387–393.
- Jaiswal SK, Bordia A. 1996. Radioprotective effect of garlic *Allium sativum* Linn. in albino rats. *Indian J Med Sci* **50**: 231–233.
- Jo EH, Hong HD, Ahn NC, et al. 2004. Modulations of Bcl2/Bax family were involved in chemopreventive effects of licorice root (*Glycyrrhiza uralensis* Fisch) in MCF-7 human breast cancer cells. *J Agric Food Chem* **52**: 1715–1719.
- Joshima H, Ohara H, Aoki Y. 1992. The effect of OK-432 upon erythropoietic recovery in sub-lethally irradiated mice: a preliminary report. *J Radiat Res* **33**: 290–300.
- Kalechman Y, Shani A, Albeck M, Sredni B. 1995a. Induction of acute phase proteins in mice and humans by treatment with AS101, an immunomodulator with radioprotective properties. *Immunopharmacology* **29**: 149–158.
- Kalechman Y, Zuloff A, Albeck M, Strassmann G. 1995b. Role of endogenous cytokine secretion in radioprotection conferred by the immunomodulator ammonium trichloro-(dioxethylene-0, 0') tellurate. *Blood* **85**: 1555–1561.
- Kamat JP, Bolor KK, Devasagayam TPA, Jayashree B, Kesavan PC. 2000a. Differential modification by caffeine of oxygen-dependent and independent effects of gamma irradiation on rat liver mitochondria. *Int J Radiat Biol* **76**: 1281–1288.
- Kamat JP, Bolor KK, Devasagayam TPA, Kesavan PC. 1999. Protection of superoxide dismutase by caffeine in rat liver mitochondria against γ -irradiation. *Curr Sci* **77**: 286–289.
- Kamat JP, Bolor KK, Devasagayam TPA, Venkatachalam SR. 2000b. Antioxidant properties of *Asparagus racemosus* induced damage induced by gamma-radiation in rat liver mitochondria. *J Ethnopharmacol* **71**: 425–435.
- Kandaswami C, Middleton E. 1994. Free Radical Scavenging and Antioxidant Activity of Plant Flavonoids. In *Free Radicals in Diagnostic Medicine* Armstrong D (ed.). Plenum Press: New York, 351–376.
- Kapil A, Sharma S. 1997. Immunopotentiating compounds from *Tinospora cordifolia*. *J Ethnopharmacol* **58**: 89–95.
- Kapoor LD. 1990. *Handbook of Ayurvedic Medicinal Plants*. CRC Press: Boca Raton, FL.
- Kapoor R, Mehta U. 1998. Supplementary effect of spirulina on haematological status of rats during pregnancy and lactation. *Plant Foods Hum Nutr* **52**: 315–324.
- Karpov LM, Brown II, Poltavtseva NV, Ershova SG, Vasileva TV, Chaban lu L. 2000. The postradiation use of vitamin-containing complexes and a phycocyanin extract in a radiation lesion in rats. *Radiat Biol Radioecol* **40**: 310–314.
- Keum YS, Park KK, Lee JM, et al. 2000. Antioxidant and anti-tumor promoting activities of the methanolic extract of heat-processed ginseng. *Cancer Lett* **150**: 41–48.
- Kim JH, Kim SH, Lee EJ. 1998a. Radioprotective effect of Lifukang, a Chinese medicinal plant prescription. *Nat Prod Sci* **4**: 26–31.
- Kim JK, Kim JH, Yoon YD. 2003. Evaluation of caffeine as a radioprotector in whole-body irradiated male mice. *In vivo* **17**: 197–200.
- Kim JY, Germolec DR, Luster MI. 1990. *Panax ginseng* as a potential immunomodulator: studies in mice. *Immunopharmacol Immunotoxicol* **12**: 257–276.
- Kim KH, Lee YS, Jung IS, et al. 1998b. Acidic polysaccharides from *Panax ginseng*, ginsan, induces Th1 cells and macrophage cytokines and generates LAK cells in synergy with rIL-2. *Planta Med* **64**: 110–115.
- Kim SH, Cho CK, Yoo SY, Koh KH, Yun HG, Ki MTH. 1993. *In vivo* radioprotective activity of *Panax ginseng* and diethyldithiocarbamate. *In Vivo* **7**: 467–470.
- Kim SH, Son CH, Nah SY, Jo SK, Byun MW, Shin DH. 2001. Modification of radiation response in mice by *Panax ginseng* and diethyldithiocarbamate. *In Vivo* **15**: 407–411.
- Kim YC, Kim SR, Markelonis GJ, Oh TH. 1998b. Ginsenosides Rb1 and Rg3 protect cultured rat cortical cells from glutamate-induced neurodegeneration. *J Neurosci Res* **53**: 426–432.
- Kim YK, Guo Q, Packer L. 2002. Free radical scavenging activity of red ginseng aqueous extracts. *Toxicology* **172**: 149–156.
- Klein AD, Penneys NS. 1988. Aloe vera. *J Am Acad Dermatol* **18**: 714–720.
- Kligerman MM, Glover DJ, Turrisi AT, et al. 1984. Toxicity of WR-2721 administered in single and multiple doses. *Int J Radiat Oncol Biol Phys* **10**: 1773–1776.
- Korina LG, Afanas'ev IB. 1997. Antioxidant and chelating properties of flavonoids. *Adv Pharmacol* **38**: 151–163.
- Kovalenko PG, Antonjuk VP, Maluita SS. 2003. Secondary metabolites production from transformed cells of *Glycyrrhiza glabra* and *Potentilla alba* producers of radioprotective compounds. *Ukrainian Biorganica Acta* **1**: 1–17.

- Kroemer G, Reed JC. 2000. Mitochondrial control of cell death. *Nat Med* **6**: 513–519.
- Kropacova K, Misurova E, Hakova H. 1998. Protective and therapeutic effect of silymarin on the development of latent liver damage. *Radiat Biol Radioecol* **38**: 411–415.
- Kulp KS, Green SL, Vulliet PR. 1996. Iron deprivation inhibits cyclin-dependent kinase activity and decreases cyclin D/CDK4 protein levels in synchronous MDA-MB-543 human breast cancer cells. *Exp Cell Res* **229**: 60–68.
- Kulp KS, Vulliet PR. 1996. Mimosine blocks cell cycle progression by chelating iron in asynchronous human breast cancer cells. *Toxicol Appl Pharmacol* **139**: 356–364.
- Kumar KS, Srinivasan V, Toles R, Jobe L, Seed TM. 2002. Nutritional approaches to radioprotection: Vitamin E. *Milit Med* **167**: 57–59.
- Kumar M, Sharma MK, Saxena PS, Kumar A. 2003. Radioprotective effect of *Panax ginseng* on the phosphatases and lipid peroxidation level in testes of Swiss albino mice. *Biol Pharm Bull* **26**: 308–312.
- Kumar MHV, Gupta VK. 2002. Effect of different extracts of *Centella asiatica* on cognition and markers of oxidative stress in rats. *J Ethnopharmacol* **79**: 253–260.
- Kumar P, Kuttan R, Kuttan G. 1996. Radioprotective effect of Rasayanas. *Indian J Exp Biol* **34**: 848–850.
- Kumar R, Singh DP, Chaturvedi VK, Pathak RC. 1997. A note on antiviral property of neem (*Melia azadirachta*) and tulsi (*Ocimum sanctum*) against Newcastle disease virus. *Indian J Comp Microbiol Immunol Infect Dis* **18**: 192.
- Kumar SS, Devasagayam TPA, Jayshree B, Kesavan PC. 2001. Mechanism of protection against radiation-induced DNA damage in plasmid pBR322 by caffeine. *Int J Radiat Biol* **77**: 617–623.
- Kure F. 1992. The radioprotective effects of methylprednisolone and Sho-Saikoto on mouse lung. *Nippon Igaku Hoshasen Gakkai Zasshi* **52**: 96–103.
- Lam SK, Ng TB. 2002. Pananotin, a potent antifungal protein from roots of the traditional Chinese medicinal herb *Panax ginseng*. *Plant Med* **68**: 1024–1028.
- Landauer MR, Davis HD, Dominitz JA, Weiss JF. 1987. Dose and time relationships of the radioprotector WR-2721 on locomotor activity in mice. *Pharmacol Biochem Behav* **27**: 573–576.
- Landauer MR, McChesney DG, Ledney GD. 1997. Synthetic trehalose dicorynomycolate (S-TDCM): behavioural effects and radioprotection. *J Radiat Res* **38**: 45–54.
- Landauer MR, Srinivasan V, Seed TM. 2003. Genistein treatment protects mice from ionizing radiation injury. *J Appl Toxicol* **21**: 25–31.
- Landauer MR, Srinivasan V, Shapiro A, Takimoto C, Seed TM. 2000. Protection against lethal irradiation by genistein. *Int J Toxicol* **19**: 37.
- Lee YS, Jung IS, Lee IR, Kim KW, Hong WS, Yun YS. 1997. Activation of multiple effector pathways of immune system by the antineoplastic immunomodulator acidic polysaccharide ginsan isolated from *Panax ginseng*. *Anticancer Res* **17**: 323–332.
- Li TSC. 1999. Sea Buckthorn: New Crop Opportunity. *Perspectives on New Crops and New Uses*, Janick I (ed.). ASHS Press: Alexandria, USA, 335–337.
- Lin IH, Hau DM, Chen WC, Chen KT, Lin JG. 1996. Effects of glycyrrhizae and glycyrrhizic acid on cellular immunocompetence of gamma-ray-irradiated mice. *Chin Med J (Engl)* **109**: 138–142.
- Links M, Lewis C. 1999. Chemoprotectants: a review of their clinical pharmacology and therapeutic efficiency. *Drugs* **57**: 293–308.
- Liu CX, Xiao PG. 1993. Danggui (*Angelica sinensis*). *An Introduction to Chinese Materia Medica*. Beijing Medicinal University and Peking Union Medical University Press: Beijing, 168.
- Liu FM, Li ZX, Shi S. 1998. Effect of total flavones of *Hippophae rhamnoides* L. on cultured rat heart cells and on cAMP level and adenylate cyclase in myocardium. *Chung-kuo-yao-li-Hsueh-Pao* **9**: 539–542.
- Livesey JC, Reed DJ. 1987. Chemical protection against ionizing radiation. *Adv Radiat Res* **13**: 285–353.
- Ma L, Zhou ZL, Yang Q. 1995. Study on effect of polysaccharide of ginseng on peripheral blood mononuclear cells induced interleukin-2 production and activity of its receptors *in vitro*. *Chung Kuo Chung Hsi Chieh Ho Tsa chih* **15**: 411–413.
- Maharwal J, Samarth RM, Saini MR. 2003. Radiomodulatory influence of Rajgira (*Amaranthus paniculatus*) leaf extract in Swiss albino mice. *Phytother Res* **17**: 1150–1154.
- Maiche AG. 1991. Effect of chamomile cream and almond ointment on acute radiation skin reaction. *Acta Oncol* **30**: 395–396.
- Maisin JR. 1998. Chemical radioprotection: past, present and future prospects. *Int J Radiat Biol* **73**: 443–450.
- Martin RF, Anderson RF. 1999. Pulse radiolysis studies indicate that electron transfer is involved in radioprotection by Hoechst 33342 and methylproamine. *Int J Radiat Oncol Biol Phys* **42**: 827–831.
- Martin RF, Broadhurst S, D'Abrew S, et al. 1996. Radioprotection by DNA ligands. *Br J Cancer Suppl* **27**: S99–S101.
- Matsubara J, Tajima Y, Karasawa M. 1987. Metallothionein induction as a potent means of radiation protection in mice. *Radiat Res* **120**: 442–455.
- Mehra KS, Mikuni I, Gupta U, Gode KD. 1984. *Curcuma longa* (Linn.) drops in corneal wound healing. *Tokai J Exp Clin Med* **9**: 27–31.
- Mei QB. 1988. Effects of *Angelica sinensis* polysaccharides on hemopoietic stem cells in irradiated mice. *Chung Kuo Yao Li Hsueh Pao* **9**: 279–282.
- Mei QB, Tao JY, Cui B. 1991. Advances in the pharmacological studies of *Radix angelica sinensis* (Oliv) diels (Chinese danggui). *Chin Med J* **104**: 776–781.
- Mettler FA, Guskova AK. 2001. Treatment of acute radiation sickness. In *Medical Management of Radiation Accidents*, 2nd edn, Gusev IA, Guskova AK, Mettler FA (eds). CRC Press: Boca Raton, FL, 53–67.
- Meyer K, Schwartz J, Crater D, Keyes B. 1995. *Zingiber officinale* (ginger) used to prevent 8-MOP associated nausea. *Dermatol Nurs* **7**: 242–244.
- Middleton E Jr, Kandaswami C, Theoharides TC. 2000. The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease and cancer. *Pharmacol Rev* **52**: 673–751.
- Miko S, Yanai T, Hasegawa H, et al. 1998. Concentration of metallothionein in mice livers after small dose of irradiation. *J Radiat Res* **39**: 239–242.
- Mitchell JB, De Graff W, Kaufman D, et al. 1991. Inhibition of oxygen-dependent radiation-induced damage by the nitroxide superoxide dismutase mimic, tempol. *Arch Biochem Biophys* **289**: 62–70.
- Mitchell JB, Russo A, Kuppusamy, Krishna MC. 2000. Radiation radicals and images. *Ann NY Acad Sci* **899**: 28–43.
- Mittal A, Pathania V, Agarwala PK, Prasad J, Singh S, Goel HC. 2002. Influence of *Podophyllum hexandrum* on endogenous antioxidant defense system in mice: possible role in radioprotection. *J Ethnopharmacol* **76**: 253–262.
- Miyanomae T, Frindel E. 1988. Radioprotection of haemopoiesis conferred by *Acanthopanax senticosus* Harms (Shigoka) administered before or after irradiation. *Exp Hematol* **16**: 801–806.
- Mizina T Yu, Sitnikova SG. 1999. Antiradiation activity of juice concentrate from *Hippophae rhamnoides* L. fruits. *Rastitel'mye Resursy* **35**: 85–92.
- Mizuno M, Yamada J, Teral H, Kozukue N, Lee YS, Tsuchida H. 1994. Differences in immunomodulatory effects between wild and cultured *Panax ginseng*. *Biochem Biophys Res Commun* **16**: 1672–1678.
- Molteni A, Moulder JE, Cohen EF, et al. 2000. Control of radiation induced pneumopathy and lung fibrosis by angiotensin converting enzyme inhibitors and angiotensin II type 1 receptor blocker. *Int J Radiat Biol* **76**: 523–532.
- Morel I, Lescoat G, Gogrel P, et al. 1993. Antioxidant and iron-chelating activities of the flavonoids catechin, quercetin, and diosmetin on iron loaded rat hepatocyte cultures. *Biochem Pharmacol* **7**: 13–19.
- Moulder JE. 2002. Radiobiology of nuclear terrorism: Report on an interagency workshop (Bethesda, Maryland, USA, December, 17–18, 2001). *Int J Radiat Oncol Biol Phys* **54**: 327–328.
- Moulder JE, Fish BL, Cohen EF. 1998a. Angiotensin II receptor antagonists in treatment and prevention of radiation nephropathy. *Int J Radiat Biol* **73**: 415–421.
- Moulder JE, Fish BL, Cohen EF. 1998b. Brief pharmacological intervention in experimental radiation nephropathy. *Radiat Res* **150**: 535–541.
- Moulder JE, Fish BL, Cohen EP. 1998c. Radiation nephropathy

- is treatable with an angiotensin converting enzyme inhibitor or an angiotensin II type I (ATI) receptor antagonist. *Radiother Oncol* **46**: 307–315.
- Nair AGR, Gunasegaran R, Joshi BS. 1982. Chemical investigation of certain south Indian plants. *Indian J Chem* **21B**: 216.
- Nair CKK, Parida DK, Nomura T. 2001. Radioprotectors in radiotherapy. *J Radiat Res* **159**: 812–834.
- Narimanov AA. 1992. The reproductive capacity of male mice protected against the superlethal action of gamma radiation by the administration of *Archangelica officinalis* and *Ledum palustre*. *Radiobiologia* **32**: 271–275.
- Narimanov AA. 1993. The antiradiation effectiveness of a mixture of *Archangelica officinalis* and *Ledum palustre* extracts in the fractionated gamma irradiation of mice. *Radiobiologia* **33**: 280–284.
- Narimanov AA, Miakisheva SN, Kuznetsova SM. 1991. The radioprotective effect of extracts of *Archangelica officinalis* Hoffm. and *Ledum palustre* L. on mice. *Radiobiologia* **31**: 391–393.
- Narimanov AA, Popova OI, Muraveva DA. 1992. Changes in the sensitivity of mice to the action of gamma irradiation by *Viscum album* L. polysaccharide. *Radiobiologia* **32**: 868.
- Ni YB, Zhao B, Hou JW, Xin WJ. 1996. Preventive effect of *Ginkgo biloba* extract on apoptosis in rat cerebellar neuronal cells induced by hydroxyl radicals. *Neurosci Lett* **214**: 115–118.
- Nikitin VA, Chistiakow AA, Bugaewa V. 1989. Therapeutic endoscopy in combined therapy of gastroduodenal ulcers. *Khirurgia-mosk* **4**: 33–35.
- Nikulin AA, Iakusheva EN, Zakharova NM. 1992. A comparative pharmacological evaluation of Sea Buckthorn, rose and plantain oil in experimental eye burn. *Eksp Klin Farmakol* **55**: 64–66.
- Norr H, Wagner H. 1982. New constituents from *Ocimum sanctum*. *Planta Med* **58**: 574.
- Ohara M, Lu H, Shiraki K, et al. 2001. Radioprotective effects of miso (fermented soy bean paste) against radiation in B6C3F1 mice: increased small intestinal crypt survival, crypt lengths and prolongation of average time to death. *Hiroshima J Med Sci* **50**: 83–86.
- Ohnishi Y, Yasumizu R, Fan HX, et al. 1990. Effects of juzentaiho-toh (TJ-48), a traditional oriental medicine, on hematopoietic recovery from radiation injury in mice. *Exp Hematol* **18**: 18–22.
- Oyama Y, Chikahisa L, Ueha T, Kancemaru K, Noda K. 1996. *Ginkgo biloba* extract protects brain neurons against oxidative stress induced by hydrogen peroxide. *Brain Res* **712**: 349–352.
- Ozaki Y, Ono K. 2002. Antiinflammatory effect of glycyrrhizin topically applied to the cotton pellet by granuloma pouch method in rats. *Nat Med* **56**: 261–263.
- Pahadiya S, Sharma J. 2003. Alteration of lethal effects of gamma rays in Swiss albino mice by *Tinospora cordifolia*. *Phytother Res* **17**: 552–554.
- Pande S, Kumar M, Kumar A. 1998a. Evaluation of radiomodifying effects of root extract of *Panax ginseng*. *Phytother Res* **12**: 13–17.
- Pande S, Kumar M, Kumar A. 1998b. Radioprotective efficacy of *Aloe vera* leaf extract. *Pharm Bull* **36**: 227–232.
- Park KK, Chun KS, Lee JM, Lee SS, Surh J. 1998. Inhibitory effects of (6) gingerol, a major pungent principle of ginger, on phorbol ester-induced inflammation, epidermal ornithine decarboxylase activity and skin tumour promotion in ICR mice. *Cancer Lett* **129**: 139–144.
- Parshad R, Sanford KK, Price FM, et al. 1998. Protective action of plant polyphenols on radiation-induced chromatid breaks in cultured human cells. *Anticancer Res* **18**: 3263–3266.
- Penna SC, Medeiros MV, Aimbrie FSC, Faria-Neto HCC, Sertie JAA, Lopes-Martins RAB. 2003. Anti-inflammatory effect of the hydroalcoholic extract of *Zingiber officinale* rhizomes on rat paw and skin edema. *Phytomedicine* **10**: 381–385.
- Phadke SA, Kulkarni SD. 1989. Screening of *in vitro* antibacterial activity of *Terminalia chebula*, *Eclipta alba* and *Ocimum sanctum*. *Indian J Med Sci* **43**: 113.
- Phillips S, Ruggier R, Hutchinson SE. 1993. *Zingiber officinale* (ginger) – an antiemetic for day case surgery. *Anaesthesia* **48**: 715–717.
- Pincemail J, Dupuis M, Nasr C, et al. 1989. Superoxide anion scavenging effect and SOD activity of *Ginkgo biloba* extract. *Experientia* **45**: 708–712.
- Prasad KN. 1999. *Handbook of Radiobiology*. CRC Press: Boca Raton, FL, 344.
- Prem Kumar I, Goel HC. 2000. Iron chelation and related properties of *Podophyllum hexandrum*, a possible role in radioprotection. *Indian J Exp Biol* **38**: 1003–1006.
- Prem Kumar I, Samanta N, Goel HC. 2002. Modulation of chromatin organization by RH-3, a preparation of *Hippophae rhamnoides*, a possible role in radioprotection. *Mol Cell Biochem* **238**: 1–9.
- Qishen P, Baojing G, Kolman A. 1989. Radioprotective effect of extract from *Spirullina platensis* in mouse bone marrow cells studied by using the micronuclei test. *Toxicol Lett* **48**: 165–169.
- Rai MK. 1996. *In vitro* evaluation of medicinal plant extracts against *Pastalopsis mangifera* Hindustan Antibiot. **86**: 53–58.
- Ramadan LA, Roushdy HM, Abu Senna GM, Amin NE, El-Deshw OA. 2002. Radioprotective effect of silymarin against radiation-induced hepatotoxicity. *Pharmacol Res* **45**: 447–454.
- Ramnath N, Lo Russo, Simon M, Martino S. 1997. Phase II evaluation of cisplatin and WR-2721 for refractory metastatic breast cancer. *Am J Clin Oncol* **20**: 368–372.
- Rao AV, Uma Devi P, Kamath R. 2001. *In vivo* radioprotective effect of *Moringa oleifera* leaves. *Indian J Exp Biol* **39**: 858–863.
- Rao Mohandras KG, Rao MS, Karanth S, Rao GM. 1999. Effect of plant extract *Centella asiatica* (Linn.) on rat CNS – a functional and morphological correlation. *Indian J Pharmacol* **31**: 56.
- Raphael TJ, Kuttan G. 2003. Effect of naturally occurring triterpenoids glycyrrhizic acid, ursolic acid, oleanolic acid and nomilin on the immune system. *Phytomedicine* **10**: 483–489.
- Rastogi RP, Mehrotra BN. 1991. *Compendium of Indian Medicinal Plants*. Central Drug Research Institute (CDRI) and Publications and Information Directorate, Council of Scientific and Industrial Research (CSIR). New Delhi, India; vol. **3**: 420.
- Rastogi RP, Mehrotra BN. 1995. *Compendium of Indian Medicinal Plants*. Central Drug Research Institute (CDRI) and Publications and Information Directorate, Council of Scientific and Industrial Research (CSIR). New Delhi, India; vol. **4**: 930.
- Real A, Guenechea G, Bueren JA, Maganto G. 1992. Radioprotection mediated by the hemopoietic stimulation conferred by AM5: a protein-associated polysaccharide. *Int J Radiat Biol* **62**: 65–72.
- Reeve VE, Bosnic M, Rosinova E, Boehm-Wilcox C. 1993. A garlic extract protects from ultraviolet B (280–320 nm) induced suppression of contact hypersensitivity. *Photochem Photobiol* **58**: 813–817.
- Reiter RJ, Tan DX. 2002. Melatonin: an antioxidant in edible plants. *Ann NY Acad Sci* **957**: 341–344.
- Reyners H, Gianfelice DE, Reyners E, Poortmans F, Crametz A, Maisin JR. 1992. Brain atrophy after fetal exposure to very low doses of ionizing radiation. *Int J Radiat Biol* **62**: 619–629.
- Riehl T, Cohen S, Tessner T, Scholemann S, Stenson WS. 2000. Lipopolysaccharide is radioprotective in mouse intestine through a prostaglandin mediated mechanism. *Gastroenterology* **118**: 1106–1116.
- Rodrigues-Bigas M, Crus NI, Suarez A. 1988. Comparative evaluation of *Aloe vera* in the management of burn wounds in guinea pigs. *Plast Reconstr Surg* **81**: 386–389.
- Romay C, Armesto J, Ramirez D, Gonzalez R, Ledon N, Garcia I. 1998. Antioxidant and anti-inflammatory properties of C-phycoerythrin from blue-green algae. *Inflamm Res* **47**: 36–41.
- Rong Y, Geng Z, Lau BHS. 1996a. *Ginkgo biloba* modulates glutathione redox cycle in vascular endothelial cells. *Nutr Res* **16**: 1913–1923.
- Rong Y, Geng ZH, Lau BHS. 1996b. *Ginkgo biloba* attenuates oxidative stress in macrophages and endothelial cells. *Free Radic Biol Med* **20**: 121–127.
- Russel N, Pessell E, Staier C, Haynes A, Das Gupta E, Byrne J. 2000. Allogenic haemopoietic stem cell transplantation for

- multiple myeloma or plasma cell leukemia using fractionated total-body radiation and high dose of melphalan conditioning. *Acta Oncol* **39**: 837–841.
- Saini MR, Kumar S, Jagetia GC, Saini N. 1985. Whole body radiation-induced damage to the peripheral blood and protection by Liv 52. *Radiobiol Radiother* **26**: 487–493.
- Sajikumar S, Goel HC. 2003. *Podophyllum hexandrum* prevents radiation induced neuronal damage in post-natal rats exposed *in utero*. *Phytother Res* **17**: 761–766.
- Sakar MK, Tamer AU. 1990. Antimicrobial activity of different extracts from *Hypericum* species. *Fitoterapia* **61**: 464–466.
- Salin CA, Samanta N, Goel HC. 2001. Protection of mouse jejunum against lethal irradiation by *Podophyllum hexandrum*. *Phytomedicine* **8**: 413–422.
- Samanta N, Goel HC. 2002. Protection against radiation induced damage to spermatogenesis by *Podophyllum hexandrum*. *J Ethnopharmacol* **81**: 217–224.
- Samarth RM, Goyal PK, Kumar A. 2001. Modulatory effect of *Mentha piperita* (Linn.) on serum phosphatases activity in Swiss albino mice against gamma irradiation. *Indian J Exp Biol* **39**: 479.
- Samarth RM, Goyal PK, Kumar A. 2002b. Modulation of serum phosphatases activity in Swiss albino mice against gamma irradiation by *Mentha piperita* Linn. *Phytother Res* **16**: 586–589.
- Samarth RM, Kumar A. 2003. *Mentha piperita* (Linn.) leaf extract provides protection against radiation induced chromosomal damage in bone marrow of mice. *Indian J Exp Biol* **41**: 229–237.
- Samarth RM, Saini MR, Maharwal J, Dhaka A, Kumar A. 2002a. *Mentha piperita* (Linn.) leaf extract provides protection against radiation induced alterations in intestinal mucosal of Swiss albino mice. *Indian J Exp Biol* **40**: 1245.
- Samman MA, Bowen ID, Taiba K, Antonius J, Hannan MA. 1998. Mint prevents shamma-induced carcinogenesis in hamster cheek pouch. *Carcinogenesis* **19**: 1795.
- Sanjay K. 2000. India's government promotes traditional healing practices. *Lancet* **355**: 1252.
- Sarma DNK, Khosa RL, Chansouria JPN, Sahai M. 1995. Antiulcer activity of *Tinospora cordifolia* Miers and *Centella asiatica* Linn. extracts. *Phytother Res* **9**: 589–590.
- Sarma L, Tiku AB, Kesavan PC, Ogaki M. 1993. Evaluation of radioprotective action of mutant (E-25) form of *Chlorella vulgaris* in mice. *J Radiat Res* **34**: 277–284.
- Sarma SP, Aithal KS, Srinivasan KK. 1990. Antiinflammatory and wound healing activities of the crude alcoholic extracts and flavonoids of *Vitex leucoxylon*. *Fitoterapia* **61**: 263–265.
- Sastre JA, Millan J, Garcia De La Asuncion, Pla R, et al. 1998. A *Ginkgo biloba* extract (Egb761) prevents mitochondrial aging by protecting against oxidative stress. *Free Rad Biol Med* **24**: 298–304.
- Sato Y. 1990. Studies on chemical protectors against radiation. XXXI. Protection effects of *Aloe arborescens* on skin injury induced by x-irradiation. *Yakugaku Zasshi* **110**: 876–884.
- Sato Y. 1991. Studies on chemical protectors against radiation. XXXIII. Protective mechanisms of various compounds against skin injury induced by radiation. *Yakugaku Zasshi* **111**: 51–58.
- Satoh M, Miura N, Naganuma M, Matsuzaki N, Kawamura E, Imura N. 1989. Prevention of adverse effects of gamma ray irradiation after metallothionein induction by bismuth subnitrate in mice. *Eur J Cancer Clin Oncol* **25**: 1729–1731.
- Satyavati GV, Gupta KA, Tandon N (eds). 1987. *Medicinal Plants of India*, vol. II Indian Council of Medical Research: New Delhi, 354.
- Scarterzzimi P, Speroni E. 2000. Review of some plants of Indian traditional medicine with antioxidant activity. *J Ethnopharmacol* **71**: 23–43.
- Schapoval EE, Alice CB, Zuanazzi JA, Silva GAAB. 1988. Determinaao da acividade antimicrobiana dos extracto de *Syzygium cumini*. *Rev Port Farm* **38**: 55–57.
- Schuchter LM, Glick JH. 1993. The current status of WR-2721 (Amifostine): a chemotherapy and radiation therapy protector. *Biol Ther Cancer Updates* **3**: 1–10.
- Sestili P, Guidarelli A, Dacha M, Cantoni O. 1998. Quercetin prevents DNA single strand breakage and cytotoxicity caused by tert-butylhydroperoxide: Free radical scavenging versus iron chelating mechanism. *Free Radic Biol Med* **25**: 196–200.
- Sgaragli GP, Valoti M, Gorellil B, Fusi F, Palmi M, Mantovani P. 1993. Calcium antagonist and antioxidant properties of some hindered phenols. *Br J Pharmacol* **119**: 369–377.
- Sharma J, Sharma R. 2002. Radioprotection of Swiss albino mouse by *Centella asiatica* extract. *Phytother Res* **16**: 785–786.
- Sharma RK, Arora R, Prasad J, et al. 2004. Radioprotective efficacy of *Hippophae rhamnoides*. In *Proceedings of the Seminar on Sea Buckthorn: A Resource for Environment, Health and Economy*. Directorate of Life Sciences, Defence Research and Development Organization: New Delhi, 21–22.
- Sharma RK, Jain V. 2002. Tackling radioresistance of hypoxic cancer cells by metabolic modulation of bioenergetics: A P-31 MRS study on perfused Ehrlich ascites tumor cells. *Indian J Physiol Pharmacol* **46**: 51–60.
- Sharma RK, Maitra A, Jain V. 2000a. Effects of 6-aminonicotinamide and 2-deoxy-D-glucose combination on the bioenergetics of perfused Ehrlich ascites tumour cells as monitored by ³¹P MR spectroscopy. *Indian J Biochem Biophys* **37**: 307–312.
- Sharma RK, Singh S, Degaonkar M, Raghunathan P, Maitra A, Jain V. 2000b. Optimization of tumour radiotherapy. Part-VI-modification of tumour glucose metabolism for increasing the bioavailability of 2-deoxy-D-glucose in murine tumour model. *Strahlenther Oncol* **176**: 135–143.
- Shen Yu, Yang XY, Zhan QM, Guo R, Liu JW, Yang CZ. 1989. The use of Chinese herb medicine in experimental radiotherapy. *Int J Radiat Oncol Biol Phys* **16**: 347–352.
- Shetty S, Udupa SL, Udupa AL, Ullas K, Bhaskar. 1999. Effect of *Ocimum sanctum* on wound healing. *Indian J Pharmacol* **31**: 78–81.
- Shetty TK, Satav JG, Nair CKK. 2002. Protection of DNA and microsomal membranes *in vitro* by *Glycyrrhiza glabra* L. against gamma radiation. *Phytother Res* **16**: 576–578.
- Shimoi K, Masuda S, Furugori M, Esaki S, Kinai N. 1994. Radioprotective effect of antioxidative plant flavonoids in γ -ray irradiated mice. *Carcinogenesis* **15**: 2669–2672.
- Shimoi K, Masuda S, Shen B, Furugori M, Kinai N. 1996. Radioprotective effect of antioxidative plant flavonoids in mice. *Mutat Res* **350**: 153–161.
- Shobi V, Goel HC. 2001. Protection against radiation-induced conditioned taste aversion by *Centella asiatica*. *Physiol Behavior* **73**: 19–23.
- Shukla A, Rasik AM, Jain GK. 1999. *In vitro* and *in vivo* wound healing activity of asiaticoside isolated from *Centella asiatica*. *J Ethnopharmacol* **65**: 1–11.
- Sidhu GS, Mani H, Gaddipate JP, et al. 1999. Curcumin enhances wound healing in streptozotocin induced diabetic rats and genetically diabetic mice. *Wound Repair Regen* **7**: 362–374.
- Singh J, Shah NC. 1994. *Podophyllum*: A Review. *Curr Res Med Arom Plants* **16**: 53–83.
- Singh RP, Banerjee S, Rao AR. 2000. Effect of *Aegle marmelos* on biotransformation enzyme systems and protection against free-radical-mediated damage in mice. *J Pharm Pharmacol* **52**: 991–1000.
- Singh S, Agarwal SS. 1991. Antiasthmatic and anti-inflammatory activity of *Ocimum sanctum*. *Int J Pharmacog* **29**: 306.
- Singh S, Majumdar DK, Rehan HM. 1996. Evaluation of antiinflammatory potential of fixed oil of *Ocimum sanctum* (holy basil) and its possible mechanism of action. *J Ethnopharmacol* **54**: 19.
- Singh SP, Abraham SK, Kesavan PC. 1995a. *In vivo* radioprotection with garlic extract. *Mutat Res* **345**: 147–153.
- Singh SP, Ashu B, Tiku AB, Kesavan PC. 1995b. Post-exposure radioprotection by *Chlorella vulgaris*. *Indian J Exp Biol* **33**: 612–615.
- Smith PF, MacLennan K, Darlington CL. 1996. The neuroprotective properties of the *Ginkgo biloba* leaf: A review of the possible relationship of platelet activating factor (PAF). *J Ethnopharmacol* **50**: 131–139.
- Smyshliaeva AV, Kudriashov lu B. 1992. The modification of a radiation lesion in animals with an aqueous extract of *Hypericum perforatum* L. *Nauchnye Doke Vyss Shkoly Biol Nauki* **4**: 9–13.
- Song JY, Han SK, Bac KG, et al. 2003. Radioprotective effects of ginsan, an immunomodulator. *Radiat Res* **159**: 768–774.

- Song JY, Han SK, Son EH, Pyo SN, Yun YS, Yi SY. 2002. Induction of secretory and tumoricidal activities in peritoneal macrophages by ginsan. *Int Immunopharmacol* **2**: 857–865.
- Sonoda Y, Kasahara T, Mukaida N, Shimizu N, Tomoda M, Takeda T. 1998. Stimulation of interleukin-8 production by acidic polysaccharides from the root of *Panax ginseng*. *Immunopharmacology* **38**: 287–294.
- Spencer CM, Goa KL. 1995. Amifostine: A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic as a radioprotector and cytotoxic chemoprotector. *Drugs* **50**: 1001–1031.
- Srivastava R, Shukla YN, Sushil Kumar. 1997. Chemistry and pharmacology of *Centella asiatica*: a review. *J Med Arom Plant Sci* **19**: 1049–1056.
- Stanley P, Mainzen P, Menon VP, Gunasekaran G. 1999a. Hypolipidaemic action of *Tinospora cordifolia* roots in alloxan-induced diabetic rats. *J Ethnopharmacol* **70**: 9–15.
- Stanley P, Mainzen P, Menon VP. 1999b. Antioxidant activity of *Tinospora cordifolia* roots in experimental diabetes. *J Ethnopharmacol* **65**: 277–281.
- Stelzer KJ, Koh WJ, Kurtz H, Greer BE, Griffin TW. 1994. Caffeine consumption is associated with decreased severe late toxicity after radiation to the pelvis. *Int J Radiat Oncol Biol Phys* **30**: 411–417.
- Stevens GR, Morris JE, Anderson E. 2000. Hemochromatosis heterozygotes may constitute a radiation-sensitive subpopulation. *Radiat Res* **153**: 844–847.
- Subramanian M, Chintalwar GJ, Chattopadhyay S. 2003. Radioprotective property of polysaccharide in *Tinospora cordifolia*. *Indian J Biochem Biophys* **40**: 22–26.
- Suguna L, Singh S, Sivakumar P, Padmavathi S, Gowri C. 2002. Influence of *Terminalia chebula* on dermal wound healing in rats. *Phytother Res* **16**: 227–231.
- Sweeney TR. 1979. *A Survey of Compounds from the Antiradiation Drug Development Program of the US Army Medical Research and Development Command*. Walter Reed Army Institute of Research: Washington DC, USA.
- Takeda A, Katoh N, Yonezawa M. 1982. Restoration of radiation injury by ginseng III. Radioprotective effect of thermostable fraction of ginseng extract on mice, rats and guinea pigs. *J Radiat Res* **23**: 150–167.
- Tan PV, Njimi CK, Ayufor JF. 1997. Screening of some African medicinal plant ulcerogenic activity-Part I. *Phytother Res* **11**: 45–47.
- Tannehill SP, Mehta MP. 1996. Amifostine and radiation therapy: past, present, and future. *Semin Oncol* **23**: 69–77.
- Taraphdar AK, Shaw BP, Bhattacharya RK, Mukherjee PK. 2002. Role of sharpunka (*Tephrosia purpurea*) in haemopoietic injury. *Antiseptic* **99**: 302–304.
- Thali S, Thatte U, Dahanukar SA. 1998. The potential of *Boerhaavia diffusa* in radiation induced haemopoietic injury. *Amala Res Bull* **18**: 20–22.
- Thatte UM, Chabria S, Karandikar SM, Dhanukar S. 1988. Immunotherapeutic modification by Indian medicinal plants. *Indian Drugs* **25**: 95–97.
- Theresiamma KC, George J, Kuttan R. 1996. Protective effect of curcumin, ellagic acid and bixin on radiation induced toxicity. *Indian J Exp Biol* **34**: 845–847.
- Thomson M, Al-Qattan KK, Al-Sarvan SM, Alnaqueeb MA, Khan I, Ali M. 2002. The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostaglandins Leukot Essent Fatty Acids* **67**: 475–478.
- Udupa SI, Udupa AL, Kulkarni DR. 1994. Anti-inflammatory and wound healing properties of *Aloe vera*. *Fitoterapia* **65**: 141–145.
- Uma Devi P. 2001. Radioprotective, anticarcinogenic and antioxidant properties of the Indian holy basil, *Ocimum sanctum*. *Indian J Exp Biol* **39**: 185–190.
- Uma Devi P, Bisht KS, Vinitha M. 1998. A comparative study of radioprotection by *Ocimum* flavonoids and synthetic aminothiol protectors in mouse. *Br J Radiol* **71**: 782–784.
- Uma Devi P, Ganasoundari A. 1995. Radioprotective effect of leaf extract of Indian medicinal plant *Ocimum sanctum*. *Indian J Exp Biol* **33**: 205–209.
- Uma Devi P, Ganasoundari A, Rao BSS, Srinivasan KK. 1999. *In vivo* radioprotection by *Ocimum sanctum* flavonoids: survival of mice. *Radiat Res* **151**: 74–78.
- Uma Devi P, Ganasoundari A, Vrinda B, Srinivasan KK, Unnikrishnan MK. 2000. Radiation protection by *Ocimum sanctum* flavonoids orientin and vicenin – mechanisms of action. *Radiat Res* **154**: 455–460.
- Uma Devi P, Kamath R, Rao BSS. 2000. Radioprotective effect of *Phyllanthus niruri* on mouse chromosome. *Curr Sci* **78**: 1245–1247.
- Upasani CD, Balaraman. 2003. Protective effect of *Spirulina* on lead-induced deleterious changes in the lipid peroxidation and endogenous antioxidants in rats. *Phytother Res* **17**: 330–334.
- Van Buul PPW, Van Duyn-Goedhart A, Shankaranarayanan K. 1999. *In vivo* and *in vitro* radioprotective effects of prostaglandin E, analogue misoprostol in DNA repair-proficient and – deficient rodent cell systems. *Radiat Res* **152**: 398–403.
- Van der Meeren A, Mouthon MA, Gaugler MH, Vandamme M, Gourmelon P. 2002. Administration of recombinant human IL11 after supralethal radiation exposure promotes survival in mice: interactive effect with thrombopoietin. *Radiat Res* **157**: 642–649.
- Vazquez B, Avila G, Segura D, Escalante B. 1996. Anti-inflammatory activity of extracts from *Aloe vera* gel. *J Ethnopharmacol* **55**: 69–75.
- Verma RK, Jain M, Saini PP, Sisodia SR, Bhatia AL. 2003. Effect of *Amaranthus paniculatus* leaf extract on radiation-induced biochemical changes in Swiss albino mice brain. *J Med Arom Plant Sci* **25**: 362–368.
- Vijaylaxmi, Herman TS, Meltz. 1996. Melatonin and radioprotection from genetic damage: *in vivo/in vitro* studies with human volunteers. *Mutat Res* **371**: 221–228.
- Vokovic-Gacic B, Simic D. 1993. Identification of natural antimutagens with modulating effects on DNA repair. *Basic Life Sci* **61**: 269.
- Vose JM, Armitage JO. 1995. Clinical applications of hematopoietic growth factors. *J Clin Oncol* **13**: 1023–1035.
- Vrinda B, Uma Devi P. 2001. Radioprotection of human lymphocyte chromosomes *in vitro* by orientin and vicenin. *Mutat Res* **498**: 39–46.
- Wang CM, Ohta S, Shimoda. 1992. Studies on chemical protectors against radiation. XXXV. Effects of radioprotective Chinese traditional medicines on radiation-induced lipid peroxidation *in vivo* and *in vitro*. *Chem Pharm Bull* **40**: 493–498.
- Wang LY 1996. A histopathologic study on recovering effect of Kangdu shengxue decoction on spleen and thymus of mice with radiation injury. *Chung Kuo Chung Hsi I Chieh Ho Tsa Chih* **16**: 489–491.
- Wasserman TH. 1994. Radiotherapeutic studies with amifostine (Ethyol). *Semin Oncol* **21**: 21–25.
- Wazir V, Maurya R, Kapil RS. 1995. Cordioside, aclerodane furano diterpene glucoside from *Tinospora cordifolia*. *Phytochemistry* **38**: 447–449.
- Weiss JF. 1997. Pharmacologic approaches to protection against radiation-induced lethality and other damage. *Environ Health Perspect* **105** (Suppl 6): 1473–1478.
- Weiss JF, Kumar KS, Walden TL, Neta R, Landauer MR, Clark EP. 1990. Advances in radioprotection through the use of combined agent regimens. *Int J Radiat Biol* **57**: 709.
- Weiss JF, Landauer MR. 2003. Protection against ionizing radiation by antioxidant nutrients and phytochemicals. *Toxicology* **189**: 1–20.
- Weiss JF, Simic MG. 1998. Introduction: perspectives in radiation protection. *Pharmacol Ther* **39**: 1–2.
- Wheeler TG, Hardy KA. 1985. Retrograde amnesia produced by electron beam exposure: causal parameters and duration of memory loss. *Radiat Res* **101**: 74–80.
- William MS, Mary Burk. Loprinzi CL, et al. 1996. Phase III double blind evaluation of an *Aloe vera* gel as a prophylactic agent for radiation induced skin toxicity. *Int J Radiat Oncol Biol Phys* **36**: 345–349.
- Wong SK, Tsui SK, Kwan SY, Su XL, Lin RC. 2000. Identification of characterization of *Podophyllum emodi* by API-LC/MS/MS. *J Mass Spectrom* **35**: 1246–1251.
- World Health Organization Monographs on Selected Medicinal Plants. 1999. Vol. 1. WHO, Geneva.
- Xiao M, Yang Z, Jiu M, You J, Xiao R. 1992. The antigastrolcerative action of β -cytosterol-D-glucoside and its aglycan in rats. *Hua-His-I-Ko-ta-Hsueh-Hsueh-Pao* **23**: 98–101.
- Yang JJ, Lin CC, Hsu HY. 1997. The possible use of Peh-Hue-Juwa-Chi-cao as an antitumor agent and radioprotector after therapeutic irradiation. *Phytother Res* **11**: 6–10.

- Yang Y, Kinoshita K, Koyama K, Takahashi K, Kondo S, Watanabe K. 2002. Structure-antiemetic-activity of some diarylheptanoids and their analogues. *Phytomedicine* **9**: 146–152.
- Yonezawa M, Katoh N, Takeda A. 1985. Restoration of radiation injury by ginseng. IV. Stimulation of recoveries in CFU and megakaryocyte counts related to the prevention of occult blood appearance in x-irradiated mice. *J Radiat Res* **26**: 436–442.
- Yonezawa M, Katoh N, Takeda A. 1989. Radiation protection by shigoka extract on split-dose in mice. *J Radiat Res* **30**: 247–254.
- Yonezawa M, Takeda A, Katoh N. 1981. Restoration of radiation injury by ginseng. II. Some properties of the radioprotective substances. *J Radiat Res* **22**: 336–343.
- Yoshioka H. 1997. Beta ray-induced scission of DNA in tritiated water and protection by a green tea percolate and epigallocatechin gallate. *Biosci Biotechnol Biochem* **61**: 1560–1563.
- Zafar R, Naaz SF. 2002. *Centella asiatica* Linn. A review. *Hamdard Medicus* **45**: 55–72.
- Zhang C, Zheng S, Zhang Y, Luo C, Guo C. 1997. The protective effects of polysaccharide and C-phycocyanin from *Spirulina platensis* on acute radiation injury in mice. *Acta Nutr Sinica* **18**: 327–331.
- Zhang JS, Sigdestad CP, Gemmell MA, Grdina DJ. 1987. Modification of radiation response in mice by fractionated extracts of *Panax ginseng*. *Radiat Res* **112**: 156–163.
- Zhang MS. 1987. A control trial of flavonoids *Hippophae rhamnoides* L. in treating ischemic heart disease. *Chung-Hua-Hsin-Hsueh-Kuan-Ping-Tsa-Chih* **15**: 97–99.
- Zhang SW, Zhang. 1990. Radiation sensitizing and protective action of *Ligusticum wallichii* Franch. *Chung His I Chieh Ho Tsa Chih* **10**: 697–698.
- Zozulia IS, Iurchenko AV. 2000. The adaptive potentials of those who worked in the cleanup of the aftermath of the accident at the Chernobyl Atomic Electric Power Station under the influence of different treatment methods. *Lik Sprava* **4**: 18–21.
- Zsebo KM, Smith KA, Hartley CA, Greenblatt M, Cooke WR, McNiece IK. 1992. Radioprotection of mice by recombinant rat stem cell factor. *Proc Natl Acad Sci USA* **89**: 9464–9468.