

# Randomized multicentre pilot study of sacubitril/valsartan versus irbesartan in patients with chronic kidney disease: United Kingdom Heart and Renal Protection (HARP)- III—rationale, trial design and baseline data

## UK HARP-III Collaborative Group

Correspondence and offprint requests to: Richard Haynes, Medical Research Council Population Health Research Unit, Nuffield Department of Population Health, Richard Doll Building, Old Road Campus, Roosevelt Drive, Oxford OX3 7LF, UK; E-mail: harp3@ndph.ox.ac.uk

### ABSTRACT

**Background.** Patients with chronic kidney disease (CKD) are at risk of progression to end-stage renal disease and cardiovascular disease. Data from other populations and animal experiments suggest that neprilysin inhibition (which augments the natriuretic peptide system) may reduce these risks, but clinical trials among patients with CKD are required to test this hypothesis.

**Methods.** UK Heart and Renal Protection III (HARP-III) is a multicentre, double-blind, randomized controlled trial comparing sacubitril/valsartan 97/103 mg two times daily (an angiotensin receptor–neprilysin inhibitor) with irbesartan 300 mg one time daily among 414 patients with CKD. Patients  $\geq 18$  years of age with an estimated glomerular filtration rate (eGFR) of  $\geq 45$  but  $< 60$  mL/min/1.73 m<sup>2</sup> and urine albumin:creatinine ratio (uACR)  $> 20$  mg/mmol or eGFR  $\geq 20$  but  $< 45$  mL/min/1.73 m<sup>2</sup> (regardless of uACR) were invited to be screened. Following a 4- to 7-week pre-randomization single-blind placebo run-in phase (during which any current renin–angiotensin system inhibitors were stopped), willing and eligible participants were randomly assigned either sacubitril/valsartan or irbesartan and followed-up for 12 months. The primary aim was to compare the effects of sacubitril/valsartan and irbesartan on measured GFR after 12 months of therapy. Important secondary outcomes include effects on albuminuria, change in eGFR over time and the safety and tolerability of sacubitril/valsartan in CKD.

**Results.** Between November 2014 and January 2016, 620 patients attended a screening visit and 566 (91%) entered the pre-randomization run-in phase. Of these, 414 (73%) participants were randomized (mean age 63 years; 72% male). The mean eGFR was 34.0 mL/min/1.73 m<sup>2</sup> and the median uACR was 58.5 mg/mmol.

**Conclusions.** UK HARP-III will provide important information on the short-term effects of sacubitril/valsartan on renal function, tolerability and safety among patients with CKD.

**Keywords:** cardiovascular disease, chronic kidney disease, neprilysin, progression

### INTRODUCTION

Chronic kidney disease (CKD) affects between 2 and 17% of the general population (depending on the country) [1, 2] and is associated with increased risks of progression to end-stage renal disease (ESRD) and morbidity and mortality from cardiovascular disease (CVD) [3, 4]. Renin–angiotensin system (RAS) inhibitors [angiotensin-converting enzyme inhibitors (ACEis) and angiotensin receptor blockers (ARBs)] have been shown to reduce the risk of ESRD in patients with proteinuric CKD [5–8], but despite such treatments, patients remain at significant risk of progression to ESRD and CVD.

The natriuretic peptide (NP) system is a neurohormonal system that has a variety of potentially beneficial functions, including natriuresis, diuresis, vasodilatation and counterregulation of RAS [9, 10]. The NP system can be augmented by inhibiting the main enzyme responsible for degrading NPs, namely neprilysin [or neutral endopeptidase (NEP)] [10]. NEP is a membrane-bound zinc-containing metalloproteinase [11] that also degrades other peptides, including angiotensin II, bradykinin, endothelin and substance P [12]. However, isolated NEP inhibition (NEPi) leads to reflex RAS activation, and inhibits angiotensin II breakdown (counteracting any potentially beneficial effects) and therefore NEPi must be combined with RAS inhibition.

As NEPi and ACEi both inhibit bradykinin degradation, their combination is associated with substantially elevated bradykinin

levels that cause unacceptable rates of angioedema [13]. ARBs do not inhibit bradykinin degradation and can be safely combined with NEPi [creating a new class of drugs called angiotensin receptor–neprilysin inhibitors (ARNis)]. Sacubitril/valsartan (previously known as LCZ696) is the first drug in this new class, combining valsartan with sacubitril [(AHU377) a prodrug that is metabolized via esterases to the active NEPi sacubitrilat (LBQ657)]. Sacubitril/valsartan 97/103 mg provides equivalent plasma concentrations of valsartan as oral valsartan 160 mg [14].

In a 5/6 nephrectomy model, treatment with combined NEP/RAS inhibition was associated with greater reductions in proteinuria and glomerulosclerosis compared with RAS inhibition alone [15, 16]. Micropuncture studies also demonstrated NEPi led to greater reductions in capillary glomerular pressure [15]. Among patients with heart failure, trials comparing sacubitril/valsartan with either ACEi or ARB have suggested that the estimated glomerular filtration rate (eGFR) of patients allocated sacubitril/valsartan declined less than those assigned ACEi or ARB [17, 18]. Sacubitril/valsartan also reduced blood pressure more than equivalent doses of valsartan in trials among patients with elevated blood pressure [19]. Trials in heart failure populations suggest NEPi might increase albuminuria [18, 20], but this effect was not observed in patients with hypertension [19] and baseline albuminuria was very low in all these trials. Overall, these data raise the hypothesis that treatment with an ARNi may be superior to either ACEi or ARB alone in slowing the progression of CKD.

The United Kingdom (UK) Heart and Renal Protection III (HARP-III) trial (ISRCTN11958993) was designed to provide information on the short-term efficacy (in terms of effect on renal function), tolerability and safety of sacubitril/valsartan among patients with CKD. The trial will also assess the effects of sacubitril/valsartan on albuminuria, blood pressure and biomarkers of kidney and cardiac damage.

## MATERIALS AND METHODS

### Study design

UK HARP-III is a double-blind, multicentre, randomized controlled trial comparing sacubitril/valsartan 97/103 mg two times daily versus irbesartan 300 mg one time daily among at least 400 participants  $\geq 18$  years of age with stages 3 and 4 CKD. Irbesartan 300 mg was selected as the comparator, as it has been shown to reduce the risk of ESRD among patients with diabetic kidney disease and is licensed for the treatment of proteinuric CKD [6, 21]. Participants were randomly allocated to receive sacubitril/valsartan or irbesartan and will be followed up for 1 year (Figure 1). The primary aim of UK HARP-III is to assess the effect of sacubitril/valsartan 97/103 mg two times daily versus irbesartan 300 mg one time daily on measured glomerular filtration rate (mGFR) at 12 months. Important secondary outcomes include the effect on urine albumin:creatinine ratio (uACR) and eGFR. All the secondary and tertiary assessments are shown in Figure 2 and further details are available in the data analysis plan (see [Supplementary data](#)). A summary of substantial amendments to the protocol is provided in the [Supplementary data](#).

### Eligibility

To fulfil the inclusion criteria, patients need to be  $\geq 18$  years of age and have either an eGFR  $\geq 45$  but  $< 60$  mL/min/1.73 m<sup>2</sup> with a uACR  $> 20$  mg/mmol or eGFR  $\geq 20$  but  $< 45$  mL/min/1.73 m<sup>2</sup> (regardless of uACR). The exclusion criteria were designed to identify patients for whom the safety of sacubitril/valsartan or irbesartan may have been a concern. The full eligibility criteria are shown in Figure 3.




FIGURE 1: UK HARP-III trial design.

<b>Primary:</b> Measured GFR (adjusted for body-surface area) at 12 months
<b>Secondary:</b>
Urine albumin:creatinine ratio at 3, 6 and 12 months
eGFR at 3, 6 and 12 months
Metabolites of sacubitril/valsartan at 3 months
<b>Tertiary:</b>
Systolic and diastolic blood pressure at 1, 3, 6, 9 and 12 months
Markers of kidney damage (kidney injury molecule 1 and neutrophil gelatinase-associated lipocalin) at 6 and 12 months
Markers of renal tubular function ( $\beta_2$ -microglobulin and retinol binding protein) at 6 and 12 months
Cardiac biomarkers (troponin I and N-terminal prohormone brain natriuretic peptide [NT-proBNP]) at 6 and 12 months
Urine cyclic guanosine monophosphate at 6 and 12 months
Rate of change of eGFR calculated from creatinine values at 0, 1, 3, 6, 9 and 12 months (overall for 0-3 and 3-12 months and separately)

FIGURE 2: UK HARP-III trial outcomes.

## Study enrolment and randomization

**Identification and invitation.** After relevant ethics [Nottingham Research Ethics Committee 2 (13/EM/0434)] and regulatory approvals had been obtained, sites were established in UK renal units. Site staff identified potentially eligible patients from hospital electronic databases, mailed these individuals an invitation letter and a copy of the patient information sheet and called them ~1 week later to discuss the trial in more detail, answer any questions they might have and to see whether they were interested in participating. Those individuals interested in participating were invited to attend a screening visit.

**Screening.** At the screening visit, eligibility was assessed and written informed consent was obtained from eligible individuals. All data were recorded directly into a bespoke Internet-based electronic case report form system. Relevant details of their medical history (including primary renal diagnosis, presence of diabetes mellitus and prior CVD) were recorded by trained research nurses and their height, weight and blood pressure were measured. Blood pressure was measured and recorded three times using an Omron M6 automated digital sphygmomanometer after sitting for at least 5 minutes. Willing and eligible patients entered the pre-randomization run-in phase. Samples of blood and urine were sent to the local hospital laboratory for confirmation of eligibility. If the results were considered inaccurate (e.g. haemolysed sample) by the local study staff the samples could be repeated once, but if the results did not confirm eligibility the participant was withdrawn from the run-in phase.

**Pre-randomization run-in.** The aims of the pre-randomization run-in phase were (i) to 'wash out' any ACEi prior to introduction of NEPi, (ii) to allow a comparison of the acute effects of the study treatments on GFR and (iii) to reduce the rate of post-randomization discontinuation of study treatment and to produce a consequent improvement in the trial's statistical sensitivity [22]. Following the screening visit, any current ACEi and/or ARB that the participant was taking was stopped and the participant entered the 4- to 7-week single-blind pre-randomization run-in phase, during which they were

asked to take one placebo sacubitril/valsartan tablet and one placebo irbesartan capsule once daily. If elevated blood pressure became a concern during the run-in phase, local investigators were advised to titrate up or start additional anti-hypertensive medications, but to avoid an ACEi, ARB or direct renin inhibitor (DRI). The choice of additional anti-hypertensive therapy remained at the discretion of the responsible clinician. Participants could withdraw from the trial for any reason during this run-in phase. Participants who did not withdraw returned 4–7 weeks later and had their GFR measured and attended a randomization visit. GFR was measured using a standard  $^{51}\text{Cr}$ -EDTA technique, although if this was not available at the site, other methods ( $^{99\text{m}}\text{Tc}$ -DTPA or iohexol) could be used with the agreement of the coordinating centre. In willing participants, a 24-hour collection of urine for albumin and sodium quantification was also obtained.

**Randomization visit.** Participants were not eligible for randomization if the mean of their second and third measurements of systolic blood pressure was <110 mmHg (or <130 mmHg with symptoms of hypotension) or if they reported an adverse event they believed to be related to their run-in treatment. Participants who remained willing and eligible were then randomly allocated in a 1:1 ratio to receive either sacubitril/valsartan or irbesartan. Participants were randomized by an Internet-based system using a minimization algorithm to ensure balance of important predictors of renal progression, including age, sex, systolic blood pressure, eGFR, uACR and the presence or absence of diabetes mellitus.

At the randomization visit, run-in treatment was collected and willing and eligible participants were issued two bottles of study treatments: one containing sacubitril/valsartan 97/103 mg or placebo tablets and the other containing irbesartan 150 mg or placebo capsules (therefore a double-dummy technique to protect blinding). Participants were initially instructed to take one tablet and one capsule daily in the morning (i.e. either sacubitril/valsartan 97/103 mg plus placebo irbesartan or placebo sacubitril/valsartan plus irbesartan 150 mg). Blood and urine samples were collected for the local analysis of creatinine, electrolytes, liver function tests and uACR and others were prepared for central analysis (Table 1).

## Post-randomization follow-up

Randomization is now complete and all participants are in follow-up. In order to check potassium and renal function after starting study treatment, participants attend their study clinic or local primary care physician at 2 weeks after randomization for a blood sample. If these results are satisfactory, study treatments are increased to either sacubitril/valsartan 97/103 mg two times daily plus two capsules of placebo irbesartan one time daily or one tablet of placebo sacubitril/valsartan two times daily plus irbesartan 300 mg one time daily.

**Follow-up assessments.** Study follow-up visits are scheduled at 1, 3, 6, 9 and 12 months after randomization. At all visits, study staff systematically seek the information on all serious adverse events, any non-serious adverse events considered by participants to be related to study treatment; and on symptoms

<p><b>Inclusion criteria</b></p> <p>Men or women aged <math>\geq 18</math> years (at screening)</p> <p>Established CKD:</p> <ul style="list-style-type: none"> <li>○ eGFR <math>\geq 20 &lt; 45</math> mL/min/1.73m<sup>2</sup>; or</li> <li>○ eGFR <math>\geq 45 &lt; 60</math> mL/min/1.73m<sup>2</sup> and uACR <math>&gt; 20</math> mg/mmol</li> </ul> <p><b>Exclusion criteria</b></p> <p>ARB therapy contraindicated e.g. bilateral renal artery stenosis</p> <p>Known intolerance of ARB</p> <p>Current treatment with aliskiren</p> <p>Mean systolic blood pressure <math>&gt; 180</math> mmHg at screening visit (or investigator unwilling to withdraw ACEi or ARB for another reason)</p> <p>Serum potassium <math>&gt; 5.5</math> mmol/L (this was increased from <math>&gt; 5.2</math> mmol/L in February 2015)</p> <p>Patients that currently have nephrotic syndrome (i.e. urine protein:creatinine ratio <math>&gt; 350</math> mg/mmol [or uACR <math>&gt; 300</math> mg/mmol] AND serum albumin <math>&lt; 30</math> g/L) or are currently receiving immunosuppression to treat the nephrotic syndrome</p> <p>Functioning renal transplant</p> <p>Acute coronary syndrome, stroke or transient ischaemic attack in 3 months prior to Screening</p> <p>Known chronic liver disease or ALT / AST <math>&gt; 2x</math> ULN at Screening visit</p> <p>History of angioedema (drug-related or otherwise)</p> <p>Use of unlicensed investigational medicinal product in previous month</p> <p>Pregnancy, lactating women, or women with child-bearing potential (refusing a reliable method of contraception)</p> <p>Medical history that might limit the patient's ability to take study treatments for the duration of the study (e.g. severe respiratory disease, or recent history of alcohol or substance misuse or history of cancer or evidence of spread in last 5 years other than non-melanoma skin cancer)</p>
--

**FIGURE 3:** Inclusion and exclusion criteria. ALT, alanine aminotransferase; AST, aspartate aminotransferase; ULN, upper limit of normal.

of hepatitis. Compliance with study treatment is assessed and participants unable to tolerate the maximum dose of study treatments are encouraged to continue on the lower dose of study drug (i.e. sacubitril/valsartan 97/103 mg or irbesartan 150 mg daily) for the remainder of the trial. If relevant, a reason for discontinuation or dose reduction is recorded. Participants prescribed contraindicated medications (ACEi, ARB or DRI) have their randomized treatment stopped. Weight and blood pressure are measured (three times after sitting for at least 5 minutes) at all visits. In both treatment groups, blood pressure is to be controlled according to the Kidney Disease: Improving Global Outcomes guidelines [23], with the initiation and choice of additional anti-hypertensive treatment being at the discretion of the responsible clinician. Within the 2 weeks before their 12-month visit participants have their second GFR measurement (using the same method as at baseline). Copies of results of both measurements of GFR are sent to the coordinating centre so the results entered by site staff can be verified by clinical study staff blind to the treatment allocation.

**Biological samples and safety monitoring.** At each follow-up visit, blood and urine samples are sent to the local hospital laboratory for creatinine, electrolytes, liver function tests (bilirubin, alanine or aspartate transaminase and alkaline phosphatase) and uACR. In addition, at the 3-, 6- and 12-month visits, samples are also taken for central analysis. EDTA samples are centrifuged and the plasma aliquoted into Cryovials, which are stored locally (with Cryovials of urine) at or below  $-20^{\circ}\text{C}$  prior to transfer to the central laboratory in Oxford, UK, where they are stored at  $-80^{\circ}\text{C}$ . The main plasma analytes measured at the central laboratory are creatinine, cardiac and inflammatory biomarkers and the urine analytes include albumin and markers of tubular damage and function [including kidney injury molecule 1, neutrophil gelatinase-associated lipocalin,  $\beta 2$ -microglobulin and retinol binding protein; Table 1]. Participants are asked not to take their morning dose of study treatment on the day of their 3-month visit (at this visit only) and the date and time of the last dose is recorded, as these samples are to be used for pharmacokinetic analyses.

**Table 1. Planned central laboratory blood and urine analyses**

Analyte	Time point			
	Randomization	3 months	6 months	12 months
<b>EDTA plasma samples</b>				
Creatinine	■	■	■	■
Albumin	■		■	■
Troponin-I	■		■	■
NT-proBNP	■		■	■
CRP	■		■	■
IL-6	■		■	■
<b>Urine samples</b>				
Albumin:creatinine ratio	■	■	■	■
KIM-1	■		■	■
NGAL	■		■	■
cGMP	■		■	■
β-2-microglobulin	■		■	■
Retinol binding protein	■		■	■

NT-proBNP, N-terminal prohormone brain natriuretic peptide; CRP, C-reactive protein; IL-6, interleukin 6; KIM-1, kidney injury molecule 1; NGAL, neutrophil gelatinase-associated lipocalin; cGMP, cyclic guanosine monophosphate.

The results of local samples are entered into the trial database once available and reviewed daily by a trained clinician at the coordinating centre. If the potassium is  $>5.5$  mmol/L, alanine or aspartate transaminase  $>2\times$  the upper limit of normal or if the eGFR has fallen  $>25\%$  from the previous value, then the trial protocol provides advice on further tests and study treatment (see [Supplementary data](#)).

### Monitoring

Prior to starting recruitment, study staff received training in the study procedures and the web-based data collection system at the coordinating centre. Recruitment rates, adherence to trial procedures and completeness of follow-up data are monitored closely by staff at the coordinating centre. All sites have at least one on-site monitoring visit, with further visits as indicated by the results of central monitoring of the data. An independent data monitoring committee (see [Supplementary data](#)) regularly reviews unblinded interim analyses of all relevant data.

### Statistical considerations

**Sample size.** The chief aim of this study is to compare mGFR between the two treatment groups at the final follow-up visit. Analysis of covariance (ANCOVA) compares mean follow-up mGFR between treatment groups after adjustment for baseline mGFR [24]. Assuming a between-person standard deviation (SD) in mGFR of  $15$  mL/min/ $1.73$  m<sup>2</sup> and a correlation between an individual's baseline and follow-up mGFR of  $0.8$ , randomization of  $400$  participants will provide at least  $80\%$  power (at  $2 P = 0.05$ ) to detect a difference in mGFR at the final follow-up (adjusted for baseline values) of  $3$  mL/min/ $1.73$  m<sup>2</sup> (the chosen minimum clinically meaningful difference), even if  $15\%$  of participants discontinue allocated study treatment [20].

**Statistical analysis.** All analyses will involve comparing outcomes during the scheduled treatment period among all those participants allocated at randomization to receive sacubitril/

valsartan  $97/103$  mg two times daily versus all those allocated to receive irbesartan  $300$  mg one time daily [i.e. intention-to-treat (ITT) analyses] [25, 26]. Comparisons of continuous outcomes (including the primary outcome) between the allocated treatment arms will be performed using ANCOVA adjusted for each patient's value at baseline [27]. If continuous outcomes are not normally distributed, then appropriate transformations (e.g. log transformation) will be made. Multiple imputation techniques will be used to account for any missing data in the primary and secondary outcomes [28]. Further details are provided in the data analysis plan (see [Supplementary data](#)).

## RESULTS

Study sites were established in 24 renal units in the UK. Between November 2014 and January 2016 a total of 620 patients attended the study screening visits and 566 (91%) entered the pre-randomization run-in (Figure 4).

### Pre-randomization run-in

A total of 138 participants withdrew from the pre-randomization run-in before attending a randomization visit (Table 2A). The most common medical reason for withdrawal from run-in was that the results from blood and urine samples taken at the screening visit did not confirm the participant's eligibility (Table 2A). Adverse events were uncommon and four participants were withdrawn because of a serious adverse event (myocardial infarction, septic shock and two cases of pneumonia).


In addition, 14 individuals attended a randomization visit but were not eligible to be randomized: the most common reason for this was their blood pressure being too low (Table 2B). Overall, 152 (27%) of the 566 individuals who entered the pre-randomization single-blind placebo run-in phase were not subsequently randomized.

### Baseline characteristics of randomized participants

A total of 414 people were randomized (Figure 4). The mean age was 63 (SD 14) years and 298 (72%) were male (Table 3). The mean systolic blood pressure was 146 (SD 16) mmHg at randomization (i.e. after 4–7 weeks of withdrawal of any prior ACEi or ARB). Based on results from the local laboratories, the mean eGFR was 34.0 (SD 10.6) mL/min/ $1.73$  m<sup>2</sup> and the median uACR was 58.5 (interquartile range 12.5–156.3) mg/mmol. Central laboratory assays will be conducted at the end of the study. About half of randomized participants had either glomerular [111 (27%)] or diabetic [83 (20%)] kidney disease and 165 (40%) patients reported diabetes mellitus at baseline. The median 5-year risk of ESRD (calculated using a validated risk calculator [29]) was 16.5%, and 62% of participants had a 5-year risk  $>10\%$ .

## DISCUSSION

The UK HARP-III trial has recruited 414 participants with CKD and will provide information on the short-term effects of sacubitril/valsartan on the change in kidney function (using mGFR) and the tolerability and safety of the drug compared



**FIGURE 4:** Trial profile: flow of participants through the trial. \*Indicates that participants may have more than one reason.

**Table 2. Reasons for (A) withdrawal during run-in and (B) ineligibility at a randomization visit**

	n (%)
(A)	
Number entering run-in	566
<i>Adverse event</i>	
Serious adverse event	4 (3)
Non-serious adverse reaction	7 (5)
<i>Other reason</i>	
Ineligible on laboratory results sent at screening visit	59 (43)
Participant wishes	16 (12)
Medical advice	13 (9)
Other non-medical reason	39 (28)
Total withdrawn during Run-in	138 (100)
(B)	
Number attending randomization visit	428
<i>Adverse event</i>	
Serious adverse event	0 (0)
Non-serious adverse reaction	3 (21)
<i>Other reason</i>	
Blood pressure too low	9 (64)
Other	2 (14)
Total ineligible at randomization visit	14 (100)

with irbesartan in people with CKD. The trial will also provide information on the effects of sacubitril/valsartan on albuminuria, blood pressure and other biomarkers of both kidney and cardiac function. These results are important because sacubitril/valsartan has now entered routine clinical practice as a treatment for heart failure with reduced ejection fraction (HFrEF)

[30], and many of these patients also have CKD. Moreover, NEPi has the potential to be a useful treatment for CKD itself.

Large randomized trials of interventions to slow the progression of CKD are required since currently available treatments do not prevent ESRD in all patients with CKD. Although ACEis and ARBs reduce the risk of progression of proteinuric diabetic and non-diabetic kidney disease, their effect (like most medical treatments) is moderate. For example, in proteinuric diabetic kidney disease, irbesartan reduced the risk of ESRD, doubling of creatinine or death from any cause by 20% compared with placebo {hazard ratio [HR] 0.80 [95% confidence interval (CI) 0.66–0.97];  $P = 0.02$ }, but this composite outcome still occurred in nearly one-third of those allocated irbesartan (and 14% reached ESRD) during the mean 2.6 years of follow-up [6]. Other strategies to reduce the risk of renal progression have either been ineffective, hazardous or both [31–33]. Neprilysin inhibition appears to be effective in rat models of CKD [15, 16, 34], but these are poorly predictive of efficacy in humans [35, 36]. In addition, sacubitril/valsartan has been shown to increase albuminuria in trials among patients with heart failure (who typically have very low baseline albuminuria) [18, 20]. NPs (particularly atrial NP) cause afferent arteriolar vasodilatation [37, 38] that may lead to increased intraglomerular pressure and hyperfiltration, which would be detrimental to the kidney. However, NEPi also disturbs degradation of other vasoactive peptides, so the net effect of NEPi on glomerular haemodynamics is uncertain, and in rat models at least, it appears to be favourable [15, 16, 34]. NPs may alter glomerular permeability and/or tubular reabsorption of protein, which may lead to albuminuria without hyperfiltration, the consequences of which are

**Table 3. Baseline characteristics of UK HARP-III participants**

Baseline characteristic	All participants (n = 414)
Age (years)	63 ± 14
<50	73 (18)
≥50–70	196 (47)
≥70	145 (35)
Gender	
Male	298 (72)
Female	116 (28)
Ethnicity	
White	377 (91)
Black	7 (2)
South Asian	18 (4)
Other	12 (3)
Prior disease	
Coronary heart disease	55 (13)
Cerebrovascular disease	31 (7)
Peripheral arterial disease	44 (11)
Heart failure	17 (4)
Diabetes	165 (40)
Systolic blood pressure (mmHg)	146 ± 16
<140	149 (36)
≥140–160	180 (43)
≥160	85 (21)
Diastolic blood pressure (mmHg)	81 ± 11
<80	191 (46)
≥80–90	133 (32)
≥90	90 (22)
Body mass index (kg/m <sup>2</sup> )	30.6 ± 6.2
<25	68 (16)
≥25–30	147 (36)
≥30	195 (47)
Not available	4 (1)
Medication	
Antiplatelet therapy	138 (33)
Oral anticoagulant	28 (7)
Diuretic	164 (40)
Calcium channel blocker	207 (50)
β-blocker	112 (27)
α-blocker	112 (27)
LDL-lowering agent	263 (64)
Prior use of ACEi or ARB	
Yes	339 (82)
No	75 (18)
CKD-EPI eGFR (mL/min/1.73 m <sup>2</sup> )	34.0 ± 10.6
<30	169 (41)
≥30–45	176 (43)
≥45	64 (15)
Not available	5 (1)
Urine albumin:creatinine ratio (mg/mmol)	58.5 (12.5–156.3)
<3	48 (12)
≥3–<30	88 (21)
≥30	251 (61)
Not available	27 (7)
Primary renal diagnosis	
Glomerular disease	111 (27)
Tubulointerstitial disease	50 (12)
Diabetic kidney disease	83 (20)
Hypertensive/renovascular disease	42 (10)
Other systemic diseases affecting the kidneys	3 (1)
Familial/hereditary nephropathies	43 (10)
Miscellaneous renal disorders	9 (2)
Unknown	73 (18)

Recorded at randomization visit unless otherwise stated. Values are given as n (%), mean ± SD or median (interquartile range).

CKD-EPI, Chronic Kidney Disease Epidemiology Collaboration; LDL, low-density lipoprotein.

uncertain. UK HARP-III is the first trial of NEPi in humans with CKD and the measurements of GFR, albuminuria and other markers of kidney function and damage will help to resolve these uncertainties.

Most patients with CKD do not progress to ESRD [39], but are at high risk of CVD [4]. Lowering low-density lipoprotein cholesterol has been shown to clearly reduce the risk of atherosclerotic vascular disease in CKD [40]. However, as renal function declines, the pattern of CVD changes from atherosclerotic disease (i.e. myocardial infarction, ischaemic stroke) to non-atherosclerotic disease (characterized by arteriosclerosis and structural heart disease, which manifests clinically similarly to heart failure, with a high incidence of sudden cardiac death) [4, 41–43], but effective treatments for non-atherosclerotic disease are not yet available. Lowering blood pressure in patients with CKD appears to reduce the risk of a wide variety of cardiovascular events, but residual risk remains [44]. The similarities in the manifestation of non-atherosclerotic disease observed in CKD and heart failure suggest that treatments that are effective in heart failure may well also be effective at reducing cardiovascular risk among patients with CKD. In the Prospective Comparison of ARNI with ACEI to Determine Impact on Global Mortality and Morbidity in Heart Failure (PARADIGM-HF) trial, sacubitril/valsartan reduced the risk of cardiovascular mortality or hospitalization for heart failure by 20% [HR 0.80 (95% CI 0.73–0.93) P < 0.001] compared with enalapril, with similar effects observed among participants with and without CKD at baseline [45]. These data suggest that NEPi would be an ideal candidate to test among patients with CKD. Nevertheless, most patients with CKD have a normal ejection fraction [41, 43], and treatments that improve outcomes in HF rEF do not necessarily improve outcomes in patients with heart failure with preserved ejection fraction [46, 47], so direct evidence is needed. NEPi has improved cardiac biomarkers (e.g. troponin, N-terminal pro-hormone brain NP) in trials in heart failure [18, 48], so the effects of NEPi on these cardiac biomarkers in people with CKD will also be of interest.

NEPi has the potential to improve both renal and cardiovascular outcomes among patients with CKD. The UK HARP-III trial will provide important information on the efficacy, safety and tolerability of sacubitril/valsartan in people with CKD. Results are anticipated in 2017.

## SUPPLEMENTARY DATA

Supplementary data are available online at <http://ndt.oxfordjournals.org>.

## ACKNOWLEDGEMENTS

The most important acknowledgement is to the participants in the study and to the collaborators listed in the appendix. The UK HARP-III Collaborative Group acknowledges the support of the National Institute for Health Research, through the Clinical Research Network. The UK HARP-III trial was funded by a grant to the University of Oxford from Novartis

(manufacturers of sacubitril/valsartan), but the study design, conduct, analysis and interpretation are the responsibility of the trial Steering Committee working with the Clinical Trial Service Unit & Epidemiological Studies Unit (CTSU). The University of Oxford is the regulatory sponsor for the trial. CTSU receives core funding from the UK Medical Research Council, British Heart Foundation and Cancer Research UK.

### CONFLICT OF INTEREST STATEMENT

CTSU has a staff policy of not accepting honoraria or other payments from the pharmaceutical industry, except for the reimbursement of costs to participate in scientific meetings ([www.ctsu.ox.ac.uk](http://www.ctsu.ox.ac.uk)).

### REFERENCES

- Bruck K, Stel VS, Gambaro G *et al*. CKD prevalence varies across the European General Population. *J Am Soc Nephrol* 2015; 27: 2135–2147
- Coresh J, Selvin E, Stevens LA *et al*. Prevalence of chronic kidney disease in the United States. *JAMA* 2007; 298: 2038–2047
- Gansevoort RT, Correa-Rotter R, Hemmelgarn BR *et al*. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; 382: 339–352
- Go AS, Chertow GM, Fan D *et al*. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004; 351: 1296–1305
- Lewis EJ, Hunsicker LG, Bain RP *et al*. The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. *N Engl J Med* 1993; 329: 1456–1462
- Lewis EJ, Hunsicker LG, Clarke WR *et al*. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001; 345: 851–860
- Brenner BM, Cooper ME, de Zeeuw D *et al*. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001; 345: 861–869
- Ruggenenti P, Perna A, Gherardi G *et al*. Renoprotective properties of ACE-inhibition in non-diabetic nephropathies with non-nephrotic proteinuria. *Lancet* 1999; 354: 359–364
- Sager G. Cyclic GMP transporters. *Neurochem Int* 2004; 45: 865–873
- Wilkins MR, Redondo J, Brown LA. The natriuretic-peptide family. *Lancet* 1997; 349: 1307–1310
- Mangiafico S, Costello-Boerrigter LC, Andersen IA *et al*. Neutral endopeptidase inhibition and the natriuretic peptide system: an evolving strategy in cardiovascular therapeutics. *Eur Heart J* 2013; 34: 886–893c
- Benigni A, Zoja C, Zatelli C *et al*. Vasopeptidase inhibitor restores the balance of vasoactive hormones in progressive nephropathy. *Kidney Int* 2004; 66: 1959–1965
- Kostis JB, Packer M, Black HR *et al*. Omapatrilat and enalapril in patients with hypertension: the Omapatrilat Cardiovascular Treatment vs. Enalapril (OCTAVE) Trial. *Am J Hypertens* 2004; 17: 103–111
- Gu J, Noe A, Chandra P *et al*. Pharmacokinetics and pharmacodynamics of LCZ696, a novel dual-acting angiotensin receptor-neprilysin inhibitor (ARNi). *J Clin Pharmacol* 2010; 50: 401–414
- Taal MW, Nenov VD, Wong W *et al*. Vasopeptidase inhibition affords greater renoprotection than angiotensin-converting enzyme inhibition alone. *J Am Soc Nephrol* 2001; 12: 2051–2059
- Roksnoer LC, van Veghel R, van Groningen MC *et al*. Blood pressure-independent renoprotection in diabetic rats treated with AT1 receptor-neprilysin inhibition compared with AT1 receptor blockade alone. *Clin Sci (Lond)* 2016; 130: 1209–1220
- Damman K, Andersen K, Belohlavek J *et al*. Angiotensin receptor neprilysin inhibition and renal function and in heart failure: results from PARADIGM-HF. *Eur Heart J* 2015; 36(Suppl 1): 545, abstract
- Solomon SD, Zile M, Pieske B *et al*. The angiotensin receptor neprilysin inhibitor LCZ696 in heart failure with preserved ejection fraction: a phase 2 double-blind randomised controlled trial. *Lancet* 2012; 380: 1387–1395
- Ruilope LM, Dukat A, Bohm M *et al*. Blood-pressure reduction with LCZ696, a novel dual-acting inhibitor of the angiotensin II receptor and neprilysin: a randomised, double-blind, placebo-controlled, active comparator study. *Lancet* 2010; 375: 1255–1266
- Voors AA, Gori M, Liu LC *et al*. Renal effects of the angiotensin receptor neprilysin inhibitor LCZ696 in patients with heart failure and preserved ejection fraction. *Eur Journal Heart Fail* 2015; 17: 510–517
- Medicines.org. Summary of Product Characteristics (SPC) Irbesartan 150 mg. Accessed 27 January 2016 <https://www.medicines.org.uk/emc/medicine/27608#ORIGINAL>
- Lang JM, Buring JE, Rosner B *et al*. Estimating the effect of the run-in on the power of the Physicians' Health Study. *Stat Med* 1991; 10: 1585–1593
- Kidney Disease: Improving Global Outcomes (KDIGO) Blood pressure Work Group. KDIGO clinical practice guidelines for the management of blood pressure in chronic kidney disease. *Kidney Int Suppl* 2012; 2: 337–414
- Borm GF, Franssen J, Lemmens WA. A simple sample size formula for analysis of covariance in randomized clinical trials. *J Clin Epidemiol* 2007; 60: 1234–1238
- Peto R, Pike MC, Armitage P *et al*. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. I. Introduction and design. *Br J Cancer* 1976; 34: 585–612
- Peto R, Pike MC, Armitage P *et al*. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. II. Analysis and examples. *Br J Cancer* 1977; 35: 1–39
- Vickers AJ, Altman DG. Statistics notes: analysing controlled trials with baseline and follow up measurements. *BMJ* 2001; 323: 1123–1124
- Rubin D. *Multiple Imputation for non-Response in Surveys*. New York: John Wiley, 1987
- Tangri N, Stevens LA, Griffith J *et al*. A predictive model for progression of chronic kidney disease to kidney failure. *JAMA* 2011; 305: 1553–1559
- Medical Letter on Drugs and Therapeutics. Sacubitril/valsartan (entresto) for heart failure. *JAMA* 2015; 314: 722–723
- Fried LF, Emanuele N, Zhang JH *et al*. Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med* 2013; 369: 1892–1903
- Parving H-H, Brenner BM, McMurray JVV *et al*. Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *N Engl J Med* 2012; 367: 2204–2213
- de Zeeuw D, Akizawa T, Audhya P *et al*. Bardoxolone methyl in type 2 diabetes and stage 4 chronic kidney disease. *N Engl J Med* 2013; 369: 2492–2503
- Cao Z, Burrell LM, Tikkanen I *et al*. Vasopeptidase inhibition attenuates the progression of renal injury in subtotal nephrectomised rats. *Kidney Int* 2001; 60: 715–721
- Perel P, Roberts I, Sena E *et al*. Comparison of treatment effects between animal experiments and clinical trials: systematic review. *BMJ* 2007; 334: 197
- Pound P, Ebrahim S, Sandercock P *et al*. Where is the evidence that animal research benefits humans? *BMJ* 2004; 328: 514–517
- Marin-Grez M, Fleming JT, Steinhausen M. Atrial natriuretic peptide causes pre-glomerular vasodilatation and post-glomerular vasoconstriction in rat kidney. *Nature* 1986; 324: 473–476
- Stockand JD, Sansom SC. Regulation of filtration rate by glomerular mesangial cells in health and diabetic renal disease. *Am J Kidney Dis* 1997; 29: 971–981
- Keith DS, Nichols GA, Gullion CM *et al*. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organization. *Arch Intern Med* 2004; 164: 659–663
- Baigent C, Landray MJ, Reith C *et al*. The effects of lowering LDL cholesterol with simvastatin plus ezetimibe in patients with chronic kidney disease (Study of Heart and Renal Protection): a randomised placebo-controlled trial. *Lancet* 2011; 377: 2181–2192
- Foley RN, Parfrey PS, Harnett JD *et al*. Clinical and echocardiographic disease in patients starting end-stage renal disease therapy. *Kidney Int* 1995; 47: 186–192
- Foley RN, Parfrey PS, Harnett JD *et al*. Impact of hypertension on cardiomyopathy, morbidity and mortality in end-stage renal disease. *Kidney Int* 1996; 49: 1379–1385



43. Patel RK, Jardine AG, Mark PB *et al.* Association of left atrial volume with mortality among ESRD patients with left ventricular hypertrophy referred for kidney transplantation. *Am J Kid Dis* 2010; 55: 1088–1096
44. Ettehad D, Emdin CA, Kiran A *et al.* Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016; 387: 957–967
45. McMurray JJ, Packer M, Desai AS *et al.* Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014; 371: 993–1004
46. Massie BM, Carson PE, McMurray JJ *et al.* Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med* 2008; 359: 2456–2467
47. Yusuf S, Pfeffer MA, Swedberg K *et al.* Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *Lancet* 2003; 362: 777–781
48. Jhund PS, Claggett BL, Voors AA *et al.* Elevation in high-sensitivity troponin T in heart failure and preserved ejection fraction and influence of treatment with the angiotensin receptor neprilysin inhibitor LCZ696. *Circ Heart Fail* 2014; 7: 953–959

Dayanandan, M. Hill, P.A. Kalra, J.J. McMurray, M. Taal, D.C. Wheeler, M.J. Landray<sup>†</sup>, C. Baigent<sup>†</sup>

<sup>\*</sup>, <sup>†</sup> Joint contribution.

#### STEERING COMMITTEE

*Chairman* C Baigent *Principal Investigators* R Haynes (Clinical Coordinator), MJ Landray *Administrative coordinator* R Dayanandan *Computing* A Baxter *Statistics* N Staplin *Other members* A Bethel, L Bowman, N Brunskill, P Cockwell, WG Herrington, M Hill, PK Judge, PA Kalra, C Knott, JJ McMurray, K Murphy, M Taal, DC Wheeler.

#### DATA MONITORING COMMITTEE

*Chairman* K Wheatley *Statistician* J Emberson *Other members* C Tomson, P Roderick

*Received:* 13.7.2016; *Editorial decision:* 24.7.2016

#### APPENDIX

#### WRITING COMMITTEE

P.K. Judge<sup>\*</sup>, R. Haynes<sup>\*</sup>, W.G. Herrington, B.C. Storey, N. Staplin, A. Bethel, L. Bowman, N. Brunskill, P. Cockwell, R.

*Nephrol Dial Transplant* (2017) 32: 2051–2058

doi: 10.1093/ndt/gfw317

Advance Access publication 16 September 2016

## An investigation of *APOL1* risk genotypes and preterm birth in African American population cohorts

Catherine C. Robertson<sup>1</sup>, Christopher E. Gillies<sup>1</sup>, Rosemary K.B. Putler<sup>1</sup>, Derek Ng<sup>2</sup>, Kimberly J. Reidy<sup>3</sup>, Brendan Crawford<sup>1</sup> and Matthew G. Sampson<sup>1</sup>

<sup>1</sup>Pediatric Nephrology, University of Michigan School of Medicine, 3560B MSRB II, 1150 West Medical Center Drive, Ann Arbor, MI 48109, USA, <sup>2</sup>Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA and <sup>3</sup>Montefiore Medical Center, Bronx, NY, USA

Correspondence and offprint requests to: Matthew G. Sampson; E-mail: mgsamps@med.umich.edu

#### ABSTRACT

**Background.** Two genetic variants in apolipoprotein L1 (*APOL1*) are associated with increased risk of focal segmental glomerulosclerosis as well as other glomerular phenotypes. These risk variants are common in individuals of African ancestry but absent in other racial groups. Yet, the majority of individuals with two *APOL1* risk alleles [high-risk (HR) genotype] do not have renal disease. It is critical to identify environmental and secondary genetic influences that, when combined

with these alleles, lead to kidney disease. In a recent study of black children with glomerular disease enrolled in the Nephrotic Syndrome Study Network (NEPTUNE) and Chronic Kidney Disease in Children Study ( $n = 104$ ), we found that subjects with an HR genotype had a 4.6-fold increase in the odds of preterm birth as compared to those with a low risk (LR) genotype [odds ratio 4.6 (CI 1.4–15.5)]. There are known racial disparities in preterm birth, which itself is a known risk factor for chronic kidney disease and focal segmental glomerulosclerosis. Thus, we questioned whether an HR *APOL1* genotype is