

Rationality for isobaric automorphic representations: the CM-case

Harald Grobner¹ 

Received: 8 May 2017 / Accepted: 9 May 2018 / Published online: 21 May 2018
© The Author(s) 2018

Abstract In this note we prove a simultaneous extension of the author’s joint result with M. Harris for critical values of Rankin–Selberg L -functions $L(s, \Pi \times \Pi')$ (Grobner and Harris in *J Inst Math Jussieu* 15:711–769, 2016, Thm. 3.9) to (i) general CM-fields F and (ii) cohomological automorphic representations $\Pi' = \Pi_1 \boxplus \dots \boxplus \Pi_k$ which are the isobaric sum of unitary cuspidal automorphic representations Π_i of general linear groups of arbitrary rank over F . In this sense, the main result of these notes, cf. Theorem 1.9, is a generalization, as well as a complement, of the main results in Raghuram (*Forum Math* 28:457–489, 2016; *Int Math Res Not* 2:334–372, 2010. <https://doi.org/10.1093/imrn/rnp127>), and Mahnkopf (*J Inst Math Jussieu* 4:553–637, 2005).

Keywords Period · L -function · Critical value · Isobaric sum · Cuspidal automorphic

Mathematics Subject Classification 11F67 (Primary) 11F70 · 11G18 · 11R39 · 22E55 (Secondary)

Communicated by A. Constantin.

Harald Grobner is supported by START-prize Y-966 of the Austrian Science Fund (FWF).

✉ Harald Grobner
harald.grobner@univie.ac.at

¹ Fakultät für Mathematik, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria

1 Rationality for isobaric automorphic representations: the general case

1.1 Introductory comments: a leitfaden for the reader

The purpose of this note is to prove a broad generalization of our own rationality-result, [12, Thm. 3.9], established *ibidem* for critical values of the Rankin–Selberg L -function $L(s, \Pi \times \Pi')$ of certain automorphic representations $\Pi \otimes \Pi'$ of $\mathrm{GL}_n \times \mathrm{GL}_{n-1}$ over an imaginary quadratic field \mathcal{K} . Our generalization of this result will be in terms of the nature of the base field \mathcal{K} , and even more importantly, of the nature of the automorphic representation Π' .

1.1.1 A short review of our result in [12]

To put ourselves *in medias res*, we will briefly recall our rationality-theorem, [12, Thm. 3.9]. It applies to a pair (Π, Π') of a cohomological cuspidal automorphic representation Π of $\mathrm{GL}_n(\mathbb{A}_{\mathcal{K}})$ and a cohomological abelian automorphic representation Π' of $\mathrm{GL}_{n-1}(\mathbb{A}_{\mathcal{K}})$, i.e., an isobaric sum of distinct unitary Hecke characters $\Pi' = \chi \boxplus \cdots \boxplus \chi_{n-1}$, over imaginary quadratic fields \mathcal{K} . By a principle found in [14, 22, 27], which works in even greater generality as exploited in the latter references, one may attach a Whittaker period $p(\Pi)$ and $p(\Pi')$ to such representations: Explained in due shortness, this period is defined by comparison of

- (i) a fixed rational structure of the (unique) Whittaker model $W(\Pi_f)$ (resp. $W(\Pi'_f)$) of the finite part of the given automorphic representation and
- (ii) a fixed rational structure on a (uniquely chosen) Π_{f-} (resp. Π'_{f-}) isotypic subspace in the cohomology $H^{b_n}(S_n, \mathcal{E}_\mu)$ (resp. $H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda)$) of the adelic “locally symmetric space” S_n (reps. S_{n-1}) in the lowest, possible degree b_n (resp. b_{n-1}).

As both, the Whittaker model and the above cohomological model, are irreducible representations, their rational structures are unique up to multiplication by non-zero complex numbers. Hence, the Whittaker periods $p(\Pi)$ and $p(\Pi')$ may simply be defined as a choice of normalization-factor, which makes the isomorphism between the Whittaker model and our cohomological model, induced from the global ψ -Fourier coefficient, respect the two fixed choices of rational structures on domain and target space.

Recall the Gauß-sum $\mathcal{G}(\omega_{\Pi'_f})$ of the central character $\omega_{\Pi'_f}$ of Π'_f and assume that the coefficient modules \mathcal{E}_μ and \mathcal{E}_λ in cohomology allow a non-trivial $\mathrm{GL}_{n-1}(\mathbb{C})$ -equivariant intertwining $\mathcal{E}_\mu \otimes \mathcal{E}_\lambda \rightarrow \mathbb{C}$. Under these assumptions the rationality-theorem [12, Thm. 3.9] asserts that for every critical point of $L(s, \Pi \times \Pi')$, i.e., for every half-integer $s_0 = \frac{1}{2} + m$, for which the archimedean L -factors on both sides of the functional equation of $L(s, \Pi \times \Pi')$ are holomorphic, there is a non-zero archimedean period $p(m, \Pi_\infty, \Pi'_\infty) \in \mathbb{C}^*$, only depending on m , Π_∞ and Π'_∞ , such that

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} p(\Pi) p(\Pi') p(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_f}). \quad (1.1)$$

In other words, the critical value $L(\frac{1}{2} + m, \Pi_f \times \Pi'_f)$ equals the product of three periods and the above Gauß-sum, up to multiplication by an element in the composition of rationality-fields $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$: These latter fields are defined by reference to the natural action of $\text{Aut}(\mathbb{C})$ on non-archimedean representations Π_f and Π'_f (see [32], §I.1), and, most importantly, they are number fields. Hence, our rationality-theorem [12, Thm. 3.9] amounts to a description of the transcendental part of $L(\frac{1}{2} + m, \Pi_f \times \Pi'_f)$, asserting that all critical values of $L(s, \Pi \times \Pi')$ are a product of transcendental periods and a Gauß-sum, up to a factor coming out of a concrete number field, namely $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$, attached to Π_f and Π'_f .

1.1.2 The main result of this paper

In this paper, we show that (1.1) is still true, if we enlarge our framework to

- (i) general CM-fields F —instead of imaginary quadratic fields \mathcal{K} and
- (ii) general cohomological isobaric automorphic representations $\Pi' = \Pi_1 \boxplus \dots \boxplus \Pi_k$, which are fully-induced from distinct unitary cuspidal automorphic representation Π_i of general linear groups $\text{GL}_{n_i}(\mathbb{A}_F)$ of arbitrary rank $n_i \geq 1$ —instead of sums of Hecke characters χ_i .

In summary, our main result is

Theorem *Let F be any CM-field. Let Π be a cuspidal automorphic representation of $\text{GL}_n(\mathbb{A}_F)$, which is cohomological with respect to \mathcal{E}_μ and let $\Pi' = \Pi_1 \boxplus \dots \boxplus \Pi_k$ by an isobaric automorphic representation of $\text{GL}_{n-1}(\mathbb{A}_F)$, fully induced from distinct unitary cuspidal automorphic representations Π_i , $1 \leq i \leq k$, which is cohomological with respect to \mathcal{E}_λ and of central character $\omega_{\Pi'}$. We assume that there is a non-trivial $\text{GL}_{n-1}(F \otimes_{\mathbb{Q}} \mathbb{R})$ -equivariant intertwining $\mathcal{E}_\mu \otimes \mathcal{E}_\lambda \rightarrow \mathbb{C}$. Then, for every critical point $\frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$, there is a non-zero archimedean period $p(m, \Pi_\infty, \Pi'_\infty) \in \mathbb{C}^*$, only depending on m, Π_∞ and Π'_∞ , such that*

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} p(\Pi) p(\Pi') p(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_f}),$$

where “ $\sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)}$ ” means up to multiplication by an element in the composition of number fields $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$.

Our main result has the following direct consequence:

Corollary *Let Π and Π' be as in the statement of the main theorem above. Let $\frac{1}{2} + m, \frac{1}{2} + \ell$ be two critical values of $L(s, \Pi \times \Pi')$ and abbreviate $\Omega_{\Pi_\infty, \Pi'_\infty}(m, \ell) := p(m, \Pi_\infty, \Pi'_\infty)p(\ell, \Pi_\infty, \Pi'_\infty)^{-1}$. Then, whenever $L^S(\frac{1}{2} + \ell, \Pi \times \Pi')$ is non-zero (e.g., if Π is unitary and $\ell \neq 0$),*

$$\frac{L^S\left(\frac{1}{2} + m, \Pi \times \Pi'\right)}{L^S\left(\frac{1}{2} + \ell, \Pi \times \Pi'\right)} \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} \Omega_{\Pi_\infty, \Pi'_\infty}(m, \ell),$$

which only depends on the archimedean components Π_∞ and Π'_∞ .

In particular, if $L^S(\frac{3}{2} + m, \Pi \times \Pi')$ is non-zero (e.g., if Π is unitary and $m \neq -1$), then the quotient of consecutive critical L -values satisfies

$$\frac{1}{\Omega_{\Pi_\infty, \Pi'_\infty}(m)} \frac{L^S(\frac{1}{2} + m, \Pi \times \Pi')}{L^S(\frac{3}{2} + m, \Pi \times \Pi')} \in \mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f).$$

Here we wrote $\Omega_{\Pi_\infty, \Pi'_\infty}(m) := \Omega_{\Pi_\infty, \Pi'_\infty}(m, m + 1)$

As its key-feature, our corollary avoids any reference to Whittaker periods and expresses quotients of critical values of $L(s, \Pi \times \Pi')$ in terms of archimedean factors only. The reader may want to compare this corollary to the main result of [15], where a similar result on quotients of consecutive critical values of Rankin–Selberg L -functions attached to cuspidal representations Π and Π' over totally real fields has been established. Our corollary hence complements this important result.

In order to keep our presentation precise, but at the same time short, we will focus on the crucial parts of the proof of our main theorem in this note and avoid repeating arguments given in [12] already, if they transfer verbatim to the more general situation here. In other words, we will only work out in details those steps of the proof, which need an extra argument, not contained in [12], and refer to precise statements in [12], if possible. *The reader is hence strongly advised to keep a copy of [12] ready at hand. Unexplained notation or references (e.g., “§2.1.1”) refer to this source [12].*

1.2 The setup

1.2.1 Algebraic data

We let F be any CM-field of dimension $2d = \dim_{\mathbb{Q}} F$ and set of archimedean places S_∞ . Each place $v \in S_\infty$ refers to a fixed pair of conjugate complex embeddings $(\iota_v, \bar{\iota}_v)$ of F , where we will drop the subscript “ v ” if it is clear from the context. We let \mathcal{O} be the ring of integers of F and for $v \notin S_\infty$, \mathcal{O}_v its local integral completion in F_v . The non-trivial additive character $\psi : F \backslash \mathbb{A} \rightarrow \mathbb{C}^*$ is defined as in §2.1.1. Throughout this note G denotes the general linear group GL_n and G' denotes the general linear group GL_{n-1} , both defined over F ($n \geq 2$).

1.2.2 Highest weight modules

We let E_μ (resp. E_λ) be an irreducible finite-dimensional representation of the *real* Lie group $G_\infty = R_{F/\mathbb{Q}}(G)(\mathbb{R})$ (resp. $G'_\infty = R_{F/\mathbb{Q}}(G')(\mathbb{R})$) on a complex vector-space, given by its highest weight $\mu = (\mu_v)_{v \in S_\infty}$ (resp. $\lambda = (\lambda_v)_{v \in S_\infty}$). Both representations are assumed to be algebraic: In terms of the standard choice of a maximal torus and positivity on the corresponding set of roots, this means that $\mu_v = (\mu_{\iota_v}, \mu_{\bar{\iota}_v}) \in \mathbb{Z}^n \times \mathbb{Z}^n$ (and the analogous assertion for λ). If $\sigma \in \mathrm{Aut}(\mathbb{C})$ is any automorphism of the field \mathbb{C} , then we define ${}^\sigma E_\mu$ to be the irreducible finite-dimensional representation of G_∞ of highest weight ${}^\sigma \mu = (({}^\sigma \mu)_v)_{v \in S_\infty}$, where at a place $v = (\iota_v, \bar{\iota}_v)$ we let $({}^\sigma \mu)_v = (\mu_{\sigma^{-1} \circ \iota_v}, \mu_{\sigma^{-1} \circ \bar{\iota}_v})$. The analogous definition yields us an irreducible finite-dimensional representation ${}^\sigma E_\lambda$ of G'_∞ .

1.2.3 Real unitary subgroups

We chose a maximal compact subgroup C_∞ (resp. C'_∞) of G_∞ (resp. G'_∞) and define real Lie subgroups $K_\infty := Z_{G_\infty} C_\infty \cong (\mathbb{R}_{>0} U(n))^d$ of G_∞ (resp. $K'_\infty := Z_{G'_\infty} C'_\infty \cong (\mathbb{R}_{>0} U(n-1))^d$ of G'_∞), where $U(k)$ denotes the usual compact unitary Lie group of rank k .

1.2.4 The cuspidal representation Π

Throughout this note, Π denotes a cuspidal automorphic representation of $G(\mathbb{A})$ with non-trivial $(\mathfrak{g}_\infty, K_\infty)$ -cohomology with respect to E_μ : This is equivalent to Π being regular algebraic in the sense of [6, Def. 3.12] (cf. [13, Thm. 6.3] for details). We do not assume Π to be unitary, but allow arbitrary integer twists $\|\det\|^m$ of unitary cuspidal automorphic representations $\tilde{\Pi}$: $\Pi = \tilde{\Pi} \cdot \|\det\|^m$. For convenience we will not distinguish between a cuspidal automorphic representation, its smooth Fréchet space completion of moderate growth and its (non-smooth) Hilbert space completion in the L^2 -spectrum. Introducing subindices “ v ”, $\Pi_\infty = \otimes_{v \in S_\infty} \Pi_v$ is hence locally of the form described in §2.4:

$$\Pi_v \cong \text{Ind}_{B(\mathbb{C})}^{G(\mathbb{C})} \left[z_1^{\ell_{v,1}+m} \bar{z}_1^{-\ell_{v,1}+m} \otimes \cdots \otimes z_n^{\ell_{v,n}+m} \bar{z}_n^{-\ell_{v,n}+m} \right],$$

where

$$\ell_{v,j} := \ell(\mu_{\iota_v}, j) := -\mu_{\iota_v, n-j+1} - m + \frac{n+1}{2} - j$$

and induction is unitary. By [6, Thm. 3.13], for each $\sigma \in \text{Aut}(\mathbb{C})$ there exists a unique cuspidal automorphic representation ${}^\sigma \Pi$ of $G(\mathbb{A})$, which is cohomological with respect to ${}^\sigma E_\mu$ and whose finite part satisfies $({}^\sigma \Pi)_f = {}^\sigma (\Pi_f) := \Pi_f \otimes_\sigma \mathbb{C}$. Since m is an integer, we have ${}^\sigma \Pi = ({}^\sigma \tilde{\Pi}) \cdot \|\det\|^m$, where ${}^\sigma \tilde{\Pi}$ is a regular algebraic, unitary cuspidal automorphic representation, defined similarly. We let $W(\Pi_f)$ be the finite part of the global Whittaker model $W(\Pi, \psi^{-1})$ defined by the ψ^{-1} -Fourier coefficient.

1.2.5 The isobaric representation Π'

Let $\sum_{i=1}^k n_i = n - 1$ be any partition of $n - 1$. As the second representation-theoretic ingredient, Π' denotes an automorphic representation of $G'(\mathbb{A})$ with non-trivial $(\mathfrak{g}'_\infty, K'_\infty)$ -cohomology with respect to E_λ , which is the isobaric sum of pairwise different, unitary cuspidal automorphic representations Π_i of $\text{GL}_{n_i}(\mathbb{A})$, $1 \leq i \leq k$,

$$\Pi' := \Pi_1 \boxplus \cdots \boxplus \Pi_k \cong \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} [\Pi_1 \otimes \cdots \otimes \Pi_k].$$

Here, P' denotes the standard parabolic subgroup of G' with Levi factor isomorphic to $\prod_{i=1}^k \text{GL}_{n_i}$ (and the latter isomorphism of representations is automatic, [1, 2, 21, 31]).

Remark 1.1 As a paradigmatic example, any representation Π' which is the cohomological quadratic base change from a quasi-split unitary group as in [7], p. 122, will be of the above form, see [7, Thm. 6.1].

Since the cuspidal representations Π_i are pairwise different, a combination of [29, Prop. 7.1.3, Thm. 3.5.12 and Rem. 3.5.14] implies that Π' is globally ψ -generic. We let $W(\Pi'_f)$ be the finite part of the global Whittaker model $W(\Pi', \psi)$ defined by the ψ -Fourier coefficient.

Abstract local genericity of the irreducible unitary representations Π'_v at an archimedean place $v \in S_\infty$ hence shows (cf., e.g., [13] §5.5) that necessarily

$$\Pi'_v \cong \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})} \left[z_1^{k_{v,1}} \bar{z}_1^{-k_{v,1}} \otimes \cdots \otimes z_{n-1}^{k_{v,n-1}} \bar{z}_{n-1}^{-k_{v,n-1}} \right],$$

where

$$k_{v,j} := k(\lambda_{l_v}, j) := -\lambda_{l_v, n-j} + \frac{n}{2} - j,$$

i.e., each Π'_v is of the form considered in §2.5.

Let $\rho_{P'}$ be the usual square-root of the modulus character of $P'(\mathbb{A})$, [5, 0, 3.5]. We write $\rho_i := \rho_{P'}|_{\text{GL}_{n_i}}$ for the restriction of $\rho_{P'}$ to the particular factor GL_{n_i} of the Levi subgroup. By [5, III, Thm. 3.3] the global representations $\Xi_i := \Pi_i \cdot \rho_i$ are regular algebraic cuspidal automorphic representations (for details see [13, pp. 1002–1003]). Hence, as for Π above, for each $\sigma \in \text{Aut}(\mathbb{C})$ and all $1 \leq i \leq k$, there are uniquely determined cuspidal automorphic representations ${}^\sigma \Xi_i$, which are cohomological with respect to the corresponding, σ -permuted coefficient module of $\text{GL}_{n_i}(\mathbb{C})$ and whose finite part satisfies $({}^\sigma \Xi_i)_f = {}^\sigma(\Xi_{i,f}) := \Xi_{i,f} \otimes_\sigma \mathbb{C}$. The representations $({}^\sigma \Xi_i) \cdot \rho_i^{-1}$ are hence pairwise different, unitary cuspidal automorphic representations. We let

$${}^\sigma \Pi' := ({}^\sigma \Xi_1) \cdot \rho_1^{-1} \boxplus \cdots \boxplus ({}^\sigma \Xi_k) \cdot \rho_k^{-1}$$

be their isobaric sum.

Lemma 1.2 *The representation ${}^\sigma \Pi'$ is fully induced, i.e.,*

$${}^\sigma \Pi' = ({}^\sigma \Xi_1) \cdot \rho_1^{-1} \boxplus \cdots \boxplus ({}^\sigma \Xi_k) \cdot \rho_k^{-1} \cong \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} \left[({}^\sigma \Xi_1) \cdot \rho_1^{-1} \otimes \cdots \otimes ({}^\sigma \Xi_k) \cdot \rho_k^{-1} \right]$$

and we have $({}^\sigma \Pi')_f \cong {}^\sigma(\Pi'_f)$.

Proof For the first assertion observe that $({}^\sigma \Xi_{i,v}) \cdot \rho_{i,v}^{-1}$ is irreducible and unitary for each $1 \leq i \leq k$ and each place v of F . Hence, $\text{Ind}_{P'(F_v)}^{G'(F_v)} [({}^\sigma \Xi_{1,v}) \cdot \rho_{1,v}^{-1} \otimes \cdots \otimes ({}^\sigma \Xi_{k,v}) \cdot \rho_{k,v}^{-1}]$ is irreducible for each v of F , see [2] (for $v \notin S_\infty$) and [1, 31] (for $v \in S_\infty$). It follows that

$$\begin{aligned} \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} [({}^\sigma \Xi_1) \cdot \rho_1^{-1} \otimes \cdots \otimes ({}^\sigma \Xi_k) \cdot \rho_k^{-1}] &\cong \otimes_v \text{Ind}_{P'(F_v)}^{G'(F_v)} [({}^\sigma \Xi_{1,v}) \\ &\cdot \rho_{1,v}^{-1} \otimes \cdots \otimes ({}^\sigma \Xi_{k,v}) \cdot \rho_{k,v}^{-1}] \end{aligned}$$

is irreducible as well. Hence, $\sigma \Pi' = (\sigma \Xi_1) \cdot \rho_1^{-1} \boxplus \dots \boxplus (\sigma \Xi_k) \cdot \rho_k^{-1}$ being isomorphic to a non-trivial subquotient of the latter global, induced representation, cf. [21], p. 208, shows that

$$\sigma \Pi' = (\sigma \Xi_1) \cdot \rho_1^{-1} \boxplus \dots \boxplus (\sigma \Xi_k) \cdot \rho_k^{-1} \cong \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} \left[(\sigma \Xi_1) \cdot \rho_1^{-1} \otimes \dots \otimes (\sigma \Xi_k) \cdot \rho_k^{-1} \right].$$

For the second claim, observe that at $v \notin S_\infty$, the action of $\sigma \in \text{Aut}(\mathbb{C})$ commutes with unnormalized, algebraic induction “ ${}^a\text{Ind}$ ”, i.e., one has

$$\begin{aligned} \sigma \text{Ind}_{P'(F_v)}^{G'(F_v)} [\Pi_{1,v} \otimes \dots \otimes \Pi_{k,v}] &= \text{Ind}_{P'(F_v)}^{G'(F_v)} [\Pi_{1,v} \otimes \dots \otimes \Pi_{k,v}] \otimes_\sigma \mathbb{C} \\ &\cong {}^a\text{Ind}_{P'(F_v)}^{G'(F_v)} [(\Xi_{1,v} \otimes_\sigma \mathbb{C}) \otimes \dots \otimes (\Xi_{k,v} \otimes_\sigma \mathbb{C})] \\ &\cong \text{Ind}_{P'(F_v)}^{G'(F_v)} [(\sigma \Xi_{1,v}) \cdot \rho_{1,v}^{-1} \otimes \dots \otimes (\sigma \Xi_{k,v}) \cdot \rho_{k,v}^{-1}], \end{aligned}$$

This completes the proof. □

As a consequence of Lemma 1.2, reading [5, III, Thm. 3.3] backwards shows that $\sigma \Pi'$ is cohomological with respect to ${}^\sigma E_\lambda$. Moreover, the same argument as above shows that $\sigma \Pi'$ is globally ψ -generic for all $\sigma \in \text{Aut}(\mathbb{C})$.

Hence, $\sigma \Pi'$ satisfies the same properties imposed on Π' above, i.e., $\text{Aut}(\mathbb{C})$ leaves the class of $(\mathfrak{g}'_\infty, K'_\infty)$ -cohomological isobaric sums of pairwise different, unitary cuspidal automorphic representations stable.

1.3 Differences to the imaginary quadratic case: archimedean considerations

1.3.1 Highest weight representations carrying cuspidal data

Let E_μ be a coefficient module as in Sect. 1.2.4, i.e., $H^*(\mathfrak{g}_\infty, K_\infty \Pi \otimes E_\mu) \neq 0$ for a cuspidal representation Π as described above. This implies strong restrictions on the highest weight $\mu = (\mu_v)_{v \in S_\infty}$ in terms of its local components at archimedean places (which we may now have in an arbitrary number $d = |S_\infty|$), which we summarize shortly as

- Lemma 1.3** (1) $\mu_{\iota_v} - \mu_{\iota_v}^\vee = (-2\mathfrak{m}, \dots, -2\mathfrak{m})$ for all $v \in S_\infty$.
 (2) $(\sigma \mu)_{\bar{\iota}_v} = \mu_{\sigma^{-1} \circ \iota_v}$ for all $v \in S_\infty$ and all $\sigma \in \text{Aut}(\mathbb{C})$.

Proof (1) By assumption E_μ supports non-zero cohomology with respect to the cuspidal representation $\Pi = \tilde{\Pi} \cdot \|\det\|^{\mathfrak{m}}$, where $\tilde{\Pi}$ is unitary. Hence, $E_{\mu+\mathfrak{m}}$ is conjugate self-dual by [5, I, Cor. 4.2] and [4, Lem. 1.3]. This implies (1).

(2) Let $\sigma \in \text{Aut}(\mathbb{C})$. The irreducible module ${}^\sigma E_\mu$ of highest weight ${}^\sigma \mu = ((\sigma \mu)_v)_{v \in S_\infty}$ supports non-zero $(\mathfrak{g}_\infty, K_\infty)$ -cohomology with respect to the cuspidal automorphic representation ${}^\sigma \Pi$. Since ${}^\sigma \Pi = (\sigma \tilde{\Pi}) \cdot \|\det\|^{\mathfrak{m}}$, our point (1) above implies that $(\sigma \mu)_{\iota_v} - (\sigma \mu)_{\iota_v}^\vee = (-2\mathfrak{m}, \dots, -2\mathfrak{m})$ for all $v \in S_\infty$ and the same integer \mathfrak{m} for all σ . Inserting the definition of ${}^\sigma \mu$ gives

$$\mu_{\sigma^{-1} \circ \iota_v, j} + \mu_{\sigma^{-1} \circ \bar{\iota}_v, n-j+1} = -2\mathfrak{m}.$$

for all $1 \leq j \leq n$. On the other hand, applying (1) to the embedding $\iota'_v := \sigma^{-1} \circ \iota_v$ of F , we obtain

$$\mu_{\sigma^{-1} \circ \iota_v, j} + \overline{\mu_{\sigma^{-1} \circ \iota_v, n-j+1}} = -2m$$

for all $1 \leq j \leq n$. Combining the latter two equations shows $\mu_{\sigma^{-1} \circ \iota_v, n-j+1} = \overline{\mu_{\sigma^{-1} \circ \iota_v, n-j+1}}$ for every j and arbitrary $v \in S_\infty$, and $\sigma \in \text{Aut}(\mathbb{C})$. This proves (2). □

1.3.2 Cohomological automorphic representations

Although maybe looking as a pure technicality at first, Lemma 1.3 (2) is an important assertion: It guarantees that the action of $\text{Aut}(\mathbb{C})$ on those coefficient modules E_μ and E_λ , which carry automorphic cohomology as in Sects. 1.2.4 and 1.2.5,—although defined abstractly as a potentially arbitrary permutation of all the embeddings $\iota : F \hookrightarrow \mathbb{C}$ —does not tear apart the data $(\mu_{\iota_v}, \mu_{\bar{\iota}_v})$ resp. $(\lambda_{\iota_v}, \lambda_{\bar{\iota}_v})$ which is attached to a pair of embeddings $(\iota_v, \bar{\iota}_v)$ forming an archimedean place v . This implies the following corollary, which says that $\text{Aut}(\mathbb{C})$ acts on Π_∞ and Π'_∞ simply as a permutation of the local factors, potentially followed by a conjugation of the characters forming the inducing data:

Corollary 1.4 *For $\sigma \in \text{Aut}(\mathbb{C})$, let ${}^\sigma \ell_v := \ell({}^\sigma \mu)_{\iota_v}, j = \ell(\mu_{\sigma^{-1} \circ \iota_v}, j)$ and ${}^\sigma k_v := k({}^\sigma \lambda)_{\iota_v}, j = k(\lambda_{\sigma^{-1} \circ \iota_v}, j)$. For the archimedean components of the automorphic representations ${}^\sigma \Pi$ and ${}^\sigma \Pi'$, we obtain*

- (1) $({}^\sigma \Pi)_\infty \cong \otimes_{v \in S_\infty} \text{Ind}_{B(\mathbb{C})}^{G(\mathbb{C})} \left[z_1^{\ell_{v,1}+m} \bar{z}_1^{-\ell_{v,1}+m} \otimes \dots \otimes z_n^{\ell_{v,n}+m} \bar{z}_n^{-\ell_{v,n}+m} \right]$
- (2) $({}^\sigma \Pi')_\infty \cong \otimes_{v \in S_\infty} \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})} \left[z_1^{{}^\sigma k_{v,1}} \bar{z}_1^{-{}^\sigma k_{v,1}} \otimes \dots \otimes z_{n-1}^{{}^\sigma k_{v,n-1}} \bar{z}_{n-1}^{-{}^\sigma k_{v,n-1}} \right]$

Proof For Π this follows from Lemma 1.3, [5, IV Lem. 4.9] and the uniqueness of irreducible unitary generic representations of $\text{GL}_r(\mathbb{C})$, $r \geq 1$, with non-trivial cohomology with respect to a given finite-dimensional coefficient module, cf. [8, Thm. 6.1] (See also [13, §5.5] for a detailed exposition of the latter assertion). For Π' one first applies what we just said about Π to the cuspidal datum Ξ_1, \dots, Ξ_k and then carefully uses [5, III, Thm. 3.3] together with induction in stages. □

As a final consequence, and this is establishes the purpose of this section, we derive the following

“Meta-Lemma” Let A_∞ be an assertion of first-order predicate calculus, involving only ${}^\sigma \Pi_\infty$ or ${}^\sigma \Pi'_\infty$ for a family of $\sigma \in \text{Aut}(\mathbb{C})$. If A_∞ is true if and only if its restriction A_v to ${}^\sigma \Pi_v$ and ${}^\sigma \Pi'_v$ is true for all $v \in S_\infty$, and A_v is shown by an argument in [12], then A_∞ holds.

1.3.3 Archimedean consequences of the Meta-Lemma

Making our choices place-by-place $v \in S_\infty$ and applying our meta-lemma, we obtain

- (1) A natural $\mathbb{Q}(E_\mu)$ -rational vector-space structure on $H^q(\mathfrak{g}_\infty, K_\infty, \Pi_\infty \otimes E_\mu)$ (resp. $\mathbb{Q}(E_\lambda)$ -rational vector-space structure on $H^q(\mathfrak{g}'_\infty, K'_\infty, \Pi'_\infty \otimes E_\lambda)$) as in §2.7.
- (2) Basis-vectors $[\Pi_\infty]$ (resp. $[\Pi'_\infty]$) of the one-dimensional spaces $H^{b_n}(\mathfrak{g}_\infty, K_\infty, W(\Pi_\infty) \otimes E_\mu)$ (resp. $H^{b_n-1}(\mathfrak{g}'_\infty, K'_\infty, W(\Pi'_\infty) \otimes E_\lambda)$), where $b_r = d \cdot \frac{r(r-1)}{2}$, as in §3.3.
- (3) A well-defined “interlacing-hypothesis” of the highest weights μ and λ as in Hypothesis 2.3: This means we assume the validity of

Hypothesis 1.5 *For all archimedean places $v = (i_v, \bar{i}_v)$ the following inequalities hold:*

$$\begin{aligned} \mu_{i_v,1} &\geq -\lambda_{i_v,n-1} \geq \mu_{i_v,2} \geq -\lambda_{i_v,n-2} \geq \dots \geq -\lambda_{i_v,1} \geq \mu_{i_v,n} \\ \mu_{\bar{i}_v,1}^v &\geq -\lambda_{\bar{i}_v,n-1}^v \geq \mu_{\bar{i}_v,2}^v \geq -\lambda_{\bar{i}_v,n-2}^v \geq \dots \geq -\lambda_{\bar{i}_v,1}^v \geq \mu_{\bar{i}_v,n}^v. \end{aligned}$$

- (4) Given (the well-definedness of) this hypothesis, a description of the set of critical points $\text{Crit}(\Pi \times \Pi') \subset \frac{1}{2} + \mathbb{Z}$ of $L(s, \Pi \times \Pi')$:

$$\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi') \iff \text{Hom}_{R_{F/\mathbb{Q}}(G')(\mathbb{C})}(E_{\mu-m} \otimes E_\lambda, \mathbb{C}) \neq 0.$$

The proof proceeds as in Lem. 3.5, though, one needs to correct a slight mistake *ibidem* first: The restriction to non-negative $m \geq 0$ there is not to be made. See also Thm. 2.21 in [25], where this has meanwhile been proved in even greater generality.

- (5) For all $\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$, compatible choices of intertwining operators $\mathcal{T}^{(m)} \in \text{Hom}_{R_{F/\mathbb{Q}}(G')(\mathbb{C})}(E_{\mu-m} \otimes E_\lambda, \mathbb{C})$ as in §3.7. Again, following the previous point, there is no restriction on m being positive or negative here.
- (6) Finally and most importantly, for all $\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$, well-defined complex numbers $c(\frac{1}{2} + m, \Pi_\infty, \Pi'_\infty)$, defined as in §3.10, and proved to be non-vanishing as in Thm. 3.8. This allows us to define archimedean periods $p(m, {}^\sigma \Pi_\infty, {}^\sigma \Pi'_\infty)$ as in §3.10, i.e., as the inverse of $c(\frac{1}{2} + m, {}^\sigma \Pi_\infty, {}^\sigma \Pi'_\infty)$, for all $\sigma \in \text{Aut}(\mathbb{C})$. As it has been discussed above, this works whether or not $m \geq 0$.

1.4 Differences to the imaginary quadratic case: non-archimedean considerations

1.4.1 Special Whittaker vectors

We will choose very particular vectors $\xi_{\Pi'_v} \in W(\Pi'_v)$, at all non-archimedean places $v \notin S_\infty$ in analogy to §3.9. Let $T' \subset B' \subset G'$ be the diagonal maximal torus in the standard Borel subgroup B' of G' and denote $T'(F_v)^+ := \{t \in T'(F_v) \mid t_i t_{i+1}^{-1} \in \mathcal{O}_v, t_{n-1} = 1\}$. Since Π'_v is the generic, the assumptions of [20, Proposition (3.2)] are satisfied. Hence, any non-vanishing functional $\xi_{\Pi'_v} \in W(\Pi'_v)$ is already non-zero on $T'(F_v)^+ \subset G'(F_v)$. As another ingredient, let $K'(m'_v)$ be the mirahoric subgroup of $G'(F_v)$ of level m'_v . If m'_v equals the conductor of Π'_v , then, by [17, Theorem (5.1)] the

space of Whittaker vectors, transforming by the central character $\omega_{\Pi'_v}$ of Π'_v under the $K'(m'_v)$ is one-dimensional, its elements being called *new vectors*. As a consequence of the above discussion, we may fix a matrix $t_{\Pi'_v} \in T'(F_v)^+$ on which all the non-trivial new vectors of Π'_v do not vanish simultaneously, where we observe that we may choose the same matrix for all σ -twists of Π'_v , i.e., such that $t_{\Pi'_v} = t_{\sigma \Pi'_v}$. Moreover, if the non-archimedean place v is outside the set of ramification of Π' and ψ , then we may take $t_{\Pi'_v} := id$. Depending on these (mild) choices, for all $v \notin S_\infty$, we define $\xi_{\Pi'_v} \in W(\Pi'_v)$ to be the unique new vector such that $\xi_{\Pi'_v}(t_{\Pi'_v}) = 1$.

As the last ingredient, we remark that we may similarly also choose particular Whittaker vectors ξ_{Π_v} for $\Pi_v, v \notin S_\infty$: These choices depend on our data fixed for Π'_v above and can be made, *mutatis mutandis*, precisely as in §3.9: First, we fix a matrix $t_{\Pi_v} \in T(F_v)^+$, analogously as for $G'(F_v)$. Now, for a non-archimedean place v outside the set of ramification of Π' and ψ , we let ξ_{Π_v} be the unique new vector of Π_v , which satisfies $\xi_{\Pi_v}(t_{\Pi_v}) = 1$. It is a certain, non-zero multiple c_{Π_v} of the essential vector, see [17, (4.1) Théorème]. If v is, however, inside the set of ramification of Π' or ψ , then we take ξ_{Π_v} to be the unique Whittaker vector, whose restriction to $G'(F_v)$ is supported on $N'(F_v)t_{\Pi'_v}K'(m'_v)$ and there equal to $\psi_v^{-1}\omega_{\Pi'_v}^{-1}$. See also [26, 3.1.4] and [22, 2.1.1], where such choices were coined first.

Finally, we observe that Lemma 3.7 still holds for these special Whittaker vectors.

1.4.2 Rational structures for Whittaker models

Keeping in mind the above considerations, we see as in Prop. 2.7 that the representations $W(\Pi_f)$ and $W(\Pi'_f)$ may be defined over the rationality fields $\mathbb{Q}(\Pi_f)$, respectively $\mathbb{Q}(\Pi'_f)$, by taking invariants of normalized new vectors in each model. Moreover, both fields $\mathbb{Q}(\Pi_f)$ and $\mathbb{Q}(\Pi'_f)$ are number fields by the regular-algebraicity of the cuspidal representations Π and Ξ_1, \dots, Ξ_k , see [6, Thm. 3.13] (or, for a detailed proof, [13, Thm. 8.1]).

1.5 Global considerations

1.5.1 Eisenstein cohomology

We let $S_n := G(F)\backslash G(\mathbb{A})/K_\infty, S_{n-1} := G'(F)\backslash G'(\mathbb{A})/K'_\infty$ and $\tilde{S}_{n-1} := G'(F)\backslash G'(\mathbb{A})/C'_\infty \cong S_{n-1} \times \mathbb{R}_+^d$, similar to §3.1. These spaces are orbifolds and we have $\dim_{\mathbb{R}}(\tilde{S}_{n-1}) = b_n + b_{n-1}$.

We define $\varphi_{P'}$ to be the associate class of cuspidal automorphic representations of $L'(\mathbb{A})$, which is defined by the unitary cuspidal $\tau := \Pi_1 \otimes \dots \otimes \Pi_k$. The space $\mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}$ of automorphic forms is then defined as in §3.1. See also the original source [10, §1.3] or [11, §2.3]. We obtain the following important result on Eisenstein cohomology:

Proposition 1.6 *The natural morphism*

$$i_{\Pi'}^{b_{n-1}} : H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \Pi' \otimes E_\lambda) \rightarrow H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_\lambda)$$

of $G(\mathbb{A}_f)$ -modules, induced by the natural injection $\iota_{\Pi'} : \Pi' \hookrightarrow \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}$, is an isomorphism. Hence, there is the following commuting triangle of natural injections of $G'(\mathbb{A}_f)$ -modules

$$\begin{array}{ccc}
 H^{b_{n-1}}(\mathfrak{g}'_{\infty}, K'_{\infty}, \Pi' \otimes E_{\lambda}) & & \\
 \downarrow \Psi_{\Pi'}^{\text{Eis}} & \searrow \begin{matrix} i_{\Pi'}^{b_{n-1}} \\ \cong \\ \tilde{\mathcal{F}}_{\lambda}^{b_{n-1}} \end{matrix} & \\
 H^{b_{n-1}}(S_{n-1}, \mathcal{E}_{\lambda}) & \longleftarrow & H^{b_{n-1}}(\mathfrak{g}'_{\infty}, K'_{\infty}, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_{\lambda})
 \end{array}$$

Proof We assume familiarity with the general results of [11]. In [11, §3.1], following [9], a filtration

$$\mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} = \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(0)} \supseteq \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(1)} \supseteq \dots \supseteq \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(m)} \tag{1.2}$$

of $\mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}$ of finite length $m = m(\{P'\})$ has been defined. The successive quotients are shown to be isomorphic to a direct sum, index by a set of (isomorphism classes) of quadruples in $M_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(j)}$, $0 \leq j \leq m$. See [11, Thm. 4] for this result and [11, §3.2] for a precise definition of $M_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(j)}$. By construction (of the filtration (1.2) and of the sets $M_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(j)}$), one necessarily finds $(P', \tau, 0, 0) \in M_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(m)}$, cf. [11, §3.1–3.2]. However, as all summands in Π' are different and unitary, the description of the residual spectrum of GL_N , cf. [24] II–III, implies that this is the only quadruple at all, i.e., $\cup_{j=0}^m M_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(j)} = \{(P', \tau, 0, 0)\}$. As a consequence, see again [11, Thm. 4] in combination with Multiplicity One for the discrete spectrum of $G'(\mathbb{A})$,

$$\begin{aligned}
 \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} &= \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(0)} = \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(1)} = \dots = \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}^{(m)} \\
 &\cong \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} \left[\tau \otimes S(\check{\mathfrak{a}}_{P', \mathbb{C}}^{G'}) \right],
 \end{aligned}$$

where $S(\check{\mathfrak{a}}_{P', \mathbb{C}}^{G'})$ is the symmetric algebra of the dual of the Lie algebra of the split component $A_{P'}$ of P' , modulo the split component of G' . Hence,

$$H^q(\mathfrak{g}'_{\infty}, K'_{\infty}, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_{\lambda}) \cong H^q(\mathfrak{g}'_{\infty}, K'_{\infty}, \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})} \left[\tau \otimes S(\check{\mathfrak{a}}_{P', \mathbb{C}}^{G'}) \right] \otimes E_{\lambda}),$$

for all degrees q , see also [11, Cor. 16]. By the minimality of the degree $q = b_{n-1}$, we obtain

$$H^{b_{n-1}}(\mathfrak{g}'_{\infty}, K'_{\infty}, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_{\lambda}) \cong \Pi'_f,$$

see [13] (7.25), revealing $H^{b_{n-1}}(\mathfrak{g}'_{\infty}, K'_{\infty}, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_{\lambda})$ as irreducible. The natural map in cohomology $\iota_{\Pi'}^{b_{n-1}}$ induced from the natural inclusion $\iota_{\Pi'} : \Pi' \hookrightarrow$

$\mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}}$ has by construction the same image as the map in cohomology induced from the Eisenstein summation map

$$\text{Eis}_0 : \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\tau] \xrightarrow{\sim} \Pi',$$

cf. [21]. Hence, recalling that all Eisenstein series attached to K_∞ -finite sections in $\text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\tau]$ are holomorphic at $\Lambda = 0$, $\iota_{\Pi'}^{b_{n-1}}$ is non-zero by [28], Satz 4.11. See also [3] 2.9. As $H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \text{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\tau] \otimes E_\lambda) \cong \Pi'_f$ is irreducible, too, by the minimality of $q = b_{n-1}$, $\iota_{\Pi'}^{b_{n-1}}$ is an isomorphism. Now define $\mathcal{F}_\lambda^{b_{n-1}}$ to be the restriction to $H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_\lambda)$ of the isomorphism of [9, Thm. 18] and $\Psi_{\Pi'}^{\text{Eis}} := \mathcal{F}_\lambda^{b_{n-1}} \circ \iota_{\Pi'}^{b_{n-1}}$. Recalling the direct sum decomposition of Eisenstein cohomology, cf. [10, Thm. 2.3] or [11, §4.1–4.3] shows that $\mathcal{F}_\lambda^{b_{n-1}}$ (and hence also $\Psi_{\Pi'}^{\text{Eis}}$) are injections. \square

1.5.2 Rational structures on submodules of automorphic cohomology and related Whittaker periods

As a consequence of the previous section, the following global results and assertions transfer from [12]: firstly, we obtain

Proposition 1.7 *For any $\sigma \in \text{Aut}(\mathbb{C})$ the natural σ -linear bijection $\tilde{\sigma} : H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) \rightarrow H^{b_{n-1}}(S_{n-1}, {}^\sigma \mathcal{E}_\lambda)$ maps the image of $\Psi_{\Pi'}^{\text{Eis}}$ onto the image of $\Psi_{\sigma\Pi'}^{\text{Eis}}$.*

Proof Let ${}^\sigma \varphi_{P'}$ be the associate class of the unitary cuspidal automorphic representation ${}^\sigma \tau := ({}^\sigma \Xi_1) \cdot \rho_i^{-1} \otimes \cdots \otimes ({}^\sigma \Xi_k) \cdot \rho_k^{-1}$. By its very definition ${}^\sigma \Pi'$ is the isobaric automorphic sum of the unitary cuspidal automorphic representations $({}^\sigma \Xi_i) \cdot \rho_{P'}^{-1}$, from which it is fully-induced, see Lemma 1.2. Applying Proposition 1.6 to Π' and ${}^\sigma \Pi'$ reduces the problem to showing that $\tilde{\sigma} : H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) \rightarrow H^{b_{n-1}}(S_{n-1}, {}^\sigma \mathcal{E}_\lambda)$ maps $\mathcal{F}_\lambda^{b_{n-1}}(H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \mathcal{A}_{\mathcal{J}', \{P'\}, \varphi_{P'}} \otimes E_\lambda))$ onto the analogously defined module $\mathcal{F}_{\sigma\lambda}^{b_{n-1}}(H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \mathcal{A}_{\mathcal{J}', \{P'\}, {}^\sigma \varphi_{P'}} \otimes {}^\sigma E_\lambda))$. However, using that $\iota_{\Pi'}^{b_{n-1}}$ and $\iota_{\sigma\Pi'}^{b_{n-1}}$ are isomorphisms, i.e., invoking Proposition 1.6 once more, exactly the same arguments as in [13, proof of Thm. 7.23] go through, where this assertion is proved for regular coefficients E_λ . This shows the claim. \square

Definition 1.8 As a consequence of Propositions 1.6 and 1.7 the composition $(\Psi_{\sigma\Pi'}^{\text{Eis}})^{-1} \circ \tilde{\sigma} \circ \Psi_{\Pi'}^{\text{Eis}}$ makes sense and we denote the resulting σ -linear bijection

$$H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \Pi' \otimes E_\lambda) \rightarrow H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, {}^\sigma \Pi' \otimes {}^\sigma E_\lambda)$$

again by $\tilde{\sigma}$.

As an immediate corollary, we obtain a $\mathbb{Q}(\Pi'_f)$ -structure on the image of the injection $\Psi_{\Pi'}^{\text{Eis}}$, which naturally extends the $\mathbb{Q}(E_\lambda)$ -structure of $H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda)$ defined by Betti-cohomology: This follows easily from Propositions 1.7 above, invoking [6, Lem.

3.2.1] (and recalling that $\mathbb{Q}(E_\lambda) \subseteq \mathbb{Q}(\Pi'_f)$, which ones concludes exactly as in the proof of [13, Cor. 8.7]). Hence, by transfer of structure along the injection $\Psi_{\Pi'}^{\text{Eis}}$, constructed in Proposition 1.6, the irreducible $G'(\mathbb{A}_f)$ -module $H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \Pi' \otimes E_\lambda)$ carries a $\mathbb{Q}(\Pi'_f)$ -structure. We assume from now on to have fixed precisely this rational structure on the cohomology of Π' (and analogously on all its σ -twists ${}^\sigma \Pi'$).

Similarly, as it is well-known, the same arguments apply for the cuspidal automorphic representation Π and its $(\mathfrak{g}_\infty, K_\infty)$ -cohomology, which injects into $H^{b_n}(S_n, \mathcal{E}_\mu)$: We obtain a $\mathbb{Q}(\Pi_f)$ -structure on $H^{b_n}(\mathfrak{g}_\infty, K_\infty, \Pi \otimes E_\mu)$, which naturally extends the $\mathbb{Q}(E_\mu)$ -structure of $H^{b_n}(S_n, \mathcal{E}_\mu)$ defined by Betti-cohomology and a natural σ -linear bijection $\tilde{\sigma} : H^{b_n}(\mathfrak{g}_\infty, K_\infty, \Pi \otimes E_\mu) \rightarrow H^{b_n}(\mathfrak{g}_\infty, K_\infty, {}^\sigma \Pi \otimes {}^\sigma E_\mu)$.

With respect to these two rational structures on relative Lie algebra cohomology and the σ -linear bijections $\tilde{\sigma}$, the proof of Prop. 3.1 goes through word-for-word, recalling the validity of [17, Theorem (5.1)] for $\Pi'_v, v \notin S_\infty$. Hence, we obtain this way Whittaker-periods $p(\Pi)$ and $p(\Pi')$, well-defined up to multiplication by $\mathbb{Q}(\Pi_f)^*$, resp. $\mathbb{Q}(\Pi'_f)^*$. In turn, again as in Prop. 3.1, these periods define rationally normalized isomorphism Θ_0^{cusp} and Θ_0^{Eis} of the corresponding Whittaker models and relative Lie algebra cohomologies.

1.6 Statement and proof of the main theorem

Theorem 1.9 *Let F be any CM-field. Let Π be a cuspidal automorphic representation of $\text{GL}_n(\mathbb{A})$ (as in Sect. 1.2.4) which is cohomological with respect to E_μ and let Π' by an isobaric automorphic representation of $\text{GL}_{n-1}(\mathbb{A})$ (as in Sect. 1.2.5) which is cohomological with respect to E_λ and of central character $\omega_{\Pi'}$. We assume that the highest weights $\mu = (\mu_v)_{v \in S_\infty}$ and $\lambda = (\lambda_v)_{v \in S_\infty}$ satisfy the interlacing-hypothesis 1.5. Then the following holds:*

- (1) For all critical values $\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$ and every $\sigma \in \text{Aut}(\mathbb{C})$,

$$\begin{aligned} & \sigma \left(\frac{L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right)}{p(\Pi) p(\Pi') p(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_f})} \right) \\ &= \frac{L\left(\frac{1}{2} + m, {}^\sigma \Pi_f \times {}^\sigma \Pi'_f\right)}{p({}^\sigma \Pi) p({}^\sigma \Pi') p(m, {}^\sigma \Pi_\infty, {}^\sigma \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_f})}. \end{aligned}$$

- (2)

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} p(\Pi) p(\Pi') p(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_f}),$$

where “ $\sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)}$ ” means up to multiplication by an element in the composition of number fields $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$.

Proof As a first step, we observe that Lemma 3.4 and the results of §3.8 transfer verbatim from [12] to our case here. Hence, recollecting all the preparatory results of

this note, the following diagram, which amplifies the main diagram of §3.2, is finally well-defined:

$$\begin{array}{ccc}
 H_c^{b_n}(S_n, \mathcal{E}_\mu) \times H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) & \longrightarrow & H_c^{b_n}(\tilde{S}_{n-1}, \mathcal{E}_\mu) \times H^{b_{n-1}}(\tilde{S}_{n-1}, \mathcal{E}_\lambda) \\
 \uparrow \downarrow & & \downarrow \wedge \\
 H_{cusp}^{b_n}(S_n, \mathcal{E}_\mu) \times H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) & & H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathcal{E}_\mu \otimes \mathcal{E}_\lambda) \\
 \uparrow \downarrow \Psi = \Psi_\Pi^{\text{cusp}} \times \Psi_\Pi^{\text{Eis}} & & \downarrow \mathcal{T}^* \\
 H^{b_n}(\mathfrak{g}_\infty, K_\infty, \Pi \otimes E_\mu) \times H^{b_{n-1}}(\mathfrak{g}'_\infty, K'_\infty, \Pi' \otimes E_\lambda) & & H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C}) \\
 \uparrow \downarrow \Theta_0 = \Theta_0^{\text{cusp}} \times \Theta_0^{\text{Eis}} & & \downarrow f \\
 W(\Pi_f) \times W(\Pi'_f) & \xrightarrow{\text{Dia}} & \mathbb{C}
 \end{array}$$

As a next step, we observe that the results of [17, 18], as well as [6, Lemme 4.6] are valid for Π_v , whenever $\psi = \otimes_v \psi_v$ is unramified at $v \notin S_\infty$, whence the proof of [23, Prop. 2.3.(c)] carries over to the situation considered here. In other words, the correction-factors c_{Π_v} of Sect. 1.4.1 satisfy $\sigma(c_{\Pi_v}) = c_{\sigma\Pi_v}$ for all $\sigma \in \text{Aut}(\mathbb{C})$ and at all non-archimedean places, where both Π' and ψ are unramified.

As a final consequence, the proof of [12, Thm. 3.9] now goes through word-for-word in our more general situation at hand and we hence obtain Theorem 1.9 (1) by chasing our special Whittaker vectors $\xi_{\Pi_f} := \otimes_{v \notin S_\infty} \xi_{\Pi_v}$ and $\xi_{\Pi'_f} := \otimes_{v \notin S_\infty} \xi_{\Pi'_v}$ through the above diagram. Assertion (2) follows from (1) applying Strong Multiplicity One for isobaric automorphic representations ([19], Thm. 4.4) together with Multiplicity One ([10] §3.3 and [16,30]). □

Remark 1.10 Theorem 1.9 represents a rather vast generalization of [26, Thm. 1.1] and [25, Thm. 1.1] over general CM-fields F : In the latter references, the analogous result has been shown for cuspidal automorphic representations Π' (over $F = \mathbb{Q}$ in [26] and over a general number field F in [25])—a condition, which we stretched to all isobaric sums Π' , which are fully-induced from cuspidal representation Π_1, \dots, Π_k (as in Sect. 1.2.5) over arbitrary CM-fields F . The situation for isobaric representations over general number fields F will be significantly more complicated, notably at infinity.

2 A consequence

2.1 Ratios of critical values

The following result is a direct consequence of our main result. It avoids any reference to Whittaker periods and expresses quotients of critical values of $L(s, \Pi \times \Pi')$ in terms of archimedean factors only. The reader may compare this corollary to the main result of [15] on quotients of consecutive critical values of Rankin–Selberg L -functions attached to cuspidal representations Π and Π' over totally real fields.

Corollary 2.1 *Let F be any CM-field. Let Π be a cuspidal automorphic representation of $GL_n(\mathbb{A})$ (as in Sect. 1.2.4) which is cohomological with respect to E_μ and let $\Pi' = \Pi_1 \boxplus \dots \boxplus \Pi_k$ by an isobaric automorphic representation of $GL_{n-1}(\mathbb{A})$ (as in Sect. 1.2.5) which is cohomological with respect to E_λ and of central character $\omega_{\Pi'}$. We assume that the highest weights $\mu = (\mu_v)_{v \in S_\infty}$ and $\lambda = (\lambda_v)_{v \in S_\infty}$ satisfy the interlacing-hypothesis 1.5. Let $\frac{1}{2} + m, \frac{1}{2} + \ell \in \text{Crit}(\Pi \times \Pi')$ be two critical values and abbreviate $\Omega_{\Pi_\infty, \Pi'_\infty}(m, \ell) := p(m, \Pi_\infty, \Pi'_\infty)p(\ell, \Pi_\infty, \Pi'_\infty)^{-1}$. Then, whenever $L^S(\frac{1}{2} + \ell, \Pi \times \Pi')$ is non-zero (e.g., if Π is unitary and $\ell \neq 0$),*

$$\frac{L^S(\frac{1}{2} + m, \Pi \times \Pi')}{L^S(\frac{1}{2} + \ell, \Pi \times \Pi')} \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} \Omega_{\Pi_\infty, \Pi'_\infty}(m, \ell),$$

and hence only depends on the archimedean components Π_∞ and Π'_∞ . In particular, if $L^S(\frac{3}{2} + m, \Pi \times \Pi')$ is non-zero (e.g., if Π is unitary and $m \neq -1$), then the quotient of consecutive critical L -values satisfies

$$\frac{1}{\Omega_{\Pi_\infty, \Pi'_\infty}(m, m+1)} \frac{L^S(\frac{1}{2} + m, \Pi \times \Pi')}{L^S(\frac{3}{2} + m, \Pi \times \Pi')} \in \mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f).$$

Acknowledgements Open access funding provided by Austrian Science Fund (FWF). I would like to thank Jie Lin, who encouraged me to write up this short note as an individual paper. I would also like to thank the referee for her/his helpful comments, which improved the presentation of this article.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Baruch, E.M.: A proof of Kirillov’s conjecture. *Ann. Math.* **158**, 207–252 (2003)
2. Bernstein, J.N.: P-invariant distributions on $GL(N)$ and the classification of unitary representations of $GL(N)$ (non-Archimedean case). In: *Lie Group Representations, II*, (College Park, Md., 1982/1983) *Lecture Notes in Mathematics*, vol. 1041, pp. 50–102 (1984)
3. Borel, A.: Cohomology and spectrum of an arithmetic group. In: *Proceedings of a Conference on Operator Algebras and Group Representations*, Neptun, Rumania (1980), Pitman, pp. 28–45 (1983)
4. Borel, A., Casselman, W.: L^2 -Cohomology of locally symmetric manifolds of finite volume. *Duke Math. J.* **50**, 625–647 (1983)
5. Borel, A., Wallach, N.: Continuous cohomology, discrete subgroups and representations of reductive groups. *Ann. Math. Stud.* **94** (Princeton Univ. Press, New Jersey) (1980)
6. Clozel, L.: Motifs et Formes Automorphes: Applications du Principe de Functorialité. In: Clozel, L., Milne, J.S. (eds.) *Automorphic Forms, Shimura Varieties, and L-functions, Vol. I*, *Perspect. Math.*, vol. 10 (Ann Arbor, MI, 1988) Academic Press, Boston, MA, pp. 77–159 (1990)
7. Cogdell, J.W., Piatetski-Shapiro, I.I., Shahidi, F.: Functoriality for the Quasisplit Classical Groups. In: Arthur, J., Cogdell, J.W., Gelbart, S., Goldberg, D., Ramakrishnan, D., Yu, J.-K. (eds.) *On Certain L-functions*, *Clay Mathematics Proceedings*, vol. 13 (West Lafayette, IN, 2007) AMS, Providence, RI, pp. 117–140 (2011)
8. Enright, T.J.: Relative Lie algebra cohomology and unitary representations of complex Lie groups. *Duke Math. J.* **46**, 513–525 (1979)
9. Franke, J.: Harmonic analysis in weighted L_2 -spaces. *Ann. Sci. de l’ENS* **31**, 181–279 (1998)

10. Franke, J., Schwermer, J.: A decomposition of spaces of automorphic forms, and the Eisenstein cohomology of arithmetic groups. *Math. Ann.* **311**, 765–790 (1998)
11. Grobner, H.: Residues of Eisenstein series and the automorphic cohomology of reductive groups. *Compos. Math.* **149**, 1061–1090 (2013)
12. Grobner, H., Harris, M.: Whittaker periods, motivic periods, and special values of tensor product L -functions. *J. Inst. Math. Jussieu* **15**, 711–769 (2016)
13. Grobner, H., Raghuram, A.: On some arithmetic properties of automorphic forms of GL_m over a division algebra. *Int. J. Number Theory* **10**, 963–1013 (2014)
14. Harder, G.: General aspects in the theory of modular symbols. In: Bertin, M.-J. (ed.) *Séminaire de Théorie des Nombres, Paris 1981–82*. Progress in Mathematics 38 (Boston, Basel, Stuttgart 1983), pp. 73–88
15. Harder, G., Raghuram, A.: Eisenstein cohomology for $GL(N)$ and ratios of critical values of Rankin–Selberg L -functions. With Appendix 1 by Uwe Weselmann and Appendix 2 by Chandrasheel Bhagwat and A. Raghuram, preprint (2017)
16. Jacquet, H., Langlands, R.P.: *Automorphic Forms on $GL(2)$* , Lecture Notes in Mathematics, vol. 114. Springer (1970)
17. Jacquet, H., Piatetski-Shapiro, I., Shalika, J.: Conducteur des représentations du groupe linéaire. *Math. Ann.* **256**, 199–214 (1981)
18. Jacquet, H., Piatetski-Shapiro, I.I., Shalika, J.: Rankin–Selberg convolutions. *Am. J. Math.* **105**, 367–464 (1983)
19. Jacquet, H., Shalika, J.: On Euler products and the classification of automorphic representations II. *Am. J. Math.* **103**, 777–815 (1981)
20. Jacquet, H., Shalika, J.: The Whittaker models of induced representations. *Pac. J. Math.* **109**, 107–120 (1983)
21. Langlands, R.P.: Automorphic representations, Shimura varieties, and motives. Ein Märchen. In: Proc. Sympos. Pure Math., Vol. XXXIII, part II, AMS, Providence, R.I., pp. 205–246 (1979)
22. Mahnkopf, J.: Cohomology of arithmetic groups, parabolic subgroups and the special values of automorphic L -Functions on $GL(n)$. *J. Inst. Math. Jussieu* **4**, 553–637 (2005)
23. Mahnkopf, J.: Modular symbols and values of L -functions on GL_3 . *J. Reine Angew. Math.* **497**, 91–112 (1998)
24. Mœglin, C., Waldspurger, J.-L.: Le Spectre Résiduel de $GL(n)$. *Ann. Sci. de l'ENS* **22**, 605–674 (1989)
25. Raghuram, A.: Critical values of Rankin–Selberg L -functions for $GL_n \times GL_{n-1}$ and the symmetric cube L -functions for GL_2 . *Forum Math.* **28**, 457–489 (2016)
26. Raghuram, A.: On the special values of certain Rankin–Selberg L -functions and applications to odd symmetric power L -functions of modular forms. *Int. Math. Res. Not.* **2**, 334–372 (2010). <https://doi.org/10.1093/imrn/rnp127>
27. Raghuram, A., Shahidi, F.: On certain period relations for cusp forms on GL_n . *Int. Math. Res. Not.* (2008). <https://doi.org/10.1093/imrn/rnn077>
28. Schwermer, J.: Kohomologie arithmetisch definierter Gruppen und Eisensteinreihen, Lecture Notes in Mathematics, vol. 988. Springer (1983)
29. Shahidi, F.: Eisenstein series and Automorphic L -functions, Colloquium publications 58, AMS (2010)
30. Shalika, J.A.: The multiplicity one theorem for GL_n . *Ann. Math.* **100**, 171–193 (1974)
31. Vogan Jr., D.A.: The unitary dual of $GL(n)$ over an Archimedean field. *Invent. Math.* **83**, 449–505 (1986)
32. Waldspurger, J.-L.: Quelques propriétés arithmétiques de certaines formes automorphes sur $GL(2)$. *Comp. Math.* **54**, 121–171 (1985)