

1 ***Re-evaluating homoploid*** 2 ***reticulate evolution in the*** 3 ***annual sunflowers.***

4 5 **Authors:**

6 Gregory L. Owens^{1*}

7 Kaichi Huang²

8 Marco Todesco²

9 Loren H. Rieseberg²

10

11 ¹: Department of Biology, University of Victoria, BC, Canada

12 ²: Department of Botany and Beaty Biodiversity Center, University of British Columbia, BC,
13 Canada

14 *Corresponding author: grego@uvic.ca

15 **Abstract:**

16 Sunflowers of the genus *Helianthus* are models for hybridization research and contain three
17 of the best studied examples of homoploid hybrid speciation. To understand the broader picture
18 of hybridization within the annual sunflowers, we used whole genome resequencing to conduct
19 a phylogenomic analysis and test for gene flow between lineages. We find that all annual
20 sunflower species tested have evidence of admixture, suggesting hybridization was common

21 during the radiation of the genus. Support for the major species tree decreases with
22 recombination rate, consistent with hybridization and introgression contributing to discordant
23 topologies. Admixture graphs found hybridization to be associated with the origins of the three
24 putative hybrid species (*H. anomalus*, *H. deserticola*, and *H. paradoxus*). However, the
25 hybridization events are more ancient than suggested by previous work. Furthermore, *H.*
26 *anomalus* and *H. deserticola* appear to have arisen from a single hybridization event involving
27 an unexpected donor, rather than through multiple independent events as previously proposed.
28 Using a broader data set that covers the whole *Helianthus* genus, including perennial species,
29 we find that signals of introgression span the genus and beyond, suggesting highly divergent
30 introgression and/or the sorting of ancient haplotypes. Thus, *Helianthus* can be viewed as a
31 syngameon in which largely reproductively isolated species are linked together by occasional or
32 frequent gene flow.

33 **Introduction**

34 The evolutionary importance of hybridization and introgression has been explored since
35 Darwin's time (Darwin 1859). While initially zoologists thought that hybridization was relatively
36 rare and unimportant (Mayr 1963), botanists have long emphasized the role of hybridization and
37 introgression in both adaptation and speciation (Anderson 1949; Heiser 1949; Anderson and
38 Stebbins 1954). Genomics eventually proved botanists correct in this regard; evidence of past
39 hybridization is common in both animals and plants (Figueiró et al. 2017; Fontaine et al. 2015;
40 Mallet et al. 2016; Calfee et al. 2021; Dagilis et al. 2021). In several cases, alleles responsible
41 for ecologically important traits have been found to be acquired through introgression (Suarez-
42 Gonzalez et al. 2018; Jones et al. 2018; Oziolor et al. 2019), supporting the notion that
43 introgression not only occurs, but can be evolutionarily important.

44 Hybridization can also result in the formation of new species through homoploid hybrid
45 speciation, in which an admixed lineage acquires reproductive isolation from its progenitors

46 through a combination of alleles or chromosomal rearrangements from both parents (Grant
47 1958; Schumer et al. 2014; Stebbins 1957). One way this can occur is if two parental species
48 differ at both habitat and mate choice. A hybrid population between these two may have
49 alternate parental alleles for these two traits, and therefore be separated from one parental
50 species by habitat and the other by mate choice (Wang et al., 2021). Hybrid speciation can also
51 occur when hybridization results in the ability to colonize a new ecological niche, reject parental
52 species as potential mates or have unique combinations of chromosomal rearrangements
53 (Gross and Rieseberg 2005; Melo et al. 2009; Rieseberg et al. 1995). Schumer et al. (2014)
54 argued that three criteria must be satisfied to demonstrate homoploid hybrid speciation: (1)
55 reproductive isolation from parental species, (2) prior admixture, and (3) evidence that
56 reproductive barriers arose via hybridization. Evaluation of the natural hybridization literature
57 indicates that while admixed lineages are relatively common (criterion 2), few case studies
58 satisfy criteria 1 and 3 (Schumer et al. 2014). Three examples of hybrid species thought to
59 address all three criteria are within the sunflower genus, *Helianthus*.

60 The widespread species *Helianthus annuus* and *H. petiolaris* are thought to be the parents
61 of three independent homoploid hybrid species, *H. anomalus*, *H. deserticola* and *H. paradoxus*.
62 These were initially identified as hybrid species through early genetic studies that found they
63 had a mixture of chloroplast DNA, allozyme and ribosomal DNA markers from both parental
64 species (Rieseberg et al. 1990a; Rieseberg 1991). The hybrid species have strong ecological
65 separation based on habitat choice; *Helianthus anomalus* occurs on sand dunes, *H. deserticola*
66 on sandy soil of the desert floor and *H. paradoxus* in brackish salt marshes (Heiser et al. 1969).
67 Comparative QTL mapping of hybrids and parental species suggested that this ecological
68 separation was achieved by combinations of parental alleles that allowed for extreme
69 (transgressive) phenotypes (Rieseberg et al. 2003). Subsequent experiments found that such
70 transgressive phenotypes were favored when segregating hybrids of *H. annuus*-*H. petiolaris*

71 were transplanted into the natural habitats of the three ancient hybrid species, with some
72 synthetic hybrids rivaling the ancient hybrids in fitness (Lexer et al. 2003; Gross et al. 2004;
73 Ludwig et al. 2004).

74 In addition to ecological isolation, the three hybrid species are partially reproductively
75 isolated by hybrid pollen sterility caused mainly by chromosomal rearrangements (Lai et al.
76 2005). This was hypothesized to be due to the hybrid species having a combination of parental
77 chromosomal rearrangements as well as unique changes (Rieseberg et al. 1995; Lai et al.,
78 2005). Synthetic hybrids of *H. annuus*-*H. petiolaris* not only recovered fertility in a small number
79 of generations, but also were more compatible with the putative hybrid species than were the
80 parental species (Rieseberg et al. 1996; Rieseberg 2000, 2006). Microsatellite, chloroplast, and
81 karyotypic analyses suggested all hybrid species originated between 63,000 and 208,000
82 generations before present and a single (*H. paradoxus*) or multiple (*H. anomalus* and *H.*
83 *deserticola*) origins depending on the species (Schwarzbach and Rieseberg 2002; Welch and
84 Rieseberg 2002; Gross et al. 2003). The strong link between admixture and reproductive
85 isolation, in terms of ecology and karyotype, comparisons with experimental hybrid lineages,
86 and partial replication across three species have made *Helianthus* a model for understanding
87 homoploid hybrid speciation.

88 Although the *Helianthus* hybrid species were probed with cutting-edge molecular markers
89 when initially identified, they have been relatively unexamined in the genomics age. Studies
90 using Genotyping-By-Sequencing grouped *H. anomalus* and *H. deserticola* together in a
91 phylogenetic tree but did not probe further (Baute et al. 2016). Moody and Rieseberg (2012)
92 sequenced ten nuclear genes for multiple individuals of most annual sunflowers, including the
93 hybrid species. They found extensive discordance between individual gene trees and notably
94 that *H. anomalus* and *H. deserticola* grouped together and that *H. paradoxus* grouped with *H.*
95 *annuus*, although the overall species tree did not agree with previous expectations. The most

96 recent phylogeny of *Helianthus* did not include any hybrid species (Stephens et al., 2015). In
97 each of these cases, modern approaches for identifying gene flow and hybrid ancestry were not
98 performed. This is important because sunflower species are both young — *Helianthus* crown
99 age is 3.6 mya — and generally have high effective population size (Strasburg and Rieseberg
100 2008). This means that incomplete lineage sorting (ILS) is likely high and discordant topologies
101 that could be produced by hybridization may also be present without any interspecies gene flow.
102 In addition, signals of hybridization can sometimes derive from unexpected sources (Owens et
103 al. 2021).

104 Beyond the three well-characterized *Helianthus* hybrid species, numerous other cases of
105 hybridization and introgression have been reported in the genus (Heiser et al. 1969; Rieseberg
106 et al. 1991), leading Rieseberg (1991) to conclude that evolution in the group “must be
107 considered reticulate, rather than exclusively dichotomous and branching.” While many of these
108 early cases – which were based on relatively limited evidence – have been subsequently
109 confirmed with genomic data (Kane et al. 2009; Owens et al. 2016; Mondon et al. 2018; Lee-
110 Yaw et al. 2019; Zhang et al. 2019), others have not (Owens et al. 2021), and several remain to
111 be tested. This highlights that the combination of modern genomic datasets and modern
112 phylogenetic techniques allow for hypotheses about hybridization to be rigorously tested
113 (Hibbins and Hahn 2022).

114 Here we use modern phylogenomic techniques to assess the extent of admixture during the
115 evolution of the genus. To do this, we first use whole genome sequencing data for most annual
116 sunflowers to infer the genome-wide phylogeny of this group. We then apply tests of
117 introgression and gene flow across the annual clade, including the three hybrid species, *H.*
118 *anomalus*, *H. deserticola* and *H. paradoxus*. This allows us to revisit criterion 2 of homoploid
119 hybrid speciation for these three species, and to gain greater insight regarding their ancestry
120 and their relationship to the broader patterns of admixture within the annual sunflower clade.

121 Lastly, using a previously published phylogenomic data (Stephens et al., 2015), we extend our
122 analyses across the entire genus.

123 **Results:**

124 **Data filtering**

125 We generated whole genome sequence data for two *Helianthus paradoxus* samples. We
126 called variants against the *H. annuus* HA412-HOV2 reference genome for these samples as well
127 as previously sequenced data for two samples of eight other annual species, or subspecies, and
128 four outgroups (supplementary table 1; Todesco et al. 2020). We retained 5,877,513 variable
129 and 28,274,146 invariant sites after filtering for quality, mappability and call rate (supplementary
130 figure 1). We divided the genome into a total of 11,797 10 kbp non-overlapping windows and
131 retained 3,267 that had $\geq 2,000$ called bases.

132 **Species and gene trees**

133 The species topology for non-hybrid taxa from ASTRAL (fig. 1A,B) was mostly consistent
134 with previous phylogenies that used fewer markers (Stephens et al. 2015; Baute et al. 2016;
135 Zhang et al. 2019). Quadripartition support at nodes ranged from 38 to 84%, suggesting
136 substantial variation in topology at gene trees. The maximum likelihood concatenated tree also
137 found the same topology as ASTRAL and monophyly for all samples of the same species (fig.
138 1B).

139 As outlined in the introduction, the three putative homoploid hybrid taxa, *H. paradoxus*, *H.*
140 *deserticola* and *H. anomalus* are thought to have originated from hybridization between *H.*
141 *annuus* and *H. petiolaris* (Rieseberg 1991). Since ASTRAL does not account for hybridization,
142 we would expect the potential hybrid species to group near to the dominant parental species.
143 We found that *H. paradoxus* grouped with *H. annuus* and its sister species *H. argophyllus*,
144 whereas *H. anomalus* and *H. deserticola* grouped with *H. niveus canescens* (fig. 1B).

145 Quadripartition support for these groupings is low (each 39%), as expected for hybrid lineages,
146 although this was not unique to branches involving the previously identified hybrid species.

147 We further examined variation in gene tree topology by calculating the gene concordance
148 factor for each branch on our concatenated tree using gene trees. We found relatively low
149 support at branches separating species compared to branches separating samples within
150 species (fig. 1C). When including all samples, each tree was unique. When we subsampled tips
151 down to a single individual per species we found higher agreement; the most common tree was
152 found 6 times out of 3,242 trees.


153 **Gene tree discordance**

154 To explore why gene concordance values were low, we compared the proportion of trees
155 matching the two alternate topologies for every quadripartition using IQ-TREE2's gene
156 concordance measure (Minh et al. 2020). Under purely incomplete lineage sorting, we expect
157 that both alternate topologies should be found at equal proportions, while introgression will lead
158 to imbalances. We found that at five nodes within the annual species, there was significant (X-
159 squared, $p < 0.05$) imbalance in alternate topologies (fig. 1C).

160 Selection against introgressed ancestry can lead to reduced introgression in regions of low
161 recombination (Schumer et al. 2018; Martin et al. 2019). We explored this by using a binomial
162 regression to quantify the relationship between support for the species topology and the
163 recombination rate (as identified from the *H. annuus* genetic map) at each node with > 10%
164 support for alternate topologies. In other words, are genomic regions with lower recombination
165 more likely to show the species topology? We found a significant negative effect of
166 recombination rate ($\beta_1 = -0.30$, $p = e^{-14}$) on support for the species topology (fig. 1C,D).
167 Interestingly, we find a significant interaction between node and recombination rate at six nodes.
168 At four nodes, most of which include the *H. anomalus/H. deserticola* clade, the relationship is

169 less negative, or is positive (fig. 1D). At the node separating the *H. annuus* samples, the
 170 relationship with recombination is most strongly negative.

171 *Helianthus niveus canescens* was previously considered a variety of *H. petiolaris*, with which
 172 it intergrades for part of its range (Heiser et al. 1969). Our species topology places *H. niveus*
 173 *canescens* with *H. anomalus* and *H. deserticola* instead of *H. petiolaris*, but with relatively low
 174 support and overall less support at low recombination regions (fig. 1D). We counted the number
 175 of trees where *H. niveus canescens* grouped with *H. petiolaris petiolaris* and *H. petiolaris fallax*
 176 to the exclusion of *H. anomalus* and *H. deserticola*, as well as the reverse (supplementary figure
 177 2). We found more support for *H. niveus canescens* together with *H. petiolaris*.



178

179 **Figure 1: Phylogenetics of annual sunflowers.** A) The prior hypothesis of annual species

180 relationships (Rieseberg 1991; Stephens et al. 2015). B) Phylogenetic tree from combined 10

181 kbp gene trees used as input for ASTRAL species topology. Quartet support plotted on
182 branches. C) Concatenated maximum likelihood tree using IQ-TREE2. Gene concordance
183 plotted on nodes. Bolded values have significantly imbalanced secondary topologies. D) The
184 relationship between gene concordance proportion and recombination quantile for each node
185 with <90% overall gene concordance. Significant interactions with node ID are highlighted here
186 and on panel C.


187 ***Admixture in the annual sunflowers***

188 To assess the extent of hybridization during the evolution of the annual sunflowers, we used
189 comprehensive approaches to test for admixture across all species. The f_{branch} analysis, which
190 uses the f_4 -ratio test to identify introgression within a phylogenetic framework, found many
191 significant signals (fig. 2A, supplementary table 2). The two strongest signals show that *H.*
192 *petiolaris* is closer to all other species, when compared to *H. debilis*, and *H. niveus canescens* is
193 closer to *H. petiolaris* and *H. debilis*, than *H. anomalus* and *H. deserticola* are. The f_{branch}
194 patterns are dependent on the underlying phylogeny, so to explore this effect we repeated f_{branch}
195 using the top nine most common species tree topologies (supplementary fig. 3). While no
196 topology completely removes significant f_{branch} signal, this shows that introgression signals from
197 f_{branch} can come from either close phylogenetic position or introgression. For example, in three of
198 the top nine trees, *H. paradoxus* is placed within the *H. petiolaris* clade, but its f_{branch} signal
199 shows it is much more similar to *H. annuus* and *H. argophyllus* than others in the *H. petiolaris*
200 clade (supplementary fig. 3). The three hybrid species do not stand out in terms of the extent of
201 admixture.

202 A different picture comes from the TreeMix analysis, which uses the covariance of
203 population allele frequency to estimate a phylogeny and migration edges. We found that three
204 was the optimal number of migration edges and explained 99.9% of variance. Above four
205 edges, there was diminishing returns in likelihood gain and variance explained (supplementary

206 figure 4). Admixture was found to be associated with the origins of the three putative hybrid
207 species in the most likely TreeMix graphs at both three and four edges (supplementary figure 5).
208 The origin of *H. paradoxus* appears to involve ancient hybridization between the ancestor of the
209 *H. annuus* clade and *H. petiolaris*, although this admixture is inherited by the other sampled
210 members of the *H. annuus* clade. Likewise, there is an admixture event between a basal lineage
211 of the *H. annuus* clade and the ancestor of *H. deserticola* and *H. anomalus*. The third signal in
212 common between the top three admixture graphs was between *H. debilis* and *H. petiolaris*.
213 However, this likely is an artifact of misplacing *H. debilis* as basal in the tree and may instead
214 suggest that it has ancestry from a lineage at the base of the annual sunflower clade. When we
215 look other graphs that are close in likelihood to the top graphs, we frequently find admixture
216 between *H. niveus canescens* and *H. petiolaris fallax*, although this results in the loss of the
217 admixture event associated with *H. anomalus* and *H. deserticola*.

218 We also used ADMIXTOOLS2, which builds an admixture graph that explains f-statistics
219 between populations. Admixture scenarios seen using ADMIXTOOLS2 have both similarities
220 and differences with the TreeMix graphs. Out-of-sample scores decreased with increasing
221 admixture edges from one to two, and from two to three, but not from three to four, suggesting
222 that three admixture edges explain much of the f_2 incongruence (supplementary figure 6). While
223 bootstrapping was unable to distinguish between the top three graphs with three admixture
224 nodes ($p > 0.1$), they each present a similar pattern that distinguishes them from the species
225 tree (fig. 2C). The strongest signal is associated with origin of two of the putative hybrid species
226 (*H. anomalus* and *H. deserticola*), but from admixture with the ancestor of *H. niveus canescens*
227 and an unsampled or extinct basal lineage. *Helianthus petiolaris* is placed at the base of the
228 *annuus* clade in the ADMIXTOOLS2 graph, implying that it shares ancestry with members of the
229 *annuus* clade, including *H. paradoxus*. In addition, recent admixture is seen between *H. niveus*
230 *canescens* and *H. petiolaris fallax* similar to that observed in the TreeMix analysis.



232 **Figure 2: Introgression across sunflowers.** A) F_{branch} statistic for annual species using the
233 ASTRAL species topology. B) The four best supported TreeMix graph with three admixture
234 edges. Higher likelihood score is better. C) The best supported admixture graph from
235 ADMIXTOOLS2 with three admixture edges. Branch lengths are not to scale. Lower score is
236 better. D) ABBA-BABA with the groupings [*H. annuus* OR *H. argophyllus*], other annual species,
237 perennial species, *Phoebanthus*. Positive values indicate more ABBA counts. E) ABBA-BABA
238 with the groupings *H. maximiliani*, other perennials, annual species, *Phoebanthus*.


239 **Triplet tests of hybrid ancestry**

240 While the admixture graphs (above) detected signals of hybridization associated with the
241 origins of the three hybrid species, there were differences relative to previous hypotheses
242 (Rieseberg 1991). Most notably, the origin of *H. anomalus* and *H. deserticola* appears to involve
243 hybridization with the ancestor of *H. niveus canescens* rather *H. petiolaris fallax*. Also, there was
244 uncertainty as to whether the signal of admixture associated with the origin of *H. paradoxus* was
245 independent from other members of the *annuus clade*. Therefore, we explicitly tested
246 hypotheses of hybrid ancestry for *H. anomalus*, *H. deserticola* and *H. paradoxus* against their
247 putative parents. In the case of *H. paradoxus*, the testing is relative to *H. annuus* and *H.*
248 *petiolaris fallax*, whereas for *H. anomalus* and *H. deserticola*, the putative parents are *H. annuus*
249 and *H. niveus canescens*. For each test, we expect to see that the hybrid species is most
250 closely related to the major parental species, and that tests of introgression show that the
251 dominant signal of introgression involves the hybrid species and its minor parental species,
252 rather than the parental species. We used four different approaches. Patterson's D compares
253 the amounts of shared derived alleles in a trio (Green et al. 2010; Malinsky et al. 2021). In this
254 case, we are using Twisst to compare the counts of different gene tree topologies (Martin and
255 Van Belleghem 2017). The branch length test compares the branch lengths in alternate
256 topology gene trees. Lastly, QuIBL models ILS and introgression based on branch lengths. We

257 note that the results of triplet tests should be interpreted with caution because of the long history
258 of introgression between the putative parental clades (Yatabe et al. 2007; Strasburg and
259 Rieseberg 2008; Kane et al. 2009). For the topology-based tests (Patterson's D and Twisst),
260 this means that the extent of introgression from the minor parent must be significantly greater
261 than that between the parental lineages. For the distance-based tests (branch length and
262 QuIBL), both the extent and the timing of introgression becomes critical, as the distance-based
263 tests have little power for detecting ancient admixture.

264 For *H. paradoxus*, we consistently find *H. annuus* is the closest species, but inconsistent
265 evidence of admixture with *H. petiolaris fallax* (fig. 3). Patterson's D finds significant *H.*
266 *paradoxus* – *H. petiolaris fallax* admixture. The results from Twisst are in the same direction, but
267 are not significant. In contrast, the branch length test supports greater *H. annuus* – *H. petiolaris*
268 *fallax* introgression and QuIBL supports neither (fig. 3). We find qualitatively similar results when
269 using *H. petiolaris petiolaris* as a potential parental species. The branch length test is more
270 sensitive to recent introgression, such as that ongoing between *H. annuus* and *H. petiolaris*
271 (Yatabe et al. 2007), whereas D has greater power to detect ancient hybridization, perhaps
272 accounting for the apparent discrepancy between the different tests. At the genomic window
273 level, there is considerable variation in D with values ranging from -1 to 1 (supplementary figure
274 7).

275 For *H. anomalus* and *H. deserticola*, we see a consistent pattern for all four tests; they are
276 more closely related to *H. niveus canescens* and there is more admixture between the potential
277 parental species, than between the hybrid species and *H. annuus* (fig. 3). This could indicate
278 that the hybrid species are ancient, and that the minor parent is at the base of the *H. annuus*
279 clade (as suggested by TreeMix), and therefore has a relatively small contribution from
280 contemporary *H. annuus*, or that the other parent of *H. anomalus* and *H. deserticola* is a now
281 extinct basal lineage, as suggested in the ADMIXTOOLS2 graph.



282

283 **Figure 3: Triplet tests of previous hypotheses of hybrid ancestry.** A-C) Counts of
 284 shared derived alleles and topological weighting for proposed hybrid species. P-values from
 285 block bootstrapping and chi-squared tests for ABBA-BABA and Twisst respectively. D-F) Branch
 286 Length Test of introgression showing genetic distance between taxa at gene trees with

287 secondary topologies. G-I) QuIBL delta BIC values and interpretation. Lower values indicate
288 more support for introgression.

289 ***Admixture across the sunflower genus***

290 To assess introgression across the rest of the genus, as well as to confirm our findings
291 within the annual clade, we expanded our analysis of gene flow using a previously published
292 sequence capture dataset spanning the genus. With this data, we used Dsuite to calculate f_{branch}
293 and D for all trios (supplementary figure 8, supplementary table 3). We found evidence of
294 introgression between perennial species, including *H. verticillatus* – *H. giganteus*, *H. divaricatus*
295 – *H. arizonensis* and *H. atrorubens* – *H. mollis*. This dataset also affirms the signal of
296 introgression between *H. petiolaris* and both *H. niveus* and *H. annuus*. However, no
297 introgression was found involving *H. debilis*. Surprisingly, we also find introgression-like signals
298 between annual and perennial species. *Helianthus maximilliani* shares more derived alleles with
299 annual species than other perennials (fig. 2E). Strangely, both *H. annuus* and *H. argophyllus*
300 share fewer derived alleles with perennials than other annual species, which is consistent with
301 widespread annual-perennial introgression excluding *H. annuus*/*H. argophyllus*, or introgression
302 between the *H. annuus*/*H. argophyllus* clade and the outgroup *Phoebanthus* (fig. 2D).


303

304

305

306

307



308

309 **Figure 4: The proposed annual sunflower phylogeny with introgression events**
310 **identified in the present study.** L1 represents an unknown lineage seen through admixture
311 analyses. Dotted lines indicate signatures of admixture seen in different analyses.

312 ***Discussion:***

313 For three-fourths of a century, *Helianthus* sunflowers have served as a model for the study
314 of hybridization and its evolutionary role (Heiser 1947; Rieseberg 1991; Todesco et al. 2020).
315 Early studies documented hybrid swarms between different sunflower species (Heiser 1947;
316 Heiser 1949; Heiser 1951), demonstrated that chromosomal rearrangements contributed to
317 hybrid sterility (Chandler et al. 1986; Lai et al. 2005), and speculated on possible ecotypes and
318 species that might be the products of hybridization (Heiser 1949; Heiser 1958; Rieseberg 1991).
319 Later molecular phylogenetic work and early low-resolution genomic studies supported some of
320 these hypotheses (Rieseberg et al. 1990a; Yatabe et al. 2005), but not others (Rieseberg et al.
321 1988), and also suggested new hypotheses, including the putative origins of three homoploid
322 hybrid species (Rieseberg 1991). Here, we have taken advantage of new high resolution

323 genomic data and computational methods to further explore the role of hybridization in
324 sunflower evolution. We pay particular attention to the three homoploid hybrid species, but we
325 also examine other well-studied cases of hybridization in the *Helianthus* (and potential new
326 cases) with the goal of reconciling the new data and analyses with previous work. Lastly, we
327 explore the effects of variation in recombination rate on tree topology and introgression.

328 ***Hybridization in sunflowers***

329 The whole genome phylogeny for sunflower is consistent with that found by previous multi-
330 locus studies (Stephens et al. 2015; Baute et al. 2016; Zhang et al. 2019). Leveraging this
331 phylogeny, we employed several different methods to search for possible cases of admixture
332 during the evolution of the genus. Remarkably, all species included in our study appear to be
333 admixed, although the level of admixture depends in part on the particular analysis performed
334 (fig. 2,4). Overall, the strongest signal of hybridization occurs between two subspecies of *H.*
335 *petiolaris* and *H. annuus*. *Helianthus petiolaris* is sympatric with *H. annuus* and contemporary
336 hybridization is well-documented (Heiser 1947; Rieseberg et al. 1998). Previous studies have
337 documented “rampant” introgression between the species (Yatabe et al. 2005) and have
338 suggested that interspecific gene flow was common in the past as well (Strasburg and
339 Rieseberg 2008).

340 Another strong signal of admixture is found between *H. petiolaris* and *H. niveus canescens*,
341 confirming a previous report (Zhang et al. 2019). *Helianthus petiolaris* intergrades with *H. niveus*
342 *canescens* phenotypically, and Heiser et al. (1969) suggested that it might be due to
343 hybridization. Although both types of phylogenetic analysis suggested *H. niveus canescens*
344 grouped with *H. anomalus* and *H. deserticola* rather than *H. petiolaris*, there were more gene
345 trees that placed *H. niveus canescens* with *H. petiolaris*. Considering the weight of evidence, we
346 support that *H. niveus canescens* was involved with admixture of the *H. anomalus/deserticola*
347 ancestor but was originally a member of the *H. petiolaris* clade. The beach sunflower, *H. debilis*,

348 is sister to *H. petiolaris* in our tree, but f_{branch} shows that it is less similar to all other annual
349 sunflower species (fig. 2A). This implies that it has ancestry from a lineage at the base of the
350 annual sunflowers, as also suggested by TreeMix. Lastly, both the TreeMix and ADMIXTOOLS2
351 graphs suggest a shared hybrid origin for *H. anomalus* and *H. deserticola*, which we will discuss
352 in more length below.

353 We used an alternate dataset generated by Stephens et al. (2015) to assess introgression
354 across the entire *Helianthus* genus. We found evidence of hybridization between *H. verticillatus*
355 and *H. giganteus*. *Helianthus verticillatus* was once thought to be a hybrid species but was later
356 disproven by molecular data (Heiser et al. 1969; Ellis et al. 2006). Interestingly, *H. verticillatus*
357 was previously thought to be a subspecies of *H. giganteus*, and the two species overlap in
358 range (Farwall 1916). Similarly, introgression between *H. mollis* and *H. atrorubens* is highly
359 plausible considering they are known to hybridize (Beatley 1969). In contrast, the proposed
360 introgression between *H. divaricatus* and *H. arizonensis* is difficult to explain because the
361 species do not share any current range and there are no plausible proxy ancestry donors.

362 Surprisingly, there was identified introgression between the annual and perennial clades,
363 despite strong post-pollination reproductive barriers between them. Most crosses between
364 annual and perennial sunflower species fail to set seed, and the few successful hybrids that
365 have been reported typically have low pollen viability (Heiser et al. 1969). Crosses involving the
366 hexaploid perennials, represent an exception to this rule (Atlagić and Terzić 2006), but we found
367 the strongest signal of introgression with a diploid perennial *H. maximiliani*, which is
368 incompatible with members of the annual clade (Henn et al. 1998). Thus, we likely are seeing
369 the outcome of ancient hybridization, which occurred before reproductive isolation was
370 complete. However, the most unexpected pattern is a signal of ancestral allele sharing between
371 *H. annuus/argophyllus* and the outgroup, *Pheobanthus* (fig. 2E). These two lineages diverged
372 2.5-5.4 mya, and are therefore expected to be completely reproductively isolated (Owens and

373 Rieseberg 2014; Mason 2018), again suggesting that the signal of admixture is from ancient
374 hybridization with a now extinct species. Nonetheless, we recommend experimental crosses be
375 undertaken between *Pheobanthus* and members of the annual clade to see if hybrids can be
376 made. It is also possible that this signal represents a change in evolutionary rate in *H. annuus*
377 and *H. argophyllus*. This has been proposed to explain similar patterns of introgression signals
378 in *Papilio* butterflies (Xiong et al. 2022). More broadly, our results suggest that admixture has
379 been common in *Helianthus* throughout its evolutionary history, and that it has had a profound
380 effect on phylogenomic relationships in the genus (fig. 4). Thus, *Helianthus* can be viewed as a
381 syngameon, in which gene flow connects otherwise distinct species (Grant 1981). Our findings
382 also address earlier concerns that hybridization in *Helianthus* might be a recent consequence of
383 anthropogenic disturbance and therefore of little significance to its long-term evolution
384 (Schemske 2000). Whether these ancient hybridization events triggered diversification, as
385 suggested in other groups (e.g., Meier et al. 2017), remains unclear.

386 ***Ancestry of putative homoploid hybrid species***

387 The TreeMix and ADMIXTOOLS2 graphs indicate that the three hybrid species are admixed,
388 thereby fulfilling the second criterion for homoploid hybrid speciation. However, there are
389 differences relative to the original scenario put forward by Rieseberg (1991). In particular, the
390 hybridization events associated with the origins of the hybrid species appear to have occurred
391 further back in time than previously believed. For *H. paradoxus*, a potential hybrid speciation
392 event is older than previously anticipated (Welch and Rieseberg 2002) since individual gene
393 trees place *H. paradoxus* most often at the base of the annuus clade, therefore at least older
394 than 1 mya. For *H. anomalus* and *H. deserticola*, our phylogenomic results suggest they are
395 sister species, and share an admixture close in time to their origin. This would also place a
396 potential hybrid speciation event further in the past than previous work suggested (Schwarzbach

397 and Rieseberg 2002; Gross et al. 2003; see supplementary discussion for reconciliation with
398 previous microsatellite dating of the hybrid species origin).

399 Another important difference is that *H. anomalus* and *H. deserticola* are sister species and
400 their major parent appears to be *H. niveus canescens* rather than *H. petiolaris fallax* as
401 originally proposed (Rieseberg 1991). While this was unexpected, *H. niveus canescens*
402 intergrades phenotypically with *H. petiolaris fallax* and the two taxa share chloroplast and
403 nuclear ribosomal DNA haplotypes, which were the markers employed to hypothesize a hybrid
404 origin in the first place (Heiser et al. 1969; Beckstrom-Sternberg et al. 1991). As a consequence,
405 some taxonomic treatments have considered *H. niveus canescens* to be a variety of *H.*
406 *petiolaris* (Blake 1942; Schilling 2020). The present study indicates that *H. niveus canescens* is
407 genetically distinct from *H. petiolaris fallax*, although there is considerable admixture between
408 them, which might account for their phenotypic resemblance (Zhang et al. 2019). Fortunately,
409 recognition that the ancestor of *H. niveus canescens* was a parent of *H. anomalus* and *H.*
410 *deserticola* has little impact on the interpretation of previous work. The population of *H.*
411 *petiolaris fallax* that was employed for most of the experimental studies of hybrid speciation
412 came from the zone of intergradation with *H. niveus canescens* (Rieseberg et al. 2003; Lexer et
413 al. 2003; Gross et al. 2004; Ludwig et al. 2004). The taxa are indistinguishable morphologically
414 and ecologically in this area, and may recapitulate the ancestor of *H. niveus canescens*.

415 Despite the power of the phylogenomic analyses, some uncertainties remain (see
416 supplementary discussion about the challenges for interpreting admixture signals in sunflower).
417 For *H. paradoxus*, it is unclear whether the signal of admixture associated with its origin is
418 independent from other members of the *annuus clade*. The triplet tests of introgression provide
419 indirect support for independence. For D, there is a significant enrichment for *H. petiolaris* – *H.*
420 *paradoxus* sharing as predicted under the hybrid origin scenario; gene tree topology counts
421 show the same pattern but are not significant. However, the branch length test, which has

422 greater power to detect recent introgression, indicates more introgression between *H. annuus* –
423 *H. petiolaris*. These results may be reconciled by the long history of introgression between the
424 *annuus* and *petiolaris* clades (Strasburg and Rieseberg 2008), with the branch length test
425 successfully detecting recent introgression between *H. annuus* and *H. petiolaris*, and D offering
426 evidence of more ancient hybridization associated with the evolution of *H. paradox*. Likewise, *H.*
427 *paradoxus* shares more derived alleles with *H. anomalus* and *H. deserticola* than do other
428 members of the *annuus* clade, offering further indirect support that *H. paradoxus* is the product
429 of an independent hybridization event (supplementary table 2).

430 There is also uncertainty regarding the parentage of *H. anomalus* and *H. deserticola*. While
431 the role of *H. niveus canescens* as one parent is strongly supported, the other hybrid parent is
432 less clear. Is it the *H. annuus* lineage? The most likely TreeMix graph suggests the ancestor of
433 the *H. annuus* clade is the other parent, but this pattern is not strongly supported by admixtools,
434 the triplet testing or D. The hybrid species share fewer derived alleles with *H. annuus* and *H.*
435 *argophyllus* when compared to *H. debilis*, *H. niveus canescens* or *H. petiolaris* (supplementary
436 table 2), but they share more derived alleles with *H. paradoxus* when compared to *H. debilis*.
437 This may be indicative of more recent introgression between the *H. annuus* lineage and
438 members of the *petiolaris* clade sampled in this study, all of which overlap in geographic
439 distribution and hybridize with *H. annuus*. Alternatively, *H. anomalus* and *H. deserticola* are a
440 combination of *H. niveus canescens* and an unsampled or extinct basal lineage, as is suggested
441 in the admixtools tree. Future comparisons with an allopatric control from the *H. petiolaris* clade
442 (e.g., *H. niveus niveus*) will be needed to distinguish between these hypotheses.

443 One perhaps surprising aspect of our results is that the amount of admixture in “hybrid
444 species” is not exceptional compared to non-hybrid species. This highlights that the amount of
445 hybrid ancestry is not critical for hybrid speciation, only that admixture was involved in
446 reproductive isolation (Rieseberg 1997; Schumer et al. 2014), which we do not directly address

447 in this study. Nonetheless, our results have implications for the interpretation of previous
448 experimental studies that have attempted to make this link (e.g., Rieseberg et al. 1995, 1996,
449 2003; Lexer et al. 2003; Gross et al. 2004; Ludwig et al. 2004). For example, evidence that the
450 homoploid hybrid species are more ancient than previously believed makes it more difficult to
451 determine which ecological, phenotypic and genomic changes are associated with hybridization
452 and which changes arose as a consequence of divergence after hybrid speciation (Ungerer et
453 al. 2006). We suspect that the clusters of parental markers found to be linked to QTLs in
454 segregating hybrids, and which were also found in the hybrid species genomes (Rieseberg et al.
455 2003), likely correspond to inversions. Such inversions are frequent in *Helianthus* (Ostevik et al.
456 2020) and often appear to be ancient, pre-dating the species they are found in (Todesco et al.
457 2020). Future studies attempting to connect reproductive barriers with admixture should be
458 conducted within a phylogenetic framework to better assess when key differences arose.

459 Previous work attempted to dissect parental ancestry across the putative hybrid genomes,
460 finding strings of linked parental markers suggestive of rapid establishment of the hybrid
461 genomes (Ungerer et al. 1998; Buerkle and Rieseberg 2008). However, incomplete lineage
462 sorting, multiple ancient admixture events, structural variation, and the auto-correlational effects
463 of recombination rate variation may all affect ancestry assignments across the genome. Future
464 attempts at identifying and interpreting ancestry tracks will have to contend with this complexity.

465 Are there other examples of homoploid hybrid speciation in *Helianthus*? Probably – although
466 the relevant data come from earlier studies rather than the present analysis (see supplementary
467 discussion about outcomes of hybridization in the annual sunflower clade). Independently
468 derived dune ecotypes of the prairie sunflower are isolated from non-dune populations by
469 multiple reproductive barriers (Heiser 1958; Ostevik et al. 2016). Recent population genomic
470 studies indicate that most of the traits underlying reproductive isolation map to ancient
471 chromosomal inversions that appear to have originated via introgression from a basal lineage in

472 the annual clade (or even earlier) (Huang et al. 2020; Todesco et al. 2020). In an example
473 involving more recent hybridization, a 77-day difference in flowering between coastal and barrier
474 island ecotypes of *H. argophyllus* was found to result from a 30 Mb introgression from *H.*
475 *annuus* containing a functional copy of *HaFT1* (Todesco et al. 2020). The dune ecotypes
476 arguably represent new species, and one of them is described as *H. neglectus* in most
477 taxonomic treatments (Heiser et al. 1969; Schilling 2020). Ecotypic differentiation in *H.*
478 *argophyllus* represents an earlier stage of the speciation process, but it illustrates how
479 introgression of a single gene can cause significant reproductive isolation. It is noteworthy that
480 all three potential new cases of homoploid hybrid speciation involve the colonization of
481 ecologically divergent habitats that provide ecogeographic isolation similar to that seen in the
482 ancient hybrids.

483 ***Recombination rate and introgression***

484 Local recombination rate is a critical population genetics parameter that can affect ILS, gene
485 flow and genetic diversity (Ortíz-Barrientos et al. 2002; Nachman and Payseur 2012; Haenel et
486 al 2018). For example, introgressed loci are more likely to persist in regions of high
487 recombination because they more quickly unlink from deleterious neighbouring loci, such as
488 genetic incompatibilities (Brandvain et al. 2012; Schumer et al. 2018). In our dataset, higher
489 recombination regions are less likely to support the inferred species topology. This is likely a
490 combination of reduced introgression in low recombination regions, as well as lower N_e , which
491 reduces ILS (Martin et al. 2019; Li et al. 2019; Pease and Hahn 2013). The effect is strongest
492 for the node separating *H. annuus* samples, where the recombination rate estimates are most
493 accurate. Sunflowers have highly labile chromosome structure and there are numerous large-
494 scale rearrangements between species in this tree, but this pattern is retained even between
495 our outgroup perennial sunflower species, suggesting that recombination rate is relatively
496 conserved even if genome structure changes (Ostevik et al. 2020). At several nodes we find

497 that the relationship is reversed, which may be because at those nodes we have the
498 introgressed topology, rather than the true species topology. Exploring the relationship between
499 recombination rate and tree topology is useful for identifying an accurate species topology.

500 **Conclusions**

501 Despite vast improvements in the quantity of genetic data available to current researchers,
502 disentangling introgression patterns is often challenging (Hibbins and Hahn 2022). When
503 multiple introgression events have occurred, this can lead to false positives or false negatives,
504 which is especially true when the true donors are not sampled or, in some cases, no longer
505 extant. Broad sampling of species can help reduce these issues and broad sampling of
506 populations within species can identify when introgression is geographically localized. Thus,
507 when evaluating the plausibility of introgression signals detected using genomic data, it is
508 important to consider factors such as reproductive compatibility and the likelihood of geographic
509 overlap over the course of evolutionary divergence. That being said, the reconstruction of
510 historical ranges is an inexact science, and information on extinct congeners is completely
511 unknown for most wild genera.

512 Phylogenomic analysis below the genus level often finds evidence of introgression,
513 especially in plants (Dagilis et al., 2021). This suggests that a significant proportion of plant
514 species have some admixed ancestry. Despite this, homoploid hybrid species are rare because
515 most studies fail to link admixture with the evolution of reproductive barriers (Schumer et al.,
516 2014). Although attempts have been made to identify admixture-derived reproductive barriers
517 through purely bioinformatic analyses (e.g. Sun et al. 2020; Wang et al. 2022), links between
518 phenotypes, reproductive isolation and introgressed ancestry are required to confirm the
519 homoploid hybrid speciation. A notable exception to this is work by Wang et al. (2021), which
520 identified an admixed species in *Ostryopsis*, quantified reproductive barriers and transgenically
521 tested candidate genes acquired through admixture. Widespread introgression across sunflower

522 species raises the possibility that admixture played a role in other speciation events. In
523 particular, chromosome rearrangements are both common and are known to cause reproductive
524 isolation in sunflowers (Ostevik et al. 2020; Lai et al. 2005), so future work should leverage
525 chromosome resolved genomes and QTL mapping to identify how whether the introgressed
526 rearrangements cause speciation.

527 **Methods:**

528 **Data generation**

529 We generated whole genome resequencing data for two *H. paradoxus* samples. DNA was
530 extracted from individual seedlings or dried leaf tissue using a modified CTAB extraction
531 protocol (Murray et al. 1980; Zeng et al. 2002), and an indexed Illumina sequencing library was
532 created, which included a Duplex Specific Nuclease (DSN) repeat depletion step (Todesco et al.
533 2020). This data was added to previously published whole genome resequencing data to create
534 a dataset of nine species, or subspecies, of annual sunflowers (Hubner et al. 2019; Todesco et
535 al. 2020; Owens et al. 2021). For each, we randomly selected two samples to be used in the
536 analysis. We only included two per species because we were interested in understanding
537 between species relationships, rather than within species diversity. Additionally, using fewer
538 samples substantially reduced the computational bottleneck in both variant calling and
539 subsequent phylogenetic analyses. To act as outgroups, we also included one sample each for
540 four diploid perennial sunflower species (See supplementary table 1).

541 For each sample, Illumina sequence data was aligned to the *Helianthus annuus* HA412-
542 HOv2 genome using NextGenMap (v0.5.3) (Sedlazeck et al. 2013), PCR duplicates were
543 marked with samtools v0.1.19 (Danecek et al. 2021), and variants were called using GATK
544 (v4.1.4.1) (Mckenna et al. 2010). We specifically included invariant sites in GATK
545 genotypeGVCFs to facilitate downstream analyses. The sunflower genome contains a large

546 proportion of transposable element repeats that hamper read alignment. To avoid those regions,
547 we used GenMap (-K 50 -E 2) (v1.3) to calculate k-mer uniqueness or mappability for all
548 positions in the genome (Pockrandt et al. 2020). We then removed all contiguous regions with
549 mappability < 1 that are greater than 100 bp. This removed 2.43 GBp out of 3.23 Gbp, but
550 should highly enrich retained regions for single copy sequences.

551 For our dataset, we filtered variant and invariant sites separately. For variant sites, we
552 removed all indels and visualized the distribution of quality metrics of SNPs using bcftools
553 (v1.14) to extract values and ggplot (v3.3.5) in R (v4.1.2) to plot (Whickham 2011; R core team
554 2021). We visually selected thresholds to remove outliers based on the distribution (bcftools
555 view -e 'INFO/DP > 1000' -e 'INFO/FS > 50' -e 'INFO/QD < 2' -e 'INFO/SOR > 4' -e 'INFO/MQ <
556 30') (supplementary figure 1). Variant and invariant sites were combined, and then filtered to
557 require ≥ 4 reads per genotype and $\geq 80\%$ of samples genotyped using VCFtools (v0.1.16)
558 (Danecek et al. 2011).

559 Code used to conduct analyses and make figures is deposited at
560 https://github.com/owensgl/helianthus_hybrid_species_2021.

561 ***Gene and species tree creation***

562 We took two approaches to explore genome-wide gene trees. In the first, we equalized
563 information across windows by requiring each genomic window have 10,000 called bp, leading
564 to variable physical size. In the second approach, we equalized physical size but not information
565 by dividing the entire genome into non-overlapping 10,000 bp windows and retained windows
566 with $\geq 2,000$ called bases. In both cases, heterozygous sites were retained and coded using
567 IUPAC coding.

568 For each of our genomic windows, we calculated a maximum likelihood phylogeny using IQ-
569 Tree (v2.0.6) (Minh et al. 2020a) with *H. giganteus*, a perennial species, as the outgroup.

570 Additionally, a single concatenated species tree using all retained bases was created using
571 maximum likelihood and model selection in IQ-TREE. We used a coalescent approach in
572 ASTRAL (v5.7.3) to estimate the species tree using both sets of gene trees separately (Zhang
573 et al. 2018).

574 We explored the diversity of gene tree topologies by using the tool findCommonTrees.py
575 (Edelman et al. 2020) to identify and count the occurrence of each topology. We found that
576 every tree was unique, so we subsampled trees down to a single sample per species and a
577 single perennial outgroup species to reduce the possible tree space. We repeated the counts of
578 trees and visualized the most common gene trees.

579 Using the concatenated phylogeny as a backbone, we measured gene concordance using
580 IQ-Tree2 (Minh et al. 2020b). This measures the percent of gene trees that support the topology
581 and the two possible alternate topologies for each branch quartet. We matched the *H. annuus*
582 recombination rate for each genomic window and calculated the percent support for each
583 topology for each recombination quintile. To determine if recombination rate affected support for
584 the species topology, we used a binomial regression with the formula:

585
$$gC \sim node_ID * cm_rate$$

586 ***Broad tests of introgression***

587 We used several methods that integrate signals of introgression between all samples. Using
588 Dsuite, we calculated the f_{branch} statistic for the species phylogeny (Malinsky et al. 2021). This
589 statistic incorporates all valid f_4 ratios (similar to Patterson's D) as well as the phylogeny to
590 identify branches that share allelic imbalance. This can help identify introgression events that
591 predate speciation times and therefore have a signal in multiple species.

592 We also used TreeMix to identify introgression events (Pickrell and Pritchard 2012). This
593 program builds a graph model of population splits based on the covariance of allele frequencies.
594 It allows for migration edges to be added to the graph that explain remaining covariance not
595 accounted for by the initial graph topology. Since TreeMix relies on covariance, we pruned our
596 dataset for SNPs in LD by only including sites with $> 0.8 r^2$ in 100,000 bp using `bcftools prune`
597 (Danecek et al. 2021), and did not include sample size correction. We ran TreeMix with 0 to 10
598 migration edges, with 99 replicates using different starting seeds, and selected the optimal
599 number of migration edges using the Evanno method in OptM (Fitak 2021). We visualized the
600 replicates with the likelihood values within 10 of the highest likelihood for each number of
601 migration edges (i.e. the models that were not the best, but were close).

602 Lastly, we built an admixture graph for the phylogeny using ADMIXTOOLS 2 (Patterson et
603 al. 2012). We first converted our VCF file to eigenvector format using a custom Perl script, and
604 then calculated all f_2 statistics using the `extract_f2` and `f2_from_precomp` functions. We used a
605 5 Mbp window for block bootstrapping. Admixture graphs were found using the `find_graphs`
606 command with `stop_gen=200`, `stop_gen2=20` and `perennials` set as the consistent outgroup.
607 This method finds graphs that fit the observed f -statistics in the data. The graph search can get
608 stuck at local optima, so we repeated each search 100 times for each number of admixture
609 events, from one to four, and retained the best fitting graph in each iteration. From this, we
610 picked the top three scoring graph for each number of admixture events. We then used
611 bootstrap-resampling of the out-of-sample scores for each graph using the function
612 `qpgraph_resample_multi` to ask whether adding additional nodes was significantly better
613 supported, by comparing the best graph with n admixture events with the best graph with $n+1$
614 admixture events, as well as comparing graphs.

615 ***Trio based tests of introgression***

616 Given the high levels of discordant gene trees, and conflicting signals from broad tests of
617 introgression, we explored trio-based tests of introgression using Dsuite to calculate Patterson's
618 D (Green et al. 2010; Malinsky et al. 2021). This test looks at quartets with the relationship
619 (((A,B),C),O) and asks if there is greater derived allele sharing between A and C or B and C.
620 Under ILS, both should share equal counts of derived alleles, but introgression will lead to
621 imbalance. Significance was tested using a block bootstrap algorithm. We used the perennial
622 species as an outgroup and tested all trio sets consistent with the species phylogeny.

623 For the putative hybrid species and their parents, we used three additional approaches, the
624 branch length test (BLT), QuIBL and Twisst. For these tests, we used genomic windows with
625 equal physical size, as described above. The BLT test compares the tip-to-tip distance between
626 inferred sister clades in the minor topologies using a Mann-Whitney test (Suvorov et al. 2022).
627 Under ILS, both minor topologies should have equal distance, but if introgression is occurring
628 then the topology it produces should have reduced distance due to its more recent coalescence.
629 These tests were done in R using TreeTools (Smith 2019), using the subsampled trees with
630 only a single sample per species. QuIBL examines trios in gene trees and asks whether the
631 branch length distribution fits better to a model that includes both ILS and later introgression, or
632 purely ILS (Edelman et al. 2020). As before, we used our perennial samples as outgroups. We
633 considered models with Bayesian information criterion score difference of ≥ 30 to be evidence of
634 introgression. We used Twisst to calculate the topological weighting for each possible tree using
635 the equal size gene trees (Martin and Van Belleghem 2017). This takes into account
636 phylogenetic position variation amongst individuals within a species and calculates a weight for
637 how common a particular topology is at each gene tree. We normalized weights such that each
638 genomic window had a total weight of one and counted the total weight for each topology.
639 Similar to the Discordant Count Test from Suvorov et al. (2020), we compared the counts of the
640 two minor topologies using a chi-squared test, under a null hypothesis that ILS should produce

641 equal counts of both minor topologies. We also calculated D across the genome using Dsuite,
642 with a window size of 50 SNPs and a step size of 25 for each possible candidate trio.

643 **Quantifying introgression across the genus**

644 Given the amount of gene flow identified within annual sunflowers, we expanded our
645 analysis of gene flow by using a previously published sequence capture set encompassing a
646 majority of diploid species in the genus (Stephens et al. 2015). This dataset included 170
647 aligned sequences totalling 106,862 bp and 11,407 parsimony informative sites. Since these
648 markers were anonymous and genome positions are required for block bootstrapping, we
649 aligned them to the *H. annuus* HA412-HOv2 reference genome using BLAST and selected the
650 location hit with the highest e-value. Sequence capture gene alignments include numerous
651 indels, so we were not able to combine this dataset with our previous WGS dataset. As before,
652 we used Dsuite to calculate D, f_4 and f_{branch} . For the species topology, we used the MP-EST tree
653 from Stephens et al. (2015), with polytomies randomly resolved.

654 **Acknowledgements:**

655 We would like to thank Sarah Yakimowski and Robert Sivinski for providing tissue used for
656 DNA extractions. This work was funded by a Banting Fellowship to GLO, as well as a Discovery
657 Grant from the Natural Sciences and Engineering Research Council of Canada to LHR.

658 **References**

- 659 Anderson E. Introgressive hybridization. New York: J. Wiley. 1949.
- 660 Anderson E, Stebbins Jr GL. Hybridization as an evolutionary stimulus. *Evolution*. 1954 378:88.
- 661 Atlagić J, Terzić S. 2006. Cytogenetic study of hexaploid species *Helianthus tuberosus* and its
662 F1 and BC1F1 hybrids with cultivated sunflower, *H. annuus*. *Genetika*. 38

- 663 Baute GJ, Owens GL, Bock DG, Rieseberg LH. 2016. Genome-wide GBS data provide a high-
664 resolution view of wild *Helianthus* diversity, genetic structure and interspecies gene flow.
665 *Am J Bot* 103:2170-2177.
- 666 Beatley JC. 1963. The sunflowers (genus *Helianthus*) in Tennessee. *J Tennessee Acad Sci* 38:
667 135-154.
- 668 Beckstrom-Sternberg S, Rieseberg LH, Doan K. 1991. Gene lineage analysis of populations of
669 *Helianthus niveus* and *H. petiolaris*. *Plant Syst Evol* 175:125-138.
- 670 Blake SF. 1942. In T. H. Kearney and R.H. Peebles. Flowering plants and ferns of Arizona.
671 U.S.D.A. Misc. Publ. 423.
- 672 Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. 2014. Speciation and introgression
673 between *Mimulus nasutus* and *Mimulus guttatus*. *PLoS genet* 10:e1004410.
- 674 Buerkle CA, Rieseberg LH. 2008. The rate of genome stabilization in homoploid hybrid species.
675 *Evolution* 62:266-275.
- 676 Calfee E, Gates D, Lorant A, Perkins MT, Coop G, Ross-Ibarra J. 2021. Selective sorting of
677 ancestral introgression in maize and teosinte along an elevational cline. *PLoS genet*
678 17:e1009810.
- 679 Chandler JM, Jan CC, Beard BH 1986. Chromosomal Differentiation among the Annual
680 *Helianthus* Species. *Syst Bot* 11:354–371.
- 681 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
682 Marth GT, Sherry ST, McVean G. 2011. The variant call format and VCFtools.
683 *Bioinformatics*. 27:2156-2158.

- 684 Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
685 McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.
686 Gigascience. 10:giab008.
- 687 Dagilis AJ, Peede D, Coughlan JM, Jofre GI, D'Agostino ER, Mavengere H, Tate AD, Matute
688 DR. 2021 15 years of introgression studies: quantifying gene flow across Eukaryotes.
689 bioRxiv.
- 690 Darwin C. On the origin of species, 1859. Routledge;
- 691 Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, García-Accinelli G, Van
692 Belleghem SM, Patterson N, Neafsey DE, Challis R. 2019. Genomic architecture and
693 introgression shape a butterfly radiation. *Science* 366:594-599.
- 694 Ellis JR, Pashley CH, Burke JM, McCauley DE. 2006. High genetic diversity in a rare and
695 endangered sunflower as compared to a common congener. *Mol Ecol* 15:2345-2355.
- 696 Farwall OA 1916. Contribution to the Botany of Michigan. No. 14, *Rep Mich Acad Sci* 17:180
- 697 Figueiró HV, Li G, Trindade FJ, Assis J, Pais F, Fernandes G, Santos SH, Hughes GM,
698 Komissarov A, Antunes A, Trinca CS. 2017 Genome-wide signatures of complex
699 introgression and adaptive evolution in the big cats. *Sci Adv* 3:e1700299.
- 700 Fitak RR. 2021. OptM: estimating the optimal number of migration edges on population trees
701 using Treemix. *Biol Methods Protoc* 6:bpab017.
- 702 Fontaine MC, Pease JB, Steele A, Waterhouse RM, Neafsey DE, Sharakhov IV, Jiang X, Hall
703 AB, Catteruccia F, Kakani E et al. 2015. Extensive introgression in a malaria vector species
704 complex revealed by phylogenomics. *Science*, 347:1258524.

- 705 Grant V, 1958. The regulation of recombination in plants. *Cold Spring Harb Symp Quant*,
706 23:337–363.
- 707 Grant V, 1981. *Plant Speciation*. New York Chichester, West Sussex: Columbia University
708 Press.
- 709 Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W,
710 Fritz MH, Hansen NF. 2010 A draft sequence of the Neandertal genome. *Science* 328:710-
711 722.
- 712 Gross BL, Rieseberg LH. 2005. The ecological genetics of homoploid hybrid speciation. *J Hered*
713 96:241-252.
- 714 Gross BL, Schwarzbach AE, Rieseberg LH. 2003. Origin(s) of the diploid hybrid species
715 *Helianthus deserticola* (Asteraceae). *Am J Bot* 90:1708-1719.
- 716 Gross BL, Kane NC, Lexer C, Ludwig F, Rosenthal DM, Donovan LA, Rieseberg LH. 2004.
717 Reconstructing the origin(s) of *Helianthus deserticola*: Survivorship and selection on the
718 desert floor. *Am Nat* 164:145-156.
- 719 Haenel Q, Laurentino TG, Roesti M, Berner D. 2018. Meta-analysis of chromosome-scale
720 crossover rate variation in eukaryotes and its significance to evolutionary genomics. *Mol*
721 *Ecol* 27:2477-2497.
- 722 Hibbins MS, Hahn MW. 2022. Phylogenomic approaches to detecting and characterizing
723 introgression. *Genetics*. 220:iyab173.
- 724 Heiser CB. 1947. Hybridization between the sunflower species *Helianthus annuus* and *H.*
725 *petiolaris*. *Evolution*, 1:249-262.

- 726 Heiser CB. 1949a Natural hybridization with particular reference to introgression. *Bot Rev*
727 15:645-687.
- 728 Heiser CB. 1949b. Study in the evolution of the sunflower species *Helianthus annuus* and *H.*
729 *bolanderi*. *UC Pub Bot*, 12:157–196.
- 730 Heiser CB. 1951. Hybridization in the annual sunflowers: *Helianthus annuus* x *H. debilis* var.
731 *cucumerifolius*. *Evolution* 5:42-51.
- 732 Heiser CB. 1958. Three new annual sunflowers from the southwestern United States. *Rhodora*
733 60:275-279.
- 734 Heiser CB, Smith DM, Clevenger SB, Martin WC. 1969. The North American sunflowers
735 (*Helianthus*) *Mem Torrey Bot Club*, 22:1-218.
- 736 Henn HJ, Wingender R, Schnabl, H. 1998. Regeneration of fertile interspecific hybrids from
737 protoplast fusions between *Helianthus annuus* L. and wild *Helianthus* species. *Plant Cell*
738 *Rep* 18:220-224
- 739 Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH. 2020. Multiple chromosomal
740 inversions contribute to adaptive divergence of a dune sunflower ecotype. *Mol Ecol*
741 29:2535-2549.
- 742 Hübner S, Bercovich N, Todesco M, Mandel JR, Odenheimer J, Ziegler JE, Lee JS, Baute GJ,
743 Owens GL, Grassa CJ, et al. 2019. Sunflower pan-genome analysis shows that
744 hybridization altered gene content and disease resistance. *Nat Plants* 5:54-62.
- 745 Jones MR, Mills LS, Alves PC, Callahan CM, Alves JM, Lafferty DJ, Jiggins FM, Jensen JD,
746 Melo-Ferreira J, Good JM. 2018. Adaptive introgression underlies polymorphic seasonal
747 camouflage in snowshoe hares. *Science*. 360:1355-1358.

- 748 Kane NC, King M, Barker MS, Raduski A, Karrenberg S, Yatabe Y, Knapp SJ, Rieseberg LH.
749 2009. Comparative genomic and population genetic analyses indicate highly porous
750 genomes and high levels of gene flow between divergent *Helianthus* species. *Evolution*
751 63:2061-2075.
- 752 Karrenberg S, Lexer C, Rieseberg LH. 2007. Reconstructing the history of selection during
753 homoploid hybrid speciation. *Am Nat* 169:725-737.
- 754 Lai Z, Nakazato T, Salmaso M, Burke JM, Tang S, Knapp SJ, Rieseberg LH. 2005. Extensive
755 chromosomal repatterning and the evolution of sterility barriers in hybrid sunflower species.
756 *Genetics* 171:291-303.
- 757 Li G, Figueiró HV, Eizirik E, Murphy WJ. 2019. Recombination-aware phylogenomics reveals
758 the structured genomic landscape of hybridizing cat species. *Mol Biol Evol* 36:2111-2126.
- 759 Lee-Yaw JA, Grassa CJ, Joly S, Andrew RL, Rieseberg LH. 2019. An evaluation of alternative
760 explanations for widespread cytonuclear discordance in annual sunflowers (*Helianthus*).
761 *New Phyt* 221:515-526.
- 762 Lexer C, Welch M, Raymond O, Rieseberg LH. 2003. The origins of ecological divergence in
763 *Helianthus paradoxus* (Asteraceae): Selection on transgressive characters in a novel hybrid
764 habitat. *Evolution* 57:1989-2000.
- 765 Ludwig F, Rosenthal D, Johnston JA, Kane N, Gross BL, Lexer C, Dudley SA, Rieseberg LH,
766 Donovan LA. 2004. Selection on leaf ecophysiological traits in a hybrid *Helianthus* species
767 and early generation hybrids in a desert dune habitat. *Evolution* 58:2682-2692.
- 768 Malinsky M, Matschiner M, Svardal H. 2021 Dsuite-fast D-statistics and related admixture
769 evidence from VCF files. *Mol Ecol Resour* 21:584-595.

- 770 Mason CM. 2018. How old are sunflowers? A molecular clock analysis of key divergences in the
771 origin and diversification of *Helianthus* (Asteraceae). *Int J Plant Sci* 179:182-191.
- 772 Mallet J, Besansky N, Hahn MW. 2016. How reticulated are species?. *BioEssays*. 38:140-149.
- 773 Martin SH, Van Belleghem SM. 2017. Exploring evolutionary relationships across the genome
774 using topology weighting. *Genetics* 206:429-438.
- 775 Martin SH, Davey JW, Salazar C, Jiggins CD. 2019. Recombination rate variation shapes
776 barriers to introgression across butterfly genomes. *PLoS Biol* 17:e2006288.
- 777 Mayr E. 1963. *Animal species and evolution*. Harvard University Press.
- 778 McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler
779 D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a MapReduce
780 framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297-
781 1303.
- 782 Meier J, Marques D, Mwaiko S, Wagner CE, Excoffier L, Seehausen O. 2017. Ancient
783 hybridization fuels rapid cichlid fish adaptive radiations. *Nat Commun* 8:14363
- 784 Melo MC, Salazar C, Jiggins CD, Linares M. 2009. Assortative mating preferences among
785 hybrids offers a route to hybrid speciation. *Evolution* 63:1660-1665.
- 786 Minh BQ, Hahn MW, Lanfear R. 2020a New methods to calculate concordance factors for
787 phylogenomic datasets. *Mol Biol Evol* 37:2727-2733.
- 788 Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, Lanfear R.
789 2020b. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the
790 genomic era. *Mol Biol Evol* 37:1530-1534.

- 791 Mondon A, Owens GL, Poverene M, Cantamutto M, Rieseberg LH. 2018. Gene flow in
792 Argentinian sunflowers as revealed by genotyping by sequencing data. *Evol Appl* 11:193-
793 204.
- 794 Moody ML, Rieseberg LH. 2012. Sorting through the chaff, nDNA gene trees for phylogenetic
795 inference and hybrid identification of annual sunflowers (*Helianthus* sect. *Helianthus*). *Mol*
796 *Phylogenet Evol* 64:145-155.
- 797 Murray MG, Thompson WF. 1980 Rapid isolation of high molecular weight plant DNA. *Nucleic*
798 *Acids Res.* 8:4321–4325
- 799 Nachman MW, Payseur BA. 2012. Recombination rate variation and speciation: theoretical
800 predictions and empirical results from rabbits and mice. *Philos Trans R Soc Lond, B, Biol*
801 *Sci* 367:409-421.
- 802 Ortíz-Barrientos D, Reiland J, Hey J, Noor MA. 2002. Recombination and the divergence of
803 hybridizing species. In: Etges WJ, Noor MAF, editors. *Genetics of Mate Choice: From*
804 *Sexual Selection to Sexual Isolation*. Kluwer Academic Publishers p. 167-78.
- 805 Ostevik KL, Andrew RL, Otto SP, Rieseberg LH. 2016. Multiple reproductive barriers separate
806 recently diverged sunflower ecotypes. *Evolution* 70:2322-2335.
- 807 Ostevik KL, Samuk K, Rieseberg LH. 2020. Ancestral reconstruction of karyotypes reveals an
808 exceptional rate of nonrandom chromosomal evolution in sunflower. *Genetics.* 214:1031-
809 1045.
- 810 Owens GL, Rieseberg LH. 2014. Hybrid incompatibility is acquired faster in annual than in
811 perennial species of sunflower and tarweed. *Evolution* 68:893-900.

- 812 Owens GL, Baute GJ, Rieseberg LH. 2016. Revisiting a classic case of introgression:
813 Hybridization and gene flow in Californian sunflowers. *Mol Ecol* 25:2630-2643.
- 814 Owens GL, Todesco M, Bercovich N, Légaré JS, Mitchell N, Whitney KD, Rieseberg LH. 2021.
815 Standing variation rather than recent adaptive introgression probably underlies
816 differentiation of the *texanus* subspecies of *Helianthus annuus*. *Mol Ecol* 30:6229-6245.
- 817 Oziolor EM, Reid NM, Yair S, Lee KM, Guberman VerPloeg S, Bruns PC, Shaw JR, Whitehead
818 A, Matson CW. 2019. Adaptive introgression enables evolutionary rescue from extreme
819 environmental pollution. *Science*. 364:455-457.
- 820 Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster T,
821 Reich D. 2012. Ancient admixture in human history. *Genetics*. 192:1065-1093.
- 822 Pease JB, Hahn MW. 2013. More accurate phylogenies inferred from low-recombination regions
823 in the presence of incomplete lineage sorting. *Evolution*. 67:2376-84.
- 824 Pickrell J, Pritchard J. 2012. Inference of population splits and mixtures from genome-wide
825 allele frequency data. *Nat Preceed*
- 826 Pockrandt C, Alzamel M, Iliopoulos CS, Reinert K. 2020 GenMap: ultra-fast computation of
827 genome mappability. *Bioinform* 36:3687-3692.
- 828 R Core Team 2021. R: A language and environment for statistical computing. R Foundation for
829 Statistical Computing, Vienna, Austria.
- 830 Rieseberg LH, Soltis DE, Palmer JD. 1988. A molecular reexamination of introgression between
831 *Helianthus annuus* and *H. bolanderi*. *Evolution* 42:227-238.

- 832 Rieseberg LH, Carter R, Zona S. 1990a. Molecular tests of the hypothesized hybrid origin of two
833 diploid *Helianthus* species (Asteraceae). *Evolution* 44:1498-1511.
- 834 Rieseberg LH, Beckstrom-Sternberg S, Doan K. 1990b. *Helianthus annuus* ssp. *texanus* has
835 chloroplast DNA and nuclear ribosomal RNA genes of *Helianthus debilis* ssp.
836 *cucumerifolius*. *PNAS* 87:593-597.
- 837 Rieseberg LH. 1991. Homoploid reticulate evolution in *Helianthus*: Evidence from ribosomal
838 genes. *Am J Bot* 78:1218-1237.
- 839 Rieseberg LH, Beckstrom-Sternberg S, Liston A, Arias D. 1991. Phylogenetic and systematic
840 inferences from chloroplast DNA and isozyme variation in *Helianthus* sect. *Helianthus*. *Syst*
841 *Bot* 16:50-76.
- 842 Rieseberg LH, Van Fossen C, Desrochers A. 1995. Hybrid speciation accompanied by genomic
843 reorganization in wild sunflowers. *Nature* 375:313-316.
- 844 Rieseberg LH, Sinervo B, Linder CR, Ungerer M, Arias DM. 1996. Role of gene interactions in
845 hybrid speciation: evidence from ancient and experimental hybrids. *Science* 272:741-745.
- 846 Rieseberg LH. 1997. Hybrid origins of plant species. *Ann Rev Ecol Syst* 28:359-389.
- 847 Rieseberg LH, Baird S, Desrochers A. 1998. Patterns of mating in wild sunflower hybrid zones.
848 *Evolution* 52:713-726.
- 849 Rieseberg LH. 2000. Crossing relationships among ancient and experimental sunflower hybrid
850 lineages. *Evolution* 54:859-865.

- 851 Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL,
852 Schwarzbach AE, Donovan LA, Lexer C. 2003. Major ecological transitions in annual
853 sunflowers facilitated by hybridization. *Science* 301:1211-1216.
- 854 Rieseberg LH. 2006. Hybrid speciation in wild sunflowers. *Ann Mo Bot Gard* 93:34-48.
- 855 Schemske DW. 2000. Understanding the origin of species. *Evolution* 54:1069-1073.
- 856 Schilling EE, Linder CR, Noyes R, Rieseberg LH. 1998. Phylogenetic relationships in *Helianthus*
857 (Asteraceae) based on nuclear ribosomal DNA internal transcribed spacer region sequence
858 data. *Syst Bot* 23:177-188.
- 859 Schilling EE. 2020 *Helianthus*. In: Flora of North America Editorial Committee, eds. 1993. Flora
860 of North America North of Mexico. 19+ vols. New York and Oxford. Vol. 21, p. 141.
- 861 Schumer M, Rosenthal GG, Andolfatto P. 2014 How common is hybrid speciation? *Evolution*,
862 68:1553-1560.
- 863 Schumer M, Xu C, Powell DL, Durvasula A, Skov L, Holland C, Blazier JC, Sankararaman S,
864 Andolfatto P, Rosenthal GG, Przeworski M. 2018. Natural selection interacts with
865 recombination to shape the evolution of hybrid genomes. *Science*. 360:656-660.
- 866 Schwarzbach AE, Rieseberg LH. 2002. Likely multiple origins of a diploid hybrid sunflower
867 species. *Mol Ecol* 11:1703-1717.
- 868 Sedlazeck FJ, Rescheneder P, Von Haeseler A. 2013. NextGenMap: fast and accurate read
869 mapping in highly polymorphic genomes. *Bioinform* 29:2790-2791.
- 870 Smith MR. 2019 TreeTools: create, modify and analyse phylogenetic trees. Comprehensive R
871 Archive Network.

- 872 Sousa VC, Grelaud A, Hey J. 2011. On the nonidentifiability of migration time estimates in
873 isolation with migration models. *Mol Ecol* 20:3956-3962
- 874 Stebbins GL 1957. The hybrid origin of microspecies in the *Elymus glaucus* complex. *Cytologia*.
875 36:336–340.
- 876 Stephens JD, Rogers WL, Mason CM, Donovan LA, Malmberg RL 2015. Species tree
877 estimation of diploid *Helianthus* (Asteraceae) using target enrichment. *Am J Bot*, 102:910-
878 920.
- 879 Strasburg JL, Rieseberg LH. 2008. Molecular demographic history of the annual sunflowers
880 *Helianthus annuus* and *H. petiolaris* – large effective population sizes and rates of long-
881 term gene flow. *Evolution* 62:1936-1950.
- 882 Suarez-Gonzalez A, Lexer C, Cronk QC. 2018. Adaptive introgression: a plant perspective. *Biol*
883 *Lett* 14:20170688.
- 884 Sun Y, Lu Z, Zhu X, Ma H. 2020. Genomic basis of homoploid hybrid speciation within chestnut
885 trees. *Nat Commun* 11:1
- 886 Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'agostino ER, Price DK, Waddell PJ,
887 Lang M, Courtier-Orgogozo V, David JR. 2022. Widespread introgression across a
888 phylogeny of 155 *Drosophila* genomes. *Curr Biol* 32:111-123.
- 889 Timme RE, Simpson BB, Linder CR. 2007. High-Resolution Phylogeny for *Helianthus*
890 (Asteraceae) Using the 18S-26S Ribosomal DNA External Transcribed Spacer. *Am J Bot*
891 94: 1837-1852.

- 892 Todesco M, Owens GL, Bercovich N, Légaré JS, Souidi S, Burge DO, Huang K, Ostevik KL,
893 Drummond EBM, et al. 2020. Massive haplotypes underlie ecotypic differentiation in
894 sunflowers. *Nature* 584:602-607.
- 895 Ungerer MC, Baird S, Pan J, Rieseberg LH. 1998. Rapid hybrid speciation in wild sunflowers.
896 *PNAS* 95:11757-11762.
- 897 Ungerer MC, Strakosh SC, Zhen Y. 2006. Genome expansion in three hybrid sunflower species
898 is associated with retrotransposon proliferation. *Curr Biol* 16:R872-873.
- 899 Wang Z, Jiang Y, Bi H, Lu Z, Ma Y, Yang X, Chen N, Tian B, Liu B, Mao X, Ma T. 2021. Hybrid
900 speciation via inheritance of alternate alleles of parental isolating genes. *Mol Plant* 14:208-
901 222.
- 902 Wang Z, Kang M, Li J, Zhang Z, Wang Y, Chen C, Yang Y, Liu J. 2022. Genomic evidence for
903 homoploid hybrid speciation between ancestors of two different genera. *Nat commun* 13:1-
904 9.
- 905 Welch ME, Rieseberg LH. 2002. Habitat divergence between a homoploid hybrid sunflower
906 species, *Helianthus paradoxus* (Asteraceae), and its progenitors. *Am J Bot* 89:472-478.
- 907 Wickham H. 2011 ggplot2. *Wiley Interdiscip Rev Comput Stat* 3:180-185.
- 908 Xiong T, Li X, Yago M, Mallet J. 2022. Admixture of evolutionary rates across a butterfly hybrid
909 zone. *eLife* 11:e78135.
- 910 Yatabe Y, Kane NC, Scotti-Saintagne C, Rieseberg LH. 2007. Rampant gene exchange across
911 a strong reproductive barrier between the annual sunflowers, *Helianthus annuus* and *H.*
912 *petiolaris*. *Genetics* 175:1883-1893.

- 913 Zeng J, Zou Y, Bai J, Zheng H. 2002 Preparation of total DNA from recalcitrant plant taxa. *Acta*
914 *Bot Sin* 44:694–697
- 915 Zhang JQ, Imerovski I, Borkowski K, Huang K, Burge D, Rieseberg LH. 2019. Intraspecific
916 genetic divergence within *Helianthus niveus* and the status of two new morphotypes from
917 Mexico. *Am J Bot* 106:1229-1239.
- 918 Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree
919 reconstruction from partially resolved gene trees. *BMC Bioinform* 19:153
- 920